

CMFRI

bulletin 42

Part One

AUGUST 1988



NATIONAL SEMINAR ON SHELLFISH RESOURCES AND FARMING

TUTICORIN

19-21 January, 1987

Session - I

CENTRAL MARINE FISHERIES RESEARCH INSTITUTE
(Indian Council of Agricultural Research)
P. B. No. 2704, E. R. G. Road, Cochin-682 031, India

40. CYCLIC ACTIVITY IN THE DIGESTIVE DIVERTICULA OF *SUNETTA SCRIPTA* IN ACCORDANCE WITH TIDES

M. Dravidamani, A. Shanmugam and R. Kasinathan
Centre of Advanced Study in Marine Biology Parangipettai - 608 502

ABSTRACT

Feeding and digestion in bivalves are considered to be continuous and simultaneous. The tide influenced physiological changes in the digestive diverticula of *Sunetta scripta* on the basis of histological observation was studied. The shape of the digestive diverticula changes in accordance with the tidal level. The maximum pH value in the mantle cavity was recorded at high tide (7.46) when the animal was covered by the almost static tidal water, the constancy in the pH of the mantle cavity was due to the renewal of the water in the mantle cavity. At low tide the pH gradually comes down to 6.87 due to the exposure of the animal and the subsequent non-renewal of water in the mantle cavity.

INTRODUCTION

The effect of tidal cycle on the physiological state of the digestive diverticula of the bivalve molluscs is known since the work of Yonge (1926a). Though the overall function of digestive diverticula of bivalves remains a matter of controversy, this organ has been the subject of investigation for over 100 years. Feeding and digestion processes in bivalves are considered to be continuous and simultane-

ous (Yonge 1923, 1926 a, b; Owen 1966; Purchon 1968). Mansour (1946) and Mansourgek (1946) described the possibility of intracellular digestion in the lamellibranchs.

The rhythmic activity of the digestive diverticula in different bivalves has been studied by different authors. Some important works are by Morton (1956, 1969, 1970a, d, 1971, 1977), McQuiston (1969) and Langton and Gabbot (1974).

Perusal of the literature clearly shows that the information on the influence of tidal cycle on the function of digestive diverticula of bivalves is very meagre except a few like Shahul Hameed (1984) in *Crassostrea madrasensis* and *Anadara rhombea*. The present attempt is aimed at studying the tide influenced physiological changes in the digestive diverticula of *Sunetta scripta* on the basis of histological observations.

MATERIAL AND METHODS

The specimens of *S. scripta* were collected at the intertidal region near the mouth of Vellar estuary. The collections were carried out at an hourly interval starting from low tide to low tide extending over a period of 12 h. For every hour two specimens were collected. Immediately the pH of the mantle cavity was measured by using a BIOCHEM pm 79 model pH meter. Then digestive diverticula was dissected out and fixed in Aqueous Bouin's solution. The materials were brought to the laboratory and embedded in paraffin wax. 8-10 μ sections were taken and stained in Delafield Haematoxylin with Eosin as a counter-stain.

RESULTS AND DISCUSSION

The maximum pH value (7.46) was recorded at hightide when the animal was covered by the almost static tidal water; the constancy in the pH of the mantle cavity was due to the renewal of the water in the mantle cavity. Similar observation was also reported by Morton (1970a) in *C. edula*. At lowtide the pH gradually comes down to 6.87 due to the exposure of the animal and the subsequent non-renewal of water in the mantle cavity (Fig 1).

During lowtide the water in the mantle cavity is not replenished and the metabolic activity depletes the O_2 and increases the CO_2 level. As a result the pH value of this non-replenished water falls down. This phenomenon was also observed by Morton (1970a) in *C. edula*. Proceeding hightide replenishes the water due to which the PH increases again.




Fig. 1 Below: Mantle cavity pH in accordance with tides. Above: Showing the different stages of the digestive diverticula; be — basophil cells; dc — digestive cells fs-fr. gm. n. tation spherule; i-ium. n ifv — large food vacuoles; n — nucleus; sfv-sm. ll food vacuoles; wm - waste material.

Inspection of sectioned digestive diverticula of *S. scripta* fixed at hourly intervals over the tidal cycle revealed that the structure of the tubule underwent rhythmic histological changes in accordance with the state of the tide,

Just before the hightide, the cells of digestive tubules were flattened (Fig 1). It showed two types of cells (a) the digestive cells; (b) the basophil cells (Sumner 1966a) or 'Young cells' (Yonge 1926). The basophil cells stained much darker than the digestive cells; nuclei were located basally in the digestive cell and the lumen of the tubule was large. During hightide the digestive cells became swollen slightly and were cuboidal (or) columnar in form due to the absorption of food material arriving in the lumen of the diverticula from the stomach (absorptive phase). The number of nuclei in the 'nest of young cells' increased apparently. After hightide the lumen of the tubule reduced and the food materials were readily observed in vacuoles of the digestive cells (Shahul Hameed 1984).

At lowtide the tubules were different in condition and the bulges at the base of the digestive cells projected into the haemocoel due to the breakdown of food materials in the digestive cells. The distal surface of the digestive cells became swollen and was loaded with indigestible wastes called

fragmentation spherules. It has been reported by Morton (1969b) in *Dreissena* that the distal portion perhaps produce a fragmentation spherules and the proximal portion passes these into the haemocoel with food and waste materials.

At a later stage spherules were being budded off from the distal surfaces of the digestive cells in this case the nests of young cells had apparently oriented towards the lumen of the tubules. As a result the length of the digestive cells was mostly reduced and the orientation towards the lumen of the tubule was lost. Approximately 4 h after low tide the digestive cells are reduced in length as well as in number and only few nuclei were seen. The basiphil cells could not be distinguished from digestive cells. The tubules were in the disintegration phase. Just before the high tide the tubules showed significant increase in the size and number of the digestive cells. The basiphil cells were recognisable in small groups with a few cells (regenerative phase).

The structure of the tubules essentially similar to that described in the early high tide of the previous tidal cycle indicating the cyclic nature of the events.

In *S. scripta* the tidal cycle is related with the activity of digestive diverticula. Morton (1969a) opined that the animal when submerged by the tide the food is absorbed in the digestive diverticula during high tide and intracellular digestion takes place within the digestive cells.

REFERENCES

- LANGTON, R. W. AND P. A. GABBOT 1974. The tidal rhythm of extracellular digestion and response to feeding in *Ostrea edulis*. *Mar. Biol.* 24 (2) : 181-187.
- MANSOUR, K. 1946. Food and digestive processes of the lamellibranchia. *Nature*, Lond., Vol. 157, pp. 482.
- MANSOUR K. AND J. J. BEK 1946. Extracellular proteolytic and lipolytic enzymes of some lamellibranchs. *Nature*, London, 158 : 378.
- MCQUISTON, R. W. 1969. Cyclic activity in the digestive diverticula of *Lasaea rubra* (Montage) Bivalvia : Eulamellibranchia). *Proc. malac. Soc. London*, 38: 483-492.
- MORTON, J.E. 1956. The tidal rhythm and action of the digestive system of the Lamellibranchia *Lasaea rubra*. *J. mar. bio/Ass. U. K.* 35 : 563-586.
- MORTON, J. E. 1969 a. Studies on the biology of *Dreissena polymorpha* pall. 1. General anatomy and morphology. *Proc. Malacol. Soc. London*, 38 : 301-321.
- MORTON, B. S. 1969 b. Studies on the biology of *Dreissena polymorpha* pall. II. Correlation of the rhythms of adductor activity, feeding digestion and excretion. *Proc. Malacol. Soc. Lond.*, 38 : pp. 401-414.
- MORTON, B. S. 1970 a. The tidal rhythm and rhythm of feeding and digestion in *Cardium edule*. *J. mar. Biol. Ass. U. K.*, 50:499-512.
- MORTON, B. S. 1970 b. A note on the cytological structure and function of the digestive diverticula of *Macoma balthica* correlated with the rhythm of *Macoma balthica* correlated with the rhythm of the tide. *Malacol. Rev.*, 3:115-119.
- MORTON, B. S. 1971. The daily rhythm and tidal rhythm of feeding and digestion in *Ostrea edulis*. *Biol. J. Linn. Soc. London*, 3: 329-342.
- MORTON, B. S. 1977. The tidal rhythm of feeding and digestion in the pacific oyster, *Crassostrea gigas* (Thunberg). *J. Exp. Mar. Biol. Ecol.*, 26 : 135-151.
- OWEN, G. 1966, 'Digestion', chapter 2 In : *physiology of Mollusca*, Vol. II, Academic Press, Newyork.
- PURCHON, R.D. 1968. The Biology of the Mollusca. Pergamon Press, Oxford, 560 pp.

- SHAHUL HAMEED, P. 1984. Studies on the crystalline style of some bivalve molluscs of Porto Novo coastal waters. Ph. D. Thesis, Annamalai University, pp. 162.
- SUMNER, A. T. 1966 a. The cytology and histochemistry of the digestive gland cells of some freshwater lamellibranchs *J. R. microscr Soc*, 85 : 201 -211.
- YONGE, C. M. 1923. Studies on the comparative physiology of digestion. I. The mechanism of feeding, digestion and assimilation in the lamellibranch *Mya*. *Soc. Ex Biol.* 1 : 15-63.
- YONGE, C. M. 1926 a. The digestive diverticula in the Lamellibranchia. *Trans. R. Soc, Edins.* 54 : 703-718:
- YONGE, C. M. 1926 b Structure and physiology of the organs of feeding and digestion in *Ostrea edulis*, *J. mar Biol. Ass. U.K.*, 14: 295-386.