VILLORITA CYPRINOIDES (HANLEY) (PELECYPODA: CORBICULIDAE) FROM COCHIN BACKWATERS, KERALA

Dissertation submitted to the Mangalore University
in partial fulfillment of the requirements
for the Degree of
Doctor of Philosophy in Biosciences

SIVAPRASAD, P. S., M. Sc.



Library of the Central Marine Fisheri Research Institute, Cochin	es
Date of receipt	****
Accession No	****
Class No	*****

Central Marine Fisheries Research Institute (Indian Council of Agricultural Research)
P. B. No. 1603, Ernakulam North P. O.
Cochin- 682 018, Kerala, India

December 2007

Dedicated to my
Parents
&
God

Declaration

I hereby declare that this thesis entitled Bioaccumulation of heavy metals and pesticides in Villorita cyprinoides (Hanley) (Pelecypoda: Corbiculidae) from Cochin backwaters, Kerala is the bona fide record of the original research work carried out by me under the supervision and guidance of Dr. P. Kaladharan, Senior Scientist, Central Marine Fisheries Research Institute, and no part of the thesis has been presented earlier for any other degree, diploma or similar titles of any University or Institution in India or abroad.

Ernakulam, December 2007 Sivaprasad P. S. (Research Student)



Phone (Off)

: 0484-2394867

2391407

Telegram

: CADALMIN EKM

Telex

: 0885-6435 MFRI IN

Fax

: 91-44-2394909

केंद्रीय समुद्री माठित्स्यकी अनुसंधान संस्थान

पोस्ट बॉक्स सं 1603; एरणाकुलम नोर्त, कोव्चिन - 682 018

CENTRAL MARINE FISHEIRES RESEARCH INSTITUTE

POST BOX No.1603; ERNAKULAM NORTH P.O., COCHIN - 682 018

(भारतीय कुषि अनुसंधान परिषद)

(Indian Council Of Agricultural Research)

Dr. P. Kaladharan, M. Sc., PhD, ARS Senior Scientist

Certificate

This is to certify that the Doctoral Dissertation entitled Bioaccumulation of heavy metals and pesticides in Villorita cyprinoides (Hanley) (Pelecypoda: Corbiculidae) from Cochin backwaters, Kerala is the authentic record of the original research work carried out by Mr. Sivaprasad P. S., M. Sc., under my guidance and supervision and that no part of this dissertation has been submitted earlier for any other degree, diploma or similar titles of any other University or Institution in India or abroad.

Ernakulam, December 2007 P. Kaladharan (Supervising Guide)

Acknowledgements

I place on record, the deepest gratitude to my supervising guide Dr. P. Kaladharan, Senior Scientist, CMFRI for suggesting the topic, unfailing guidance, constant encouragement and constructive criticism I received from him throughout the course of my study.

I am grateful to Dr. N. G. K. Pillai, Director and Prof. (Dr.) Mohan Joseph Modayil, Former Director, CMFRI for providing necessary facilities to carryout the work.

I would like to express my thanks to the authorities of ICAR, New Delhi for granting Senior Research Fellowship in connection with the AP Cess Funded National Network Project on National Risk Assessment Programme for Fish and Fish Products for Domestic and International Markets.

I am deeply obliged to Dr. D. Prema, Senior Scientist and Shri. N. K, Sanil, Scientist (Senior Scale), CMFRI for the timely helps and the enthusiastic encouragements extended throughout my study.

I express my sincere thanks to Dr. M. Rajagopalan, Principal Scientist (Former HOD) and Dr. M. Lakshminarayana, Head, Fisheries Environment Management Division (FEMD) for providing laboratory facilities and timely support.

I am extremely grateful to Dr. P. K, Surendran, Principal Scientist (Retd.), Microbiology Fermentation and Biotechnology Division and Dr. T. V. Sankar, Head, Biochemistry and Nutrition Division (BND), CIFT for the generous helps and encouragements during the entire course of study.

I am indebted to Dr. P. K, Krishnakumar, Principal Scientist, CMFRI for kindly going through the thesis and for offering valuable suggestions.

I am much thankful to Smt. Leelabai, Smt. K, K, Valsala, Shri. Palaniswami, Shri. Nandakumar, Shri. Kambatkar, FEM Division, CMFRI and Dr. Usha Rani, Shri. Mathai, BND, CIFT for the helps and assistance received during the period of study.

I am thankful to my colleagues at CIFT, Cochin (Shri. Sivaperumal and Dr. O. K, Sindhu), CIBA, Chennai (Shri. Abey Abrham and Shri. Abhilash) and CMFRI, Cochin (Shri. Anoop Krishnan, Shri. Anoop B., Shri. Afsal, Miss. Anjana Mohan, Miss. Neetha Susan and Smt. Dhanya Lenin) for their kind co-operation and help.

I express my deep sense of gratitude to the Director of Fisheries, Government of Kerala for granting me study leave to pursue the doctoral programme.

I am deeply indebted to my parents, wife, daughters, brother and others in the family for their unstinted and wholehearted support and help throughout the study.

Above all, I bow my head before God almighty.

Contents

		Page
Dec	laration	
Cer	tificate	
Ack	nowledgments	
	onyms	
	tract	
Intr	oduction	1-12
Rev	iew of Literature	13-28
Mat	erials and Methods	29-36
1.	Collection site	29
2.	Collection of samples	29-31
3.	Analytical methods	31-35
4.	Bioaccumulation of heavy metals and pesticides	35
5.	Statistical analysis	35-36
Results		37-77
1.	Hydrographic parameters	37
2.	Fat	37
3.	Heavy metals	37-40
4.	Organochlorine pesticides	41-45
5.	Bioaccumulation of heavy metals and pesticides	46-49
6.	Hazard levels in Villorita cyprinoides	49-50
Disc	cussion	78-109
Sun	nmary	110-115
Refe	erences	116-148

Abstract

Bivalves are notorious for their ability to bioconcentrate contaminants and hence they can be used as effective indicators for monitoring trace toxic contaminants in coastal waters due to their wide distribution, sessile life style, euryhaline nature, resistance to stress and high accumulation of wide range of chemicals and ease of sampling. Bivalve resources of Kerala State mainly include different species of clams, oysters and mussels. During 2005- 2006 period, the total bivalve landing was about 61280 tones, of which 75% was contributed by a single species, *Villorita cyprinoides* (black clam). More than 90% of the total black clam production of the state is supported by Vembanad Lake system. Cochin backwaters, a part of Vembanad Lake system, one of the largest tropical estuaries along the west coast of India, is considered to be highly polluted due to effluent discharge from various sources.

The samples (water, sediment and black clam) for the present study were collected from one of the major canals of Cochin backwaters (Lat. 9 ⁰ 40' & 10 ⁰ 12' N and Long. 76 ⁰ 10' & 76 ⁰ 30' E) namely the Vemblai Canal. Samples for heavy metal analysis (water, sediment and black clam) were collected for a period of two years (January 2004 to December 2005). In the case of Organochlorine pesticides (OCPs), samples of water and clam meat were collected from the same site for a period of one year (January 2004 to December 2004).

The study revealed that the concentration of six heavy metals (As, Cd, Cr, Hg, Pb, Zn) and thirteen organochlorine pesticides (α BHC, γ BHC, β BHC, heptachlor, heptachlor epoxide, aldrin, dieldrin, endrin, ppDDE, opDDD, ppDDD, opDDT, ppDDT) found in *V. cyprinoides*, water and sediment are comparable to the earlier reports. Seasonal differences in the occurrence of heavy metal and pesticides are evident in the study. However, consistent statistically significant variations are rare. Environmental parameters like temperature, salinity and pH showed statistically significant correlation with few heavy metals such as Cd and Cr and OCPs like γ BHC, β BHC, aldrin, heptachlor epoxide, dieldrin and ppDDT. Higher values of

Bioconcentration factor (BCF) were obtained for almost all the heavy metals and organochlorine pesticide in the study. Values of Biotic sediment accumulator factor (BSAF) calculated for various heavy metals were moderately high for few metals. Seasonal difference is also evident in BCF and BSAF. Residual concentration of all the heavy metals and OCPs studied are within the safe limits prescribed by WHO and USFDA.

The results of this study enabled to conclude that *V. cyprinoides* possess the capabilities for accumulating heavy metals and OCPs from the surrounding medium. However, the bioaccumulation potential of the animal was found to vary with contaminants, seasons, levels in the surrounding medium and other unknown intrinsic and extrinsic factors. Therefore, the black clam can be used as sentinel organism in monitoring heavy metal and pesticide pollution in inland waters of Vembanad Lake systems. The residual limits of all the toxic contaminants are within the safe limits, and therefore the bivalve clam from Cochin backwater is safe and suitable for human consumption and for aquaculture as nutrient rich wet feed.

Acronyms

AAS Atomic Absorption Spectrophotometer

Anon. Anonymous

ANOVA Analysis of variance

As Arsenic

BCF Bioconcentration factor

BDL Below detection limits

BHC Benzene hexachloride

BSAF Biotic sediment accumulator factor

CBZ Cominco Binani Zinc

Cd Cadmium

CF Concentration factor

Con. Concentrated

Cr Chromium

DDD Dichloro diphenyl dichloroethane

DDE Dichloro diphenyl dichloroethylene

DDT Dichloro diphenyl trichloroethane

DO Dissolved oxygen

ECD Electron capture detector

EPA Environmental Protection Agency

FACT Fertilizer and Chemicals Travancore

FAO Food and Agriculture Organization

FDA Food and Drug Administration

GC Gas Chromatorgraphy

H₂SO₄ Sulphuric acid

HCH Hexa chlorohexane

HCl Hydrochloric acid

Hg Mecury

HIL Hindustan Insecticides Limited

HNO₃ Nitric acid

IRE Indian Rare Earths

MPEDA Marine Products Export Development Authority

MRL Maximum residual limit

ND Not detected

OCPs Organochlorine pesticides

op ortho para

PAH Polyaromatic hydrocarbons

Pb Lead

PCB Polychlorinated biphenyls

pp para para

ppb parts per billion

ppm parts per million

ppt parts per thousand

SD Standard deviation

SPSS Statistical Package for the Social Sciences

TCC Travancore Cocki Chemicals

TCMC Travancore Codin Notufacturing Company

UCI United Satal st India

WHO World Health Organization

Zn Zinc

α **W**pha

β Beta

γ yama