

Indian J. Fish., 59(3) : 153-155, 2012



Note

Occurrence of isopod parasites in clupeids off Chennai coast, India

SATYANARAYAN SETHI

Madras Research Centre of Central Marine Fisheries Research Institute, 75, Santhom High Road

R. A. Puram, Chennai - 600 028, Tamil Nadu, India

e-mail: sethisatyanarayana@yahoo.co.in

ABSTRACT

The study was carried out to investigate the incidence of isopod parasites in clupeids, with special emphasis on the Indian pellona fish, *Pellona ditchela*. Two species of isopods, *Joryma hilsae* and *Joryma sawayah* were found infecting this fish. The isopods of the genus *Joryma* are typical Indo-Pacific genus, found infecting the gills of marine fishes. However, this is the first report of these two species in Indian pellona, *P. ditchela* along the Bay of Bengal off Chennai coast, India. The prevalence (%) of *J. hilsae* and *J. sawayah* in *P. ditchela* recorded during the present investigation were 17.7 and 32.3 respectively. Cymothoids are common isopod parasites of marine fishes and their infection can cause serious damages to the fishes.

Keywords: Cymothoid, Indian pellona, *Joryma hilsae*, *Joryma sawayah*, *Pellona ditchela*

Cymothoids are obligatory parasites infesting many of the commercially important fishes. They are protandric hermaphrodites and bloodsuckers, living on the skin, gill filaments, or in the mouth of the fishes. These parasites retard growth and cause emaciation followed by death. Pathological conditions resulting from parasitic diseases in fishes reaches a high magnitude of epidemics under crowded and other unnatural conditions (Ravichandran *et al.*, 2007). Isopod parasite of the family *Cymothoidae* has been reported in about 350 species of fishes. Over 80% of these are from tropical and subtropical seas, many are from the Indo-Malaysian archipelago (Lester and Roubal, 1995). Their life cycle involves only one host (holoxenic cycle) (Trilles, 1994) and usually these are large sized parasites, which can cause deleterious effects on the host fishes (Trilles, 1996).

The information regarding cymothoid fauna of marine fishes from the Indian coasts is scanty (Pillai, 1954; Bal and Joshi, 1959; Veerapan and Ravichandran 2000). Most of the studies were from the east coast of India (Ravichandran *et al.*, 1999, 2009; Ravichandran and Rameshkumar, 2004; Rajkumar *et al.*, 2004, 2005; Ravichandran 2007; Rameshkumar and Ravichandran, 2010 a, b, Rameshkumar *et al.*, 2011). The present study reports the infection of two isopod parasites (*J. hilsae* and *J. sawayah*) in the Indian pellona, *Pellona ditchela* (Clupeidae) from Chennai coast.

Samples of freshly landed clupeid fishes viz., *Pellona ditchela*, *Thryssa mystax*, *Thryssa setirostris* and *Tenualosa ilisha* captured in gillnets from the inshore waters of Bay of Bengal off Chennai (lat. 13° 03' 974" N;

long 80° 17' 362" E, south-east coast of India) from March, 2009 to May, 2012 were collected and transported to the laboratory for the study. The fishes were weighed, measured, and investigated for the presence of isopods with the help of a dissecting microscope. The isopods were isolated, cleaned, fixed in 4% formaldehyde solution in physiological saline and preserved in 70% ethanol for taxonomic identification. Photomicrographs were taken using an Olympus digital camera C7070 fitted to the Olympus CX41 microscope. The isopod parasites were identified according to Bruce (1986) and Trilles (1994). The taxonomic classification of the Indian pellona fish host was carried out following Fischer and Whitehead (1974) and Froese and Pauly (2012). The total length of the fish hosts and isopods parasites were measured.

During the present study, four species of clupeids including 130 numbers of Indian pellona, (184.4±27.26 mm in length and 56.94±8.28 g in weight) were examined for presence of isopods. Among the various species of clupeids examined, only the Indian pellona, *P. ditchela* were found infected with two species of isopod parasites. The parasites were identified as *Joryma hilsae* Bowman and Tareen, 1983 and *Joryma sawayah* Rameshkumar, Ravichandran and Trilles, 2011 (Crustacea; Isopoda; Cymothoidae).

J. hilsae (Fig. 1) was collected from 23 fishes and *J. sawayah* (Fig. 2) was found infected in 42 fishes. *J. sawayah* showed higher prevalence (32.3%) in *P. ditchela* followed by *J. hilsae* (17.7%) (Table 1). The number of parasites per fish ranged from one to four with average of 1.028 ± 0.70 and were found mostly from gill chambers.



Fig. 1. *J. hilsae* attached to the gill region of *P. ditchela*



Fig. 2. *J. sawayah* attached to the gill region of *P. ditchela*

Body colour of *J. hilsae* (Fig. 3) was pale tan with large dark stripes on lateral parts of pereonites 6 and 7 and pleonites 1 – 5. The dorsal posterior side of *J. hilsae* showed clear blackish bands. The average length and weight of *J. hilsae* was 22.8 ± 4.13 mm and 0.405 ± 0.196 g respectively. In *J. sawayah* (Fig. 4) body colour was whitish



Fig. 3. Gravid females of *J. hilsae* collected from *P. ditchela* (dorsal & ventral view)

with a slight greenish tinge on the dorsal side. The mean length and weight of *J. sawayah* was 24.05 ± 2.37 mm and 0.4194 ± 0.104 g respectively.




Fig. 4. Gravid females of *J. sawayah* collected from *P. ditchela* (dorsal and ventral view)

Rameshkumar and Ravichandran (2010) reported infection of two species of isopods *Cymothoa indica* and *Alitropus typus* in *Tilapia mossambica* from Vellar Estuary, Tamil Nadu, India. Ravichandran *et al.* (2009) reported infestation of *Rastrelliger kanagurta* with cymothoid isopod, *Joryma brachysoma* from Colachel, south-west coast of India. Ravichandran *et al.* (2007) studied the histopathological changes associated with *Joryma tartoor* infestation in *Parastromateus niger*. Mance (juvenile parasitic stages of cymothoid) feed voraciously and kill fry and fingerlings of several species of fishes. Permanently attached adults parasites stunt the growth of fish and retard reproduction process. Parasites in the gill chamber usually lead to stunted and anaemic gill conditions. Isopod infections can lead to severe economic loss in culture operations (Bragoni, *et al.*, 1984).

In the present study, two isopods parasites (*J. hilsae* and *J. sawayah*) are reported in a single species of are *P. ditchela* along Chennai coast. According to Rameshkumar *et al.* (2011) all the species of *Joryma* are reported from the North-western Indian Ocean. *J. hilsae*, *J. engraulidis*, *J. tartoor* and *J. brachysoma*, from the South-western coasts of India and only one species *J. sawayah* from Kuwait. In the present study, *J. sawayah* which was earlier reported only from Kuwait waters is reported for the first time from the Indian pellona, *P. ditchela* along Chennai coast of India.

Table 1. Nature of isopod infection in clupeid fishes of Chennai coast

Fish species	Number of fishes examined	Number of fishes infected with <i>J. hilsae</i>	Number of fishes infected with <i>J. sawayah</i>
<i>Pellona ditchela</i>	130	23 (17.7)*	42 (32.3)*
<i>Thryssa mystax</i>	30	0	0
<i>Thryssa setirostris</i>	50	0	0
<i>Tenualosa ilisha</i>	40	9	0

*Prevalence (%)

Acknowledgements

Author is thankful to Dr. Kalaiselvam, Assistant Professor, CAS in Marine Biology, Annamalai University, Parangipettai, Tamil Nadu, India for identification of the parasites and thanks are also due to Shri. S. Rajapackiam, Technical Officer, MRC of CMFRI, Chennai for his constant support and encouragements.

References

- Bal, D. V. and Joshi, U. N. 1959. Some new isopod parasites on fishes. *J. Bombay Nat. Hist. Soc.*, 56 : 563–569.
- Bowman, T. E. and Tareen, I. U. 1983. Cymothoidae from fishes of Kuwait (Arabian Gulf) (Crustacea: Isopoda). *Smithson. Contrib. Zool.*, 382: 1–30.
- Bragoni, G., Romestand, B. and Trilles, J. P. 1984. Parasitoses a Cymothoadien chez le loup *Dicentrarchus labrax* (Linnaeus, 1758) en élevage. I. Ecologie parasitaire dans le cas de l'étang de Diana (Isopoda, Cymothoidae). *Crustaceana*, 47(1): 44–51.
- Bruce, N. L. 1986. Australian *Pleopodias* Richardson, 1910, and *Anilocra* Leach, 1818 (Isopoda, Cymothoidae), Crustacean Parasites of Marine Fishes. *Rec. Aus. Mus.*, 39: 85–130.
- Fischer, W. and Whitehead P. J. P. 1974. *FAO Species identification sheets for fishery purposes. Eastern Indian Ocean (Fishery Area 57) and Western Central Pacific (Fishery Area 71)*, Rome, Food and Agricultural Organization of the United Nations, Vol. I-V.
- Froese, R. and Pauly, D. 2012. Fishbase: a global information system on fishes. World Wide Web electronic publication. Available from: <http://www.fishbase.org>.
- Lester, R. J. G. and Roubal, F. R. 1995. Phylum Arthropoda. In: Woo P. T. K. (Ed.), *Fish diseases and disorders*, 1: *Protozoan and Metazoan infections*. CAB International, Wallingford, p. 475–598.
- Pillai, N. K. 1954. A preliminary note on the Tanaidacea and Isopoda of Travancore. *Bull. Cent. Res. Inst., Univ. Kerala, Ser. C, Natl Sci.*, 3: 1–21.
- Rajkumar, M., Perumal, P. and Trilles, J. P. 2005. *Cymothoa indica* (Crustacea, Isopoda, Cymothoidae) parasitizes the cultured larvae of the Asian Seabass *Lates calcarifer* under laboratory conditions. *Dis. Aquat. Org.*, 66: 87–90.
- Rajkumar, M., Santhanam, P. and Perumal, P. 2004. Report on new host record of *Cymothoa indica* Schioedte & Meinert, 1884 (Crustacea: Isopoda) from Parangipettai coastal waters, south-east coast of India. *J. Aquat. Biol.*, 19: 113–114.
- Rameshkumar, G. and Ravichandran, S. 2010a. *Cymothoa indica* (Isopoda; Cymothoidae) and *Alitropus typus* (Isopoda; Aegidae) on freshwater fish *Tilapia mossambica* (Cichlidae) in Vellar estuary, South-east coast of India, *Biotemas*, 23 (3): 67–70.
- Rameshkumar, G. and Ravichandran, S. 2010b. New host record, *Rastrelliger kanagurta*, for *Nerocila phaeopleura* parasites (Crustacea, Isopoda, Cymothoidae). *Middle-East J. Sci. Res.*, 5: 54–56.
- Rameshkumar, G., Ravichandran, S. and Trilles, J. P. 2011. Cymothoidae (Crustacea, Isopoda) from Indian fishes, *Acta Parasit.*, 56 (1): 78–91.
- Ravichandran, S. 2007. Infestation of isopod parasite *Lironeca puhi* in slender needle fish *Strongylura leiura*. *Res. J. Parasit.*, 2: 87–93.
- Ravichandran, S., Ajitha Kumar, T. T., Ronald, Ross, P. and Muthulingam, M. 2007. Histopathology of the infestation of parasite isopod *Joryma tartoor* of the host fish, *Parastromateus niger*. *Res. J. Parasit.*, 2(1): 68–71.
- Ravichandran, S., Balasubramanian, T. and Kannupandi, T. 2007. Incidence of parasitic isopods on the fish *Sphyraena obtusata*. *Res. J. Parasit.*, 2 (1): 45–50.
- Ravichandran, S. 2009. Invasion of gill region of *Ilisha melastoma* by isopod parasites. *ICFAI Univ. J. Life Sci.*, 3: 65–71.
- Ravichandran, S. and Rameshkumar, T. 2004. Infestation of isopod parasite *Nerocila phaeopleura* on *Chirocentus dorab*. *Proceedings National Seminar on Marine Resources*, ST Hindu College, Nagercoil, p. 45–48.
- Ravichandran, S., Rameshkumar, G., Mahesh Babu, B. and Kumaravel, K. 2009. Infestation of *Rastrelliger kanagurta* with cymothoid isopod, *Joryma brachysoma* in the Colachel environment of South-west coast of India. *J. Fish. Mar. Sci.*, 1: 80–84.
- Ravichandran, S., Ranjith Singh, A. J. A., Veerappan, N. and Kannupandi, T. 1999. Effect of isopod parasite *Joryma brachysoma* from Parangipettai coastal waters. *Ecol. Env. Conser.*, 5: 95–101.
- Trilles, J. P. and Hipeau-Jacquotte, R. 1996. Associations et parasitisme chez les Crustacés. In: de J. Forest, Masson (Ed.), *Traité de Zoologie, Anatomie, Systématique, Biologie*, publié sous la direction de Pierre-P. Grassé, Tome VII, *Crustacés, Fascicule 2, Généralités (Suite) et Systématique, sous la direction de*, Paris, p. 187–234.
- Trilles, J. P. 1994. Les Cymothoidae du Monde. Prodrôme pour une Faune. *Stud. Mar.*, (21/22): 1–288.
- Veerapan, N. and Ravichandran, S. 2000. *Isopod parasites from marine fishes of Parangipettai coast*. Publication of the Centre of Advanced Study in Marine Biology, Annamalai University, Parangipettai, 24 pp., pls 1–3.

Date of Receipt : 08.05.2012

Date of Acceptance : 01.09.2012