

Indian J. Fish., 61(3) : 29-34, 2014



## Seamount associated fishery of south-west coast of India - a preliminary assessment

K. K. BINEESH, K. V. AKHILESH, E. M. ABDUSSAMAD AND D. PRAKASAN

Central Marine Fisheries Research Institute, P. B. No. 1603, Ernakulam North P. O., Cochin - 682 018, Kerala, India

e-mail: [kbbineesh@gmail.com](mailto:kbbineesh@gmail.com)

### ABSTRACT

West coast of India accounts for 55% of the total marine capture fishery production of India. Wider continental shelf with high productivity due to coastal upwelling is attributed to this large contribution. Present study shows that fishing in the seamounts, knolls, guyots and ridges in the Arabian Sea too contribute considerably to the landings of west coast. During 2010 and 2011, an estimated 2314 and 1934 t of fishes were landed from seamounts and associated knolls at Kochi (Kerala) in the west coast. Landings were supported mostly by commercially important high value large pelagics, elasmobranchs, carangids and perches. Present study indicates that these areas are under intensive exploitation and more detailed studies are needed for critical status assessment of resources in these vulnerable habitats.

Keywords: Arabian Sea, Fishery, India, Knolls, Seamounts Species diversity, West coast

### Introduction

Seamounts are unique environments which support high level biodiversity. Several thousands of seamounts are distributed throughout the world's oceans and their numbers are increasing in each marine survey (Wessel, 2001; Kitchingman and Lai, 2004; Hillier and Watts, 2007; Allain *et al.*, 2006, 2008; Etnoyer *et al.*, 2010). The seamounts of Indian Ocean are the most poorly studied, virtually nothing is known about its biota, habitats and characteristics (Ingole and Koslow, 2005). Satellite data indicate that Indian Ocean has numerous small to moderate sized seamounts, mainly associated with its ridge systems (Craig and Sandwell, 1998; Iyer *et al.*, 2012).

Seamounts attract increasing interest from biogeographic, ecological, economic, fishery and conservation points (Glover and Smith, 2003), due to its diverse fauna. Eight hundred fish species have been recorded from seamounts (Froese and Sampang, 2004; Morato *et al.*, 2006; Morato and Clark, 2007), of which many have high commercial value. Seamounts are well known for fish aggregations (for spawning or feeding), making them highly vulnerable to fishing activities (Morato *et al.*, 2006). Large aggregations of pelagic fishes are often found in association with seamounts throughout the tropical oceans and are often exploited by fishing fleets (Fonteneau, 1991).

Since early 1980's fisheries have developed for both teleost and crustaceans around the seamounts in the world

oceans, particularly due to the discovery of fishing grounds of orange roughy, *Hoplostethus atlanticus*. During 1980, the Soviets developed fisheries across the seamounts of Madagascar, South-West Indian and Mid Indian Ridges, Ninety-East and Broken Ridges in the Indian Ocean, and targeted alfonso ( *Beryx splendens* ), rubyfish (*Emmelichthys* spp.) and butterfish (Centrolophidae) (Romanov, 2002).

In the Indian EEZ, seamounts are located along the west coast mostly as part of Chagos-Laccadive Ridge (CLR) (Kitchingman and Lai, 2004; Iyer *et al.*, 2012), but information available about their ecology, bathymetry and biodiversity are scanty (Qasim and Wafar, 1979; Untawale *et al.*, 1989; Parulekar, 1990; Ambiye and Untawale, 1992). The islands of Chagos, the Maldives and the Lakshadweep are above-water parts of the Chagos-Laccadive Ridge. The present article provides information on the distribution and fishery on seamounts, knolls, guyots and ridges off the west coast of India.

### Materials and methods

Weekly surveys at the Cochin Fisheries Harbour (CFH), Kochi, Kerala in south-west India were conducted between 2010 and 2011 to determine the catch, species and size composition of the landings. The daily catch data was estimated based on personnel observations carried out during one to three trips per week. The annual estimates of catch were made following the stratified random sampling design, as adopted by the Central Marine Fisheries

Research Institute (CMFRI), Kochi based on Srinath *et al.* (2005). Personnel interviews with fishermen were conducted to understand the geographical locations and to ensure that the catches are from seamounts (between 12°N-14° N and 71°E -72 °E) off south-west India. Seamount distribution map (Fig. 1) was downloaded from GEBCO undersea features names (<http://www.ngdc.noaa.gov/gazetteer>). Catch composition and size ranges were monitored. Fishes were identified up to species level using Talwar and Kacker (1984) and recent taxonomic keys.

## Results and discussion

### Fishing areas and methods

Three seamounts locally known as “*Manchappara* (Bassas de ‘Pedro Bank’), *Kora* (Cora Divh Bank) and *Bank* (Sesostris Bank) (Fig. 1), and several other knolls and ridges called “*paru, thitta*” off the west coast are intensively fished throughout the year and the catches are landed mainly at Cochin Fisheries Harbour, a major landing centre in the west coast of India. These seamounts are recognised off Karnataka coast with varying size and at different depths ranging from 43 m – 2300 m (Iyer *et al.*, 2012). These seamounts are intensively fished throughout the year by approximately 100 - 130 mechanised fishing vessels (42 to 72 feet OAL), which operate drift gillnets, longlines, and handlines depending on season and resource availability. They undertake mainly multiday fishing (8-22 days) operations. The operational depth over the seamounts ranged between 16-300 m. Fishing is carried out both during day and night hours. The surface fishing drift gillnets have size of 1000 to 2500 m length, 15 m height and mesh size of 11 inches. They operate during night hours and targets tunas, pelagic sharks, rainbow runner, groupers and sword fish. Longlines which target sharks, rays, tunas, billfishes and other medium sized fishes employ hooks No. 0-3 and use about 500-2500 hooks at a time depending on ground conditions. The common baits are scads (*Decapterus russelli*,




Fig. 1. Map showing the positions of seamounts along the south-west coast of India

*Decapterus macarellus* and *Selar crumenophthalmus*), sardines (*Sardinella* spp., *Dussumieria* spp.), flying fishes (*Exocoetus* spp.) and squids. Live flying fishes (*Exocoetus* spp.) kept in containers are used while targeting sailfishes. Hand lines with smaller hooks (hooks no. 8, 9 and 10) mainly targeted groupers, snappers, pigface brems and trevallies along rocky areas of 20-70 m depth. Fishing on seamounts depends on the availability of high value fishes (sharks, tunas, groupers, billfishes and marlins) and fishing operation generally ceases during monsoon (June-September).

### Fish landings

During 2010 and 2011, a total of 2314 and 1934 t of fishes were landed respectively from seamounts and associated knolls at Cochin Fisheries Harbour (Kochi), Kerala. Major groups contributing to the fishery were tunas, sailfishes, carangids, elasmobranchs, swordfishes, groupers and snappers. Estimated landings of major groups (Fig. 2) indicate that major portion of commercially important demersal groups and large pelagic fishes landed at Cochin are from knolls and seamounts.




Fig. 2. Estimated landings of major groups exploited from seamounts and knolls off south-west coast of India, landed at Cochin Fisheries Harbour during 2010 and 2011

Month-wise catch data of important groups and their contribution to fishery at Cochin (Fig. 3) indicates a clear seasonal pattern for the seamount fishery. During 2010, major landings were observed during February-March, with 39% of total landings followed by September-October (19%). In 2011, major landings recorded during February-March (35%) followed by May-June (25 %) and September-October (19%).

### Species composition

Landings from seamounts comprised 103 species belonging to 31 families and 64 genera (Table 1). Out of this, 53 species were abundant in the landings. The most frequently caught species were *Istiophorus platypterus*, *Katsuwonus pelamis*, *Thunnus albacares*, *Aprion*




Fig. 3. Monthly landings (%) of major groups exploited from seamounts and knolls off south-west coast of India, landed at Cochin Fisheries Harbour during 2010 and 2011

*virescens*, *Pinjalo pinjalo*, *Epinephelus* spp., *Lutjanus* spp. and *Acanthocybium solandri*, *Coryphaena hippurus*, sharks *Carcharhinus* spp. and rays *Taeniurops meyeri* and *Himantura fai*.

Tunas supported a good fishery from seamounts. Their average annual contribution was 533 and 489 t during 2010 and 2011 respectively, which represented 23-25% of the total landings from seamounts. Tunas in the landings were represented mainly by yellow fin, *Thunnus albacares* (59-69%) and skipjack, *Katsuwonus pelamis* (23-39%) and small portion of longtail tuna *Thunnus tonggol* (7%).

Landing of sailfish, *Istiophorus platypterus* from seamounts was 487 t during 2010 and 416 t during 2011. Marlins were represented by *Istiompax indica* with 186 t and 159 t during 2010 and 2011 respectively and swordfish *Xiphias gladius* with 172 and 215 t.

An estimated 313 and 195 t of carangids were landed from seamounts at Cochin during 2010 and 2011 respectively. During 2010, *Caranx* spp. dominated with average landings of 165 t followed by *Elegatis bipinnulata*, with 122 t. Contribution of *Caranx* spp. declined to 150 t followed by *E. bipinnulata* with 38 t during 2011.

Elasmobranch fishery was supported by more than 30 species belonging to 13 families (Table 1). Maximum diversity was observed in Carcharhiniformes and Myliobatiformes, which contributed major portion of fishery at Cochin. *T. meyeri*, *Carcharhinus limbatus*, *Carcharhinus brevipinna*, *Alopias superciliosus*, *Alopias pelagicus*, *Sphyrna lewini* and *Himantura fai* were the most abundant species landed. Elasmobranch landings during the period ranged between 242 and 144 t. Sharks dominated the landings representing 70% and 55 % during 2010 and 2011 respectively, followed by rays.

During 2010, snappers, rock cods, pigface brems and perches accounted for 191 t, 100 t, 59 t and 18.3 t respectively from seamounts, while during 2011 landing

from seamounts were dominated by snappers 150 t, rock cods 109 t, pigface brems 26 t and perches 6 t (Fig. 2).

Fishing gears like longlines, gillnets and handlines operating in the seamounts associated ridge areas primarily target high value fish resources like groupers, perches, large carangids and tunas (*Katsuwonus pelamis*, *Thunnus albacares*, *Thunnus tonggol* and *Gymnosarda unicolor*) where they aggregate for spawning and feeding. Similar aggregations of elasmobranchs, billfishes and tunas in association with seamounts are found in all tropical oceans and these are exploited by fishing fleets (Fonteneau, 1991; Itano and Holland, 2000).

The seamounts along the west coast of India consistently aggregates commercially important fish groups and are intensively exploited. Abdussamad *et al.* (2012) stated that, tunas are mainly caught from seamounts along the south-west coast of India. Large aggregation of myctophids have been reported near Angria Bank and Ratnagiri knolls (16° 19' N-72° 05' E) off Maharashtra (Kumaran, 1978) which forms a major food item of commercially important large pelagics. Silas (1985) commented that Angria Bank in northern Arabian Sea is an important fish spawning ground. The knolls off Beypore (11° 22' N- 74° 21' E), Periyapani (12° 24' N-71° 52' E), Ezhukalpanathitta (11° 26' N-74° 08' E), Chinnapani, Vizhinjam paru (off Kerala) are also known to support good fishery.

Seamounts and associated knolls are rich areas of biodiversity. Present study confirmed the presence of new species like Indian Bunquelovely, *Symphysanodon xanthopterygion*, Pillai's anthias, *Pseudanthias pillai* and *Liopropoma randalli* from the seamounts of west coast of India. Occurrence of elasmobranchs like *Rhynchobatus australiae*, *Dasyatis microps*, *Aetomylaeus vespertilio*, *Mobula tarapacana*, *Mobula kuhlii*, and *Hexanchus griseus* reported from these areas call for an in depth study on fish diversity of seamounts. An early biogeographic study of 92 seamounts suggested that 15% of the species collected were endemic to individual seamounts (Wilson and Kaufmann, 1987). More research is needed to know the actual diversity of these seamounts.

Results of the present study indicate an increasing trend in the fishing efforts in areas around seamounts along the south-west coast of India, during recent years especially for elasmobranchs, large pelagics and billfishes. This may ultimately lead to overexploitation of species, as reported from several parts of the world. Considering the boom and bust nature of many seamount fisheries (Koslow *et al.*, 2000), a cautious approach is needed to manage the seamount fishery of India. Seamounts and their associated ecosystems are considered as important component in the fishery along the west coast, as they act as breeding, feeding and nursery grounds of the

Table 1. List of commercially important fish species landed at Cochin Fisheries Harbour (Kochi), Kerala from seamounts, off south-west coast of India

Family	Species	Size range (cm)	Status in fishery	IUCN status
Hexanchidae	<i>Hexanchus griseus</i>	87-260	Occasional	Near Threatened
	<i>Heptanchias perlo</i>	80-140	Occasional	Near Threatened
Ginglymostomatidae	<i>Nebrius ferrugineus</i>	120-300	Occasional	Vulnerable
Stegostomatidae	<i>Stegostoma fasciatum</i>	80-160	Frequent	Vulnerable
Rhincodontidae	<i>Rhincodon typus</i>	95-260	Rare	Vulnerable
Alopiidae	<i>Alopias pelagicus</i>	110-226	Abundant	Vulnerable
	<i>Alopias superciliosus</i>	112-335	Abundant	Vulnerable
	<i>Alopias vulpinus</i>	126-312	Abundant	Vulnerable
Lamnidae	<i>Isurus oxyrinchus</i>	112-207	Frequent	Near Threatened
Carcharhinidae	<i>Carcharhinus albimarginatus</i>	92-217	Frequent	Near Threatened
	<i>Carcharhinus altimus</i>	140-187	Frequent	Data Deficient
	<i>Carcharhinus amboinensis</i>	104-260	Frequent	Data Deficient
	<i>Carcharhinus leucas</i>	90-340	Frequent	Near Threatened
	<i>Carcharhinus limbatus</i>	107-286	Frequent	Near Threatened
	<i>Carcharhinus longimanus</i>	100-140	Frequent	Near Threatened
	<i>Galeocerdo cuvier</i>	106-440	Abundant	Near Threatened
	<i>Prionace glauca</i>	180-342	Occasional	Near Threatened
	<i>Triaenodon obesus</i>	80-148	Rare	Near Threatened
	Sphyrnidae	<i>Sphyrna zygaena</i>	120-322	Rare
<i>Sphyrna lewini</i>		120-387	Abundant	Near Threatened
Rhinobatidae	<i>Rhina ancylostoma</i>	120-175	Occasional	Vulnerable
Rhynchobatidae	<i>Rhynchobatus australiae</i>	180-270	Occasional	Vulnerable
Dasyatidae	<i>Pteroplatytrygon violacea</i>	38-78	Abundant	Least Concern
	<i>Dasyatis microps</i>	120-186	Occasional	Data Deficient
	<i>Taeniurops meyeri</i>	102-180	Abundant	Vulnerable
	<i>Himantura granulata</i>	106-120	Rare	Near Threatened
	<i>Himantura fai</i>	90-165	Abundant	Least Concern
Myliobatidae	<i>Aetomylaeus vespertilio</i>	160-220	Rare	Endangered
	<i>Manta birostris</i>	320-440	Occasional	Vulnerable
	<i>Mobula tarapacana</i>	110-260	Occasional	Data Deficient
	<i>Mobula kuhlii</i>	90-118	Occasional	Data Deficient
	<i>Mobula japonica</i>	120-314	Abundant	Near Threatened
Istiophoridae	<i>Istiophorus platypterus</i>	160-340	Abundant	Least Concern
	<i>Istiompax indica</i>	260-410	Abundant	Data Deficient
Xiphiidae	<i>Xiphias gladius</i>	120-210	Occasional	Least Concern
Coryphaenidae	<i>Coryphaena hippurus</i>	80-110	Abundant	Least Concern
Menidae	<i>Mene maculata</i>	16-28	Occasional	Not Evaluated
Rachycentridae	<i>Rachycentron canadum</i>	60-163	Abundant	Least Concern
Scombridae	<i>Acanthocybium solandri</i>	130-202	Occasional	Least Concern
	<i>Scomberomorus commerson</i>	82-144	Abundant	Near Threatened
	<i>Thunnus albacares</i>	64-168	Abundant	Near Threatened
	<i>Gymnosarda unicolor</i>	110-228	Frequent	Least Concern
	<i>Katsuwonus pelamis</i>	52-86	Abundant	Least Concern
	<i>Thunnus tonggol</i>	56-98	Abundant	Data Deficient
	<i>Auxis thazard</i>	52-62	Occasional	Least Concern
Gempylidae	<i>Ruvettus pretiosus</i>	83-114	Occasional	Data Deficient
Fistulariidae	<i>Fistularia commersonii</i>	74-110	Occasional	Not Evaluated
Zeidae	<i>Zenopsis conchifer</i>	34-56	Occasional	Not Evaluated
Lutjanidae	<i>Aprion virescens</i>	72-98	Abundant	Not Evaluated



	<i>Pristipomoides multidens</i>	48-92	Abundant	Not Evaluated
	<i>Lutjanus bohar</i>	62-84	Abundant	Not Evaluated
	<i>Lutjanus kasmira</i>	14-22	Abundant	Not Evaluated
	<i>Lutjanus lutjanus</i>	16-30	Abundant	Not Evaluated
	<i>Pinjalo pinjalo</i>	36-76	Abundant	Not Evaluated
Lethrinidae	<i>Lethrinus mahsena</i>	22-48	Abundant	Not Evaluated
Emmelichthyidae	<i>Erythrocles acarina</i>	26-68	Abundant	Not Evaluated
Priacanthidae	<i>Pristigenys refulgens</i>	14-28	Occasional	Not Evaluated
Berycidae	<i>Beryx splendens</i>	16-22	Occasional	Not Evaluated
Carangidae	<i>Selar crumenophthalmus</i>	13-28	Occasional	Not Evaluated
	<i>Trachinotus mookalee</i>	64-92	Occasional	Not Evaluated
	<i>Decapterus macarellus</i>	26-38	Abundant	Least Concern
	<i>Decapterus macrosoma</i>	24-36	Abundant	Not Evaluated
	<i>Megalaspis cordyla</i>	38-52	Occasional	Not Evaluated
	<i>Caranx ignobilis</i>	60-145	Abundant	Not Evaluated
	<i>Caranx sexfasciatus</i>	30-102	Abundant	Least Concern
	<i>Caranx melampygus</i>	64-116	Abundant	Not Evaluated
	<i>Caranx tille</i>	30-90	Abundant	Not Evaluated
	<i>Carangoides fulvoguttatus</i>	84-110	Occasional	Not Evaluated
	<i>Carangoides gymnothethus</i>	48-90	Occasional	Not Evaluated
	<i>Seriolina nigrofasciata</i>	38-44	Occasional	Not Evaluated
	<i>Naucrates ductor</i>	13-18	Occasional	Not Evaluated
	<i>Alectis indicus</i>	36-128	Abundant	Not Evaluated
	<i>Seriola dumerili</i>	46-82	Occasional	Not Evaluated
	<i>Seriola rivoliana</i>	68-96	Occasional	Not Evaluated
	<i>Scomberoides lysan</i>	44-96	Occasional	Not Evaluated
	<i>Scomberoides commersonianus</i>	56-110	Abundant	Not Evaluated
	<i>Elagatis bipinnulata</i>	90-160	Abundant	Not Evaluated
Serranidae	<i>Variola louti</i>	13-76	Abundant	Least Concern
	<i>Variola albimarginata</i>	44-78	Abundant	Least Concern
	<i>Plectropomus leopardus</i>	46-70	Abundant	Near Threatened
	<i>Plectropomus areolatus</i>	44-70	Abundant	Vulnerable
	<i>Epinephelus radiatus</i>	44-67	Abundant	Least Concern
	<i>Epinephelus poecilnotus</i>	44-62	Abundant	Least Concern
	<i>Epinephelus longispinis</i>	40-60	Abundant	Least Concern
	<i>Epinephelus latifasciatus</i>	42-60	Abundant	Data Deficient
	<i>Epinephelus fuscoguttatus</i>	46-86	Abundant	Near Threatened
	<i>Epinephelus flavocaeruleus</i>	40-73	Abundant	Least Concern
	<i>Epinephelus fasciatus</i>	22-36	Abundant	Least Concern
	<i>Epinephelus epistictus</i>	44-60	Rare	Data Deficient
	<i>Epinephelus diacanthus</i>	30-48	Abundant	Near Threatened
	<i>Epinephelus chlorostigma</i>	47-72	Abundant	Least Concern
	<i>Epinephelus areolatus</i>	22-32	Abundant	Least Concern
	<i>Cephalopholis urodeta</i>	20-28	Abundant	Least Concern
	<i>Cephalopholis sonnerati</i>	44-56	Abundant	Least Concern
	<i>Cephalopholis miniata</i>	31-40	Abundant	Least Concern
	<i>Cephalopholis formosa</i>	26-31	Abundant	Least Concern
	<i>Aethaloperca rogae</i>	36-56	Occasional	Not Evaluated
Kyphosidae	<i>Kyphosus vaigiensis</i>	34-58	Occasional	Not Evaluated
Acanthuridae	<i>Naso brevirostris</i>	30-46	Occasional	Least Concern
	<i>Naso</i> sp. 1		Occasional	
	<i>Naso</i> sp. 2		Occasional	
Mullidae	<i>Parupeneus</i> sp. 1		Rare	
	<i>Parupeneus</i> sp. 2		Rare	

major fishery resources. Continuous monitoring of the fishery and resources from these areas is needed to ensure sustainable fishery.

### Acknowledgements

We are grateful to the Director, Central Marine Fisheries Research Institute, Kochi for the support and facilities provided.

### References

- Abdussamad, E. M., Koya, K. P. S., Rohit, P., Joshi, K. K., Ghosh, S., Elayathu, M. N. K., Prakasan, D., Sebastine, M., Beni, M and Rao, G. S. 2012. Fishery of yellowfin tuna *Thunnus albacares* (Bonnaterre, 1788) in the Indian EEZ with special reference to their biology and population characteristics. *Indian J. Fish.*, 59(3): 43-51.
- Allain, V., Kirby, D. and Kerandel, J. 2006. *Seamount research planning workshop final report*. Report of the Seamount Research Planning Workshop, 20-21 March 2006, Secretariat of the Pacific Community, Noumea, New Caledonia, p. 1-57.
- Allain, V., Kerandel, J. A., Andrefouet, S., Magron, F., Clark, M., Kirby, D. S and Muller-Karger, F. E. 2008. Enhanced seamount location database for the Western and Central Pacific Ocean: Screening and crosschecking of 20 existing datasets. *Deep-Sea Res., Part I*, 55(8): 1035- 1047.
- Ambiye, V. and Untawale, A. G. 1992. Deep water marine algal flora of the submerged banks off west coast of India. In: Desai, B. N. (Ed.), *Oceanography of the Indian Ocean*. Oxford and IBH, New Delhi, p. 247-252.
- Craig, C. H. and Sandwell, D. T. 1988. Global distribution of seamounts from Seasat profiles. *J. Geophys. Res.*, 93(B9): 10408-10420.
- Etnoyer, P. J., Wood, J. and Shirley, T. C. 2010. How large is the Seamount Biome? *Oceanography*, 23(1): 206-209.
- Fonteneau, A. 1991. Seamounts and tuna in the tropical Atlantic. *Aquat. Living Resour.*, 4:13-25.
- Froese, R. and Sampang, A. 2004. Taxonomy and biology of seamount fishes. In: Morato, T. and Pauly, D. (Eds.), *Seamounts: biodiversity and fisheries*. University of British Columbia. Fisheries Centre Research Report, 12: 25-31.
- Glover, A. G. and Smith, C. R. 2003. The deep-sea floor ecosystem: current status and prospects of anthropogenic change by the year 2025. *Env. Cons.*, 30(3): 219-241.
- Hillier, J. K. and Watts, A. B. 2007. Global distribution of seamounts from ship-track bathymetry data. *Geo-phys. Res. Lett.*, 34: L 13304, 1-5.
- Ingle, B. and Koslow, J. A. 2005. Deep-sea ecosystems of the Indian Ocean. *Indian J. Mar. Sci.*, 34: 27-34.
- Itano, D. G. and Holland, K. N. 2000. Movement and vulnerability of bigeye (*Thunnus obesus*) and yellowfin tuna (*Thunnus albacares*) in relation to FADs and natural aggregation points. *Aquat. Living Resour.*, 13: 213-223.
- Iyer, S. D., Das, P., Kalangutkar, N. G. and Mehta, C. M. 2012. Seamounts – windows of opportunities and the Indian scenario. *Curr. Sci.*, 102(10): 1382-1391.
- Kitchingman, A. and Lai, S. 2004. Inferences on potential seamount locations from mid-resolution bathymetric data. In: Morato, T. and Pauly, D. (Eds.), *Seamounts: biodiversity and fisheries*. University of British Columbia Fish. Centre Res. Rep., 12: 7-12.
- Koslow, J. A., Boehlert, G. W., Gordon, J. D. M., Haedrich, M., R. L., Lorange, P. and Parin, N. 2000. Continental slope and deep-seawater fisheries: implications for a fragile ecosystem. *ICES J. Mar. Sci.*, 57: 548-557.
- Kumaran, M. 1978. Our fisheries resources and the role of upwelling in their fluctuations. Part II: Primary productivity and fisheries potential. *Seafood Exp. J.*, 10 (2): 9-19.
- Morato, T., Cheung, W. W. L. and Pitcher, T. J. 2006. Vulnerability of seamount fish to fishing: fuzzy analysis of life history attributes. *J. Fish. Biol.*, 68(1): 209-221.
- Morato, T. and Clark, M. R. 2008. Seamount fishes: ecology and life histories. In: Pitcher T. J., Morato, T., Hart, P. J. B., Clark M. R., Haggan N. and Santos R. S. (Eds.), *Seamounts: ecology, conservation and management*. Blackwell, Oxford, p. 170-188.
- Parulekar, A. H. 1990. Bio-indications of sunken ships and ship wrecks. In: Rao, S. R. (Ed.), *Recent advances in marine archaeology. Proceedings of the second Indian conference on marine archaeology of Indian Ocean countries, January 1990*, p. 95-101.
- Qasim, S. Z. and Wafar, M. V. M. 1979. Occurrence of living corals at several places along the west coast of India. *Mahasagar*, 12(1): 53-58.
- Romanov, E. 2002. Bycatch in the tuna purse-seine fisheries of the western Indian Ocean. *Fish. Bull.*, 100(1): 90-105.
- Silas, E. 1985. *Tuna fisheries of the Exclusive Economic Zone of India: biology and stock assessment*. Central Marine Fisheries Research Institute, Kochi, India, 216 pp.
- Srinath, M., Kuriakose, S. and Mini, K. G. 2005. Methodology for the estimation of marine fish landings in India. *CMFRI Special Publication*, 86: 1-57.
- Talwar, P. K. and Kacker, R. K. 1984. *Commercial sea fishes of India*. Zoological Survey of India, Calcutta, 997 pp.
- Untawale, A., Reddy, C. R. K. and Ambiye, V. D. 1989. Marine algal flora of submerged Angria Bank (Arabian Sea). *Indian J. Mar. Sci.*, 18: 207-209.
- Wessel, P. 2001. Global distribution of seamounts inferred from gridded Geosat/ERS-1 altimetry. *J. Geophys. Res.*, 106 (B9): 19431-19441.
- Wilson, R. R. and Kaufmann, R. S. 1987. Seamount biota and biogeography. In: Keating, B. H., Fryer, P., Batiza, R. and Boehlert, G. W. (Eds.), *Seamounts, islands and atolls*. *Geophys. Monogr.*, 43: 319-334.

Date of Receipt : 09.08.2012

Date of Acceptance : 12.02.2014