



SHORT COMMUNICATION

Molecular and Cytogenetic Characterization of Fish Cell Lines and its Application in Aquatic Research

Naresh Sahebrao Nagpure · Akhilesh Kr. Mishra · A. S. Ninawe ·
Avinash Rasal · Akhilesh Dubey · Amit Kumar · M. Goswami ·
Ravindra Kumar · J. K. Jena

Received: 22 January 2014/Revised: 8 August 2014/Accepted: 6 January 2015/Published online: 22 December 2015
© The National Academy of Sciences, India 2015

Abstract Fish cell line has emerged as an important tool in fishery biotechnology. In recent years, various fish cell lines have been developed by different researchers across the country. National Repository on Fish cell lines, established with the aim to preserve fish cell lines for training and education to stakeholders, has started functioning at National Bureau of Fish Genetic Resources, Lucknow. This repository is supposed to characterize and preserve the fish cell lines developed across the country and serve as a national referral centre for Indian and exotic fish cell lines. Currently, the repository is maintaining 50 fish cell lines deposited by various research institutes in India, including the cell lines developed at cell culture facility of National Bureau of Fish Genetic Resources. The cell lines have been successfully cryopreserved after verifying its authenticity by sequence analysis of two mitochondrial genes, viz. 16S rRNA and cytochrome c oxidase sub-unit I. Chromosomal analysis, transfection efficiency and immunocytochemistry are also being used to characterize the cell lines. The facility is serviceable for the collection, deposition and

distribution of fish cell lines. This paper discusses the status as well as the methodology adopted for fish cell lines development, characterization and storage at NRFC.

Keywords Cell line repository · Cryopreservation · Fish cell line · Molecular characterization

Cell culture is the most important means available in life sciences to study the biological activities occurring inside cells of a living body and it refers to the *in vitro* growth of cells taken from tissue of multicellular organisms. The cells are bathed in a culture medium containing essential nutrients and energy sources necessary for the cells survival [1] The culture environment usually consists of a suitable glass or plastic culture vessel containing a liquid or semisolid medium that supplies nutrients to the growing cells.

Fish cell lines have increased tremendously in number (283), since the development of the first permanent fish cell

N. S. Nagpure · A. Kr. Mishra · A. Rasal · A. Dubey ·
A. Kumar · M. Goswami · R. Kumar (✉)
Division of Molecular Biology & Biotechnology, National
Bureau of Fish Genetic Resources (Indian Council of
Agricultural Research), Canal Ring Road, P.O. Dilkusha,
Lucknow 226 002, India
e-mail: ravindra.scientist@gmail.com

N. S. Nagpure
e-mail: nagpurens@yahoo.co.in

A. Kr. Mishra
e-mail: am.mishra70@gmail.com

A. Rasal
e-mail: avinashrasal44@gmail.com

A. Dubey
e-mail: engg.akhil111@gmail.com

A. Kumar
e-mail: amitbios@gmail.com

M. Goswami
e-mail: mukugoswami@gmail.com

A. S. Ninawe
Department of Biotechnology, Ministry of Science and
Technology, CGO Complex, New Delhi, India
e-mail: ninawe@gmail.com; ninawe@dbt.nic.in

J. K. Jena
National Bureau of Fish Genetic Resources (Indian Council of
Agricultural Research), Canal Ring Road, P.O. Dilkusha,
Lucknow 226 002, India
e-mail: jkjena2@rediffmail.com

line in 1962 from rainbow trout [2]. Although, a large number of fish cell lines have been established from freshwater and anadromous species, only a few cell lines are established from commercial marine fishes [3]. Embryonic cell lines have also been developed from some selected fishes, such as zebrafish (*Danio rerio*) [4, 5], medaka (*Oryzias latipes*) [6], sea perch (*Lateolabrax japonicus*) [7] Japanese flounder (*Paralichthys olivaceus*) [8], turbot (*Scophthalmus maximus*) [9] and sea bass (*Dicentrarchus labrax*) [10].

The efforts of Dr. Dilip Kumar in early 1980s to develop fish cell lines at CIFA, Bhubaneswar and primary cultures developed from gill tissue of mrigal (*Cirrhinus mrigala*) [11], kidney of stinging catfish (*Heteropneustes fossilis*) [12] followed by caudal fin of rohu (*L. rohita*) [13] had given momentum to fish cell culture work in India. 50 fish cell lines, developed by different groups in the country, have been deposited to NRFC till date (Table 1). In recent years, a few embryonic stem-like cell lines also have been established by various researchers in fisheries [14, 15]. The crustacean cell culture has gained potential scope for development of diagnostic reagents and probes for use in shrimp, crayfish and lobster industries. Despite several attempts to develop cell culture systems from crustaceans [16–20] in the country, no established cell line has been reported from crustaceans till date. The morphology, growth mode, subculture routine, DNA profile and other details of the cell line are available at: <http://mail.nbfgf.res.in/nrfc/cellline.php>.

The fish cell line repositories ensure availability of fish cell lines which is requisite to address certain key issues, like diagnosis of viral diseases, development of preventive as well as therapeutic measures, cytotoxicity measurements, detecting environmental changes and genotoxic effects etc. The authentication of cell lines by such repositories further certifies many things, like original source, passage number and information regarding submission to the scientific publications etc.

Major cell line repositories, including American Type Culture Collection (ATCC), European Collections of Cell Cultures (ECACC), German Collection of Microorganisms and Cell Cultures (DSMZ), have received cell line submissions from researchers across the world and authenticated all cell line submissions. The European Collections of Cell Cultures (ECACC) currently holds over 40,000 cell lines representing 45 different species and 50 tissue types, including the cell lines from different tissues of 21 fish species. Till date, out of over 3,400 cell lines deposited at the American Type Culture Collection (ATCC), 43 cell lines are from aquatic animals with 19 being fish cell lines available globally for dissemination to the researchers. A fish cell line repository exists at Department of Biology, University of Waterloo, Canada where studies on

application of fish cell lines in the toxicology and ecotoxicology of fish have been done [21].

The National Bureau of Fish Genetic Resources (NBFGR), Lucknow is the premium organization mandated for fish germplasm collection, classification, cataloguing and its conservation. The Institute has been designated as a National Repository of Fish Germplasm by ICAR, Department of Agricultural Research and Education, Ministry of Agriculture and National Biodiversity Authority (NBA), Ministry of Environment and Forests. Regrettably, the cell lines developed are not available to researchers across the country after completion of time bound projects. A lot of time and resources are spent to develop the cell lines for fresh experiments. To circumvent this, National Repository of Fish Cell Lines (NRFC) has been established at NBFGR for long-term preservation and redistribution of deposited fish cell lines to interested researchers. At NRFC, methodologies for effective long-term maintenance, preservation and characterization of fish cell lines are developed and documented so that these may be reproduced in the recipient's laboratory. Currently, NRFC is maintaining 50 fish cell lines which are available to researchers for R&D work (Table 1). Photomicrograph of the cell lines is given in Fig. 1.

The objective of the repository is to acquire the fish cell lines developed by various national research laboratories for its long term preservation and distribution of the cell lines as and when required by the research community of India. Cell line deposition and request form for depositing and requesting cell lines from the repository along with many other related information are currently being available at the NRFC website hosted at NBFGR homepage.

NRFC is involved in development and characterization of new fish cell lines. So far 13 cell lines from 11 different fish species have been contributed by NBFGR cell culture facility. The National repository developed will help in preventing different research groups across the country from developing fish cell lines from different species and organs which have already been developed. Such incidents might have occurred in the past due to lack of a national level fish cell line repository in India. The establishment of national fish cell line repository in India is, thus, a meaningful step towards R&D work to enhance aquaculture production in the country.

The cell lines developed at NRFC are being characterized for growth studies, chromosomal analysis, 16S rRNA and COI sequence analysis, immunocytochemistry, transfection studies, cryopreservation and revival efficiency. The effects of different temperatures, FBS and basic Fibroblast Growth Factor (bFGF) concentrations on cell growth are examined by seeding cells at a concentration of 1×10^5 cells in 25 cm² tissue culture flasks and incubated at 18, 20, 24, 28 and 32 °C for 7 days. Cells from triplicate

Table 1 Fish cell lines available at NRFC-NBFG, Lucknow

S. no.	Name of cell line	Fish species	Organ	NRFC code
1.	PCF	<i>Puntius chelynoides</i>	Fin	NRFC001
2.	SRF	<i>Schizothorax richardsonii</i>	Fin	NRFC002
3.	TTCF	<i>Tor tor</i>	Fin	NRFC003
4.	CCF	<i>Cyprinus carpio</i>	Fin	NRFC004
5.	WAF	<i>Wallago attu</i>	Fin	NRFC005
6.	RF	<i>Labeo rohita</i>	Fin	NRFC006
7.	KCF	<i>Koi carp</i>	Fin	NRFC007
8.	HBF	<i>Horabragus brachysoma</i>	Fin	NRFC008
9.	PDF	<i>Puntius denisonii</i>	Fin	NRFC009
10.	CFFN2	<i>Amphiprion sebae</i>	Dorsal fin	NRFC010
11.	CFBR	<i>Amphiprion sebae</i>	Brain	NRFC011
12.	CFSP	<i>Amphiprion sebae</i>	Spleen	NRFC012
13.	CFCP1	<i>Amphiprion sebae</i>	Caudal peduncle	NRFC013
14.	SISK	<i>Lates calcarifer</i>	Kidney	NRFC014
15.	SISS	<i>Lates calcarifer</i>	Spleen	NRFC015
16.	SIGE	<i>Lates calcarifer</i>	Eye muscle	NRFC016
17.	IGK	<i>Epinephelus coioides</i>	Kidney	NRFC017
18.	SICE	<i>Catla catla</i>	Eye muscle	NRFC018
19.	SICH	<i>Catla catla</i>	Heart	NRFC019
20.	CB	<i>Catla catla</i>	Brain	NRFC020
21.	ICG	<i>Catla catla</i>	Gill	NRFC021
22.	ICF	<i>Clarius batrachus</i>	Fin tissue	NRFC022
23.	LRG	<i>Labeo rohita</i>	Gill tissue	NRFC023
24.	DT1CPEx	<i>Dascyllus trimaculatus</i>	Caudal peduncle explant	NRFC024
25.	DT1F4Ex	<i>Dascyllus trimaculatus</i>	Fin explant	NRFC025
26.	DT1CPTr	<i>Dascyllus trimaculatus</i>	Trypsinized caudal peduncle	NRFC026
27.	RC4H1Tr	<i>Rachycentron canadum</i>	Trypsinized heart	NRFC027
28.	CTM	<i>Catla catla</i>	Thymus (macrophage)	NRFC028
29.	CTE	<i>Catla catla</i>	Thymus (epithelial)	NRFC029
30.	EM2HTTr	<i>Epinephelus malabaricus</i>	Trypsinized heart	NRFC030
31.	EM2GEx	<i>Epinephelus malabaricus</i>	Gill	NRFC031
32.	EM3GEx	<i>Epinephelus malabaricus</i>	Gill	NRFC032
33.	EM4SPEx	<i>Epinephelus malabaricus</i>	Spleen	NRFC033
34.	CTB	<i>Catla catla</i>	Blood	NRFC034
35.	PC1CpTr	<i>Pomacentrus caeruleus</i>	Trypsinized caudal peduncle	NRFC035
36.	PC1F1Ex	<i>Pomacentrus caeruleus</i>	Fin explant	NRFC036
37.	PC1L1Tr	<i>Pomacentrus caeruleus</i>	Trypsinized liver	NRFC037
38.	HC2SPEx	<i>Epinephelus merra</i>	Spleen	NRFC038
39.	CFF	<i>Pristolepis rubripinnis</i>	Fin	NRFC039
40.	IEE	<i>Etroplus suratensis</i>	Eye muscle	NRFC040
41.	IEK	<i>Etroplus suratensis</i>	Kidney	NRFC041
42.	IEG	<i>Etroplus suratensis</i>	Gill	NRFC042
43.	IEB	<i>Etroplus suratensis</i>	Brain	NRFC043
44.	RE	<i>Labeo rohita</i>	Eye muscle	NRFC044
45.	CSK	<i>Channa striatus</i>	Kidney	NRFC045
46.	CSG	<i>Channa striatus</i>	Gill	NRFC046
47.	WAM	<i>Wallago attu</i>	Muscle	NRFC047
48.	WAG	<i>Wallago attu</i>	Gill	NRFC048
49.	CPG	<i>Channa punctatus</i>	Gill	NRFC049
50.	DRM	<i>Danio rerio</i>	Muscle	NRFC050

flasks at each temperature are trypsinized and counted using a haemocytometer for a period of 1 week. Cell growth at different FBS and bFGF concentrations are studied at optimum temperature. Immuno-typing of the fish cell lines are performed with monoclonal antibodies directed against Vimentin and Cytokeratin (C-18) at different passages.

Chromosomal analysis is done by seeding cells in 75 cm² tissue culture flasks having L-15 medium with 20 % FBS. After 24 h incubation, medium is replaced with 10 ml of fresh medium containing 0.1 ml colcemid solution (1 µg ml⁻¹) (Sigma-Aldrich) for arresting the cell division at metaphase and incubated at 28 °C for 2 h. After harvesting by centrifugation (700 g, 5 min), the cells are suspended in a hypotonic solution consisting of 0.5 % KCl for 10 min and fixed in methanol: acetic acid (3:1). Slides are prepared following the conventional drop-splash technique [22]. The chromosomes are counted under a microscope after staining with 5 % Giemsa for 10 min.

Template DNA for PCR assays is extracted from cultured cell lines following the method described by [23]. An average 575 bp long fragment of mitochondrial 16S rRNA gene and 655 bp long fragment of cytochrome c oxidase subunit I (COI) genes are amplified using universal primers [24, 25] and sequenced. The obtained sequences of PCR fragments are compared to known sequences of the species using BLASTn for confirmation.

Transfection is an important technique to introduce nucleic acid into the target cells and observing changes occurring in its morphology. At NRFC Sub confluent monolayers (with 70–80 % confluency) of cell lines are usually transfected with pEGFP-C1 plasmid using LTX and Plus Reagents (Invitrogen) using manufacturer's protocol. The green fluorescence signals are observed after 18 h under fluorescent microscope (Olympus).

Cryopreservation of the fish cell lines is a vital technique that allows them to remain alive for a longer period. They can be revived as and when required if maintained suitably under liquid nitrogen. At NRFC the viability of cells following storage in liquid nitrogen (LN₂) is evaluated in freezing medium at different passages. In brief, cells growing logarithmically are harvested by trypsinization and concentrated by centrifugation. The pellet is washed with PBS and cell count is adjusted to 3 × 10⁶ cells per ml of L-15 medium with 20 % FBS and 10 % dimethyl sulfoxide (DMSO) of cell culture grade. Aliquots of 1 ml are dispensed into 2 ml sterile cryo-vials (10 numbers) (Nunc) hold at 4 °C for 2 h, -20 °C for 1 h, -80 °C overnight and then transferred into LN₂ containers at -196 °C. After 6 months of storage, the cryopreserved vials are revived to check the revival efficiency. For revival, the cryo-vials containing preserved cell lines are thawed quickly in a water bath at 28 °C. Freezing medium

is removed by centrifugation. The cells are suspended in L-15 medium with 20 % FBS and seeded into 25 cm² tissue culture flasks. The viability of the cells is measured by trypan blue staining and the number of cells is counted using haemocytometer. The revived viable cells are used as a normal fish cell line for cellular and molecular research in fisheries.

Fish cell lines have enormous applications in biomedical research, toxicology, gene regulation, gene expressions and gene transfer [26–28]. Fish cell lines represent an essential component of diagnostic procedures for viral and bacterial pathogens [29]. The most widely employed application of fish cell cultures is the isolation of fish viruses that are agents of epizootics of commercially important aquaculture fish species. In addition to serving solely as a means of virus isolation, fish cell cultures are very useful for studying the replication and genetics of these viruses, the effect of antiviral drugs and the production of experimental vaccines. The cultured fish cells produce a wide range of biological products of commercial interest including immune-regulators, antibodies, polypeptide growth factors, enzymes, hormones etc. They are used in the development of tissue plasminogen activator, interferon, monoclonal antibodies, tumour specific antigens and for in vitro study of gene expression, transplantation analysis and cancer biology. Cell cultures are being used to identify fish cytokines and antibacterial compounds. The main advantage of cell culture is that the cell lines allow higher control of experimental conditions and at the same time reduce the variability of in vivo responses that arise due to the responses of fish to stress and environmental influences.

Fish cell cultures are also being developed as in vitro models for determining karyotypes and other aspects of cytogenetics including chromosome polymorphism, chromosome abnormalities, speciation and evolution [30]. Cell cultures of fish and shellfish can contribute indirectly to their successful farming by providing basic insights into how their growth, reproduction and health can be understood and manipulated [31]. Cell cultures are being used to understand the hormonal regulation of fish reproductive cycles. Primary pituitary cell cultures are being used to study the regulation of gonadotropin hormone (GTH) secretion in responses to gonadotropin releasing hormone (GnRH) [32, 33]. Recently, fish has emerged as a suitable model and a promising alternative to the classical mammalian systems to study vertebrate developments, in general, and skeletogenesis, in particular. To complement in vivo developmental studies and identifying signalling pathways involved in developmental processes, the fish cell lines have been developed in particular bone-derived cells [34]. Fish stem cells have the potential for use in various biotechnological works. Among them, gene targeting, germ cell transplantation and semi-cloning by nuclear transfer




Fig. 1 Photomicrograph of few cell lines available at NRFC: **i** PCF **ii** SRF **iii** TTCF **iv** CCF **v** WAF **vi** RF **vii** KCF **viii** HBF **ix** PDF **x** CFFN **xi** CFBR **xii** CFSP **xiii** CFCP1 **xiv** SISK **xv** SISS **xvi** SIGE **xvii** IGK **xviii** SICE **xix** SICH **xx** CB **xxi** ICF **xxii** ICG **xxiii** LRG **xxiv** DT1CPEx **xxv** DTIF4Ex **xxvi** DTICPTr **xxvii** RC4H1Tr. Lineage of the cell line have been mentioned in Table 1

have attracted considerable interest and progress [35]. Using advanced genetic tools, the researchers from Howard Hughes Medical Institute and their collaborators have now identified key cells involved in zebrafish heart regeneration and begun to decipher the instructions to the cells used to carry out their repair work. This may have new perspective on which cells might be taught to regenerate in human hearts.

National Repository established at NBFGR, Lucknow will, thus, provide the required impetus to accelerate research in the field of fish cellular and reproductive biology. The cell lines available at the facility will be supplied to the research community upon request.

Acknowledgments The Department of Biotechnology, Government of India is thankfully acknowledged for the financial support. The authors also thank the Director, NBFGR for providing research facilities.

References

- Martin BM (eds) (1994) Routine cell culture. Tissue culture techniques: an introduction, pp. 29–30
- Wolf K, Quimby MC (1962) Established eurythermic line of fish cells in vitro. *Science* 135:1065–1066
- Wei Y-B et al (2010) A novel heart-cell line from brown-marbled grouper *Epinephelus fuscoguttatus* and its susceptibility to iridovirus. *J Fish Biol* 76(5):1149–1158
- Collodi P et al (1992) Culture of cells from zebrafish (*Brachydanio rerio*) embryo and adult tissues. *Cell Biol Toxicol* 8:43–61
- Sun L et al (1995) ES-like cell cultures derived from early zebrafish embryos. *Mol Mar Biol Biotech* 4:193–199
- Hong Y et al (1996) Pluripotency and differentiation of embryonic stem cell lines from the medakafish (*Oryzias latipes*). *Mech Dev* 60:33–44
- Chen S-L et al (2003) Establishment of a pluripotent embryonic cell line from sea perch blastula embryo. *Aquaculture* 218:141–151
- Chen et al (2004) Establishment of a continuous embryonic cell line from Japanese flounder *Paralichthys olivaceus* for virus isolation. *Dis Aquat Organ* 60:241–246
- Chen et al (2005) Development and characterization of a continuous embryonic cell line from turbot (*Scophthalmus maximus*). *Aquaculture* 249:63–68
- Buonocore F et al (2006) Production and characterization of a continuous embryonic cell line from sea bass (*Dicentrarchus labrax* L.). *Mar Biotechnol* 8:80–85
- Sathe PS et al (1995) Establishment and characterization of a new fish cell line, MG-3, from the gills of mrigal *Cirrhinus mrigala*. *Indan J Exp Biol*. 33:589–594
- Singh et al (1995) Development of primary cell cultures from kidney of freshwater fish *Heteropneustes fossilis*. *Indian J Exp Biol* 33:595–599
- Lakra WS, Bhonde RR (1996) Development of primary cell culture from the caudal fin of an Indian major carp, *Labeo rohita* (Ham). *Asian Fish Sci* 9:149–152
- Parameswaran V, Ahmed VPI, Shukla R, Bhonde RR, Hameed ASS (2007) Development and characterization of two new cell lines from milkfish (*Chanos chanos*) and grouper (*Epinephelus coioides*) for virus isolation. *Mar Biotechnol* 9:281–291
- Lakra WS et al (2010) Development and characterization of two new cell lines from common carp, *Cyprinus carpio* (Linn). *Biol Res* 43:385–392
- Purushothaman V et al (1998) Development of in vitro primary cell culture system from penaeid shrimp, *Penaeus indicus*, *Penaeus monodon* and sand crab *Emerita asiatica*. *Indian J Animal Sci* 68(10):1097–1099
- Kumar et al (2001) Development of a cell culture system from the ovarian tissue of African catfish (*Clarias gariepinus*). *Aquaculture* 194:51–62
- Uma A et al (2002) Establishment of primary cell culture from hepatopancreas of *Penaeus monodon* for the study of whitespot syndrome virus (WSSV). *Asian Fish Sci* 15:365–370
- Shashikumar A, Desai P (2008) Development of primary cell culture from *Scylla serrata*. *Cytotechnology* 56:161–169
- Goswami et al (2010) Development of cell culture system from the giant freshwater prawn *Macrobrachium rosenbergii* (de Man). *Mol Biol Rep* 37(4):2043–2048
- Bols NC, Dayeh VR, Lee LEJ, Schirmer K (2005) Use of fish cell lines in the toxicology and ecotoxicology of fish. In: Moon TW, Mommsen TP (eds) *Biochemistry and molecular Biology of fishes—environmental toxicology*, vol 6. Elsevier Science, Amsterdam, pp 43–84
- Freshney RI (1994) *Culture of animal cells: a manual of basic technique*. Wiley, New York
- Lo CF, Leu JH, Ho CH, Chen CH, Peng SE, Chen YT, Yeh PY, Huang CJ, Wang CH, Kou GH (1996) Detection of baculovirus associated with white spot syndrome (WSSV) in penaeid shrimps using polymerase chain reaction. *Dis Aquat Org* 25:133–141
- Palumbi S, Martin A, Romano S, Mcmillan WO, Stice L, Grabowski G (1991) *The simple fool's guide to PCR*. Version 2.0. Honolulu, HL 96822. Department of Zoology and Kewalo Marine Laboratory, University of Hawaii
- Ward RD, Zemlak TS, Innes BH, Last PR, Hebert PDN (2005) DNA barcoding Australia's fish species. *Proc R Soc London Ser B* 360:1847–1857
- Babich H et al (1986) *In vitro* cytotoxicity testing of aquatic pollutants (Cd, Cu, Zn, Ni) using established fish cell lines. *Ecotoxicol Environ Safety* 11:91–99
- Hightower LE, Renfro JL (1988) Recent applications of fish cell culture to biomedical research. *J Exp Zool* 248:290–302
- Driever W, Rangini Z (1993) Characterization of a cell line derived from zebrafish (*brachydanio rerio*) embryos. *In Vitro Cell Dev Biol Animal* 29A(9):749–754
- Villena AJ (2003) Applications and needs of fish and shellfish cell culture for disease control in aquaculture. *Rev Fish Biol Fish* 13:111–140
- Robert F (1970) Atlantic salmon (*Salmo salar*) chromosomes and speciation. *Trans Am Fish Soc* 99:105–111
- Bols NC (1991) Biotechnology and aquaculture: the role of cell cultures. *Biotech Adv* 9:31–49
- Fahraeus-van Ree GE et al (1982) The fine structure and function of isolated gonadotropic cells as revealed from pituitaries of immature rainbow trout, *Salmo gairdneri*, by means of enzymatic dispersion technique. *Cell Tissue Res* 226:641–653
- Weil C et al (1986) Use of pituitary cells in primary culture to study the regulation of gonadotropin hormone (GtH) secretion in rainbow trout; setting up and validating the system as assessed by its responsiveness to mammalian and salmon gonadotropin releasing hormone. *General Comp Endocrinol* 62:202–209
- Rafael MS et al (2010) Fish bone-derived cell lines: an alternative in vitro cell system to study bone biology. *J Appl Ichthyol* 26:230–234
- Hong N, Li Z, Hong Y (2011) Fish stem cell cultures. *Int J Biol Sci* 7(4):392–402