

# Pelvic Morphology Differs in Rotation and Obliquity Between Developmental Dysplasia of the Hip and Retroversion

Moritz Tannast MD, Peter Pfannebecker MD,  
Joseph M. Schwab MD, Christoph E. Albers MD,  
Klaus A. Siebenrock MD, Lorenz Büchler MD

Published online: 14 July 2012

© The Association of Bone and Joint Surgeons® 2012

## Abstract

**Background** Developmental dysplasia of the hip (DDH) and acetabular retroversion represent distinct acetabular pathomorphologies. Both are associated with alterations in pelvic morphology. In cases where direct radiographic assessment of the acetabulum is difficult or impossible or in mixed cases of DDH and retroversion, additional indirect pelvimetric parameters would help identify the major underlying structural abnormality.

**Questions/Purposes** We asked: How does DDH and retroversion differ with respect to rotation and coronal obliquity as measured by the pelvic width index, anterior inferior iliac spine (AIIS) sign, ilioischial angle, and obturator index? And what is the predictive value of each variable in detecting acetabular retroversion?

**Methods** We reviewed AP pelvis radiographs for 51 dysplastic and 51 retroverted hips. Dysplasia was diagnosed

based on a lateral center-edge angle of less than 20° and an acetabular index of greater than 14°. Retroversion was diagnosed based on a lateral center-edge angle of greater than 25° and concomitant presence of the crossover/ischial spine/posterior wall signs. We calculated sensitivity, specificity, and area under the receiver operating characteristic (ROC) curve for each variable used to diagnose acetabular retroversion.

**Results** We found a lower pelvic width index, higher prevalence of the AIIS sign, higher ilioischial angle, and lower obturator index in acetabular retroversion. The entire innominate bone is internally rotated in DDH and externally rotated in retroversion. The areas under the ROC curve were 0.969 (pelvic width index), 0.776 (AIIS sign), 0.971 (ilioischial angle), and 0.925 (obturator index).

**Conclusions** Pelvic morphology is associated with acetabular pathomorphology. Our measurements, except the AIIS sign, are indirect indicators of acetabular retroversion. The data suggest they can be used when the acetabular rim is not clearly visible and retroversion is not obvious.

**Level of Evidence** Level III, diagnostic study. See Guidelines for Authors for a complete description of levels of evidence.

---

One or more of the authors (JMS) has received fellowship funding from the Maurice E. Müller Foundation of North America. Each author certifies that he or she, or a member of his or her immediate family, has no commercial associations (eg, consultancies, stock ownership, equity interest, patent/licensing arrangements, etc) that might pose a conflict of interest in connection with the submitted article.

All ICMJE Conflict of Interest Forms for authors and *Clinical Orthopaedics and Related Research* editors and board members are on file with the publication and can be viewed on request.

Each author certifies that his or her institution approved the human protocol for this investigation, that all investigations were conducted in conformity with ethical principles of research, and that informed consent for participation in the study was obtained.

---

M. Tannast (✉), P. Pfannebecker, J. M. Schwab,  
C. E. Albers, K. A. Siebenrock, L. Büchler  
Department of Orthopaedic Surgery, Murtenstrasse,  
Inselspital, University of Bern, 3010 Bern, Switzerland  
e-mail: moritz.tannast@insel.ch

## Introduction

Developmental dysplasia of the hip (DDH) and acetabular retroversion represent distinct acetabular pathomorphologies. In DDH, the acetabulum is undercovered and often excessively anteverted [7, 8, 16, 21]. In acetabular retroversion, coverage is typically excessive, especially anteriorly [18]. Both conditions lead to distinct pathomechanical problems: a dysplastic acetabulum leads to static overload of the articular cartilage while a retroverted

acetabulum leads to dynamic impingement between the prominent anterosuperior aspect of the acetabulum and the femoral head-neck junction.

Based on a number of reports and consistent with our clinical observation, there are indicators that the pathologic acetabular morphology in DDH and retroversion is associated with alterations in pelvic morphology [1–7, 14, 15, 26]. Kojima et al. [14, 15], using three-dimensional (3D) CT, showed a decrease in the transverse diameter of the pelvic inlet and outlet in DDH, suggesting a general narrowing of the bony pelvis. Fujii et al. [7] used CT to examine rotational deformity of the innominate bone in DDH. They noted internal rotation of the innominate bone in dysplastic hips compared with controls. In addition, hips with acetabular retroversion, both in the control group ( $n = 4$ ) and in the setting of DDH ( $n = 9$ ), had externally rotated innominate bones. While this suggests opposing rotational abnormalities of the innominate bone between acetabular retroversion and DDH, these two groups were not compared exclusively.

By directly comparing the pelvic anatomy in DDH and acetabular retroversion, we can establish association between pelvic morphology and acetabular pathomorphology. This would allow us to diagnose acetabular pathomorphology in cases where direct radiographic assessment of the acetabulum is difficult or impossible. Furthermore, using radiographic measurements on plain AP pelvis radiographs would eliminate the need for additional, expensive, and sometimes radiation-intense imaging studies.

Our goal, therefore, was to devise a method for assessing pelvic morphology in the presence of acetabular dysplasia and DDH using plain radiographs. To this end, we developed four radiographic parameters: pelvic width index, radiographic appearance of the anterior inferior iliac spine (AIIS) sign, ilioischial angle, and obturator index. For each of these four key measurement variables, we

asked two questions: (1) How do these variables differ between dysplastic hips and retroverted hips? And (2) what is the predictive value of each variable to detect acetabular retroversion?

## Patients and Methods

We performed a retrospective comparative study of radiographic pelvic morphology between dysplastic and retroverted hips. We reviewed radiographs from our institutional database of patients who underwent periacetabular osteotomy (PAO) between March 2004 and March 2011 ( $n = 86$ ) or surgical hip dislocation between March 2007 and March 2011 ( $n = 300$ ). We defined the dysplasia group using the following parameters: lateral center-edge angle of less than  $20^\circ$  [17] and acetabular index of more than  $14^\circ$  [25]. We identified a total of 86 patients fitting these inclusion criteria. We excluded patients undergoing anteverting PAO ( $n = 17$ ), incomplete radiographic documentation ( $n = 10$ ), or previous hip or pelvis surgery ( $n = 8$ ). This left 51 patients in the dysplasia group. We defined the retroversion group as a lateral center-edge angle of more than  $25^\circ$  and the presence of three radiographic signs: the crossover sign [10, 18], ischial spine sign [12, 13], and posterior wall sign [18, 23]. We identified a total of 317 patients fitting these inclusion criteria. We excluded patients in whom not all three radiographic signs were positive ( $n = 170$ ) and with a previous history of hip trauma ( $n = 39$ ), previous hip or pelvis surgery ( $n = 25$ ), incomplete radiographic documentation ( $n = 14$ ), Legg-Calvé-Perthes disease ( $n = 10$ ), and protrusio acetabuli ( $n = 8$ ). This left 51 patients in the retroversion group. The two groups were similar in terms of age, affected side, weight, height, and BMI (Table 1). Our hospital's institutional review board approved the study.

**Table 1.** Demographic information

Parameter	Females			Males		
	Dysplasia group	Retroversion group	p value	Dysplasia group	Retroversion group	p value
Number of hips	39	23		12	28	
Age (years)*	29 ± 9.0 (15–45)	29 ± 12.0 (13–59)	0.985	26 ± 6.8 (16–36)	24 ± 8.0 (13–42)	0.369
Side (% right of all hips)	54	65	0.272	58	64	0.763
Weight (kg)*	64 ± 8.9 (48–82)	71 ± 13.9 (51–107)	0.098	75 ± 11.6 (65–101)	80 ± 14.1 (62–107)	0.381
Height (cm)*	164 ± 5.4 (155–178)	166 ± 9.1 (140–184)	0.452	174 ± 10.0 (157–186)	180 ± 9.1 (165–203)	0.180
BMI (kg/m <sup>2</sup> )*	24 ± 3.3 (18–32)	26 ± 6.1 (18–40)	0.215	25 ± 2.2 (23–30)	25 ± 3.2 (20–32)	0.922
Symphysis-sacrococcygeal distance (mm)	57 ± 13 (35–82)	69 ± 11 (47–93)	< 0.001	38 ± 10 (26–55)	54 ± 12 (30–75)	< 0.001

\* Values are expressed as mean ± SD, with the range in parentheses.

**Table 2.** Descriptions and reliability/reproducibility of the four key measurement variables


Variable	Description	Intrarater ICC/kappa	Interrater ICC/kappa
Pelvic width index (Fig. 1A)	A line (Line a) is drawn from the pubic symphysis to the most lateral edge of the ischial tuberosity, parallel to the interteardrop line. A parallel line is drawn between the midpoint of the sacrum and the most lateral point on the iliac wing (Line b). The index is calculated as the ratio of the length of Line a to the length of Line b and is expressed as a percentage.	0.98 (0.96–0.99)	0.89 (0.77–0.95)
Anterior inferior iliac spine sign (Fig. 1B)	The anterior inferior iliac spine is given a grade of 0 (indicating it is not fully seen in profile) or 1 (indicating it is seen fully in profile).	0.83 (0.6–1.0)*	0.69 (0.42–0.96)*
Ilioischial angle (Fig. 1C)	A line is drawn connecting the base of each radiographic teardrop (the interteardrop line). A second line is then drawn from the intersection of the ilioischial line and the iliopectineal line to the lateral-most point on the ipsilateral obturator foramen. The inner angle between these two lines is then measured.	0.92 (0.81–0.96)	0.92 (0.81–0.96)
Obturator index (Fig. 1D)	A line (Line c) is drawn to the maximum width of the obturator foramen. The index is calculated as the ratio of Line c to 1/2 of the length of the interteardrop line (Line d) and is expressed as a percentage.	0.97 (0.94–0.98)	0.97 (0.94–0.99)

\* The kappa value was calculated for this variable; the ICC was calculated for other remaining variables; values are expressed as mean, with 95% CI in parentheses; ICC = intraclass correlation coefficient.

Since a substantial number of dysplastic hips have a positive crossover sign [16], we further subdivided our DDH group into those with and without a crossover sign to evaluate how this affected our measurements. Twenty of the 39 (51%) hips in the female DDH group and 11 of the 12 (92%) hips in the male DDH group had a positive crossover sign.

A standardized radiographic technique was performed for all reviewed AP pelvis radiographs. All radiographs were performed in the supine position. A film focus distance of 1.2 m was used with the beam centered between the pubic symphysis and a line connecting the anterior superior iliac spine with the pelvis in neutral rotation [12, 22, 23]. The longitudinal rotation of the pelvis was verified as correct when the tip of the coccyx was in line with pubic symphysis. Images were not specifically corrected for tilt; however, we recorded the distance between the superior pubic symphysis and sacrococcygeal junction for each patient (Table 1).

Two of us (PP, CEA) independently evaluated the following key measurement variables on each radiograph (Table 2): (1) pelvic width index (Fig. 1A), (2) presence of the AIIS sign (Fig. 1B), (3) ilioischial angle (Fig. 1C), and (4) obturator index (Fig. 1D). The two observers each performed two separate sets of measurements on deidentified preoperative plain AP pelvis radiographs. The two sets of measurements were performed a minimum of 7 days apart. Inter- and intraobserver reliabilities were tested for each parameter using the intraclass correlation coefficient (ICC) for continuous value measurements and the kappa value for ordinal measurements (Table 2).



**Fig. 1A–D** The diagrams illustrate how to (A) calculate the pelvic width index ( $a/b$ ), (B) determine the AIIS sign, (C) determine the ilioischial angle, and (D) calculate the obturator index ( $c/d$ ). For a detailed explanation of these calculations, see Table 2.

Due to the lack of available comparable data in the literature, a pilot set of measurements was taken on a subset of our patients to perform a power analysis. Based on these, we detected a mean difference of 10% in the pelvic width index with an estimated SD of 10%. Using these numbers, a power analysis was performed and a minimum sample size of 32 hips for each group was required to provide an  $\alpha$  of 0.01 and a  $\beta$  of 0.10.

Results from the complete set of measurements were collected and stratified by group and sex. Normal distribution was confirmed using the Kolmogorov-Smirnov test. We compared groups using a paired t-test for continuous variables and Fisher's exact test for categorical variables. To evaluate the overall predictive performance of our key measurement variables, a receiver operating characteristic (ROC) curve was calculated for each variable. We then calculated the sensitivity and specificity of each test based on thresholds detected for the ROC curve.

## Results

The pelvic width index was smaller (both sexes:  $p < 0.001$ ) in the retroversion group than in the dysplastic group (Table 3). There was a higher (females:  $p < 0.001$ ; males:  $p = 0.013$ ) prevalence of the AIIS sign in the

retroversion group than in the dysplastic group. The ilioischial angle was higher (females:  $p < 0.001$ ; males:  $p = 0.009$ ) in the retroversion group than in the dysplastic group. The obturator index was lower (both sexes:  $p < 0.001$ ) in the retroversion group than in the dysplastic group. When we subdivided our DDH hips into those with and without a crossover sign, there was no difference between the two groups for females (Table 4).

The greatest area under the ROC curve was found for the ilioischial angle with a cutoff of  $100^\circ$  (0.971), followed by the pelvic width index  $< 56\%$  (0.969) and the obturator index  $< 40\%$  (0.925) (Fig. 2, Table 5). Compared to the other key measurement variables, the pelvic width index had the highest sensitivity (100% when pelvic width index  $< 56\%$ ), and the AIIS sign had the lowest sensitivity (59%). The highest specificity was found in the AIIS sign (96%), and the lowest specificity was found in the obturator index (86%).

## Discussion

DDH and acetabular retroversion represent two distinct acetabular pathomorphologies. As these disease processes have become better understood, it is clear they include not just alterations to the acetabulum but also distinct

**Table 3.** Results comparing the four key measurement variables for the dysplasia and retroversion groups

Variable	Females			Males		
	Dysplasia group	Retroversion group	p value	Dysplasia group	Retroversion group	p value
Pelvic width index (%)*	62 ± 5 (54–81)	50 ± 6 (30–56)	< 0.001	57 ± 4 (50–61)	49 ± 3 (44–55)	< 0.001
Anterior inferior iliac spine sign (% positive)	3	57	< 0.001	8	50	0.013
Ilioischial angle (°)*	96 ± 4 (84–102)	104 ± 3 (97–109)	< 0.001	97 ± 3 (92–102)	104 ± 3 (98–110)	0.009
Obturator index (%)*	45 ± 7 (32–69)	28 ± 10 (7–44)	< 0.001	44 ± 5 (37–50)	33 ± 6 (21–44)	< 0.001


\* Values are expressed as mean ± SD, with the range in parentheses.

**Table 4.** Results comparing the four key measurement variables for dysplasia group with and without the COS

Variable	Females			Males		
	Dysplasia + COS (n = 19)	Dysplasia – COS (n = 20)	p value	Dysplasia + COS (n = 11)	Dysplasia – COS (n = 1)	p value
Pelvic width index (%)*	62 ± 4 (54–73)	63 ± 6 (56–81)	0.689	57 ± 4 (50–61)	52	NA
Anterior inferior iliac spine sign (% positive)	5	0	0.5	9	0	NA
Ilioischial angle (°)*	95 ± 5 (84–102)	96 ± 3 (89–100)	0.274	97 ± 3 (92–102)	98	NA
Obturator index (%)*	44 ± 8 (32–69)	46 ± 5 (35–55)	0.771	44 ± 5 (37–50)	38	NA

\* Values are expressed as mean ± SD, with the range in parentheses; COS = crossover sign; NA = not applicable.

**Fig. 2A–D** The ROC curves used to determine the predictive value for (A) the pelvic width index < 56%, (B) a positive AIIS sign, (C) an ilioischial angle > 100°, and (D) an obturator index < 40%, are shown.




**Table 5.** Predictive value of the key measurement variables for detection of acetabular retroversion

Variable	Sensitivity (%)	Specificity (%)	Area under ROC curve
Pelvic width index < 56%	100 (85–100)	90 (79–97)	0.969
Positive anterior inferior iliac spine sign	59 (36–79)	96 (87–100)	0.776
Ilioischial angle > 100°	95 (77–100)	94 (84–99)	0.971
Obturator index < 40%	89 (78–97)	86 (73–94)	0.925

Values are expressed as mean, with 95% CI in parentheses; ROC = receiver operating characteristic.

morphologic changes to the entire pelvis. To date, no direct comparison has been made between the morphologic pelvic changes in these two diseases. However, there is anthropologic evidence that hip function and pelvic morphology are directly related [9]. We observed a similar pattern of pelvic morphology in our human subjects,



**Fig. 3A–B** (A) The right ½ of an AP pelvis radiograph of a dysplastic hip is compared to (B) an AP pelvis radiograph of a left hip with acetabular retroversion. In both radiographs, the coccyx is in line with the pubic symphysis, indicating no pelvic malrotation. In addition, the vertical distance between the symphysis and the sacrococcygeal joint is similar in both patients, indicating no substantially different pelvic tilt. The right hemipelvis appears similar to an obturator oblique view while the left hemipelvis appears similar to an iliac oblique view.

**Table 6.** Pelvic measurement data

Variable	Definition	Flückiger et al. [6]	Kojima et al. [14]	Trousdale et al. [26]	Fujii et al. [7]	Current study
Year		2000	2001	2002	2011	2011
Hip morphology		DDH (n = 17)	DDH (n = 40)	DDH (n = 7)	DDH (n = 50)	DDH (n = 51)
Imaging modality		Radiographs	3D CT	MRI	CT	Radiographs
Transverse pelvic inlet (cm)*	Widest area of pelvis between iliac arcuate lines	15.4 (15.0–16.1)	12.3 (10.9–13.7)	12.92 (12.6–13.7)	NA	14.6 (12.3–19.7)
Transverse pelvic expansion (cm)*	Bilateral centers of the internal surface of the acetabulum	NA	11.3 (9.5–12.8)	NA	NA	NA
Pelvic interspinous distance (cm)*	Line between the two ischial spines	11.8 (11.4–12.3)	11.2 (9.7–12.4)	11.97 (11.0–13.3)	NA	10.7 (7.7–14.3)
Transverse pelvic outlet (cm)*	Bilateral internal surface of inferior ischial tuberosity	14.2 (13.6–14.5)	12.8 (10.9–14.9)	NA	NA	13.4 (10.7–17.7)
AP inlet (cm)*	Posterior-superior pubic symphysis to sacral promontory	NA	12.9 (10.7–15.2)	12.54 (11.2–13.9)	NA	NA
AP midpelvis (cm)*	Posterior-inferior pubic symphysis to sacrum, passing through midpoint of bispinous line	NA	11.6 (9.6–13.3)	12.24 (10.5–14.5)	NA	NA
Sagittal diameter of the pelvic expansion (cm)*	S2-S3 vertebral junction and the middle posterior pubic symphysis	NA	13.4 (11.1–15.2)	NA	NA	NA
Superior iliac angle (°)†	Opening angles of the ilium at the anterior superior iliac spine	NA	NA	NA	57.0 ± 6.1	NA
Inferior iliac angle (°)†	Opening angles of the ilium at the anterior inferior iliac spine	NA	NA	NA	72.0 ± 4.3	NA
Ischiopubic angle (°)†	Closing angle of the ischiopubic pelvis	NA	NA	NA	30.6 ± 2.7	NA

\* Values are expressed as mean, with the range in parentheses; † values are expressed as mean ± SD; DDH = developmental dysplasia of the hip; NA = not available; 3D = three-dimensional.

specifically associated with their hip disease. Our goal was to directly compare four key measurement variables (pelvic width index, presence of AIIIS sign, ilioischial angle, obturator index) and evaluate their predictive value for these two different morphologies.


The major limitation to our study is that we only compared two morphologic extremes. Clearly DDH and acetabular retroversion exist on a spectrum, and mixed-type pathology is common [16]. While our key measurement variables show good predictive value and reproducibility between the two extremes, with the exception of DDH with a radiographic crossover sign, we have not explicitly evaluated them in the setting of mixed morphology. Second, we were unable to measure the actual rotation of the hemipelvis. CT, and specifically 3D CT, can provide precise measurements to determine rotational properties of the pelvis in both the coronal and sagittal planes. However, despite an inability to measure rotation directly, plain radiographs are most frequently the first, and sometimes the only, imaging studies performed on patients seeking treatment for hip disease. While using plain radiographs limits our ability to directly compare our values to previously published rotational measurements performed on CT or MRI [7, 11, 20], we believe the use of an AP pelvis radiograph is faster and more universally accessible. Finally, we did not specifically adjust for tilt when evaluating these radiographs. When we evaluated each group (DDH and retroversion) subdivided by sex, we saw the difference in means for the symphysis-sacrococcygeal distance was 12 mm in females and 16 mm in males (Table 1). This translated to a difference in pelvic tilt of 5° in females and 7° in males [24]. We do not know the extent to which this variation in tilt affected our measurement parameters.

Our results support the theory that the entire hemipelvis is involved in both acetabular dysplasia and retroversion. In DDH, the pelvis appears to be internally rotated around a sagittal axis. This leads to the radiographic appearance of the hemipelvis mimicking an obturator oblique view (Fig. 3A). Focal acetabular dysplasia has been described in cases of neuromuscular hip disease [8], but our cohort of patients and multiple other reports [2, 7, 11, 20] confirm the entire hemipelvis is involved. Even in cases of hip dysplasia with a concomitant crossover sign, our measurements do not show any difference from those performed in hips without a crossover sign [8]. In acetabular retroversion, the hemipelvis appears to be externally rotated around a sagittal axis, leading to the radiographic appearance of an iliac oblique view (Fig. 3B). Previously, only the inferior hemipelvis was implicated in acetabular retroversion [13].

As part of our radiographic review protocol, we also measured several standard pelvic measurements that have been reported previously (Table 6). The fact that our

standard measurements correlate with the literature increases the validity of our key measurement variables. Therefore, we believe these key measurement variables allow detection of hemipelvis version on a plain AP pelvis radiograph, independent of sex.

Based on the predictive value of each key measurement variable, and independent of sex, a retroverted hip is likely to be present if the pelvic width index is less than 50%, the AIIIS outline is clearly visible, the ilioischial angle is greater than 100°, and the obturator index is less than 40%. These guidelines are helpful for several reasons. First, these measurements allow one to infer acetabular morphology from existing pelvic morphology, especially when clear radiographic visualization of the acetabulum may be difficult. It has been established that pathomorphologic pelvic changes relate to acetabular disease manifesting at an early age before ossification of the acetabulum is complete [2, 20]. Therefore, it is reasonable to assume a pediatric patient who does not yet display a fully ossified acetabular rim may have occult acetabular retroversion that can be detected by our pelvic measurements (Fig. 4). It should be noted, however, we have not specifically studied these measurements in the setting of skeletally immature pelvises. Similarly, in the setting of a THA, where the normal acetabular rim may be obscured by osteophytes, these measurements may help preoperatively identify a retroverted native acetabulum (Fig. 5). Finally, these measurements can assist in decision making between reorientation and rim-trimming procedures. For instance, when our four key variables indicate acetabular



**Fig. 4A–B** (A) A radiograph of the left hip of a child at age 13 years shows indistinct anterior and posterior acetabular walls, an elevated pelvic width index, positive AIIIS sign, elevated ilioischial angle, and decreased obturator index, all indicating likely acetabular retroversion. (B) By age 15 years, the acetabular walls have ossified and retroversion is plainly evident.



**Fig. 5** A radiograph of a 56-year old man with left hip degeneration shows indistinct acetabular walls, an elevated pelvic width index, elevated ilioischial angle, and decreased obturator index. The patient was noted during subsequent THA to have a retroverted acetabulum.

retroversion with associated pelvic pathomorphology, one should consider acetabular reorientation over a rim-trimming procedure [19]. In our experience, we use these measurements as an adjunct to the standard radiographic parameters to identify the major pathology to avoid inappropriate treatment, but further clinical studies could help clarify an algorithm for using these parameters.

Distinct pelvic morphology is present in both DDH and acetabular retroversion. Based on our measurements, the sagittal rotation and coronal obliquity of the entire innominate bone are directly related to these two acetabular pathomorphologies. We presented indirect, sex-independent pelvimetric parameters indicating DDH and acetabular retroversion on an AP pelvis radiograph. Recognition of these parameters will help to understand complex morphology in hips where direct radiographic assessment of the acetabulum is difficult or impossible.

## References

- Ait-Ameur A, Wakim A, Dubousset J, Kalifa G, Adamsbaum C. The AP diameter of the pelvis: a new criterion for continence in the exstrophy complex? *Pediatr Radiol*. 2001;31:640–645.
- Albiñana J, Morcuende JA, Delgado E, Weinstein SL. Radiologic pelvic asymmetry in unilateral late-diagnosed developmental dysplasia of the hip. *J Pediatr Orthop*. 1995;15:753–762.
- Albiñana-Cilveti J, Delgado-Baeza E, Miralles-Flores C. Pelvic deformity in experimental dislocation of the growing hip. *Int Orthop*. 1992;16:317–321.
- Delgado-Baeza E, Albiñana-Cilveti J, Miralles-Flores C. Why does pelvic deformity occur in experimental dislocation of the growing hip? *J Pediatr Orthop*. 1992;12:376–383.
- Dora C, Zurbach J, Hersche O, Ganz R. Pathomorphologic characteristics of posttraumatic acetabular dysplasia. *J Orthop Trauma*. 2000;14:483–489.
- Flückiger G, Egli S, Kosina J, Ganz R. [Birth after peri-acetabular osteotomy] [in German]. *Orthopade*. 2000;29:63–67.
- Fujii M, Nakashima Y, Sato T, Akiyama M, Iwamoto Y. Pelvic deformity influences acetabular version and coverage in hip dysplasia. *Clin Orthop Relat Res*. 2011;469:1735–1742.
- Gugenheim JJ, Gerson LP, Sadler C, Tullos HS. Pathologic morphology of the acetabulum in paralytic and congenital hip instability. *J Pediatr Orthop*. 1982;2:397–400.
- Hogervorst T, Bouma H, de Boer SF, de Vos J. Human hip impingement morphology: an evolutionary explanation. *J Bone Joint Surg Br*. 2011;93:769–776.
- Jamali AA, Mladenov K, Meyer DC, Martinez A, Beck M, Ganz R, Leunig M. Anteroposterior pelvic radiographs to assess acetabular retroversion: high validity of the “cross-over-sign.” *J Orthop Res*. 2007;25:758–765.
- Jia J, Zhang L, Zhao Q, Li L, Liu X. Does medial rotational deformity of the whole pelvis universally exist in unilateral DDH? *Arch Orthop Trauma Surg*. 2011;131:1383–1388.
- Kakaty DK, Fischer AF, Hosalkar HS, Siebenrock KA, Tannast M. The ischial spine sign: does pelvic tilt and rotation matter? *Clin Orthop Relat Res*. 2010;468:769–774.
- Kalberer F, Sierra RJ, Madan SS, Ganz R, Leunig M. Ischial spine projection into the pelvis: a new sign for acetabular retroversion. *Clin Orthop Relat Res*. 2008;466:677–683.
- Kojima S, Kobayashi S, Saito N, Nawata M, Horiuchi H, Takaoka K. Morphological characteristics of the bony birth canal in patients with developmental dysplasia of the hip (DDH): investigation by three-dimensional CT. *J Orthop Sci*. 2001;6:217–222.
- Kojima S, Kobayashi S, Saito N, Nawata M, Horiuchi H, Takaoka K. Three-dimensional computed tomography evaluation of bony birth canal morphologic deformity (small pelvic cavity) after dome pelvic osteotomy for developmental dysplasia of the hip. *Am J Obstet Gynecol*. 2002;187:1591–1595.
- Li PL, Ganz R. Morphologic features of congenital acetabular dysplasia: one in six is retroverted. *Clin Orthop Relat Res*. 2003;245–253.
- Murphy SB, Ganz R, Müller ME. The prognosis in untreated dysplasia of the hip: a study of radiographic factors that predict the outcome. *J Bone Joint Surg Am*. 1995;77:985–989.
- Reynolds D, Lucas J, Klaue K. Retroversion of the acetabulum: a cause of hip pain. *J Bone Joint Surg Br*. 1999;81:281–288.
- Siebenrock KA, Schoeniger R, Ganz R. Anterior femoroacetabular impingement due to acetabular retroversion: treatment with periacetabular osteotomy. *J Bone Joint Surg Am*. 2003;85:278–286.
- Suzuki S. Deformity of the pelvis in developmental dysplasia of the hip: three-dimensional evaluation by means of magnetic resonance image. *J Pediatr Orthop*. 1995;15:812–816.
- Tannast M, Albers CE, Steppacher SD, Siebenrock KA. Hip pain in the young adult. In: Bentley G, ed. *European Instructional Lectures*. Berlin, Germany: Springer; 2011:141–154.
- Tannast M, Mistry S, Steppacher SD, Reichenbach S, Langlotz F, Siebenrock KA, Zheng G. Radiographic analysis of femoroacetabular impingement with Hip2Norm—reliable and validated. *J Orthop Res*. 2008;26:1199–1205.
- Tannast M, Siebenrock KA, Anderson SE. Femoroacetabular impingement: radiographic diagnosis—what the radi-



- ologist should know. *AJR Am J Roentgenol.* 2007;188:1540–1552.
24. Tannast M, Zheng G, Anderegg C, Burckhardt K, Langlotz F, Ganz R, Siebenrock KA. Tilt and rotation correction of acetabular version on pelvic radiographs. *Clin Orthop Relat Res.* 2005;438:182–190.
  25. Tönnis D, Heinecke A. Acetabular and femoral anteversion: relationship with osteoarthritis of the hip. *J Bone Joint Surg Am.* 1999;81:1747–1770.
  26. Trousdale RT, Cabanela ME, Berry DJ, Wenger DE. Magnetic resonance imaging pelvimetry before and after a periacetabular osteotomy. *J Bone Joint Surg Am.* 2002;84:552–556.