

Stroke

JOURNAL OF THE AMERICAN HEART ASSOCIATION



Differences and Similarities Between Spontaneous Dissections of the Internal Carotid Artery and the Vertebral Artery

Michelle von Babo, Gian Marco De Marchis, Hakan Sarikaya, Christian Stapf, Frédérique Buffon, Urs Fischer, Mirjam R. Heldner, Jan Gralla, Simon Jung, Barbara Goeggel Simonetti, Heinrich P. Mattle, Ralf W. Baumgartner, Marie-Germaine Bousser and Marcel Arnold

Stroke. 2013;44:1537-1542; originally published online April 30, 2013;
doi: 10.1161/STROKEAHA.113.001057

Stroke is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231
Copyright © 2013 American Heart Association, Inc. All rights reserved.
Print ISSN: 0039-2499. Online ISSN: 1524-4628

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://stroke.ahajournals.org/content/44/6/1537>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Stroke* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Stroke* is online at:
<http://stroke.ahajournals.org/subscriptions/>

Differences and Similarities Between Spontaneous Dissections of the Internal Carotid Artery and the Vertebral Artery

Michelle von Babo, MD*; Gian Marco De Marchis, MD*; Hakan Sarikaya, MD; Christian Stapf, MD; Frédérique Buffon, MD; Urs Fischer, MD; Mirjam R. Heldner, MD; Jan Gralla, MD; Simon Jung, MD; Barbara Goeggel Simonetti, MD; Heinrich P. Mattle, MD; Ralf W. Baumgartner, MD; Marie-Germaine Bousser, MD; Marcel Arnold, MD

Background and Purpose—To compare potential risk factors, clinical symptoms, diagnostic delay, and 3-month outcome between spontaneous internal carotid artery dissection (sICAD) and spontaneous vertebral artery dissection (sVAD).

Methods—We compared patients with sICAD (n=668) and sVAD (n=302) treated in 3 university hospitals.

Results—Patients with sICAD were older (46.3±9.6 versus 42.0±10.2 years; $P<0.001$), more often men (62.7% versus 53.0%; $P=0.004$), and presented more frequently with tinnitus (10.9% versus 3.4%; $P<0.001$) and more severe ischemic strokes (median National Institutes of Health Stroke Scale, 10±7.1 versus 5±5.9; $P<0.001$). Patients with sVAD had more often bilateral dissections (15.2% versus 7.6%; $P<0.001$) and were more often smokers (36.0% versus 28.7%; $P=0.007$). Thunderclap headache (9.2% versus 3.6%; $P=0.001$) and neck pain were more common (65.8% versus 33.5%; $P<0.001$) in sVAD. Subarachnoid hemorrhage (6.0% versus 0.6%; $P<0.001$) and ischemic stroke (69.5% versus 52.2%; $P<0.001$) were more frequent in sVAD. After multivariate analysis, sex difference lost its significance ($P=0.21$), and all other variables remained significant. Time to diagnosis was similar in sICAD and sVAD and improved between 2001 and 2012 compared with the previous 10-year period (8.0±10.5 days versus 10.7±13.2 days; $P=0.004$). In sVAD, favorable outcome 3 months after ischemic stroke (modified Rankin Scale, 0–2: 88.8% versus 58.4%; $P<0.001$), recurrent transient ischemic attack (4.8% versus 1.1%; $P=0.001$), and recurrent ischemic stroke (2.8% versus 0.7%; $P=0.02$) within 3 months were more frequent.

Conclusions—sICAD and sVAD patients differ in many aspects. Future studies should perform separate analyses of these 2 entities. (*Stroke*. 2013;44:1537-1542.)

Key Words: dissection ■ outcome ■ risk factor ■ stroke

Spontaneous cervical artery dissection (sCAD) is a major cause of stroke in young and middle-aged adults, causing up to 25% of all ischemic strokes in patients 15 to 49 years of age.^{1–3} Depending on the affected vessel, sCAD is divided into spontaneous internal carotid artery dissection (sICAD) and spontaneous vertebral artery dissection (sVAD). The causes of sCAD are poorly understood. Both constitutional and environmental factors play a role. In most studies, sICAD and sVAD were analyzed as one entity. Analogous to aortic dissection, in which pathologies differ between vessel segments,⁴ there might be differences in the mechanisms leading to sICAD and sVAD.

The Cervical Artery Dissection Ischemic Stroke Patients (CADISP) group recently reported significant differences between sICAD and sVAD.⁵ In light of these new findings, we aimed to compare the frequency of potential risk factors,

presenting clinical characteristics, and outcome after 3 months in patients with sICAD and sVAD. Furthermore, we investigated whether time to diagnosis differed between sVAD and sICAD and changed over time. To answer these questions, we retrospectively compared patients with sICAD and sVAD enrolled in our ongoing database of 1027 patients with cervicocerebral artery dissection from 3 stroke centers.

Patients and Methods

Patients

In this observational study, we analyzed prospectively collected data of 1027 consecutive patients with a first-ever sCAD who were treated in 1 of the following 3 tertiary referral hospitals: University Hospital Berne, Switzerland, from October 1990 to January 2012 (n=359);

Continuing medical education (CME) credit is available for this article. Go to <http://cme.ahajournals.org> to take the quiz.

Received October 24, 2012; accepted February 21, 2013.

From the Department of Neurology (M.v.B., G.M.D.M., H.S., U.F., M.R.H., S.J., B.G.S., H.P.M., M.A.) and Neuroradiology (J.G.), University Hospital of Bern, Bern, Switzerland; Department of Neurology, University Hospital Zurich, Zurich, Switzerland (H.S., R.W.B.); and Department of Neurology, Hôpital Lariboisière, APHP, Université Paris-Diderot, Paris, France (C.S., F.B., M.-G.B.).

*Drs von Babo and De Marchis contributed equally to this article.

Oh Young Bang, MD, PhD, was the guest editor for this article.

Correspondence to Marcel Arnold, MD, Department of Neurology, University Hospital Bern, Freiburgstrasse 10, 3010 Bern, Switzerland. E-mail marcel.arnold@insel.ch

© 2013 American Heart Association, Inc.

Stroke is available at <http://stroke.ahajournals.org>

DOI: 10.1161/STROKEAHA.113.001057

University Hospital Zurich, Switzerland, from October 1990 to July 2011 (n=268); and Lariboisière University Hospital, Paris, France, from January 1997 to December 2011 (n=400). Spontaneous dissection was defined as occurring spontaneously, after physical exertion, or a so-called minor trauma.⁶ Patients with previous sCAD were not included.

Diagnosis of sCAD was made by (1) cervical or cerebral MRI or computed tomography showing a wall hematoma or (2) computed tomography angiography, ultrasound, or intra-arterial digital subtraction angiography demonstrating a double lumen, a string sign, an intimal flap, or a dissecting aneurysm at a nonbifurcation site.⁷ On hospital admission, all images were reviewed at least by a certified neuroradiologist and a neurologist, and only patients with unequivocal dissection diagnosis were approached for study enrollment. Patients with extracranial and intracranial sCAD and sVAD were included, and patients who presented with both sCAD and sVAD (n=57) were excluded. Risk factors and clinical symptoms of patients with multiple dissections of the same artery type (affecting only sCAD or only sVAD) were evaluated only once per patient.

Baseline Characteristics and Potential Risk Factors

Baseline characteristics and potential risk factors were assessed by a structured patient interview or reconstructed on the basis of the hospital reports. The following potential risk factors were collected: arterial hypertension, diabetes mellitus, hypercholesterolemia, former or present smoking, migraine with or without aura, current use of anticonception or hormone substitution, known connective tissue disorder, febrile infection or minor trauma within the past 4 weeks before sCAD, and family history of stroke, sCAD, or connective tissue disorder.

Arterial hypertension was defined as a positive history of treated or untreated hypertension (before 2000: systolic blood pressure >160 mmHg or diastolic blood pressure >95 mmHg; after 2000 systolic blood pressure >140 mmHg or a diastolic blood pressure >90 mmHg).

Hypercholesterolemia was defined by total venous fasting cholesterol values >5.0 mmol/L or treatment with cholesterol-lowering medication. Current smoking was defined as regular smoking within the past 5 years; former smokers abstained for ≥5 years. History of migraine with or without aura was assessed by a neurologist on the basis of International Headache Society criteria.⁸ Assessed connective tissue disorders included Ehlers–Danlos syndrome type IV, Marfan syndrome, and osteogenesis imperfecta type I. Family history of connective tissue disorder, sCAD, or ischemic stroke was considered positive if ≥1 first-degree relative was affected. Febrile infection was defined as ≥1 clinical symptom of infection and body temperature >38°C. Fibromuscular dysplasia was only assessed in patients who underwent digital subtraction angiography.

Clinical Presentation

At initial encounter, all patients underwent a neurological and physical examination by a trained neurologist (including a routine blood sampling and an ECG). Patients were screened for ischemic events (transient ischemic attack [TIA], stroke, amaurosis fugax, and retinal infarction) and local symptoms (neck pain, headache, thunderclap headache, Horner syndrome, pulsatile tinnitus, cranial nerve palsy, cervical radiculopathy, and cervical spinal cord ischemia). Severity of head or neck pain was assessed using the visual analogue scale.⁹ Horner syndrome and cranial nerve palsy were only recorded when not occurring in the context of a brain stem infarction. Ischemic symptoms were classified according to the duration of clinical symptoms: TIA and amaurosis fugax if <24 hours, stroke and retinal infarction if >24 hours.

In case of a cerebral stroke, the National Institutes of Health Stroke Scale (NIHSS) score was assessed and a cerebral MRI or computed tomography was performed. If the NIHSS score was not assessed, it was reconstructed from hospital reports. All patients received an evaluation of the cerebral and cervical arteries by MRI/MR angiography,

ultrasound examination or digital subtraction angiography. Reversible cerebral vasoconstriction was defined according to Ducros et al,¹⁰ and its previously reported association with sCAD^{10,11} was further investigated in the present study.

Time to Diagnosis

We defined the onset of clinical symptoms on the basis of the first symptom, which the treating neurologist could associate with an sCAD. Time to diagnosis was defined as the time interval between the date of symptom onset and the date on which the neurovascular imaging leading to the diagnosis was performed.

Three-Month Follow-Up

A clinical follow-up (n=725) or structured telephone interview (n=82), which included the assessment of the modified Rankin Scale (mRS), was performed at 3 months. A total of 163 follow-ups (17%) were missing (proportionally equally in 112 patients with sCAD [17%] and 50 patients with sVAD [17%]). Recurrent events (dissection, stroke, TIA, and death) and current medication were registered.

Statistical Analysis

Demographic variables, potential risk factors, and clinical parameters were compared between sCAD (or multiple sCAD) and sVAD (or multiple sVAD). Mean and SD of continuous variables were calculated. Differences of continuous variables were assessed with Mann–Whitney *U*-test, differences of categorical variables with χ^2 test, or Fisher exact test, if appropriate. Hormonal anticonception was only assessed in women <50 years of age. Multivariate logistic regression analysis was performed to evaluate an independent association among baseline characteristics, potential risk factors, clinical symptoms, and diagnosis of sVAD versus sCAD. The following variables were included: age, sex, center of inclusion, NIHSS score, neck pain, thunderclap headache, pulsatile tinnitus, and subarachnoid hemorrhage (SAH). The cutoff for inclusion in the multivariate analysis was $P<0.1$ in the univariate analysis. Confidence intervals (95%) were calculated.

To determine the diagnostic delay, we performed a subgroup analysis dividing all sCAD into 2 groups: (1) diagnosis of sCAD between 1991 and 2000, and (2) diagnosis of sCAD between 2001 and 2012.

To compare the 3-month follow-up, we dichotomized the clinical outcome measured by the mRS into 2 groups: good outcome (mRS, 0–2) and poor outcome (mRS, 3–6). We ran a logistic regression analysis correcting for center of inclusion, age, sex, NIHSS score, and site of dissection to determine an independent association between the affected vessel and the outcome.

Results

Baseline Characteristics and Potential Risk Factors

Of 970 patients (59.7% men), 668 (68.9%) presented with unilateral or bilateral sCAD and 302 (31.1%) with unilateral or bilateral sVAD (Table 1).

Patients with sCAD were more often men (62.7% versus 53.0%; $P=0.004$) and older (46.3±9.6 years versus 42.0±10.2 years; $P<0.001$; Table 1). Bilateral dissection occurred half as often in patients with sCAD compared with sVAD (7.6% versus 15.2%; $P<0.001$). Current smoking was significantly less frequent in sCAD than in sVAD (28.7% versus 36.0%; $P=0.02$). No other investigated potential risk factor showed a different association between sCAD and sVAD in univariate analysis. A total of 35 sCAD were isolated intracranial dissections. Of all 668 sCAD, 6 were isolated intracranial (0.9%); of all 302 sVAD, 29 were isolated intracranial (9.6%).

Table 1. Comparison of Baseline Characteristics and Potential Risk Factors Between sICAD and sVAD

	All Patients	sICAD	sVAD	<i>P</i> Univariate	Odds Ratio (95% Confidence Interval)*	<i>P</i> Multivariate*
No. (%)	970	668 (68.9)	302 (31.1)
Age, y, mean±SD	45.0±10.0	46.3±9.6	42.0±10.2	<0.001	0.97 (0.95–0.98)	<0.001
Male sex, n (%)	579 (59.7)	419 (62.7)	160 (53.0)	0.004	1.27 (0.88–1.83)	0.211
Hypertension, n (%)	245 (25.4)	175 (26.3)	70 (23.3)	0.331
Diabetes mellitus, n (%)	22 (2.3)	12 (1.8)	10 (3.3)	0.142
Current smoking, n (%)	298 (31.0)	191 (28.7)	107 (36.0)	0.024	1.69 (1.16–2.47)	0.007
Past smoking, n (%)	150 (15.7)	111 (16.8)	39 (13.2)	0.150
Hypercholesterolemia, n (%)	414 (51.0)	271 (49.3)	143 (54.8)	0.142
Family history of stroke, n (%)	133 (14.0)	97 (14.7)	36 (12.4)	0.345
Migraine with aura, n (%)	97 (10.5)	65 (10.3)	32 (11.0)	0.725
Migraine without aura, n (%)	196 (21.2)	136 (21.5)	60 (20.6)	0.756
Hormonal anticonception, n (%)†	100 (27.3)	54 (24.1)	46 (32.4)	0.083
Minor cervical trauma, n (%)	186 (19.2)	125 (18.7)	61 (20.3)	0.561
Febrile infection, n (%)	76 (9.7)	56 (10.5)	20 (7.9)	0.262
Fibromuscular dysplasia, n (%)	18 (7.9)	9 (6.6)	9 (10)	0.349
CTD, n (%)	9 (0.9)	5 (0.8)	4 (1.4)	0.469	/	/
Family history of sCAD/CTD, n (%)	18 (1.9)	15 (2.3)	3 (1.0)	0.202	/	/
Time to diagnosis, y, mean±SD	9.2±12.2	9.0±11.6	9.8±13.5	0.586	/	/

CTD indicates connective tissue disorder; sCAD, spontaneous cervical artery dissection; sICAD, spontaneous internal carotid artery dissection; and sVAD, spontaneous vertebral artery dissection.

*Multivariate logistic regression for prediction of sVAD adjusted for age, sex, center of inclusion, multiple dissection, current smoking, cervical pain, thunderclap headache, pulsatile tinnitus, ischemic stroke, subarachnoid hemorrhage, and National Institutes of Health Stroke Scale score.

†Hormonal anticonception: only women <50 years of age.

After performing multivariate analysis, the higher frequency of sICAD in men compared with women lost its significance (*P*=0.21); all other variables showed an independent, significant association (Table 1).

Clinical Presentation

Patients with sVAD presented twice as often with neck pain (65.8% versus 33.5%; *P*<0.001), but neck pain intensity was not significantly higher compared with sICAD (visual analogue scale, 6.5±1.9 versus 6.0±1.7; *P*=0.12; Table 2). However, headache was equally common in both (sVAD 70.4% versus sICAD 71.4; *P*=0.77) but significantly stronger in its intensity in sVAD (visual analogue scale, 7.5±2.1 versus 6.7±1.8; *P*<0.001). Thunderclap headache occurred more than twice as often in sVAD than in sICAD (9.2% versus 3.6%; *P*=0.001). Of the 24, 10 patients with thunderclap headache had an SAH. Patients with sVAD were 10 times more often affected by SAH than patients with sICAD (6.0% versus 0.6%; *P*<0.001). Of the 18, 4 patients with sVAD-related SAH were associated with reversible cerebral vasoconstriction syndrome. Apart from cranial nerve palsy and Horner syndrome, which only appear in carotid artery dissections, pulsatile tinnitus was the only symptom that presented significantly more often in patients with sICAD than in sVAD (10.9% versus 3.4%; *P*<0.001).

Cerebral ischemic events (ie, stroke, TIA, retinal infarction, or amaurosis fugax, combined) affected patients with sVAD significantly more often than patients with sICAD (84.4% versus 70.4%; *P*<0.001). Ischemic stroke was significantly

more frequent in sVAD (69.5% versus 52.2%; *P*<0.001), but less severe than in sICAD (mean NIHSS_{VAD} 5±5.9 versus NIHSS_{ICAD} 10±7.1; *P*<0.001). Rare findings were spinal infarction (C2-4) in 1 patient and cervical radiculopathy (C5-7) in another, both associated with sVAD. One patient with sICAD was asymptomatic.

The most common first symptom was in both sICAD and sVAD, a local symptom (66.2% versus 65.6%; *P*=0.85). Both had equally often a TIA (10.6% versus 13.0%; *P*=0.28) or an ischemic stroke (17.2% versus 21.0%, *P*=0.18) as initial symptom. All independent associations remained significant after multivariate regression analysis (Table 2).

Time to Diagnosis

During the whole observation period, time to diagnosis did not differ between sICAD and sVAD (9.0±11.6 days versus 9.8±13.5 days; *P*=0.59). However, subgroup analysis showed that patients with sICAD treated between 2001 and 2012 were diagnosed >2 days earlier compared with those treated between 1991 and 2000 (8.0±10.5 days versus 10.7±13.2 days; *P*=0.004). In sVAD, time to diagnosis did not show a significant difference between the 2 time periods (9.3±13.2 days in 2001–2012 versus 11.8±14.7 days in 1990–2000; *P*=0.10).

Clinical Outcome and Recurrence

Clinical outcome measured by mRS was significantly better in patients after sVAD than after sICAD (mRS, 0–2: sVAD 88.8%, sICAD 58.4%; *P*<0.001; Table 3). There

Table 2. Clinical Presentation According to Dissected Vessels (ICAD vs VAD)

	All Patients	sICAD	sVAD	<i>P</i> Univariate	Odds Ratio (95% Confidence Interval)*	<i>P</i> Multivariate*
No.	970	668 (68.9)	302 (31.1)
Cervical pain, n (%)	421 (43.5)	223 (33.5)	198 (65.8)	<0.001	4.23 (2.91–6.15)	<0.001
VAS cervical pain, mean±SD	6.2±1.8	6.0±1.7	6.5±1.9	0.116
Headache, n (%)	688 (71.1)	476 (71.4)	212 (70.4)	0.767
VAS headache, mean±SD†	6.9±1.9	6.7±1.8	7.5±2.1	<0.001	1.28 (1.09–1.51)	0.002
Thunderclap headache, n (%)	44 (5.4)	20 (3.6)	24 (9.2)	0.001	2.26 (1.01–5.05)	0.048
Cranial nerve palsy, n (%)	60 (6.2)	60 (9.0)	0	0.000
Horner syndrome, n (%)	315 (32.5)	315 (47.2)	0	0.000
Pulsatile tinnitus, n (%)	83 (8.6)	73 (10.9)	10 (3.4)	<0.001	0.26 (0.11–0.63)	0.003
Cervical radiculopathy, n (%)	1 (0.1)	0	1 (0.3)	0.329
Spinal infarction, n (%)	1 (0.1)	0	1 (0.3)	0.312
Cerebral ischemia n (%)	725 (74.7)	470 (70.4)	255 (84.4)	<0.001
Ischemic stroke, n (%)	558 (57.6)	348 (52.2)	210 (69.5)	<0.001	6.60 (4.21–10.34)	<0.001
Transient ischemic attack, n (%)	279 (28.8)	190 (28.4)	89 (29.5)	0.743
No. of TIA, mean±SD	1.5±1.0	1.5±1.0	1.5±0.9	0.109
Amaurosis fugax, n (%)	80 (8.3)	80 (12.0)	0	<0.001
Retinal ischemia, n (%)	8 (0.8)	8 (1.2)	0	0.064
Subarachnoid hemorrhage, n (%)	22 (2.3)	4 (0.6)	18 (6.0)	<0.001	11.13 (2.78–44.56)	0.001
NIHSS score‡, mean±SD	5±7.3	10±7.1	5±5.9	<0.001	0.87 (0.84–0.91)	<0.001

NIHSS indicates National Institutes of Health Stroke Scale; sICAD, spontaneous internal carotid artery dissection; sVAD, spontaneous vertebral artery dissection; TIA, transient ischemic attack; and VAS, visual analogue scale.

*Multivariate logistic regression adjusted for age, sex, center of inclusion, multiple dissection, current smoking, cervical pain, thunderclap headache, pulsatile tinnitus, ischemic stroke, subarachnoid hemorrhage, and NIHSS score.

†Multivariate logistic regression for sCAD adjusted for age, sex, center of inclusion, multiple dissection, current smoking, cervical pain, headache intensity (VAS), thunderclap headache, pulsatile tinnitus, ischemic stroke, cerebral ischemia, subarachnoid hemorrhage, and NIHSS score.

‡Multivariate logistic regression adjusted for age, sex, multiple dissection, current smoking, cervical pain, thunderclap headache, pulsatile tinnitus, cerebral ischemia, subarachnoid hemorrhage, and NIHSS score.

was a trend toward higher mortality after sICAD within the first 3 months after dissection ($P=0.059$) in univariate analysis.

Seven patients (0.9%) had a recurrent isolated dissection, none of them after a multiple dissection. The amount of recurrent dissections did not differ between sICAD and sVAD, but 4 of the 5 recurrent dissections after sICAD affected the vertebral artery, whereas 1 of the 2 recurrent dissections after sVAD affected the contralateral vertebral artery, and the other one affected both the internal carotid and the contralateral vertebral artery. Recurrent TIA and stroke were more common after sVAD (TIA: sVAD 4.8%, sICAD 1.1%, $P=0.001$; stroke: sVAD 2.8%, sICAD 0.7%, $P=0.02$).

Multivariate analysis was performed adjusting for age, sex, center of inclusion, NIHSS score, and site of dissection to determine an independent association between the affected vessel and outcome. Only the higher baseline NIHSS score was highly associated with a poor outcome ($P<0.001$) and higher mortality ($P<0.001$).

Discussion

This is the second large study analyzing differences in baseline characteristics, clinical symptoms, potential risk factors, and outcome between sICAD and sVAD.

Baseline Characteristics and Risk Factors

The frequency of sICAD was about twice as high compared with sVAD, which is in line with previous findings.^{12,13} In agreement with previous studies, bilateral dissection affected the vertebral artery twice as often as the internal carotid artery.^{5,14} Patients with sICAD were older and more often men, compared with patients with sVAD, which is consistent with previous findings.^{5,13–15} These age and sex differences are difficult to explain. However, because patients with sVAD were younger compared with patients with sICAD, it is possible that risk factors or triggering events, such as traumas that are less related to advanced age, are involved more in the pathogenesis of sVAD than sICAD. Of interest, younger age at CAD onset and multiple CADs have been reported to be more common in women.^{16–18} These differences in age and sex may suggest that genetic differences, sexual hormones, or even sex-related differences in arterial wall properties may play a role.^{19,20} Moreover, the vertebral artery and the carotid artery do have a different embryological origin.^{4,21} Of interest, current smoking was more frequent in patients with sVAD. This result is in line with the observation that smoking was associated with dilatation of only the thoracic aorta, which has the same embryological origin as the vertebral artery, and not of the ascending aorta, which has the same roots as the internal

Table 3. Three-Month Follow-Up: Outcome and Events According to Dissection Site (sICAD vs sVAD)

	All Patients	sICAD	sVAD	P Univariate
Modified Rankin Scale 0–2 (patients with ischemic stroke), n (%)	293 (70.1)	150 (58.4)	143 (88.8)	<0.001
Recurrent dissection, n (%)	7 (0.9)	5 (0.9)	2 (0.8)	0.999
Recurrent TIA, n (%)	18 (2.3)	6 (1.1)	12 (4.8)	0.001
Recurrent stroke, n (%)	11 (1.4)	4 (0.7)	7 (2.8)	0.021
Death, n (%)	15 (1.9)	12 (2.4)	2 (0.8)	0.162
Death (patients with ischemic stroke), n (%)	14 (3.3)	12 (4.7)	2 (1.2)	0.058

sICAD indicates spontaneous internal carotid artery dissection; sVAD, spontaneous vertebral artery dissection; and TIA, transient ischemic attack.

carotid artery.^{4,21} Diverging from previous findings, no difference in previous febrile infection or minor cervical trauma could be found.^{5,14}

Clinical Presentation

A new finding is that thunderclap headache was more than twice as common in sVAD compared with sICAD, which is only partly explained by a higher amount of SAH in sVAD. The reasons for this association remain unknown.

Pulsatile tinnitus, which corresponds to noise evoked by turbulent, nonlaminar blood flow and transmitted to the inner ear,²² occurred thrice more often in sICAD. This may be explained by the anatomic proximity of the typical location of ICAD to the inner ear. Two patients with sVAD presented with rare manifestations: 1 patient with a cervical radiculopathy and another with a cervical spine infarction.

In agreement with the CADISP study,⁵ ischemic stroke was less frequent but more severe in patients with sICAD compared with sVAD. Ischemic strokes in the anterior circulation of undefined pathogenesis have previously been reported to cause a more severe impairment compared with ischemic strokes in the posterior circulation.^{23–25}

A 10-fold amount of patients with sVAD experienced an SAH compared with patients with sICAD. Most of them affected either only the intracranial segment of the vertebral artery or extended to the intracranial segments. Of the 18, 4 patients with sVAD-related SAH were associated with a reversible cerebral vasoconstriction syndrome, which is known to cause SAH in some patients without evidence for intracranial extension of the cervical artery dissections and which was diagnosed in 12 patients in total.¹¹

Time to Diagnosis

Local clinical symptoms of sVAD are not specific compared with those of sICAD, which are typically a painful Horner syndrome or lower cranial nerve palsies. Hence, the diagnosis of sVAD poses a challenge. However, we could not find a significant diagnostic delay of sVAD compared with sICAD. An interesting observation is the earlier diagnosis of sICAD between 2001 and 2012 compared with the diagnosis made between 1990 and 2000. This indicates a diagnostic improvement, probably because of a progress in imaging modalities and an increased awareness of the disease. In sVAD, no diagnostic improvement in the later time period could be shown. This points out that there is still

much room for improvement of the clinical and diagnostic work-up in sCAD.

Clinical Outcome and Recurrence

In univariate analysis, a good outcome (mRS, 0–2) in patients who experienced an sICAD-related stroke was significantly less common compared with patients with an sVAD-related stroke. After multivariate analyses, the only parameter independently predicting functional outcome was the NIHSS score, which was higher in patients with stroke because of sICAD as opposed to sVAD.

Recurrent dissections within the first 3 months were as rare in sICAD as in sVAD, and there were no recurrent dissections after multiple dissections. Interestingly, recurrent dissections were mostly located in the vertebral artery, no matter whether the initial dissection affected the vertebral or carotid artery, which to our knowledge, has not been previously reported. Only 1 recurrent dissection affected the same vessel on the same side again.

Both recurrent TIA and recurrent ischemic stroke were more common after sVAD. We observed a nonsignificant trend toward higher mortality in patients with an sICAD-related ischemic stroke.

Strengths and Limitations

Strengths of this study are the large sample size and the significant results after correction for multiple comparisons. Because of the low incidence of sCAD in the general population, we had to include patients over a long time period and from 3 different centers. The centers of inclusions consisted only of university hospitals. Thus, a selection bias with more severe cases, especially more patients with strokes, might have occurred and patients with only local symptoms might have been underrepresented. In addition, one-sixth of the patients had no 3-month follow-up examination. However, the rate of patients with lost follow-up did not differ between sVAD and sICAD patients.

Conclusions

Although patients with sICAD and sVAD have many features in common, they differ in age, clinical presentation, risk factors, outcome, and rates of early recurrent ischemic events. The observed differences do not provide enough evidence to propose different treatment strategies for sVAD and sICAD. However, future studies on sCAD should perform separate analyses of the 2 entities.

Acknowledgments

We thank Pietro Ballinari, PhD, University of Bern, for statistical analyses.

Disclosures

Drs De Marchis, Fischer, Simonetti, Sarikaya, and Jung are supported in the frame of the Special Program for University Medicine (SPUM-Grant 33CM30-124119), which is funded by the Swiss National Science Foundation (SNSF). The other authors have no conflicts to report.

References

- Chan MT, Nadareishvili ZG, Norris JW. Diagnostic strategies in young patients with ischemic stroke in Canada. *Can J Neurol Sci.* 2000;27:120–124.
- Putala J, Metso AJ, Metso TM, Konkola N, Kraemer Y, Haapaniemi E, et al. Analysis of 1008 consecutive patients aged 15 to 49 with first-ever ischemic stroke: the Helsinki young stroke registry. *Stroke.* 2009;40:1195–1203.
- Schievink WI. Spontaneous dissection of the carotid and vertebral arteries. *N Engl J Med.* 2001;344:898–906.
- Allaire E, Schneider F, Saucy F, Dai J, Cochenne F, Michineau S, et al. New insight in aetiopathogenesis of aortic diseases. *Eur J Vasc Endovasc Surg.* 2009;37:531–537.
- Debette S, Grond-Ginsbach C, Bodenat M, Kloss M, Engelter S, Metso T, et al.; Cervical Artery Dissection Ischemic Stroke Patients (CADISP) Group. Differential features of carotid and vertebral artery dissections: the CADISP study. *Neurology.* 2011;77:1174–1181.
- Dittrich R, Rohsbach D, Heidbreder A, Heuschmann P, Nassenstein I, Bachmann R, et al. Mild mechanical traumas are possible risk factors for cervical artery dissection. *Cerebrovasc Dis.* 2007;23:275–281.
- Arnold M, Cumurciuc R, Stapf C, Favrole P, Berthet K, Boussier MG. Pain as the only symptom of cervical artery dissection. *J Neurol Neurosurg Psychiatry.* 2006;77:1021–1024.
- Headache Classification Subcommittee of the International Headache Society. The International Classifications of Headache Disorders. *Cephalalgia.* 2004;24:9–160.
- Bijur PE, Silver W, Gallagher EJ. Reliability of the visual analog scale for measurement of acute pain. *Acad Emerg Med.* 2001;8:1153–1157.
- Ducros A, Boukobza M, Porcher R, Sarov M, Valade D, Boussier MG. The clinical and radiological spectrum of reversible cerebral vasoconstriction syndrome. A prospective series of 67 patients. *Brain.* 2007;130(Pt 12):3091–3101.
- Arnold M, Camus-Jacquin M, Stapf C, Ducros A, Viswanathan A, Berthet K, et al. Postpartum cervicocephalic artery dissection. *Stroke.* 2008;39:2377–2379.
- Redekop GJ. Extracranial carotid and vertebral artery dissection: a review. *Can J Neurol Sci.* 2008;35:146–152.
- Lee VH, Brown RD Jr, Mandrekar JN, Mokri B. Incidence and outcome of cervical artery dissection: a population-based study. *Neurology.* 2006;67:1809–1812.
- Dziewas R, Konrad C, Dräger B, Evers S, Besselmann M, Lüdemann P, et al. Cervical artery dissection—clinical features, risk factors, therapy and outcome in 126 patients. *J Neurol.* 2003;250:1179–1184.
- Metso TM, Debette S, Grond-Ginsbach C, Engelter ST, Leys D, Brandt T, et al. Age-dependent differences in cervical artery dissection. *J Neurol.* 2012;259:2202–2210.
- Arnold M, Kappeler L, Georgiadis D, Berthet K, Keserue B, Boussier MG, et al. Gender differences in spontaneous cervical artery dissection. *Neurology.* 2006;67:1050–1052.
- Arnold M, De Marchis GM, Stapf C, Baumgartner RW, Nedeltchev K, Buffon F, et al. Triple and quadruple spontaneous cervical artery dissection: presenting characteristics and long-term outcome. *J Neurol Neurosurg Psychiatry.* 2009;80:171–174.
- Schievink WI, Mokri B, O'Fallon WM. Recurrent spontaneous cervical-artery dissection. *N Engl J Med.* 1994;330:393–397.
- Krejza J, Arkuszewski M, Kasner SE, Weigle J, Ustymowicz A, Hurst RW, et al. Carotid artery diameter in men and women and the relation to body and neck size. *Stroke.* 2006;37:1103–1105.
- Hansen F, Mangell P, Sonesson B, Länne T. Diameter and compliance in the human common carotid artery—variations with age and sex. *Ultrasound Med Biol.* 1995;21:1–9.
- Satti SR, Cerniglia CA, Koenigsberg RA. Cervical vertebral artery variations: an anatomic study. *AJNR Am J Neuroradiol.* 2007;28:976–980.
- Pelkonen O, Tikkakoski T, Luotonen J, Sotaniemi K. Pulsatile tinnitus as a symptom of cervicocephalic arterial dissection. *J Laryngol Otol.* 2004;118:193–198.
- De Marchis GM, Kohler A, Renz N, Arnold M, Mono ML, Jung S, et al. Posterior versus anterior circulation strokes: comparison of clinical, radiological and outcome characteristics. *J Neurol Neurosurg Psychiatry.* 2011;82:33–37.
- Libman RB, Kwiatkowski TG, Hansen MD, Clarke WR, Woolson RF, Adams HP. Differences between anterior and posterior circulation stroke in TOAST. *Cerebrovasc Dis.* 2001;11:311–316.
- Förster A, Gass A, Kern R, Griebel M, Hennerici MG, Szabo K. Thrombolysis in posterior circulation stroke: stroke subtypes and patterns, complications and outcome. *Cerebrovasc Dis.* 2011;32:349–353.