

*Ergod. Th. & Dynam. Sys.* (2008), **28**, 1465–1478 © 2008 Cambridge University Press  
doi:10.1017/S0143385707000880 Printed in the United Kingdom

## Laminated currents

JOHN ERIK FORNÆSS<sup>†</sup>, YINXIA WANG<sup>‡</sup> and ERLEND FORNÆSS WOLD<sup>§</sup>

<sup>†</sup> *Mathematics Department, The University of Michigan, East Hall, Ann Arbor,  
MI 48109, USA*

(*e-mail: fornaess@umich.edu*)

<sup>‡</sup> *Department of Mathematics, Henan Polytechnic University, Jiaozuo, 454000, China*

(*e-mail: yinxiawang@gmail.com*)

<sup>§</sup> *Mathematisches Institut, Universität Bern, Sidlerstr. 5, CH-3012 Bern, Switzerland*

(*e-mail: erlendfw@math.uio.no*)

(Received 9 March 2007 and accepted in revised form 26 October 2007)

*Abstract.* In this paper we prove the equivalence of two definitions of laminated currents.

### 1. Introduction

Let  $K$  be a relatively-closed subset of the bidisc  $\Delta^2(z, w) = \{(z, w); |z|, |w| < 1\}$ . We suppose that  $K$  is a disjoint union of holomorphic graphs,  $w = f_\alpha(z)$ , where  $f_\alpha$  is a holomorphic function on the unit disc with  $f_\alpha(0) = \alpha$  and  $|f_\alpha(z)| < 1$ . We let  $\mathcal{L}$  denote the lamination of  $K$ .

There are two notions of laminated currents that we will discuss. Let  $T$  be a positive closed  $(1, 1)$ -current supported on  $K$ . We assume that  $T$  is the restriction of a positive closed current defined on a neighborhood of  $\overline{\Delta}^2$ . We denote by  $[V_\alpha]$  the current of integration along the graph of  $f_\alpha$ . Let  $\lambda$  denote a continuous  $(1, 0)$ -form which at  $(z, f_\alpha(z))$  equals a non-zero multiple of  $dw - f'_\alpha(z) dz$ .

*Definition 1.* We say that  $T$  is a *laminated current directed by  $\mathcal{L}$*  if  $\lambda \wedge T = 0$  for any such  $\lambda$ .

These are the same as Sullivan's *structure currents* [10]. The present terminology was introduced by Berndtsson and Sibony in [1], and such currents were treated further in [4]. In accordance with Dujardin [3] we also define the following.

*Definition 2.* We say that  $T$  is a *laminated current subordinate to  $\mathcal{L}$*  if there is a positive measure  $\mu$  such that  $T = \int_\alpha [V_\alpha] d\mu(\alpha)$ .

Our main result is the following.

**MAIN THEOREM.** *The current  $T$  is subordinate to  $\mathcal{L}$  if and only if it is directed by  $\mathcal{L}$ .*

We note that this is a result by Sullivan in the case of the lamination being smooth, i.e. the graphs vary smoothly with  $\alpha$  [10]. In the continuous setting Dujardin has shown that if a current  $T$  is dominated by a current subordinate to  $\mathcal{L}$  then  $T$  is subordinate to  $\mathcal{L}$ .

The part of Sullivan’s proof that does not go through automatically in the non-smooth case is a certain approximation step, and so in the present article we are concerned with approximation of partially-smooth functions. In [5] the authors proved such an approximation theorem in the case of laminations in  $\mathbb{R}^2$  and in  $\mathbb{R}^3$ . In the last section we show that the main theorem breaks down for Riemann-surface laminations in higher dimension.

For related material on laminated currents the reader may consult the paper of Bedford et al [2].

2. *Holomorphic motions and preliminary estimates for slopes of holomorphic graphs*

We need to know how the lamination  $\mathcal{L}$  defined above varies with the parameter  $\lambda$ , and we use the fact that it defines a holomorphic motion. Let  $\Delta := \{z \in \mathbb{C} : |z| < 1\}$  denote the unit disc in  $\mathbb{C}$ . A holomorphic motion is a subset  $E$  of the complex plane  $\mathbb{C}$  (or the Riemann sphere  $\widehat{\mathbb{C}}$ ) and a map  $f : \Delta \times E \rightarrow \mathbb{C}$  (or  $\widehat{\mathbb{C}}$ ) such that  $f(0, \cdot) = \text{id}$ ,  $f(\lambda, \cdot)$  is injective for each  $\lambda$ , and  $f(\cdot, z)$  is holomorphic for each  $z$ . The lamination  $\mathcal{L}$  defines a holomorphic motion.

Let us briefly recall some facts. It is known [9] that any holomorphic motion has an extension to a holomorphic motion  $f : \Delta \times \mathbb{C} \rightarrow \mathbb{C}$ . This means that we may regard  $K$  as a subset of a lamination of  $\Delta \times \mathbb{C}$ . From [8] we have that  $f$  is automatically jointly continuous in  $(\lambda, z)$ ; in fact the map  $(\lambda, z) \mapsto (\lambda, f_\lambda(z))$  is a homeomorphism onto  $\Delta \times \mathbb{C}$ . Moreover,  $f(\lambda, \cdot)$  is quasi-conformal for each  $\lambda$ , and  $f(\lambda, \cdot)$  distorts cross-ratios by a bounded amount depending on  $|\lambda|$ . In particular we have the following. If  $C$  is compact in  $\mathbb{C}^*$  and  $x, y, z$  are three distinct points in  $\mathbb{C}$  with  $c_0 = (x - y)/(z - y) \in C$ , then  $(f_\lambda(x) - f_\lambda(y))/(f_\lambda(z) - f_\lambda(y))$  is close to  $c_0$  depending only on  $|\lambda|$  (for a fixed  $C$ ). To see this one can consider the map  $\lambda \mapsto (f_\lambda(x) - f_\lambda(y))/(f_\lambda(z) - f_\lambda(y))$ , a map from the unit disk to  $\mathbb{C} \setminus \{0, 1\}$ , and use the fact that it has to be distance-decreasing in the Poincaré metric. Finally we recall that  $f(\lambda, \cdot)$  is Hölder continuous with exponent  $1 + \epsilon(|\lambda|)$ .

Next we need a basic estimate on slopes of the graphs. For the benefit of the reader we include the details of this well-known fact. We denote by  $\mathcal{O}(\Omega)$  the space of holomorphic functions on  $\Omega$ . Let  $\|\cdot\|_\infty$  denote the sup norm. Set

$$H^\infty = H^\infty(\Delta) = \{f \in \mathcal{O}(\Delta) : \|f\|_\infty < \infty\}.$$

Also, if  $0 < C < \infty$  we set

$$H_C^\infty = H_C^\infty(\Delta) = \{f \in \mathcal{O}(\Delta) : \|f\|_\infty < C\}.$$

LEMMA 1. *If  $f \in H_1^\infty(\Delta)$  and  $f(z) \neq 0$  for all  $z \in \Delta$ , then*

$$|f'(0)| \leq 2|f(0)| \log(1/|f(0)|).$$

*Proof.* Pick a holomorphic function  $f(z)$  on the unit disc such that  $0 \neq |f(z)| < 1$  for all  $z \in \Delta$ . We can replace  $f(z)$  by  $e^{i\theta} f(z)$  for any real  $\theta$ . This does not change  $|f(0)|$  or  $|f'(0)|$ . Hence we can assume that  $f(0) > 0$ .

We set  $h(z) := \log f(z)$ . Then  $h(z)$  is a holomorphic function on the unit disc and  $\operatorname{Re}(h(z)) < 0$ . We can also choose a branch of the logarithm so that  $\log(f(0)) = -a < 0$ . If  $k(z) = h(z)/a$ , then  $k(z)$  is a holomorphic function on the unit disc and  $k(0) = -1$ ,  $\operatorname{Re}(k(z)) < 0$ . We define  $L(w) = (w + 1)/(w - 1)$ . Then  $L(-1) = 0$  and if  $\operatorname{Re}(w) < 0$  then  $|L(w)| < 1$ . Then  $\Gamma(z) := L(k(z))$  is a holomorphic function from the unit disc to the unit disc. Moreover  $\Gamma(0) = L(k(0)) = L(-1) = 0$ . Since  $\Gamma(0) = 0$  and  $|\Gamma(z)| < 1$  we can apply the Schwarz lemma. So we can conclude that  $|\Gamma'(0)| \leq 1$ . By the chain rule,  $\Gamma'(0) = L'(k(0))k'(0) = L'(-1)k'(0)$ . Since  $L'(w) = -2/(w - 1)^2$  we get  $\Gamma'(0) = -2/(-1 - 1)^2 k'(0)$  and therefore  $k'(0) = -2\Gamma'(0)$ . Hence we get  $|k'(0)| \leq 2$ . Since  $k(z) = h(z)/a$ , we can conclude next that  $|k'(0)| = |h'(0)|/a$ . Hence  $|h'(0)| = a|k'(0)| \leq a \cdot 2$ , so  $|h'(0)| \leq 2a$ . Next recall that  $h(z) = \log f(z)$ , so  $f(z) = e^{h(z)}$ . Hence  $f'(z) = e^{h(z)}h'(z)$ . Therefore  $f'(0) = e^{h(0)}h'(0) = f(0)h'(0)$ . Hence  $|f'(0)| \leq |f(0)||h'(0)|$ . This implies that  $|f'(0)| \leq 2a|f(0)|$ . Now recall that  $\log f(0) = -a$ . But we have set this up so that  $\log f(0) = \log |f(0)| + i \arg f(0)$  is real-valued. So  $\log |f(0)| = -a$ , i.e.  $\log(1/|f(0)|) = a$ . Therefore  $|f'(0)| \leq 2a|f(0)| = 2|f(0)| \log(1/|f(0)|)$ . This concludes the proof of the lemma.  $\square$

**COROLLARY 1.** *Suppose that we have two functions  $f$  and  $g$  holomorphic on the unit disc with  $f - g \in H_1^\infty(\Delta)$ . Suppose that  $f(z) \neq g(z)$  for each  $z \in \Delta$ . We then have the estimate  $|f'(z) - g'(z)| \leq 4|f(z) - g(z)| \log(1/|f(z) - g(z)|)$  for all  $z \in \Delta$ ,  $|z| < 1/2$ .*

*Proof.* Pick  $z$ ,  $|z| < 1/2$ . We define  $G(w) = f(z + w/2) - g(z + w/2)$ . Then  $G(w)$  satisfies the conditions of Lemma 1. Hence  $|G'(0)| \leq 2|G(0)| \log(1/|G(0)|)$ . Therefore,

$$\frac{1}{2}|f'(z) - g'(z)| \leq 2|f(z) - g(z)| \log \frac{1}{|f(z) - g(z)|}. \quad \square$$

### 3. Approximation for complex curves in $\mathbb{C}^2$

We assume that for every  $c = (a, b) = (a + ib) \in \mathbb{C}$  we have a holomorphic graph  $\Gamma_c$  given by  $w = y_1 + iy_2 = f_c(z)$ ,  $z = x_1 + ix_2 \in \Delta$ . We assume that all surfaces are disjoint and that there is a surface through every point in  $\Delta \times \mathbb{C}$ . We assume that  $f_c(0) = c$ .

Let  $\pi : \Delta \times \mathbb{C} \rightarrow \mathbb{C}$  be defined by  $\pi(z, f_c(z)) = c$ . By the discussion in the previous section the function  $\pi$  is continuous.

Fix a positive constant  $R$ . By Corollary 1 there exists a positive real number  $\delta_0 > 0$  such that if  $z \in (1/2)\Delta$  and if  $c, c' \in R\Delta$  with  $|c - c'| < \delta_0$  then

$$\left| \frac{\partial}{\partial z} f_{c'}(z) - \frac{\partial}{\partial z} f_c(z) \right| \leq 4 \cdot |f_{c'}(z) - f_c(z)| \log \frac{1}{|f_{c'}(z) - f_c(z)|}. \quad (1)$$

We define a class of partially-smooth functions:

$$\mathcal{A} := \left\{ \phi \in \mathcal{C}(\Delta \times \mathbb{C}) : \phi(z, f_c(z)) \in \mathcal{C}^1(\Gamma_c), \right. \\ \left. \Phi(x_1, x_2, w) := \frac{\partial}{\partial x_1} \phi(x_1, x_2, f_c(x_1, x_2)), w = f_c(x_1, x_2) \in \mathcal{C}(\Delta \times \mathbb{C}), \right. \\ \left. \Psi(x_1, x_2, w) := \frac{\partial}{\partial x_2} \phi(x_1, x_2, f_c(x_1, x_2)), w = f_c(x_1, x_2) \in \mathcal{C}(\Delta \times \mathbb{C}) \right\}.$$

**THEOREM 1.** *Let  $\phi \in \mathcal{A}$ , let  $R$  be a positive real number and let  $\epsilon > 0$ . Then there exists a function  $\psi \in C^1(\Delta \times R\Delta)$  such that for every point  $(x_1, x_2, w) = (x_1, x_2, f_c(x_1, x_2)) \in \Delta \times R\Delta$ :*

$$\begin{aligned}
 &|\psi(x_1, x_2, w) - \phi(x_1, x_2, w)| < \epsilon, \\
 &\left| \frac{\partial}{\partial x_1}[\psi(x_1, x_2, f_c(x_1, x_2))] - \frac{\partial}{\partial x_1}[\phi(x_1, x_2, f_c(x_1, x_2))] \right| < \epsilon, \\
 &\left| \frac{\partial}{\partial x_2}[\psi(x_1, x_2, f_c(x_1, x_2))] - \frac{\partial}{\partial x_2}[\phi(x_1, x_2, f_c(x_1, x_2))] \right| < \epsilon.
 \end{aligned}$$

We will prove the theorem using the following result.

**PROPOSITION 1.** *Let  $g \in \mathcal{A}$ ,  $g(x_1, x_2, f_{a+ib}(x_1, x_2)) = a$ , and let  $R$  be a positive real number. There exists a positive real number  $t_0$  such that the following holds. For all  $\epsilon > 0$  there exists a function  $h \in C^1(t_0\Delta \times R\Delta)$  such that for every point  $(x_1, x_2, w) = (x_1, x_2, f_c(x_1, x_2)) \in t_0\Delta \times R\Delta$ :*

$$\begin{aligned}
 &|h(x_1, x_2, w) - g(x_1, x_2, w)| < \epsilon, \\
 &\left| \frac{\partial}{\partial x_1}[h(x_1, x_2, f_c(x_1, x_2))] \right| < \epsilon, \\
 &\left| \frac{\partial}{\partial x_2}[h(x_1, x_2, f_c(x_1, x_2))] \right| < \epsilon.
 \end{aligned}$$

The same result holds if we replace  $a$  by  $b$  in the definition of  $g$ .

*Proof of Theorem 1 from Proposition 1.*

**LEMMA 2.** *Let  $p \in \Delta$  be a point, and let  $R, t_0$  be positive real numbers such that  $\Delta_{t_0}(p) \subset\subset \Delta$ . Consider the lamination restricted to  $\Delta_{t_0}(p) \times \mathbb{C}$ . If the conclusion of Proposition 1 holds on  $\Delta_{t_0}(p) \times R\Delta$  (with respect to projection onto  $\{p\} \times \mathbb{C}$ ), then the conclusion of Theorem 1 holds on  $\Delta_{t_0}(p) \times R\Delta$ .*

*Proof.* Let  $\pi = (\pi_1, \pi_2)$  denote the projection onto  $\{p\} \times \mathbb{C}$ . For each  $j, k \in \mathbb{Z}$  and  $\delta > 0$  we let  $c^\delta(j, k)$  denote the point  $(p, j\delta + k\delta i)$ . Let  $\Lambda_j^\delta$  denote the  $C^1$ -smooth function defined by  $\Lambda_j^\delta(t) = \cos^2[\pi/2\delta(t - j\delta)]$  when  $(j - 1)\delta \leq t \leq (j + 1)\delta$  and 0 otherwise. For each  $c^\delta(j, k)$  we first define a function

$$\psi_{jk}^\delta(z) := \phi(z, f_{c^\delta(j,k)}(z)),$$

and then we define a preliminary approximation

$$\psi^\delta(z, w) = \sum_{j,k} \psi_{jk}^\delta(z) \Lambda_j(\pi_1(z, w)) \Lambda_k(\pi_2(z, w)).$$

Let  $(z_0, w_0) \in \Delta_{t_0}(p) \times R\Delta$ . Then  $\pi(z_0, w_0)$  is contained in a square with corners  $c^\delta(j, k), c^\delta(j + 1, k), c^\delta(j, k + 1)$  and  $c^\delta(j + 1, k + 1)$ , and

$$\psi^\delta(z_0, w_0) = \sum_{m=j, j+1, n=k, k+1} \psi_{mn}^\delta(z_0) \Lambda_m(\pi_1(z_0, w_0)) \Lambda_n(\pi_2(z_0, w_0)).$$

We have

$$\begin{aligned}
 |\psi^\delta(z_0, w_0) - \phi(z_0, w_0)| &= \left| \sum_{m=j, j+1, n=k, k+1} [\psi_{mn}^\delta(z_0) - \phi(z_0, w_0)] \right. \\
 &\quad \left. \times \Lambda_m^\delta(\pi_1(z_0, w_0)) \cdot \Lambda_n^\delta(\pi_2(z_0, w_0)) \right| \\
 &\leq \max_{m=j, j+1, n=k, k+1} |\psi_{mn}^\delta(z_0) - \phi(z_0, w_0)|.
 \end{aligned}$$

Since the map from  $\overline{\Delta}_{t_0}(p) \times \mathbb{C}$  defined by  $(z, \alpha) \mapsto (z, f_\alpha(z))$  is a homeomorphism it follows that  $\psi^\delta \rightarrow \phi$  uniformly as  $\delta \rightarrow 0$ .

Next we approximate derivatives along leaves. Let  $\alpha$  be such that  $(z_0, w_0) = (z_0, f_\alpha(z_0))$ . Since the functions  $\Lambda_j^\delta \circ \pi_i$  are constant along leaves,

$$\begin{aligned}
 &\left| \frac{\partial}{\partial x_i} [\psi^\delta(z_0, f_\alpha(z_0)) - \phi(z_0, f_\alpha(z_0))] \right| \\
 &= \left| \sum_{m=j, j+1, n=k, k+1} \left[ \frac{\partial}{\partial x_i} [\psi_{mn}^\delta(z_0) - \phi(z_0, f_\alpha(z_0))] \right] \right. \\
 &\quad \left. \times \Lambda_m^\delta(\pi_1(z_0, f_\alpha(z_0))) \cdot \Lambda_n^\delta(\pi_2(z_0, f_\alpha(z_0))) \right| \\
 &\leq \max_{m=j, j+1, n=k, k+1} \left| \frac{\partial}{\partial x_i} [\psi_{mn}^\delta(z_0) - \phi(z_0, f_\alpha(z_0))] \right|.
 \end{aligned}$$

It follows that  $\psi^\delta \rightarrow \phi$  also in  $C^1$ -norm on leaves.

Now the conclusion of Lemma 2 follows because the functions  $\pi_j$  can be approximated uniformly and in  $C^1$ -norm on leaves. □

For each point  $p \in \Delta$  there exists by Proposition 1 a positive real number  $t_p$  such that constant approximation is possible on  $\Delta_{t_p}(p) \times R\Delta$ . Hence by Lemma 2 approximation of functions in  $\mathcal{A}$  is possible.

We may then choose a locally-finite cover  $\{U_\alpha\}_{\alpha \in \mathbb{N}}$  of  $\Delta$  by disks such that approximation by functions in  $\mathcal{A}$  is possible on each  $U_\alpha \times R\Delta$ . Let  $\{\varphi_\alpha\}$  be a partition of unity subordinate to  $\{U_\alpha\}$ . For each  $\alpha$  let  $C_\alpha = \|\nabla \varphi_\alpha\|$ .

For a given  $\epsilon_\alpha$  let  $g_{\epsilon_\alpha}$  be an  $\epsilon_\alpha$ -approximating function of  $\phi$  on  $U_\alpha \times R\Delta$ . We will show that there is a sequence  $\{\epsilon_\alpha\}$  such that the function

$$\psi = \sum_{\alpha} \varphi_\alpha \cdot g_{\epsilon_\alpha}$$

satisfies the claims of the theorem.

Let  $z_0 \in U_\alpha$ , and let  $\{\alpha_1, \dots, \alpha_m\}$  be the finite set of  $\alpha$ 's, such that the support of  $\phi_\alpha$  intersects  $U_\alpha$ . Then

$$\psi(z, f_c(z)) = \sum_{i=1}^m \varphi_{\alpha_i}(z) \cdot g_{\epsilon_{\alpha_i}}(z, f_c(z)),$$

for all  $z$  near  $z_0$ . Then

$$\begin{aligned} & |\psi(z_0, f_c(z_0)) - \phi(z_0, f_c(z_0))| \\ &= \left| \left[ \sum_{i=1}^m \varphi_{\alpha_i}(z_0) \cdot g_{\epsilon_{\alpha_i}}(z_0, f_c(z_0)) \right] - \phi(z_0, f_c(z_0)) \right| \\ &\leq \sum_{i=1}^m \varphi_{\alpha_i}(z_0) \cdot |g_{\epsilon_{\alpha_i}}(z_0, f_c(z_0)) - \phi(z_0, f_c(z_0))| \\ &\leq \max\{\epsilon_{\alpha_i}\}. \end{aligned}$$

Further

$$\begin{aligned} & \left| \frac{\partial}{\partial x_1} [\psi(z_0, f_c(z_0)) - \phi(z_0, f_c(z_0))] \right| \\ &= \left| \frac{\partial}{\partial x_1} \left[ \left[ \sum_{i=1}^m \varphi_{\alpha_i}(z) \cdot g_{\epsilon_{\alpha_i}}(z, f_c(z_0)) \right] - \phi(z_0, f_c(z_0)) \right] \right| \\ &= \left| \sum_{i=1}^m \frac{\partial}{\partial x_1} [\varphi_{\alpha_i}(z_0) \cdot (g_{\epsilon_{\alpha_i}}(z_0, f_c(z_0)) - \phi(z_0, f_c(z_0)))] \right| \\ &= \left| \sum_{i=1}^m \frac{\partial}{\partial x_1} [\varphi_{\alpha_i}(z_0)] \cdot (g_{\epsilon_{\alpha_i}}(z_0, f_c(z_0)) - \phi(z_0, f_c(z_0))) \right. \\ &\quad \left. + \sum_{i=1}^m \varphi_{\alpha_i}(z_0) \cdot \frac{\partial}{\partial x_1} [g_{\epsilon_{\alpha_i}}(z_0, f_c(z_0)) - \phi(z_0, f_c(z_0))] \right| \\ &\leq m \cdot \max\{C_{\alpha_i}\} \cdot \max\{\epsilon_{\alpha_i}\} + \max\{\epsilon_{\alpha_i}\}. \end{aligned}$$

Similarly we get that

$$\left| \frac{\partial}{\partial x_2} [\psi(z_0, f_c(z_0)) - \phi(z, f_c(z_0))] \right| \leq m \cdot \max\{C_{\alpha_i}\} \cdot \max\{\epsilon_{\alpha_i}\} + \max\{\epsilon_{\alpha_i}\}.$$

It is clear that we may choose  $\epsilon_{\alpha_i}$  for  $i = 1, \dots, m$  to get the desired estimate for all points  $z_0 \in U_\alpha$  for this particular  $\alpha$ . Running through all  $\alpha$  we find that any particular  $\alpha_i$  will only come under consideration a finite number of times. Hence we may choose the sequence  $\{\epsilon_\alpha\}$ . □

We proceed to prove the proposition.

Fix  $\delta_0$  to get the estimate (1) (in the beginning of §3) for all  $|c - c'| < \delta_0$  with  $|c|, |c'| \leq 2R$ . For any  $\delta$  with  $0 < \delta < \delta_0$  we let  $c^\delta(j, k) = (j + k \cdot i) \cdot \delta$  for  $j, k \in \mathbb{Z}$ . Let  $\chi : [0, 1] \rightarrow \mathbb{R}$  be a smooth function such that  $\chi(t) = 0$  for  $0 \leq t \leq 1/4$  and  $\chi(t) = 1$  for  $3/4 \leq t \leq 1$ . Let  $C$  be a constant such that  $|\chi'(t)| \leq C$  for all  $t \in [0, 1]$ .

We first define a function  $h_\delta$  on the surfaces  $\Gamma_{c^\delta(j,k)}$  simply by  $h_\delta|_{\Gamma_{c^\delta(j,k)}} \equiv j\delta$ . We want to interpolate this function between the surfaces.

For a fixed  $z$  consider the sets of points

$$Q_{c^\delta(j,k)}(z) := \{f_{c^\delta(j,k)}(z), f_{c^\delta(j+1,k)}(z), f_{c^\delta(j,k+1)}(z), f_{c^\delta(j+1,k+1)}(z)\}.$$

We first show that these sets move nicely with  $z$  for small enough  $|z|$  and independent of  $\delta$ . In particular we want to know that we may define quadrilateral regions  $R_{\delta,j,k}(z)$ , with straight edges and corners  $Q_{c^\delta(j,k)}(z)$ , and that these sets have disjoint interior.

We make the change of coordinates in the  $w$  variable, by setting

$$\tilde{w}(z, w) = \tilde{w}_{jk}(z, w) = \frac{w - f_{c^\delta(j,k)}(z)}{f_{c^\delta(j+1,k)}(z) - f_{c^\delta(j,k)}(z)}.$$

We get

$$\begin{aligned} \tilde{w}(z, f_{c^\delta(j,k)}(z)) &\equiv 0, \\ \tilde{w}(z, f_{c^\delta(j+1,k)}(z)) &\equiv 1. \end{aligned}$$

From the discussion on holomorphic motions in §2 we get the following.

LEMMA 3. Fix  $N$ . Then there exists a real number  $t_0 > 0$  independent of  $\delta$  such that if  $|l|, |m| < N$  then  $|\tilde{w}_{jk}(z, f_{c^\delta(j+l,k+m)}(z)) - \tilde{w}_{jk}(z, f_{c^\delta(j+l,k+m)}(0))| < 1/10$  for all  $|z| < t_0$  and any  $j, k$ .

From now on we assume that  $|z| \leq t_0$ .

LEMMA 4. The quadrilaterals have disjoint interiors.

*Proof.* Pick  $(j, k)$ . We use the linear change of coordinates in the  $w$  direction for fixed  $z$ :

$$\tilde{w}_{jk}(z, w) = \frac{w - f_{c^\delta(j,k)}(z)}{f_{c^\delta(j+1,k)}(z) - f_{c^\delta(j,k)}(z)}.$$

This sends  $f_{c^\delta(j+l,k+m)}(z)$  close to  $(j+l, k+m)$  on a small disc in the  $z$  direction for uniformly bounded  $(l, m)$ . Hence it is clear that the quadrilaterals are disjoint.  $\square$

Next we define preliminary functions  $h_{jk}^\delta$  on the respective quadrilaterals. First we define a function  $t_z(y_1, y_2)$  to be constant equal to 0 on the line between  $f_{c^\delta(j,k)}(z)$  and  $f_{c^\delta(j,k+1)}(z)$ , and constant equal to 1 on the line between  $f_{c^\delta(j+1,k)}(z)$  and  $f_{c^\delta(j+1,k+1)}(z)$ . We extend  $t_z$  continuously to be affine on the two other edges, and then we extend  $t_z$  to be constant equal to  $v$  on the line between  $f_{c^\delta(j,k)}(z) + v \cdot (f_{c^\delta(j+1,k)}(z) - f_{c^\delta(j,k)}(z))$  and  $f_{c^\delta(j,k+1)}(z) + v \cdot (f_{c^\delta(j+1,k+1)}(z) - f_{c^\delta(j,k+1)}(z))$ . Finally we define  $h_{jk}^\delta$  by

$$h_{jk}^\delta(z, y_1, y_2) = j\delta + \delta \cdot (\chi \circ t_z)(y_1, y_2).$$

The  $h_{jk}^\delta$  patch up smoothly along the vertical sides of the quadrilaterals where the functions are constant. To be able to patch them together in the ‘horizontal’ directions we first extend each  $h_{jk}^\delta$  across the ‘horizontal’ edges.

To do this we use the coordinates defined by  $\tilde{w}$ . Consider the normalization

$$\tilde{w}_{jk}(z, w) = \frac{w - f_{c^\delta(j,k)}(z)}{f_{c^\delta(j+1,k)}(z) - f_{c^\delta(j,k)}(z)}.$$

Let  $\tilde{h}_{jk}^\delta$  be defined by  $\tilde{h}_{jk}^\delta \circ \tilde{w} = h_{jk}^\delta$ . We want to glue together the two functions on the quadrilaterals sharing (in the new coordinates) the line segment  $\gamma$  between  $(0, 0)$  and  $(1, 0)$ , i.e. the function  $\tilde{h}_{jk}^\delta$  defined above  $\gamma$  and the function  $\tilde{h}_{j(k-1)}^\delta$  below  $\gamma$ .

We start by extending the function  $\tilde{h}_{jk}^\delta$ . Note first that by Lemma 3 the quadrilaterals  $R_{\delta,j,k}$  and  $R_{\delta,j,k-1}$  in the new coordinates – henceforth denoted  $\tilde{R}_{\delta,j,k}$  and

$\tilde{R}_{\delta,j,k-1}$  – have corners within  $(1/10)$ -distance from the points  $(l, m)$  for  $l, m \in \{0, 1, -1\}$ . Note also that if we define a function  $\tilde{t}_z(\tilde{y}_1, \tilde{y}_2)$  ( $\tilde{w} = \tilde{y}_1 + i\tilde{y}_2$ ) along lines in the quadrilateral  $\tilde{R}_{\delta,j,k}(z)$  in the new coordinates as we did when we defined  $t_z(y_1, y_2)$  above, then  $h_{jk}^\delta = (j\delta + \delta(\chi \circ \tilde{t})) \circ \tilde{w}$ . Because of the placing of the corners we see that there exists a constant  $K$  independent of  $\delta, j, k$  such that  $\|\nabla_{\tilde{w}}(j\delta + \delta(\chi \circ \tilde{t}))\| \leq K\delta$ .

Continue the lines in  $\tilde{R}_{\delta,j,k}$  that pass through the interval  $[(1/8), 1 - (1/8)]$  and extend  $\tilde{h}_{jk}^\delta$  to be constant on these lines. By the placing of the corners there is a constant  $\mu$  – independent of  $\delta$  and  $j, k$  – such that these lines can be extended to the line between  $(0, -\mu)$  and  $(1, -\mu)$ . Let  $\tilde{P}_{\delta,j,k}$  denote the extended set  $\tilde{R}_{\delta,j,k} \cup (\tilde{R}_{\delta,j,k-1} \cap \{y_2 \geq -\mu\})$ ; we see that  $\tilde{h}_{jk}^\delta$  extends to be constant on the part of  $\tilde{P}_{\delta,j,k}$  where it is not already defined. Extend  $\tilde{h}_{j(k-1)}^\delta$  similarly in the other direction.

To glue the functions together we choose a smooth function  $\varphi(z, \tilde{y}_1, \tilde{y}_2) = \varphi(\tilde{y}_2)$  such that  $\varphi(\tilde{y}_2) = 1$  if  $y_2 \geq \mu$  and such that  $\varphi(\tilde{y}_2) = 0$  if  $y_2 \leq -\mu$ . We define our final function

$$h_\delta(z, w) := (\varphi \circ \tilde{w}_{jk})(z, w) \cdot h_{jk}^\delta(z, w) + (1 - \varphi \circ \tilde{w}_{jk})(z, w) \cdot h_{j(k-1)}^\delta(z, w). \tag{2}$$

Fix a constant  $M$  such that  $\|\partial\varphi/\partial\tilde{y}_2\| = M$ .

LEMMA 5. *There are constants  $N_1$  and  $N_2$  such that for each  $j, k, \delta$  we have  $h_{jk}^\delta(z, w) = j\delta$  if  $|w - f_{c^\delta(j,k)}(z)| \leq N_1|f_{c^\delta(j+1,k)}(z) - f_{c^\delta(j,k)}(z)|$ . Moreover there is a smooth function  $\tilde{g}_{jk}^\delta(z, \tilde{y}_1, \tilde{y}_2)$  such that  $h_{jk}^\delta = \tilde{g}_{jk}^\delta \circ \tilde{w}$  and  $\|\nabla_{\tilde{w}}\tilde{g}_{jk}^\delta\| \leq N_2\delta$ .*

*Proof.* The existence of the constant  $N_1$  can be seen by our description of the function in local coordinates where we used Lemma 3. To see the rest let us give the function  $\tilde{g}_{jk}^\delta$  explicitly.

Fix  $z$ . Let  $(a_1, a_2)$  denote the corner of  $\tilde{R}_{j,k}^\delta$  that is close to  $(0, 1)$ , and define a map  $A_z(\tilde{y}_1, \tilde{y}_2) := (\tilde{y}_1 - \tilde{y}_2(a_1/a_2), \tilde{y}_2(1/a_2))$ . Then  $A_z$  changes smoothly with  $z$  and  $\|A_z\| < 2$  for all the possibilities of  $(a_1, a_2)$  we are considering.

Next we define a function  $\hat{t}$  on the quadrilateral  $A_z(\tilde{R}_{j,k}^\delta)$  along lines as above. Let  $(b_1, b_2)$  denote the corner close to  $(1, 1)$  and fix  $\hat{y} = (\hat{y}_1, \hat{y}_2)$ . We have that the two vertical sides of  $A_z(\tilde{R}_{j,k}^\delta)$  meet at the point  $(0, -L)$  where  $L = b_2/(b_1 - 1)$ . Calculating the slope of the line from the point  $\hat{y}$  to the point  $(\hat{t}(\hat{y}), 0)$ , we get that  $\hat{y}_1/(L + \hat{y}_2) = \hat{t}(\hat{y})/L$ , which gives us

$$\hat{t}(\hat{y}) = \frac{\hat{y}_1 \cdot L}{L + \hat{y}_2} = \frac{\hat{y}_1 \cdot b_2}{b_2 + \hat{y}_2(b_1 - 1)}.$$

We have that  $\hat{t}$  varies smoothly with  $(b_1, b_2)$  and we see that  $\hat{t}$  has bounded derivatives for the cases of  $(b_1, b_2)$  we are considering. Define  $\tilde{g}_{jk}^\delta$  by

$$\tilde{g}_{jk}^\delta = j\delta + \delta(\chi \circ \hat{t} \circ A_z),$$

and the function  $h_{jk}^\delta$  is given by  $h_{jk}^\delta = \tilde{g}_{jk}^\delta \circ \tilde{w}$ . □

LEMMA 6.  $h_\delta \rightarrow g$  in sup norm on  $\Delta_{t_0} \times R\Delta$ .

*Proof.* It is clear that  $h_\delta(0, \cdot) \rightarrow g(0, \cdot)$  uniformly. The claim then follows from Lemma 8 below. □



LEMMA 7. *If  $t_0$  and  $\delta$  are small enough, then  $|f_{c^\delta(j,k)}(z) - f_{c^\delta(j+1,k)}(z)| \geq \delta^2$  for all  $z$  with  $|z| \leq t_0$  and all  $j, k$  such that  $|c^\delta(j, k)| \leq 2R$ .*

*Proof.* This follows from the Hölder continuity of the holomorphic motion. □

LEMMA 8. *Let  $c \in R\bar{\Delta}$ . The function  $h_\delta(z, f_c(z))$  is small in  $\mathcal{C}^1$ -norm along the graph  $\Gamma_c$ .*

*Proof.* We need to estimate the derivatives of the function  $h_\delta(z, f_c(z))$  at an arbitrary point  $(z_0, f_c(z_0))$ , and this point is contained in some extended quadrilateral  $P_{\delta,j,k}$ . We estimate  $\partial/\partial x = \partial/\partial x_1$  – the case of  $\partial/\partial x_2$  is similar. Since we are working on lines we use the notation  $(x, y_1, y_2)$  for coordinates.

If the point is close to the vertical edges, then the function  $h_\delta$  is locally constant, so we are done. We can assume that also  $(z_0, f_c(z_0)) \in P_{\delta,j,k} \setminus P_{\delta,j,k+1}$ . We divide the proof into two cases. Assume first that  $(z_0, f_c(z_0))$  is not in  $P_{\delta,j,k-1}$ . Then the function  $h_\delta$  is simply equal to the function  $h_{jk}^\delta$  (see (2)).

We have that

$$\begin{aligned} \frac{\partial}{\partial x}(h_{jk}^\delta(x, f(x))) &= \left( \frac{\partial h_{jk}^\delta}{\partial x}, \frac{\partial h_{jk}^\delta}{\partial y_1}, \frac{\partial h_{jk}^\delta}{\partial y_2} \right) (x, f(x)) \cdot \left( 1, \frac{\partial f_1}{\partial x}, \frac{\partial f_2}{\partial x} \right) (x) \\ &= \frac{\partial h_{jk}^\delta}{\partial x}(x, f(x)) + \left( \frac{\partial h_{jk}^\delta}{\partial y_1}, \frac{\partial h_{jk}^\delta}{\partial y_2} \right) (x, f(x)) \cdot \left( \frac{\partial f_1}{\partial x}, \frac{\partial f_2}{\partial x} \right) (x). \end{aligned} \tag{3}$$

For fixed  $s, v$  we may define a curve  $(x, g(x))$ :

$$\begin{aligned} g(x) &= (1 - s) [(1 - v)f_{c^\delta(j,k)}(x) + vf_{c^\delta(j+1,k)}(x)] \\ &\quad + s[(1 - v)f_{c^\delta(j,k+1)}(x) + vf_{c^\delta(j+1,k+1)}(x)]. \end{aligned}$$

Then  $h_{jk}^\delta(x, g(x)) \equiv j\delta + \chi(v)\delta$ . Choose  $s$  and  $v$  so that  $(x_0, g(x_0)) = (x_0, f_c(x_0))$ . We get that

$$\begin{aligned} 0 &= \frac{\partial}{\partial x}(h_{jk}^\delta(x, g(x))) \\ &= \frac{\partial h_{jk}^\delta}{\partial x}(x, g(x)) + \left( \frac{\partial h_{jk}^\delta}{\partial y_1}, \frac{\partial h_{jk}^\delta}{\partial y_2} \right) (x, g(x)) \cdot \left( \frac{\partial g_1}{\partial x}, \frac{\partial g_2}{\partial x} \right) (x), \end{aligned} \tag{4}$$

and so subtracting (4) from (3) we get

$$\frac{\partial}{\partial x}(h_{jk}^\delta(x_0, f(x_0))) = \left( \frac{\partial h_{jk}^\delta}{\partial y_1}, \frac{\partial h_{jk}^\delta}{\partial y_2} \right) (x_0, g(x_0)) \cdot \left( \frac{\partial f_1}{\partial x} - \frac{\partial g_1}{\partial x}, \frac{\partial f_2}{\partial x} - \frac{\partial g_2}{\partial x} \right) (x_0).$$

Using Lemma 3 we see that  $\|f_c(x_0) - f_{c^\delta(j+l,k+m)}(x_0)\| \leq 2\|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)\|$  for  $l, m \in \{0, 1\}$ , and so

$$\begin{aligned} & \left\| \frac{\partial}{\partial x} (f_c - f_{c^\delta(j+l,k+m)}) (x_0) \right\| \\ & \leq 4 \| (f_c - f_{c^\delta(j+l,k+m)}) (x_0) \| \log \frac{1}{\| (f_c - f_{c^\delta(j+l,k+m)}) (x_0) \|} \\ & \leq 8 \| (f_{c^\delta(j+1,k)} - f_{c^\delta(j,k)}) (x_0) \| \log \frac{1}{2 \| (f_{c^\delta(j+1,k)} - f_{c^\delta(j,k)}) (x_0) \|}. \end{aligned}$$

It follows that

$$\begin{aligned} \left\| \frac{\partial}{\partial x} (h_\delta(x, f(x))) \right\| & \leq 8 \cdot \left\| \left( \frac{\partial h_\delta}{\partial y_1}, \frac{\partial h_\delta}{\partial y_2} \right) \right\| \cdot \| (f_{c^\delta(j+1,k)} - f_{c^\delta(j,k)}) (x_0) \| \\ & \quad \times \log \frac{1}{2 \| (f_{c^\delta(j+1,k)} - f_{c^\delta(j,k)}) (x_0) \|}. \end{aligned}$$

We proceed to estimate  $\|(\partial h_\delta/\partial y_1, \partial h_\delta/\partial y_2)\|$ . We change coordinates according to Lemma 5 and write  $h_\delta$  as a composition  $\tilde{g}_\delta \circ \tilde{w}(y)$ . We get  $\|D_w \tilde{w}\| = 1/(\|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)\|)$ , and we have that  $\|\nabla_{\tilde{w}} \tilde{g}_\delta\| \leq N_2 \delta$ . This shows that

$$\left\| \left( \frac{\partial h_\delta}{\partial y_1}, \frac{\partial h_\delta}{\partial y_2} \right) \right\| \leq N_2 \delta \frac{1}{\|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)\|}.$$

This gives

$$\left\| \frac{\partial}{\partial x} (h_\delta(x, f(x))) \right\| \leq 8N_2 \delta \log \frac{1}{\|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)\|}.$$

We have by Lemma 7 that  $\|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)\| \geq \delta^2$ , and so

$$\left\| \frac{\partial}{\partial x} (h_\delta(x_0, f(x_0))) \right\| \leq 8N_2 \delta \log \frac{1}{2\delta^2} \rightarrow 0 \quad \text{as } \delta \rightarrow 0.$$

The other case we have to consider is when  $(z_0, f_c(z_0))$  is contained in an overlap where we glued our functions together. In that case we may assume that  $(z_0, f_c(z_0))$  is also contained in  $P_{j(k-1)}^\delta$  (see (2)).

Let  $\vec{v}$  denote the vector  $\vec{v} = \partial/\partial x(x_0, f_c(x_0))$ . We have that

$$\begin{aligned} \nabla h_\delta(x_0, f_c(x_0)) \cdot \vec{v} & = \nabla[\varphi \circ \tilde{w} \cdot h_{jk}^\delta] (x_0, f_c(x_0)) \cdot \vec{v} \\ & \quad + \nabla[(1 - \varphi) \circ \tilde{w} \cdot h_{j(k-1)}^\delta] (x_0, f_c(x_0)) \cdot \vec{v} \\ & = h_{jk}^\delta(x_0, f_c(x_0)) \cdot \nabla[\varphi \circ \tilde{w}] (x, f_c(x_0)) \cdot \vec{v} \\ & \quad + (\varphi \circ \tilde{w})(x_0, f_c(x_0)) \cdot \nabla[h_{jk}^\delta] (x_0, f_c(x_0)) \cdot \vec{v} \\ & \quad + h_{j(k-1)}^\delta(x_0, f_c(x_0)) \cdot \nabla[(1 - \varphi) \circ \tilde{w}] (x, f_c(x_0)) \cdot \vec{v} \\ & \quad + ((1 - \varphi) \circ \tilde{w})(x_0, f_c(x_0)) \cdot \nabla[h_{j(k-1)}^\delta] (x_0, f_c(x_0)) \cdot \vec{v}. \end{aligned}$$

By the above calculations we need not worry about the second and fourth term in this sum so we have to check that

$$(h_{jk}^\delta(x_0, f_c(x_0)) - h_{j(k-1)}^\delta(x_0, f_c(x_0))) \cdot \nabla[\varphi \circ \tilde{w}] (x_0, f_c(x_0)) \cdot \vec{v} \rightarrow 0$$

as  $\delta \rightarrow 0$ .

First of all we have that  $|h_{jk}^\delta(x_0, f_c(x_0)) - h_{j(k-1)}^\delta(x_0, f_c(x_0))| \leq 2\delta$ . Further,  $|\nabla[\varphi \circ \tilde{w}](x_0, f_c(x_0)) \cdot \vec{v}| \leq M \cdot \|D[\tilde{w}](x_0, f_c(x_0))(\vec{v})\|$ .

Now

$$D[\tilde{w}](x_0, f_c(x_0))(\vec{v}) = \frac{\partial}{\partial x} \left[ \left( x, \frac{f_c(x) - f_{c^\delta(j,k)}(x)}{f_{c^\delta(j+1,k)}(x) - f_{c^\delta(j,k)}(x)} \right) \right] (x_0).$$

Ignoring the constant term (it gets killed by  $\delta$ ), we get that

$$\begin{aligned} \|D[\tilde{w}](x_0, f_c(x_0))(\vec{v})\| &\leq \frac{|f'_c(x_0) - f'_{c^\delta(j,k)}(x_0)|}{|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)|} \\ &\quad + \frac{|f_c(x_0) - f_{c^\delta(j,k)}(x_0)| \cdot |f'_{c^\delta(j+1,k)}(x_0) - f'_{c^\delta(j,k)}(x_0)|}{|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)|^2} \\ &\leq \frac{|f_c(x_0) - f_{c^\delta(j,k)}(x_0)|}{|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)|} \log \frac{1}{|f_c(x_0) - f_{c^\delta(j,k)}(x_0)|} \\ &\quad + \frac{|f_c(x_0) - f_{c^\delta(j,k)}(x_0)| \cdot |f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)|}{|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)|^2} \\ &\quad \times \log \frac{1}{|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)|}. \end{aligned}$$

By Lemma 3,  $|f_c(x_0) - f_{c^\delta(j,k)}(x_0)|/|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)| \leq 2$ , and so

$$\begin{aligned} \|D[\tilde{w}](x_0, f_c(x_0))(\vec{v})\| &\leq 2 \cdot \log \frac{1}{|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)|} \\ &\quad + 2 \log \frac{1}{|f_c(x_0) - f_{c^\delta(j,k)}(x_0)|}. \end{aligned}$$

By Lemma 5, our function is constant unless  $|f_c(x_0) - f_{c^\delta(j,k)}(x_0)| \geq N_1 |f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)| \geq N_1 \delta^2$  (by Lemma 7), and so we may assume that

$$\|D[\tilde{w}](x_0, f_c(x_0))(\vec{v})\| \leq 2 \log \frac{1}{\delta^2} + 2 \log \frac{1}{N_1 \delta^2}.$$

All in all:

$$\begin{aligned} &|(h_{jk}^\delta(x_0, f_c(x_0)) - h_{j(k-1)}^\delta(x_0, f_c(x_0))) \cdot \nabla[\varphi \circ \tilde{w}](x_0, f_c(x_0)) \cdot \vec{v}| \\ &\leq 4M\delta \left( \log \frac{1}{\delta^2} + \log \frac{1}{N_1 \delta^2} \right) \rightarrow 0 \quad \text{as } \delta \rightarrow 0. \quad \square \end{aligned}$$

#### 4. Proof of the main theorem

We are ready to prove the main theorem. As pointed out in §2, by the theorem of Słodkowski [9, 11], we can assume that  $\mathcal{L}$  is a lamination of  $\Delta \times \mathbb{C}$  as in the previous section.

*Proof of the main theorem.* Suppose that  $T$  is a positive closed  $(1, 1)$ -current on  $\Delta^2(0, 1)$ , supported on the laminated set  $K$  described in the introduction. We assume that  $T$  is subordinate to the lamination  $\mathcal{L}$  of  $K$ . Hence there is a positive measure  $\mu$  such that

$T = \int [V_\alpha] d\mu(\alpha)$ . Suppose that  $\lambda = dw - f'_\alpha(z) dz$ . We want to show that  $\lambda \wedge T = 0$ . Let  $\phi$  be any smooth  $(1, 0)$  test form. We need to show that  $\langle \lambda \wedge T, \phi \rangle = 0$ . This follows since

$$\begin{aligned} \langle \lambda \wedge T, \phi \rangle &= \int (\lambda \wedge T) \wedge \phi \\ &= \int T \wedge (\lambda \wedge \phi) \\ &= \int_\alpha \left( \int_{V_\alpha} \lambda \wedge \phi \right) d\mu(\alpha) \\ &= \int_\alpha 0 = 0. \end{aligned}$$

Assume next that  $T$  is directed by  $\mathcal{L}$ . Since  $\mathcal{L}$  is a lamination of  $\Delta \times \mathbb{C}$  we may invoke the approximation result from the previous section. With the approximation result at hand the implication follows from Sullivan’s proof of the smooth case [10]. We include the proof for the benefit of the reader.

Step 1 is to show that there exists a family of probability measures  $\sigma_\alpha$  such that  $\sigma_\alpha$  is supported on  $\Gamma_\alpha$ , and a measure  $\mu'$  on the  $\alpha$ -plane such that for all test forms  $\omega$ ,

$$T(\omega) = \int \left( \int_{\Gamma_\alpha} \omega d\sigma_\alpha \right) d\mu'.$$

Let  $\omega$  be a  $(1, 1)$  test form and let  $\lambda(z, w) = dw - f'_\alpha(z) dz$  for  $w = f_\alpha(z)$ . Let  $\vec{v}_1(z, w) = (1, f'_\alpha(z))$  and let  $\vec{v}_2(z, w) = (i, i \cdot f'_\alpha(z))$  for  $w = f_\alpha(z)$ , and define the 2-tangent field  $v(z, w) = (\vec{v}_1(z, w), \vec{v}_2(z, w))$ .

Switching basis,

$$\omega = \psi_1 dz \wedge d\bar{z} + \psi_2 dz \wedge \bar{\lambda} + \psi_3 d\bar{z} \wedge \lambda + \psi_4 \lambda \wedge \bar{\lambda},$$

for some functions  $\psi_i$ , and by assumption,  $T(\omega) = T(\psi_1 dz \wedge d\bar{z})$ . The function  $\psi_1$  is given by  $\psi_1 = (1/2i)\omega(v)$ , and so

$$T(\omega) = T\left(\frac{1}{2i}\omega(v) dz \wedge d\bar{z}\right).$$

On the other hand we may use  $T$  to define a linear functional  $L$  on  $\mathcal{C}_0(\Delta \times \mathbb{C})$  by  $L(\psi) = T(\psi dz \wedge d\bar{z})$ , and so by the Riesz representation theorem there is a measure  $\nu$  such that

$$L(\psi) = \int \psi d\nu.$$

This means that

$$T(\omega) = \int \frac{1}{2i}\omega(v) d\nu.$$

Now the measure  $\nu$  disintegrates [6]: there exists a family of probability measures  $\sigma_\alpha$  such that  $\sigma_\alpha$  is supported on  $\Gamma_\alpha$ , and a measure  $\mu'$  on the  $\alpha$ -plane such that for all  $\psi \in \mathcal{C}_0(\Delta \times \mathbb{C})$ ,

$$\int \psi d\nu = \int \left( \int_{\Gamma_\alpha} \psi d\sigma_\alpha \right) d\mu'.$$

We define currents  $T_\alpha$  by  $T_\alpha(\omega) = \int_{\Gamma_\alpha} (1/2i)\omega(v) d\sigma_\alpha$ , and we get that

$$T(\omega) = \int T_\alpha(\omega) d\mu'.$$

The next step is to show that  $T_\alpha$  is closed for  $\mu'$ -almost all  $\alpha$ . Let  $\{\omega_j\}$  be a dense set of  $C^1$ -smooth  $(0, 1)$  test forms and fix a  $j \in \mathbb{N}$ . Let  $g$  be a continuous function in the  $\alpha$ -variable and extend  $g$  constantly along leaves. We want to show that

$$\int g \cdot T_\alpha(\partial\omega) d\mu' = 0,$$

because this would imply that  $\partial T_\alpha = 0$  for  $\mu'$ -almost all  $\alpha$  (since  $g$  is arbitrary).

By Theorem 1 there exists a sequence  $g_i$  of smooth functions such that  $g_i \rightarrow g$  uniformly and in  $C^1$ -norm on leaves. Since  $T$  is closed,

$$0 = \int T_\alpha(\partial(g\omega_j)) d\mu' = \int T_\alpha(\partial g_i \wedge \omega_j) d\mu' + \int g_i \cdot T_\alpha(\partial\omega_j) d\mu'.$$

Since  $T_\alpha(\partial g_i \wedge \omega) \rightarrow 0$  we get that

$$\int g \cdot T_\alpha(\partial\omega_j) d\mu' = \lim_{i \rightarrow \infty} \int g_i \cdot T_\alpha(\partial\omega_j) d\mu' = 0.$$

Running through all  $\omega_j$  we see that  $T_\alpha$  is closed for  $\mu'$ -almost all  $\alpha$ . The only possibility then is that the measures  $\sigma_\alpha$  are constant multiples of  $dz \wedge d\bar{z}$ , i.e.  $\sigma_\alpha = \varphi(\alpha) dz \wedge d\bar{z}$  where  $\varphi$  is a measurable function [7]. Define  $\mu := \varphi \cdot \mu'$ .

### 5. Two counterexamples

In [5] the authors proved versions of the main theorem for real laminations in  $\mathbb{R}^2$  and  $\mathbb{R}^3$ . In those results we added an extra slope condition on the laminations which is analogous to the estimate in Corollary 1. We give here a simple example of a lamination of curves in  $\mathbb{R}^2$  where the slope condition is not satisfied. Also, the conclusion of the main theorem fails. The analogue of Theorem 1, i.e. approximation of partially-smooth functions, fails as well.

For each  $t \in \mathbb{R}$ , we let  $\gamma_t$  be the curve  $y = f_t(x) = (x - t)^3$  in  $\mathbb{R}^2$ . Clearly this gives a continuous lamination of  $\mathbb{R}^2$  by curves. The curves are all tangent to the  $x$ -axis. This implies that the current of integration of the  $x$ -axis is annihilated by the 1-form  $\lambda$  defined by  $dy - f'_t(x) dx$  on  $\gamma_t$ . However, this current is not an integral of currents  $[\gamma_t]$ . We also observe that the function  $a(x, y)$  defined by  $a(x, f_t(x)) = t$  cannot be approximated by  $C^1$  functions, because any such approximation will have to have a small derivative along the  $x$ -axis.

We can also modify this example so that we have a Riemann surface lamination in  $\mathbb{C}^3$ . For  $t \in \mathbb{C}$ , let  $\gamma_t$  be the complex curve  $\gamma_t(s) = (z, w, \tau) = (s, (s - t)^2, (s - t)^3)$ . These curves laminate  $\mathbb{C}^3$ , and  $\gamma_t$  is tangent to the  $z$ -axis at  $(t, 0, 0)$ . Hence the  $z$ -axis is annihilated by any continuous 1-forms defining the lamination. Hence the current of integration of the  $z$ -axis is directed. But clearly it is not subordinate to the lamination. Again the function  $a(z, w, \tau)$  defined by  $a|_{\gamma_t} = t$  cannot be approximated by  $C^1$  functions.

*Acknowledgements.* The first author is supported by an NSF grant. The third author is supported by Schweizerische Nationalfonds grant 200021-116165/1.

## REFERENCES

- [1] B. Berndtsson and N. Sibony. The  $\bar{\partial}$ -equation on a positive closed current. *Invent. Math.* **147** (2002), 371–428.
- [2] E. Bedford, M. Lyubich and J. Smillie. Polynomial diffeomorphisms of  $\mathbb{C}^2$ , IV. *Invent. Math.* **112** (1993), 77–125.
- [3] R. Dujardin. Approximation des fonctions lisses sur certaines laminations. *Indiana Univ. Math. J.* **55** (2006), 579–592.
- [4] J. E. Fornæss and N. Sibony. Harmonic currents and finite energy of laminations. *Geom. Funct. Anal.* **15** (2005), 962–1003.
- [5] J. E. Fornæss, Y. Wang and E. F. Wold. Approximation of partially smooth functions. *Proceedings of Qikeng Lu Conference (June 2006)*. *Sci. China A: Math.* **51** (4) (2008), 553–561.
- [6] P. R. Halmos. *Measure Theory*. Van Nostrand, New York, 1950.
- [7] P. Lelong. *Fonctions Plurisubharmoniques et Formes Différentielles Positives*. Gordon and Breach, Paris, London, New York, 1968.
- [8] P. Mañé, P. Sad and D. Sullivan. On the dynamics of rational maps. *Ann. Sci. École Norm. Sup.* **16** (1983), 193–217.
- [9] Z. Stodkowski. Holomorphic motions and polynomial hulls. *Proc. Amer. Math. Soc.* **111** (1991), 347–355.
- [10] D. Sullivan. Cycles for the dynamical study of foliated manifolds and complex manifolds. *Invent. Math.* **36** (1976), 225–255.
- [11] K. Astala and G. J. Martin. Holomorphic motions. Papers on Analysis (A volume dedicated to Olli Martio on the occasion of his 60th birthday). *Report*, No. 83, University of Jyväskylä, Jyväskylä, 2001, pp. 27–40.