



^b
UNIVERSITÄT
BERN

Flexible varieties and automorphism groups

I. Arzhantsev, H. Flenner, S. Kaliman, F. Kutzschebauch,
and M. Zaidenberg

Research Report 2010-14

23.12.2010

Mathematics Institute
University of Bern
Sidlerstrasse 5
CH-3012 Bern
Switzerland

www.math.unibe.ch

FLEXIBLE VARIETIES AND AUTOMORPHISM GROUPS

I. ARZHANTSEV, H. FLENNER, S. KALIMAN, F. KUTZSCHEBAUCH, M. ZAIDENBERG

ABSTRACT. Given an affine algebraic variety X of dimension $n \geq 2$, we let $\text{SAut}(X)$ denote the special automorphism group of X i.e., the subgroup of the full automorphism group $\text{Aut}(X)$ generated by all one-parameter unipotent subgroups. We show that if $\text{SAut}(X)$ is transitive on the smooth locus X_{reg} then it is infinitely transitive on X_{reg} . In turn, the transitivity is equivalent to the flexibility of X . The latter means that for every smooth point $x \in X_{\text{reg}}$ the tangent space $T_x X$ is spanned by the velocity vectors at x of one-parameter unipotent subgroups of $\text{Aut}(X)$. We provide also different variations and applications.

CONTENTS

Introduction	2
1. Flexibility versus transitivity	4
1.1. Algebraically generated groups of automorphisms	4
1.2. Transversality	9
1.3. Special subgroups	11
2. Infinite transitivity	13
2.1. Main theorem	13
2.2. Proof of Theorem 2.5	14
2.3. Examples of non-separation of orbits	17
2.4. G_Y -orbits	19
3. Collective infinite transitivity	20
3.1. Collective transitivity on G -varieties	20
3.2. Infinite transitivity on matrix varieties	20
3.3. The case of symmetric and skew-symmetric matrices	21
4. Tangential flexibility, interpolation by automorphisms, and \mathbb{A}^1 -richness	22
4.1. Flexibility of the tangent bundle	22
4.2. Prescribed jets of automorphisms	25
4.3. \mathbb{A}^1 -richness	29
5. Some applications	30
5.1. Unirationality, flexibility, and triviality of the Makar-Limanov invariant	30
5.2. Flexible quasihomogeneous varieties	31
6. Appendix: Holomorphic flexibility	36
6.1. Oka-Grauert-Gromov Principle for flexible varieties	36
6.2. Volume density property	38
References	39

The research of the third author was supported by an NSA grant No. H98230-10-1-0185. The research of the fourth author was partially supported by Schweizerische Nationalfonds grant No. 200020-124668/1. This work was done during a stay of the second, third, and fifth authors at the Max Planck Institut für Mathematik at Bonn, a stay of the fourth author at the University of Miami, and a stay of the first and the second authors at the Institut Fourier, Grenoble. The authors thank these institutions for hospitality.

INTRODUCTION

All algebraic varieties and algebraic groups in this paper are supposed to be defined over an algebraically closed field \mathbb{k} of characteristic zero. Unless we explicitly mention the opposite, the varieties are supposed to be reduced and irreducible. For such a variety X we let $\text{SAut}(X)$ denote the subgroup of the automorphism group $\text{Aut}(X)$ generated by all one-parameter unipotent subgroups of $\text{Aut}(X)$ i.e., subgroups isomorphic to the additive group \mathbb{G}_a of the field. We call $\text{SAut}(X)$ the *special automorphism group* of X . In this paper we study transitivity properties of the action of this group on X .

As an example, let us consider the special automorphism group $\text{SAut}(\mathbb{A}^n)$ of the affine space $\mathbb{A}^n = \mathbb{A}_{\mathbb{k}}^n$. In the case $n = 1$ the only automorphisms in $\text{SAut}(\mathbb{A}^1)$ are the translations, so the group acts transitively but not 2-transitively. However, for $n \geq 2$ the situation is completely different. Here $\text{SAut}(\mathbb{A}^n)$ is no longer an algebraic group, e.g. for $n \geq 2$ it contains the shears

$$(x, y) \mapsto (x, y + P(x)),$$

where $P \in \mathbb{k}[x]$ is a polynomial, which form a family of infinite dimension. It is a well known and elementary fact that $\text{SAut}(\mathbb{A}^n)$, $n \geq 2$, acts even *infinitely transitively* on \mathbb{A}^n that is, m -transitively for any $m \geq 1$.

There is a number of further cases, where $\text{SAut}(X)$ acts infinitely transitively. Consider, for instance, an equivariant projective embedding $Y \hookrightarrow \mathbb{P}^n$ of a flag variety $Y = G/P$. Then the special automorphism group of the affine cone X over Y acts infinitely transitively on the smooth locus X_{reg} of X [1]. For any non-degenerate toric affine variety of dimension ≥ 2 a similar result is true [1]. If Y is an affine variety, on which $\text{SAut}(Y)$ acts infinitely transitively, then the same holds for the suspension $X = \{uv - f(y) = 0\} \subseteq \mathbb{A}^2 \times Y$ over Y , where $f \in \mathcal{O}(Y)$ is a non-constant function ([1]; see also [22] for the case where $Y = \mathbb{A}^n$).

One of the central results of this paper is the following general theorem (cf. Theorem 2.2). It confirms a conjecture formulated in [1, §4].

Theorem 0.1. *For an affine variety X of dimension ≥ 2 , the following conditions are equivalent.*

- (i) *The group $\text{SAut}(X)$ acts transitively on X_{reg} .*
- (ii) *The group $\text{SAut}(X)$ acts infinitely transitively on X_{reg} .*

Transitivity properties of the special automorphism group are closely related to the flexibility of a variety, which was studied in the algebraic context in [1]¹. We say that a point $x \in X$ is *flexible* if the tangent space $T_x X$ is spanned by the tangent vectors to the orbits $H.x$ of one-parameter unipotent subgroups $H \subseteq \text{Aut}(X)$. The variety X is called *flexible* if every smooth point $x \in X_{\text{reg}}$ is. Clearly, X is flexible if one point of X is and the group $\text{Aut}(X)$ acts transitively on X_{reg} . With this notation we can show in Corollary 1.21 that condition (i) and then also (ii) in Theorem 0.1 is equivalent to

- (iii) *X is a flexible variety.*

¹In the analytic context, several other flexibility properties are surveyed in [10].

For the examples mentioned above flexibility was verified in [1]. As a further example, the total space of a homogeneous vector bundle over a flexible affine variety is flexible (Corollary 4.5). Furthermore, every semisimple algebraic group is well known to be generated by its unipotent 1-parameter subgroups. This implies that G itself and any affine homogeneous space G/H are flexible. Hence in case $\dim G/H \geq 2$ their special automorphism group act infinitely transitively. More generally, if a semi-simple algebraic group acts with an open orbit on a smooth affine variety X then X is homogeneous itself and so is flexible (see Theorem 5.5).

In contrast to such results, according to A. Borel a real Lie group cannot act 4-transitively on a simply connected manifold (see Theorems 5 and 6 in [4]). The latter remains true (without the assumption of simple connectedness) for the actions of algebraic groups over algebraically closed fields [25]. This shows that the special automorphisms of an affine homogeneous space of dimension ≥ 2 form always a ‘large’ group.

Let us mention several applications. As an almost immediate consequence it follows that *in a flexible affine variety X any finite subset $Z \subseteq X_{\text{reg}}$ can be interpolated by a polynomial curve*, that is by a curve $C \cong \mathbb{A}^1$ in X_{reg} (see Corollary 4.18 below for a more general statement). Indeed, given a one-dimensional \mathbb{G}_a -orbit O in X_{reg} and a finite subset $Z' \subseteq O$ of the same cardinality as that of Z by infinite transitivity there is an automorphism $g \in \text{SAut}(X)$ which sends Z' to Z . Then $g(O) \cong \mathbb{A}^1$ is a \mathbb{G}_a -orbit passing through every point of Z .

This interpolation by \mathbb{A}^1 -curves is related to the property of \mathbb{A}^1 -richness [23]. A smooth variety X is called \mathbb{A}^1 -rich if, given any two disjoint closed subvarieties Y, Z of X with $\text{codim}_X Y \geq 2$ and $\dim Z = 0$, there exists a polynomial curve in $X \setminus Y$ passing through every point of Z . For instance, \mathbb{A}^n for $n \geq 2$ is \mathbb{A}^1 -rich by the Gromov-Winkelmann theorem, see [40]. In Corollary 4.18 we show more generally that *any smooth flexible affine variety is \mathbb{A}^1 -rich by means of \mathbb{G}_a -orbits*.

An interesting case of flexible varieties are the degeneracy loci of generic matrices, which are the varieties say $X_r \subseteq \mathbb{A}^{mn}$ consisting of $m \times n$ -matrices of rank $\leq r$. It is a standard fact of linear algebra that $\text{SAut}(X_r)$ acts transitively on $X_r \setminus X_{r-1}$ unless $r = m = n$. Indeed, one can transform each matrix to a normal form by a sequence of elementary transformations, which replace row i by row $i + t \cdot \text{row } j$ ($i \neq j, t \in \mathbb{k}$), and similarly for columns. Since these transformations constitute \mathbb{G}_a -actions, $\text{SAut}(X_r)$ acts transitively and hence infinitely transitively on $X_r \setminus X_{r-1}$ unless $\dim X_r \leq 1$ or $r = m = n$.

We can prove this infinite transitivity even simultaneously for matrices of different ranks, see Theorem 3.3. This shows that any finite collection of $m \times n$ matrices can be diagonalized simultaneously by means of elementary row- and column transformations depending polynomially on the matrix entries. Similar statements also hold for symmetric and skew-symmetric matrices, see Theorems 3.5 and 3.6. Such a collective infinite transitivity for conjugacy classes of matrices was established earlier by Z. Reichstein [35] using different methods.

The *Gizatullin surfaces* represent another interesting class of examples. These are normal affine surfaces which admit a completion by a chain of smooth rational curves. Due to Gizatullin’s Theorem [15] (see also [8]), a normal affine surface X different from

$\mathbb{A}^1 \times (\mathbb{A}^1 \setminus \{0\})$ is Gizatullin if and only if the special automorphism group $\text{SAut}(X)$ has an open orbit with a finite complement. It follows from Theorem 2.2 below that the group $\text{SAut}(X)$ acts infinitely transitively on this open orbit. It is unknown however (to the best of our knowledge) whether this orbit coincides with X_{reg} i.e., whether every Gizatullin surface is flexible. This is definitely not true in positive characteristic, where the special automorphism group $\text{SAut}(X)$ of a Gizatullin surface X can have fixed points that are regular points of X [7]. We refer the reader e.g., to [9] and the references therein for a study of one-parameter groups acting on Gizatullin surfaces.

We can also prove a version of infinite transitivity including infinitesimal near points. More precisely we will show (see Theorem 4.14 for a slightly more general version):

Theorem 0.2. *Let X be a flexible affine variety of dimension $n \geq 2$ equipped with an algebraic volume form² ω . Then for every $m \geq 0$ and every finite subset $Z \subseteq X_{\text{reg}}$ there exists an automorphism $g \in G$ with prescribed m -jets at the points $p \in Z$, provided these jets preserve ω and inject Z into X_{reg} .*

In the holomorphic context such results were shown in [5] and [19].

Let us give a short overview of the content of the various sections.

In Section 1 we deal with a more general class of groups of automorphisms, namely, with groups generated by a family of algebraic subgroups. We study the orbits of such groups, and give a generalization of Kleiman's Transversality Theorem in this context (see Theorem 1.15). It is remarkable that for such actions the Rosenlicht Theorem on the separation of generic orbits by rational invariants remains true (see Theorem 1.12). As a consequence we are able to confirm a conjecture of [27] concerning the field ML-invariant.

In Section 2 we deduce Theorem 0.1 (cf. the more general Theorems 2.2 and 2.5). The methods developed there will be applied in Section 3 to show infinite transitivity on several orbits simultaneously, see Theorem 3.1. In particular, we deduce the applications to matrix varieties mentioned before.

Section 4 contains the results on the interpolation of curves and automorphisms as described above. In Section 5 we apply our techniques to homogeneous spaces and their affine embeddings. Finally in the Appendix 6 we compare our results with similar facts in complex analytic geometry. We deduce, in particular, that the Oka-Grauert-Gromov Principle is available for smooth G -fibrations with flexible fibers, where G is an algebraically generated group of automorphisms (cf. Proposition 6.3 and Corollary 6.7).

1. FLEXIBILITY VERSUS TRANSITIVITY

1.1. Algebraically generated groups of automorphisms. Let X be an algebraic variety over \mathbb{k} .

Definition 1.1. A subgroup G of the automorphism group $\text{Aut}(X)$ is said to be *algebraically generated* if it is generated by a family \mathcal{G} of connected algebraic subgroups of $\text{Aut}(X)$. More precisely, every $H \in \mathcal{G}$ is a connected algebraic group over \mathbb{k} , not

²By this we mean a nowhere vanishing n -form defined on X_{reg} .

necessarily affine, contained in G , and $G = \langle H \mid H \in \mathcal{G} \rangle$ is generated by these subgroups as an abstract group.

Proposition 1.2. *If the subgroup $G \subseteq \text{Aut}(X)$ is algebraically generated then the following hold.*

- (a) *For every point $x \in X$ the orbit $G.x$ is locally closed.*
- (b) *For every $x \in X$ there are (not necessarily distinct) subgroups $H_1, \dots, H_s \in \mathcal{G}$ such that*

$$G.x = H_1.(H_2.\cdots.(H_s.x)\cdots).$$

Proof. Replacing X by the Zariski closure of the orbit $G.x$ we may assume that $X = \overline{G.x}$ i.e., the orbit of x is dense in X . Notice that for every finite sequence $\mathcal{H} = (H_1, \dots, H_s)$ in \mathcal{G} the subset $X_{\mathcal{H},x} = H_1.(H_2.\cdots.(H_s.x)\cdots) \subseteq X$ is constructible and irreducible, being the image of the irreducible variety $H_1 \times \dots \times H_s$ under the regular map

$$(1) \quad \Phi_{\mathcal{H},x} : H_1 \times \dots \times H_s \rightarrow X, \quad (h_1, \dots, h_s) \mapsto (h_1 \cdots h_s).x.$$

Observe that enlarging \mathcal{H} we enlarge $X_{\mathcal{H},x}$ too. By assumption the union of all such sets $X_{\mathcal{H},x}$ is dense in X , hence also the union of the closures $\bar{X}_{\mathcal{H},x}$ is. Since an increasing sequence of closed irreducible subsets becomes stationary, $X = \bar{X}_{\mathcal{H},x}$ for some \mathcal{H} . In particular, the interior $\overset{\circ}{X}_{\mathcal{H},x}$ is nonempty. The union $\Omega = \bigcup_{\mathcal{H}} \overset{\circ}{X}_{\mathcal{H},x}$ over all such sequences $\mathcal{H} \subseteq \mathcal{G}$ is G -invariant and thus equal to the whole orbit $G.x$, which shows

- (a). Since an increasing sequence of open subsets of X becomes stationary we have $\overset{\circ}{X}_{\mathcal{H},x} = \Omega$ for some sequence \mathcal{H} in \mathcal{G} and so (b) follows as well. \square

We can strengthen (b) as follows.

Proposition 1.3. *There are (not necessarily distinct) subgroups $H_1, \dots, H_s \in \mathcal{G}$ such that*

$$(2) \quad G.x = H_1.(H_2.\cdots.(H_s.x)\cdots) \quad \forall x \in X.$$

Proof. Let us introduce a partial order on the set of sequences in \mathcal{G} via

$$(H_1, \dots, H_m) \succcurlyeq (H'_1, \dots, H'_s) \iff \exists i_1 < \dots < i_s : (H'_1, \dots, H'_s) = (H_{i_1}, \dots, H_{i_s}).$$

Obviously any two sequences are dominated by a third one. Given a sequence $\mathcal{H} = (H_1, \dots, H_s)$ in \mathcal{G} we consider the map

$$(3) \quad \Phi_{\mathcal{H}} : H_1 \times \dots \times H_s \times X \longrightarrow X \times X, \quad (h_1, \dots, h_s, x) \mapsto (x, (h_1 \cdots h_s).x).$$

The image $Z_{\mathcal{H}} = \Phi_{\mathcal{H}}(H_1 \times \dots \times H_s \times X)$ is constructible and irreducible. In particular, the union of closures $Z = \bigcup_{\mathcal{H}} \bar{Z}_{\mathcal{H}}$ stabilizes in $X \times X$ ³ and so is closed.

Let $\overset{\circ}{Z}_{\mathcal{H}}$ be the interior of $Z_{\mathcal{H}}$ in Z . It follows as before that also $\overset{\circ}{Z}_{\mathcal{H}}$ becomes stationary and that the union $Z' = \bigcup_{\mathcal{H}} \overset{\circ}{Z}_{\mathcal{H}}$ is an open dense subset of Z .

Suppose that G acts on $X \times X$ via $g.(x, y) = (g.x, y)$. If $\mathcal{H} = (H_1, \dots, H_s)$ and $H \in \mathcal{G}$ then for any $(h_1, \dots, h_s) \in H_1 \times \dots \times H_s$ and $h \in H$ we have

$$h.\Phi_{\mathcal{H}}(h_1, \dots, h_s, x) = h.(x, (h_1 \cdots h_s).x) = \Phi_{(\mathcal{H}, H)}(h_1, \dots, h_s, h^{-1}, hx).$$

Hence $h.Z_{\mathcal{H}} \subseteq Z_{(\mathcal{H}, H)}$. It follows that Z and Z' are G -invariant.

³ I.e., it coincides with $\bar{Z}_{\mathcal{H}}$ for \mathcal{H} sufficiently large.

Consider now for \mathcal{H} sufficiently large the sets $Z_{\mathcal{H}}$, $Z' = \overset{\circ}{Z}_{\mathcal{H}}$, and $Z = \bar{Z}_{\mathcal{H}}$ as families over X via the first projection $p : (x, y) \mapsto x$. By [17, 9.5.3] there is an open dense subset V of X such that $Z'(x)$ is dense in $Z(x)$ for all $x \in V$, where for a subset $M \subseteq X \times X$ we denote by $M(x)$ the fiber of $p|M : M \rightarrow X$ over x . Since Z and Z' are invariant under the action of G and the projection p is equivariant, we may suppose that V is as well G -invariant.

In particular there is a sequence \mathcal{H}_0 such that $Z_{\mathcal{H}}(x) = (H_1 \cdots H_s).x$ is dense in $Z(x)$ for all $x \in V$ and all sequences $\mathcal{H} = (H_1, \dots, H_s)$ dominating \mathcal{H}_0 . It follows that $Z(x)$ is the orbit closure of $G.x$ and so $(H_1 \cdots H_s).x$ is dense in the orbit $G.x$ for all $x \in V$.

We claim that for every point $x \in V$

$$(H_s \cdots H_1 \cdot H_1 \cdots H_s).x = G.x.$$

Indeed, for any $y \in G.x$ the sets $(H_1 \cdots H_s).x$ and $(H_1 \cdots H_s).y$ are both dense in the orbit $G.x = G.y$. Hence they have a point, say, z in common. Thus

$$y \in (H_s \cdots H_1).z \subseteq (H_s \cdots H_1 \cdot H_1 \cdots H_s).x.$$

Replacing \mathcal{H} by the larger sequence $(H_s, \dots, H_1, H_1, \dots, H_s)$ it follows that

$$(H_1 \cdots H_s).x = G.x \text{ for all } x \in V \text{ simultaneously.}$$

The complement $Y = X \setminus V$ is closed, and all its irreducible components are of dimension $< \dim X$. Using induction on the dimension of X it follows that (2) holds for \mathcal{H} sufficiently large and all $x \in X$ simultaneously, concluding the proof. \square

Remark 1.4. We note that Propositions 1.2 and 1.3 remain true with the same proofs for varieties over algebraically closed fields of arbitrary characteristic.

Definition 1.5. A sequence $\mathcal{H} = (H_1, \dots, H_s)$ in \mathcal{G} satisfying condition (2) of 1.3 will be called *maximal*.

Remark 1.6. It is not true in general that the ‘orbit’ $G.Y$ of a Zariski closed subset $Y \subseteq X$ under an algebraic group action is locally closed. Nevertheless, applying the same kind of arguments as in the proofs of 1.2 and 1.3 above one can show that for any algebraically generated subgroup $G \subseteq \text{Aut}(X)$ and a constructible subset Y of X the orbit $G.Y$ is a constructible subset of X .

Proposition 1.7. *Assume that the generating family \mathcal{G} of connected algebraic subgroups is closed under conjugation in G , i.e., $gHg^{-1} \in \mathcal{G}$ for all $g \in G$ and $H \in \mathcal{G}$. Then there is a sequence $\mathcal{H} = (H_1, \dots, H_s)$ in \mathcal{G} such that for all $x \in X$ the tangent space $T_x(G.x)$ of the orbit $G.x$ is spanned by the tangent spaces*

$$T_x(H_1.x), \dots, T_x(H_s.x).$$

Proof. We claim that $T_x(G.x)$ is spanned by the tangent spaces $T_x(H.x)$, where $H \in \mathcal{G}$. Indeed, consider a maximal sequence $H_1, \dots, H_s \in \mathcal{G}$ such that the map $\Phi_{\mathcal{H},x} : H_1 \times \dots \times H_s \rightarrow G.x$ in (1) is surjective. Its generic rank is maximal and so for some point $y = (h_1 \cdots h_s).x \in G.x$ the tangent map

$$d\Phi_{\mathcal{H},x} : T_{(h_1, \dots, h_s)}(H_1 \times \dots \times H_s) \longrightarrow T_y(G.x)$$

is surjective. Multiplication by $g = (h_1 \cdots h_s)^{-1}$ yields an isomorphism $\mu_g : G.x \rightarrow G.x$ which sends y to x . Hence the composition $\mu_g \circ \Phi_{\mathcal{H},x}$ has a surjective tangent map

$$d(\mu_g \circ \Phi_{\mathcal{H},x}) : T_{(h_1, \dots, h_s)}(H_1 \times \dots \times H_s) \longrightarrow T_x(G.x).$$

The restriction of $\mu_g \circ \Phi_{\mathcal{H},x}$ to the factor

$$h_1 \times \dots \times h_{\sigma-1} \times H_\sigma \times h_{\sigma+1} \times \dots \times h_s$$

can be identified with the map

$$H'_\sigma = g_\sigma^{-1} H_\sigma g_\sigma \rightarrow G.x, \quad h' \mapsto h'.x,$$

where $g_\sigma = h_{\sigma+1} \cdots h_s \in G$. Taking the tangent maps provides the claim.

Consider further the map $\Phi_{\mathcal{H}} : H_1 \times \dots \times H_s \times X \rightarrow Z$ as in (3) associated with a maximal sequence \mathcal{H} , where $Z \subseteq X \times X$ is the closure of the image of $\Phi_{\mathcal{H}}$ as in the proof of 1.3. Choose an invariant open subset $V \subseteq X_{\text{reg}}$ such that the first projection $p : Z \rightarrow X$ is smooth over V . Note that the fiber of $Z_V = p^{-1}(V) \rightarrow V$ over x is just the orbit $G.x$. Let us consider the map of relative tangent bundles

$$d\Phi_{\mathcal{H}} : T(H_1 \times \dots \times H_s \times V/V) \rightarrow \Phi_{\mathcal{H}}^*(T(Z_V/V))$$

and its restriction to $(1, \dots, 1) \times V \cong V$,

$$d\Phi_{\mathcal{H}} : T_1 H_1 \times \dots \times T_1 H_s \times V \rightarrow \Phi_{\mathcal{H}}^*(T(Z_V/V))|V.$$

The set $U_{\mathcal{H}}$ of points in V where this map is surjective, is open. By the above claim, their union $\bigcup_{\mathcal{H}} U_{\mathcal{H}}$ coincides with V . Since an increasing union of open subsets stabilizes, we obtain that $V = U_{\mathcal{H}}$ for \mathcal{H} sufficiently large. Induction on the dimension of X as in the proof of Proposition 1.3 ends the proof. \square

Remark 1.8. It may happen for a family \mathcal{G} which is not closed under conjugation that for some point $x \in X$ the tangent spaces

$$T_x(H_1.x), \dots, T_x(H_s.x)$$

do not span $T_x(G.x)$, whatever is the sequence $\mathcal{H} = (H_1, \dots, H_s)$ in \mathcal{G} . For instance, the group $G = \text{SL}_2$ is generated by the family $\mathcal{G} = \{U^+, U^-\}$, where U^\pm are the subgroups of upper and lower triangular unipotent matrices. Letting SL_2 act on itself by left multiplication the tangent space $T_1 G$ of the orbit $G = G.1$ is \mathfrak{sl}_2 , while for any sequence $\mathcal{H} = (H_1, \dots, H_s)$ in \mathcal{G} the tangent spaces $T_1(H_1), \dots, T_1(H_s)$ are contained in the 2-dimensional subspace $T_1(U^+) + T_1(U^-)$.

Definition 1.9. Let $G \subseteq \text{Aut}(X)$ be algebraically generated by a family \mathcal{G} of connected algebraic subgroups, which is closed under conjugation. We say that a point $p \in X_{\text{reg}}$ is *G-flexible* if the tangent space $T_p X$ at p is generated by the subspaces $T_p(H.p)$, where $H \in \mathcal{G}$.

Corollary 1.10. *With G and \mathcal{G} as in Definition 1.9 the following hold.*

- (a) *A point $p \in X_{\text{reg}}$ is G-flexible if and only if the orbit $G.p$ is open.*
- (b) *An open G-orbit (if it exists) is unique and consists of all G-flexible points in X_{reg} .*

Proof. (a) is immediate from Proposition 1.7. Moreover (b) follows from (a) since any two open G -orbits overlap and so must coincide. \square

Let us note that by Corollary 1.10(a) the definition of a G -flexible point only depends on G and not on the choice of the generating set \mathcal{G} .

Using the semicontinuity of the fiber dimension we can deduce the following semicontinuity result for orbits of groups that are algebraically generated.

Corollary 1.11. *If a group $G \subseteq \text{Aut}(X)$ is algebraically generated then the function $x \mapsto \dim G.x$ is lower semicontinuous on X . In particular, there is a Zariski open subset $U \subseteq X$ filled in by orbits of maximal dimension.*

Proof. We may suppose that $G = \langle \mathcal{G} \rangle$, where \mathcal{G} is a family of connected algebraic subgroups of $\text{Aut}(X)$ closed under conjugation in G . For a maximal sequence $\mathcal{H} = (H_1, \dots, H_s)$ consider the map $\Phi_{\mathcal{H}}$ from (3). By the semicontinuity of fiber dimension the function

$$X \ni x \longmapsto \dim_{\tau(x)} \Phi_{\mathcal{H}}^{-1}(x, x)$$

is upper semicontinuous on X , where $\tau(x) = (1, \dots, 1, x) \in H_1 \times \dots \times H_s \times X$. Here $\Phi_{\mathcal{H}}^{-1}(x, x)$ is just the fiber of the map $\Phi_{\mathcal{H},x} : H_1 \times \dots \times H_s \rightarrow G.x$ over x .

Fix a point $x_0 \in X$. Enlarging \mathcal{H} we may assume that $\Phi_{\mathcal{H},x_0}$ is a submersion. Thus for x in a suitable neighborhood U of x_0

$$\begin{aligned} \dim G.x_0 &= \sum_{\sigma=1}^s \dim H_{\sigma} - \dim \Phi^{-1}(x_0, x_0) \\ &\leq \sum_{\sigma=1}^s \dim H_{\sigma} - \dim \Phi^{-1}(x, x) \\ &\leq \dim G.x. \end{aligned}$$

It follows that $\dim G.x \geq \dim G.x_0$ for $x \in U$, as required. \square

In view of our preceding results the following analog of the Rosenlicht Theorem on rational invariants holds in our setting with an almost identical proof, see e.g. [33, Theorem 2.3]. For the reader's convenience we add the argument.

Theorem 1.12. *Let G be an algebraically generated group acting on a variety X . Then there exists a finite collection of rational G -invariants which separate G -orbits in general position.*

Proof. Replacing X by a subset U as in Corollary 1.11 we may assume that all orbits of G are of maximal dimension. In particular then all G -orbits are closed in X . Let $\Gamma \subseteq X \times X$ consist of all pairs (x, x') such that x and x' are in the same G -orbit. Note that this is just the the image of the map $G \times X \rightarrow X \times X$ with $(g, x) \mapsto (g.x, x)$. As we have seen in the proof of Proposition 1.3, Γ contains an open dense subset, say Γ_0 , of the closure $\bar{\Gamma}$ in $X \times X$.

Letting G act on the first component of $X \times X$ we may assume that Γ_0 is G -invariant, since otherwise we can replace it by the union of all translates of Γ_0 . If $p_2 : \Gamma_0 \rightarrow X$ denotes the second projection then for a general point $x \in X$ the fibre $p_2^{-1}(x) = G.x \times \{x\}$ is closed in $X \times \{x\}$. Hence there is an open dense subset $U \subseteq X$ such that $\Gamma_0 \cap p_2^{-1}(U)$ is closed in $X \times U$. In particular it follows that $\Gamma \cap X \times U$ is closed in $X \times U$. Shrinking U we may also assume that U is affine.

Let $\mathcal{I} \subseteq \mathcal{O}_{X \times U}$ be the ideal sheaf of $\Gamma \cap X \times U$ and let J be the ideal generated by \mathcal{I} in the algebra $\mathbb{k}(X) \otimes \mathbb{k}[U]$ ⁴. The ideal J is G -invariant assuming that G acts on the first factor of $X \times X$. Moreover, J is generated as a $\mathbb{k}(X)$ -vector subspace by G -invariant elements (see [33, Lemma 2.4]). We can find a finite set of generators of J , say F_1, \dots, F_p , among these elements. We have

$$F_i = \sum_s f_{is} \otimes u_{is}, \quad \text{where } f_{is} \in \mathbb{k}(X)^G \quad \text{and} \quad u_{is} \in \mathbb{k}[U].$$

Let us show that the functions f_{is} separate orbits in general position.

Shrinking U once again we may assume that all the f_{is} are regular functions on U and that the elements F_i generate the ideal I . Then the orbit of a point $x \in U$ is defined by the equations $F_i(x, y) = \sum_s f_{is}(x)u_{is}(y) = 0$, $i = 1, \dots, p$. Consequently, the equalities $f_{is}(x_1) = f_{is}(x_2)$ for all i and s imply that $G.x_1 = G.x_2$ on U . \square

As in [33, Corollary on p. 156] this theorem has the following consequence.

Corollary 1.13. *Let G be an algebraically generated group acting on a variety X of dimension n . Then*

$$\text{trdeg}(\mathbb{k}(X)^G : \mathbb{k}) = \min_{x \in X} \{\text{codim}_X G.x\}.$$

In particular, there is an open orbit of G if and only if $\mathbb{k}(X)^G = \mathbb{k}$.

Remark 1.14. It may happen that all G -orbits in X have a common point in their closures and so the only *regular* G -invariants are the constants. This is the case, for instance, for the group $G = \mathbb{G}_m$ acting on \mathbb{A}^n by homotheties. Cf. also an example in [27, §4.2], where the group G is generated by its one-parameter unipotent subgroups.

1.2. Transversality. If an algebraic group G acts transitively on an algebraic variety X and Z, Y are smooth subvarieties of X then by Kleiman's Transversality Theorem [24] a general g -translate $g.Z$ ($g \in G$) meets Y transversally. In this subsection we establish the following analogue of Kleiman's Theorem for an arbitrary algebraically generated group (which might be of infinite dimension).

Theorem 1.15. *Let a subgroup $G \subseteq \text{Aut}(X)$ be algebraically generated by a system \mathcal{G} of connected algebraic subgroups closed under conjugation in G . Suppose that G acts with an open orbit $O \subseteq X$.*

Then there exist subgroups $H_1, \dots, H_s \in \mathcal{G}$ such that for any locally closed reduced subschemes Y and Z in O one can find a Zariski dense open subset $U = U(Y, Z) \subseteq H_1 \times \dots \times H_s$ such that every element $(h_1, \dots, h_s) \in U$ satisfies the following.

- (a) *The translate $(h_1 \dots h_s).Z_{\text{reg}}$ meets Y_{reg} transversally.*
- (b) $\dim(Y \cap (h_1 \dots h_s).Z) \leq \dim Y + \dim Z - \dim X$.

In particular $Y \cap (h_1 \dots h_s).Z = \emptyset$ if $\dim Y + \dim Z < \dim X$.

The proof is based on the following auxiliary result, which is complementary to Proposition 1.7.

⁴Here $\mathbb{k}(X)$ denotes the function field of X .

Proposition 1.16. *Let the assumption of Theorem 1.15 hold. Then there is a sequence $\mathcal{H} = (H_1, \dots, H_s)$ in \mathcal{G} so that for a suitable open dense subset $U \subseteq H_s \times \dots \times H_1$, the map*

$$(4) \quad \Phi_s : H_s \times \dots \times H_1 \times O \longrightarrow O \times O \quad \text{with} \quad (h_s, \dots, h_1, x) \mapsto ((h_s \cdot \dots \cdot h_1).x, x)$$

is smooth on $U \times O$.

Proof. According to Proposition 1.3 there are subgroups $H_1, \dots, H_s \subseteq G$ in \mathcal{G} such that Φ_s is surjective. Hence there is an open dense subset $U_s \subseteq H_s \times \dots \times H_1 \times O$ on which Φ_s is smooth. Assuming that U_s is maximal with this property we consider the complement $A_s = (H_s \times \dots \times H_1 \times O) \setminus U_s$.

Let us study the effect of increasing the number of factors, i.e., passing to

$$\Phi_{s+1} : H_{s+1} \times \dots \times H_1 \times O \longrightarrow O \times O$$

The map Φ_{s+1} is smooth on $H_{s+1} \times U_s$. Indeed, for every $h_{s+1} \in H_{s+1}$ we have a commutative diagram

$$\begin{array}{ccc} H_{s+1} \times \dots \times H_1 \times O & \xrightarrow{\Phi_{s+1}} & O \times O \\ \uparrow h_{s+1} \times \text{id} & & \uparrow h_{s+1} \times \text{id} \\ \{1\} \times H_s \times \dots \times H_1 \times O & \xrightarrow{\Phi_s} & O \times O \end{array}$$

where the lower horizontal map is smooth on U_s . In other words, $U_{s+1} \supseteq H_{s+1} \times U_s$ or, equivalently, $A_{s+1} \subseteq H_{s+1} \times A_s$. We claim that increasing the number of factors by H_{s+1}, \dots, H_{s+t} in a suitable way, we can achieve that

$$(5) \quad \dim A_{s+t} < \dim(H_{s+t} \times \dots \times H_{s+1} \times A_s).$$

If $(h_s, \dots, h_1, x) \in A_s$ and $y = (h_s \cdot \dots \cdot h_1).x$ then for suitable H_{s+t}, \dots, H_{s+1} the map

$$H_{s+t} \times \dots \times H_{s+1} \times O \longrightarrow O \times O$$

is smooth in all points $(1, \dots, 1, y)$, see Proposition 1.7. In particular Φ_{s+t} is smooth in all points $(1, \dots, 1, h_s, \dots, h_1, x)$ with $x \in O$, i.e.

$$(1, \dots, 1) \times A_s \cap A_{s+t} = \emptyset.$$

Now (5) follows.

Thus increasing the number of factors suitably we can achieve that⁵

$$\dim A_s < \dim(H_s \times \dots \times H_1).$$

In particular, the image of A_s under the projection

$$\pi : H_s \times \dots \times H_1 \times O \longrightarrow H_s \times \dots \times H_1$$

is nowhere dense. Hence there is an open dense subset $U \subseteq H_s \times \dots \times H_1$ such that $\Phi_s : U \times O \rightarrow O \times O$ is smooth. \square

⁵In fact we can make the difference $\dim(H_s \times \dots \times H_1) - \dim A_s$ arbitrarily large.

Proof of Theorem 1.15. Let us first show (a). Replacing Y and Z by Y_{reg} and Z_{reg} , respectively, we may assume that Y and Z are smooth. Applying Proposition 1.16 there are subgroups H_1, \dots, H_s in \mathcal{G} such that $\Phi_s : U \times O \rightarrow O \times O$ is smooth for some open subset $U \subseteq H_1 \times \dots \times H_s$. In particular $\mathcal{Y} = \Phi_s^{-1}(Y \times Z) \cap (U \times O) \subseteq U \times Z$ is smooth. By Sard's Theorem the general fiber of the projection $\mathcal{Y} \rightarrow U$ is smooth as well. In other words, shrinking U we may assume that all fibers of this projection are smooth. Since for a point $h = (h_1, \dots, h_s) \in U$ the fiber $\mathcal{Y} \cap \pi^{-1}(h)$ maps bijectively via Φ_s onto $Y \cap (h_1 \cdot \dots \cdot h_s).Z$, (a) follows.

Now (b) follows by an easy induction on $l = \dim Y + \dim Z$, the case of $l = 0$ being trivial. Indeed, applying (a) and the induction hypothesis to Y_{sing} and Z and also to Y and Z_{sing} , for suitable connected algebraic subgroups H_1, \dots, H_s and general $(h_1, \dots, h_s) \in H_1 \times \dots \times H_s$ we have that Y_{reg} and $(h_1 \cdot \dots \cdot h_s).Z_{\text{reg}}$ meet transversally and that

$$\begin{aligned} \dim(Y_{\text{sing}} \cap (h_1 \cdot \dots \cdot h_s).Z) &\leq \dim Y_{\text{sing}} + \dim Z - \dim X; \\ \dim(Y \cap (h_1 \cdot \dots \cdot h_s).Z_{\text{sing}}) &\leq \dim Y + \dim Z_{\text{sing}} - \dim X. \end{aligned}$$

This immediately implies the desired result. \square

1.3. Special subgroups. Let X be an algebraic variety. The following notion is central in the sequel.

Definition 1.17. A subgroup G of the automorphism group $\text{Aut}(X)$ will be called *special*⁶ if it is generated by a family of one-parameter unipotent subgroups i.e., subgroups isomorphic to \mathbb{G}_a .

We give two simple examples.

Example 1.18. (1) The group $\text{SAut}(X)$ is special. The image of $\text{SAut}(X)$ under the diagonal embedding $\text{SAut}(X) \hookrightarrow \text{SAut}(X^m)$ is also a special subgroup.

(2) A connected affine algebraic group is special if and only if it does not admit nontrivial characters [32]. In particular, every semi-simple algebraic group is special.

In the sequel it will be important to deal with the infinitesimal generators of subgroups of $\text{Aut}(X)$ isomorphic to \mathbb{G}_a . Let us collect the necessary facts.

1.19. (1) ([36]) If the group \mathbb{G}_a acts on an affine variety $X = \text{Spec } A$ then the associated derivation ∂ on A is locally nilpotent, i.e. for every $f \in A$ we can find $n \in \mathbb{N}$ such that $\partial^n(f) = 0$.

(2) Conversely, given a locally nilpotent \mathbb{k} -linear derivation⁷ $\partial : A \rightarrow A$ and $t \in \mathbb{k}$, the map $\exp(t\partial) : A \rightarrow A$ is an automorphism of A or, equivalently, of X . Thus for $\partial \neq 0$, $H = \exp(\mathbb{k}\partial)$ can be regarded as a subgroup of $\text{Aut}(X)$ isomorphic to \mathbb{G}_a (see [12]). Considering ∂ as a vector field on X the action of $H \cong \mathbb{G}_a$ on X is just the associated phase flow.

(3) The ring of invariants $\mathbb{k}[X]^H = \ker \partial$ has transcendence degree⁸ $\dim X - 1$. Furthermore, for any H -invariant function $f \in \mathbb{k}[X]^H$ the one-parameter unipotent

⁶ In the case where $X = \mathbb{A}^n$ these groups were called ∂ -generated in [31, Definition 2.1].

⁷Or LND, for short.

⁸Or the Gelfand-Kirillov dimension.

subgroup $H_f = \exp(\mathbb{k}f\partial)$ will play an important role in the sequel. It will be called a *replica* of H .

It acts on X in the same direction as H does, but with a different speed along the orbits. In this way $\ker \partial$ is presented as an abelian subalgebra⁹ of the Lie algebra of all regular vector fields on X .

For a special group $G \subseteq \text{SAut}(X)$ we let $\text{LND}(G)$ denote the set of all locally nilpotent vector fields on X which generate the one-parameter unipotent subgroups of G . This set is stable under conjugation in G and is a cone, i.e., $\mathbb{k} \cdot \text{LND}(G) \subseteq \text{LND}(G)$. In the sequel we consider often subsets $\mathcal{N} \subseteq \text{LND}(G)$ of locally nilpotent vector fields such that the associated set of one-parameter subgroups $\mathcal{G} = \{\exp(\mathbb{k}\partial) : \partial \in \mathcal{N}\}$ form a generating set of algebraic subgroups for G . By abuse of language, we often say that \mathcal{N} is a generating set of G , and we write $G = \langle \mathcal{N} \rangle$.

From Proposition 1.7 we deduce the following result.

Corollary 1.20. *Given a special subgroup $G = \langle \mathcal{N} \rangle$ of $\text{Aut}(X)$, where $\mathcal{N} \subseteq \text{LND}(G)$ is stable under conjugation in G , there are locally nilpotent vector fields $\partial_1, \dots, \partial_s \in \mathcal{N}$ which span the tangent space $T_p(G.p)$ at every point $p \in X$.*

For a point $p \in X$ we let $\text{LND}_p(G) \subseteq T_pX$ denote the *nilpotent cone* of all tangent vectors $\partial(p)$, where ∂ runs over $\text{LND}(G)$. By Corollary 1.20 we have $T_p(G.p) = \text{Span LND}_p(G)$.¹⁰ Thus a point $p \in X_{\text{reg}}$ is G -flexible (see Definition 1.9) if and only if the cone $\text{LND}_p(G)$ spans the whole tangent space T_pX at p .

Applying Corollary 1.10 to the special automorphism group $G = \text{SAut}(X)$ yields the equivalence (i) \Leftrightarrow (iii) in the introduction, cf. Theorem 0.1.

Corollary 1.21. *Given an affine variety X the action of $\text{SAut}(X)$ on X_{reg} is transitive if and only if X is flexible.*

In the following example we illustrate the notions of replica and of a special group in the case of the special automorphism group $\text{SAut}(\mathbb{A}^n)$ of an affine space \mathbb{A}^n over \mathbb{k} .

Example 1.22. This group contains the one-parameter unipotent subgroup of translations in any given direction. The infinitesimal generator of such a subgroup is a directional partial derivative. Such a derivative defines a locally nilpotent derivation of the polynomial ring in n variables, with the associated phase flow being the group of translations in this direction. Its replicas are the one parameter groups of shears in the same direction.

As another example, consider the locally nilpotent derivation $\partial = X \frac{\partial}{\partial Y} + Y \frac{\partial}{\partial Z}$ of the polynomial ring $\mathbb{k}[X, Y, Z]$ and an invariant function $f = Y^2 - 2XZ \in \ker \partial$. The corresponding replica H_f contains in particular the famous Nagata automorphism $H_f(1) = \exp(f \cdot \partial) \in \text{SAut}(\mathbb{A}^3)$, see [37].

Notice that any automorphism $\alpha \in \text{SAut}(\mathbb{A}^n)$ preserves the usual volume form on \mathbb{A}^n (see [31] or Lemma 4.10 below). Hence $\text{SAut}(\mathbb{A}^n) \subseteq G_n$, where G_n denotes the subgroup of $\text{Aut}(\mathbb{A}^n)$ consisting of all automorphisms with Jacobian determinant 1.

⁹It is contained in the centralizer of ∂ . So the latter is infinite dimensional provided that $\dim X \geq 2$.

¹⁰Cf. Corollary 4.3 below.

Recall that the tame subgroup $T_n \subseteq \text{Aut}(\mathbb{A}^n)$ is the subgroup generated by all *elementary* automorphisms $\sigma \in \text{Aut} \mathbb{A}^n$ of the form

$$\sigma = \sigma_{i,\alpha,f} : (x_1, \dots, x_{i-1}, x_i, x_{i+1}, \dots, x_n) \mapsto (x_1, \dots, x_{i-1}, \alpha x_i + f, x_{i+1}, \dots, x_n),$$

where $f \in \mathbb{k}[x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n]$ and $\alpha \in \mathbb{G}_m$. An elementary automorphism σ with $\alpha = 1$ can be included into a one parameter unipotent subgroup $\{\sigma_{i,1,t,f}\}_{t \in \mathbb{k}}$. Since for any $\lambda \in \mathbb{G}_m$ the conjugate $h_\lambda \circ \sigma \circ h_\lambda^{-1}$, where $h_\lambda(x) = \lambda x$, is again elementary, any tame automorphism β with Jacobian determinant 1 can be written as a product of elementary automorphisms with Jacobian determinant 1 (cf. [31, Examples 2.3 and 2.5] or [13, Proposition 9]). Hence

$$(6) \quad T_n \cap G_n \subseteq \text{SAut}(\mathbb{A}^n) \subseteq G_n.$$

For $n = 3$ the first inclusion is proper. Indeed, due to the well known Shestakov-Umirbaev Theorem [37] the Nagata automorphism $H_f(1)$ is wild. Hence $H_f(1) \in \text{SAut}(\mathbb{A}^3) \setminus (T_3 \cap G_3)$.

However, for $n = 2$ by the Jung-van der Kulk Theorem $\text{Aut}(\mathbb{A}^2) = T_2$ and so by (6) $\text{SAut}(\mathbb{A}^2) = G_2$. It is known that G_2 is perfect and is the commutator subgroup of the group $\text{Aut}(\mathbb{A}^2)$ (see e.g., [13, Proposition 10]). The question arises whether the equality $\text{SAut}(\mathbb{A}^n) = G_n$ holds as well for $n \geq 3$ (cf. [31, Problem 2.1]).

2. INFINITE TRANSITIVITY

2.1. Main theorem. In this section we show that the special automorphism group of a flexible variety X acts infinitely transitively on X_{reg} . We state this in a more general setup which turns out to be necessary for later applications. Let us first fix some notation and assumptions for this and the next subsection.

2.1. Let X be an affine algebraic variety over \mathbb{k} . Let $G \subseteq \text{SAut}(X)$ be a subgroup generated by a given set, say, \mathcal{N} of locally nilpotent vector fields such that the following conditions are satisfied.

- (1) \mathcal{N} is closed under conjugation by elements in G .
- (2) \mathcal{N} is closed under taking *replicas*, i.e. for all $\partial \in \mathcal{N}$ and $f \in \ker \partial$ we also have $f\partial \in \mathcal{N}$.

We call such a generating set *saturated*¹¹. We note that the important condition here is (2), since starting from a set of locally nilpotent vector fields \mathcal{N}_0 generating G and only satisfying (2) we can add all conjugates of elements in \mathcal{N}_0 and obtain thereby a saturated set \mathcal{N} generating the same group G .

By Corollary 1.20 there are finitely many vector fields $\partial_1, \dots, \partial_s$ in \mathcal{N} that span the tangent space $T_x(G.x)$ at every point $x \in X$. Throughout this section these vector fields will be fixed.

The next result implies Theorem 0.1 in the Introduction.

Theorem 2.2. *Let X be an affine algebraic variety of dimension ≥ 2 and let $G \subseteq \text{SAut}(X)$ be a subgroup generated by a saturated set \mathcal{N} of locally nilpotent vector fields, which acts with an open orbit $O \subseteq X$. Then G acts infinitely transitively on O .*

¹¹ See [31, Definition 2.2] for a closely related notion of a *finitely ∂ -generated* group of automorphisms of $X = \mathbb{A}^n$.

Remark 2.3. In fact, the equivalence (i) \Leftrightarrow (ii) \Leftrightarrow (iii) of Theorem 0.1 is valid for any quasi-affine variety U which is the complement to a codimension ≥ 2 subvariety in a normal affine variety $X = \text{Spec } A$. This can be deduced from Theorem 2.2 by a straightforward argument since the group $\text{SAut}(U)$ of an open subset $U \subseteq X$ acts also on $A = \Gamma(U, \mathcal{O}_U)$.

2.4. For a subset $Z \subseteq X$ we let $\mathcal{N}_Z = \{\partial \in \mathcal{N} : \partial|_Z = 0\}$ be the set of derivations in \mathcal{N} vanishing on Z , and let

$$G_{\mathcal{N}, Z} = \langle H = \exp(\mathbb{k}\partial) : \partial \in \mathcal{N}_Z \rangle$$

be the subgroup of G generated by all exponentials in \mathcal{N}_Z . Clearly the automorphisms in $G_{\mathcal{N}, Z}$ fix the set Z pointwise. In the case $\mathcal{N} = \text{LND}(G)$ we simply write G_Z instead of $G_{\mathcal{N}, Z}$. Since \mathcal{N}_Z is $G_{\mathcal{N}, Z}$ -saturated the group $G_{\mathcal{N}, Z} = \langle \mathcal{N}_Z \rangle$ is again generated by a saturated set of locally nilpotent derivations.

With these notation and assumptions our main technical result can be formulated as follows.

Theorem 2.5. *Let X be an affine algebraic variety of dimension ≥ 2 and let $G \subseteq \text{SAut}(X)$ be a subgroup generated by a saturated set \mathcal{N} of locally nilpotent vector fields, which acts with an open orbit $O \subseteq X$. Then for every finite subset $Z \subseteq O$ the group $G_{\mathcal{N}, Z}$ acts transitively on $O \setminus Z$.*

Before embarking on the proof let us show how Theorem 2.2 follows.

Proof of Theorem 2.2. Let x_1, \dots, x_m and x'_1, \dots, x'_m be sequences of points in O with $x_i \neq x_j$ and $x'_i \neq x'_j$ for $i \neq j$. Let us show by induction on m that there is an automorphism $g \in G$ with $g.x_i = x'_i$ for all $i = 1, \dots, m$. As G acts transitively on O this is certainly true for $m = 1$. For the induction step suppose that there is already an automorphism $\alpha \in G$ with $\alpha.x_i = x'_i$ for $i = 1, \dots, m-1$. Applying Theorem 2.5 to $Z = \{x'_1, \dots, x'_{m-1}\}$ we can also find an automorphism $\beta \in G_{\mathcal{N}, Z}$ with $\beta(\alpha(x_m)) = x'_m$. Clearly then $g = \beta \circ \alpha$ satisfies $g.x_i = x'_i$ for all $i = 1, \dots, m$. \square

2.2. Proof of Theorem 2.5. To deduce Theorem 2.5 we need a few preparations. As before X stands for an affine algebraic variety over \mathbb{k} . Let us introduce the following technical notion.

Definition 2.6. Let $G \subseteq \text{SAut}(X)$ be a special subgroup and let $\Omega \subseteq X$ be a subset stabilized by G . We say that a locally nilpotent vector field $\partial \in \text{LND}(G)$ with associated one-parameter subgroup $H = \exp(\mathbb{k}\partial)$ satisfies the *orbit separation property* on Ω , if there is an H -stable subset $U(H) \subseteq \Omega$ such that

- (a) for each G -orbit O contained in Ω , the intersection $U(H) \cap O$ is open and dense in O , and
- (b) the global H -invariants $\mathcal{O}_X(X)^H$ separate all one-dimensional H -orbits in $U(H)$.

The reader should note that we allow $U(H) \cap O$ to contain or even to consist of 0-dimensional H -orbits. We also emphasize that Ω can be *any* union of orbits and can e.g. contain orbits in the singular part of X .

Similarly we say that a set of locally nilpotent vector fields \mathcal{N} satisfies the *orbit separation property* on Ω if this holds for every $\partial \in \mathcal{N}$.

Remarks 2.7. 1. Let ∂ be a locally nilpotent vector field on X and let $H = \exp(\mathbb{k}\partial)$ be the subgroup of $\text{SAut}(X)$ generated by ∂ . By a corollary of the Rosenlicht Theorem on rational invariants [33, Proposition 3.4] there exists an H -invariant Zariski open set $U(H)$ such that the restriction $H|U(H)$ admits a geometric quotient

$$\pi : U(H) \rightarrow U(H)/H$$

given by a finite set of regular H -invariant functions on X . If G possesses an open orbit O then we can choose such an open set $U(H)$ contained in O . Letting $\Omega = O$ every $\partial \in \text{LND}(G)$ satisfies the orbit separation property on Ω .

2. Given a locally nilpotent derivation $\partial \in \text{LND}(G)$, it satisfies the orbit separation property on any set Ω which is a union of G -orbits meeting $U(H)$. In particular this property holds for general G -orbits (cf. Corollary 1.11).

3. As we shall see in Example 2.14 the orbit separation property is not necessarily satisfied on any G -orbit. However, there are interesting geometric situations where this property holds for arbitrary G -orbits, see Subsection 3.2.

We need the following simple Lemma.

Lemma 2.8. *If a locally nilpotent vector field $\partial \in \text{LND}(G)$ satisfies the orbit separation property on a G -stable subset $\Omega \subseteq X$ then also every replica $f\partial$, $f \in \ker \partial$, and every g -conjugate $g^*(\partial) = g \circ \partial \circ g^{-1}$, $g \in G$, has this property.*

Proof. Let $\partial' = f\partial$ be a replica of ∂ with associated one-parameter subgroup H' . In the case $f = 0$ the assertion is obvious. Otherwise the one dimensional orbits of H' are also one dimensional orbits of H , and the H and H' invariant functions are the same. Hence setting $U(H') = U(H)$, (a) and (b) in Definition 2.6 are again satisfied for H' . The fact that any g -conjugate of ∂ has again the orbit separation property, is easy and can be left to the reader. \square

For the remaining part of this subsection we fix the following notation.

2.9. Let $G \subseteq \text{SAut}(X)$ be a special subgroup generated by a saturated set \mathcal{N} of locally nilpotent vector fields. Let $\Omega \subseteq X$ a G -stable subset. We choose $\partial_1, \dots, \partial_s \in \mathcal{N}$ with associated one-parameter subgroups $H_\sigma = \exp(\mathbb{k}\partial_\sigma)$ and assume that the following two conditions are satisfied.

- (1) $\partial_1, \dots, \partial_s \in \mathcal{N}$ span $T_x(G.x)$ for every point $x \in \Omega$ (see 1.20), and
- (2) ∂_σ has the orbit separation property on Ω for all $\sigma = 1, \dots, s$.

Consequently there are subsets $U(H_\sigma) \subseteq \Omega$ such that conditions (a) and (b) in Definition 2.6 are satisfied with $H = H_\sigma$. We let

$$V = \bigcap_{\sigma=1}^s U(H_\sigma).$$

In particular,

- (i) $V \cap O$ is open and dense in O for every orbit O contained in Ω , and
- (ii) any two points in V in different one dimensional H_σ -orbits can be separated by an H_σ -invariant function on X for all $\sigma = 1, \dots, s$.

Lemma 2.10. *Under the assumptions of 2.9 let $x, y \in \Omega$ be distinct points lying in G -orbits of dimension ≥ 2 . Then there is an automorphism $g \in G$ such that*

- (a) $g.x, g.y \in V$, and
- (b) $g.x$ and $g.y$ are lying in different H_σ -orbits¹² for all $\sigma = 1, \dots, s$.

Proof. (a) Since G acts transitively on every G -orbit O in Ω and $V \cap O$ is dense in O , we can find $g \in G$ with $g.x \in V$. Replacing x by $g.x$ we may assume that $x \in V$. For some $\sigma \in \{1, \dots, s\}$ we have $H_\sigma.y \cap V \neq \emptyset$. Taking $h \in H_\sigma$ general we have $h.x \in V$ and $h.y \in V$, as required.

(b) By (a) we may assume that $x, y \in V$. The property that $g.x$ and $g.y$ are in different H_σ -orbits is an open condition. Thus by recursion it suffices to find $g \in G$ such that (b) is satisfied for a fixed σ . If x and y are already in different H_σ -orbits then there is nothing to show.

So suppose that this is not the case and so x, y are sitting on the same H_σ -orbit, which is then necessarily one dimensional. By assumption the vector fields $\partial_1, \dots, \partial_s$ span the tangent space $T_x(G.x)$ at x , and the G -orbit of x has dimension ≥ 2 . Hence ∂_τ is not tangent to $H_\sigma.x$ at x for some τ . In particular the orbits $H_\sigma.x$ and $H_\tau.x$ are both of dimension one and have only finitely many points in common.

If x and y are in different H_τ -orbits then we can choose a global H_τ -invariant f with $f(x) = 1$ and $f(y) = 0$. The group $H = \exp(\mathbb{k}f\partial_\tau)$ is contained in G , fixes y and moves x along $H_\tau.x$. Hence for a general $g \in H_\tau$ the points $g.x$ and $g.y = y$ lie on different H_σ -orbits.

Assume now that x and y belong to the same H_τ -orbit. We claim that again $g.x$ and $g.y$ are in different H_σ -orbits for a general $g \in H_\tau$.

To show this claim we consider $h_t = \exp(t\partial_\tau) \in H_\tau$. By assumption $h_a.x = y$ for some $a \in \mathbb{k}$, $a \neq 0$. We can find an H_σ -invariant function f on X , which induces a polynomial $p(t) = f(h_t.x)$ of positive degree in $t \in \mathbb{k}$. If $h_t.x$ and $h_t.y$ are in the same H_σ -orbits for a general $t \in \mathbb{k}$ then

$$p(t) = f(h_t.x) = f(h_t.y) = f(h_t.(h_a.x)) = f(h_{a+t}.x) = p(t + a),$$

which is impossible. Hence for a general $g = h_t \in H_\tau$ the points $g.x$ and $g.y$ are in different H_σ -orbits, as desired. \square

Lemma 2.11. *With the notations as in 2.9 assume that $x, y \in V$ are distinct points lying in different (possibly zero dimensional) H_σ -orbits for all $\sigma = 1, \dots, s$. Then the vector fields $\partial \in \mathcal{N}$ vanishing at x span $T_y(G.y)$.*

Proof. The vectors $\partial_\sigma(y)$ with $1 \leq \sigma \leq s$ span the tangent space $T_y(G.y)$. Thus it suffices to find replicas $\partial'_1, \dots, \partial'_s$ of $\partial_1, \dots, \partial_s$, which vanish at x and are equal to ∂_σ at the point y .

If the H_σ -orbit of x is a point, then necessarily ∂_σ vanishes at x and we can choose $\partial'_\sigma = \partial_\sigma$. If the H_σ -orbit of y is a point then $\partial_\sigma(y) = 0$ and so we can take $\partial'_\sigma = 0$. Assume now that both $H_\sigma.x$ and $H_\sigma.y$ are one dimensional. By our construction of V there is an H_σ -invariant function f_σ on X with $f_\sigma(x) = 0$ and $f_\sigma(y) = 1$. Hence $\partial'_\sigma = f_\sigma\partial_\sigma$ is a locally nilpotent vector field on X vanishing in x and equal to ∂_σ at y . \square

¹²Possibly of dimension 0.

Corollary 2.12. *For each $x \in \Omega$ and every G -orbit $O \subseteq \Omega$ the group $G_{\mathcal{N},x}$ as in 2.4 acts transitively on $O \setminus \{x\}$.¹³*

Proof. Let y be a point in $O \setminus \{x\}$. With the notations as in 2.9, according to Lemma 2.10 there is an automorphism $g \in G$ with $g.x, g.y \in V$ such that $g.x, g.y$ are in different (possibly 0-dimensional) H_σ -orbits for $i = 1, \dots, s$. By Lemma 2.11 the vector fields $\partial \in \mathcal{N}$ vanishing at $g.x$ span $T_{g.y}(O)$. Using the fact that \mathcal{N} is stable under conjugation by elements $g \in G$ it follows that the vector fields in \mathcal{N} vanishing at x span the tangent space $T_y(O)$. In other words, y is a $G_{\mathcal{N},x}$ -flexible point on the orbit closure \bar{O} . Applying 1.10 we obtain that $G_{\mathcal{N},x}$ acts transitively on $O \setminus \{x\}$. \square

Proof of Theorem 2.5. By Remark 2.7(1) the orbit separation property is satisfied on the open orbit $\Omega = O$. Given a set $Z = \{x_1, \dots, x_m\} \subseteq O$ of m distinct points we consider $Z_\mu = \{x_1, \dots, x_\mu\}$ for $\mu = 1, \dots, m$. Let us show by induction on μ that $G_{\mathcal{N},Z_\mu}$ acts transitively on $O \setminus Z_\mu$. For $\mu = 1$ this follows from Corollary 2.12. Assuming for some $\mu < m$ that $G_{\mathcal{N},Z_\mu}$ acts transitively on $O \setminus Z_\mu$, Corollary 2.12 implies that $(G_{\mathcal{N},Z_\mu})_{x_{\mu+1}} = G_{\mathcal{N},Z_{\mu+1}}$ acts transitively on $O \setminus Z_{\mu+1}$. \square

2.3. Examples of non-separation of orbits. Suppose as before that a subgroup $G \subseteq \text{SAut}(X)$ is generated by a saturated set \mathcal{N} of locally nilpotent vector fields. While \mathcal{N} satisfies the orbit separation property 2.6 on a general G -orbit (see Remark 2.7(2)), this is not always true on an arbitrary G -orbit. Furthermore, the following example shows that on the union of two G -orbits this property might fail although it is satisfied on every single orbit.

Example 2.13. On the affine 4-space $X = \mathbb{A}^4 = \text{Spec } \mathbb{k}[X, Y, Z, U]$ let us consider the locally nilpotent vector fields

$$\partial_1 = Y \frac{\partial}{\partial X} + Z \frac{\partial}{\partial Y} \quad \text{and} \quad \partial_2 = \frac{\partial}{\partial U}.$$

Let $G \subseteq \text{SAut}(X)$ be the special subgroup generated by ∂_1, ∂_2 and all their replicas, and let $\mathcal{N} \subseteq \text{LND}(G)$ denote the saturated set generated by ∂_1 and ∂_2 . It is easily seen that $\ker \partial_1 = \mathbb{k}[Z, Y^2 - 2XZ, U]$ and $\ker \partial_2 = \mathbb{k}[X, Y, Z]$. Hence the G -orbits $O_\pm = \{Y = \pm 1, Z = 0\}$ of dimension two are not separated by H_1 -invariants, where $H_i = \exp(\mathbb{k}\partial_i)$, $i = 1, 2$. In particular, \mathcal{N} does not satisfy the orbit separation property on $O_+ \cup O_-$. However, this property is satisfied on $\Omega = O_+$ and also on $\Omega = O_-$ separately as this is the case for ∂_1 and ∂_2 (cf. Lemma 2.8).

We note also that the isomorphism $\sigma : O_+ \rightarrow O_-$ with $\sigma(x, 1, 0, u) = (-x, -1, 0, u)$ commutes with the actions of H_1 and H_2 . Hence there is no collective transitivity on $O_+ \cup O_-$ in the sense of Theorem 3.1 below while G acts on every single orbit O_\pm indeed infinitely transitively.

According to our next example one cannot expect infinite transitivity of G on an arbitrary G -orbit O of dimension ≥ 2 without assuming the orbit separation property on O . However, compare Theorem 3.1 below for a positive result.

¹³In particular, it is transitive on O if $x \notin O$.

Example 2.14. Consider the locally nilpotent derivations

$$\partial_1 = Y \frac{\partial}{\partial X} + Z \frac{\partial}{\partial Y} + U \frac{\partial}{\partial Z} \quad \text{and} \quad \partial_2 = Z \frac{\partial}{\partial X} + X \frac{\partial}{\partial Y} + U \frac{\partial}{\partial Z}$$

of the polynomial ring $\mathbb{k}[X, Y, Z, U]$. We claim that $\ker \partial_1 = \mathbb{k}[p_1, p_2, p_3, p_4]$, where

$$p_1 = U, \quad p_2 = Z^2 - 2YU, \quad p_3 = Z^3 - 3YZU + 3XU^2, \quad \text{and} \\ p_4 = \frac{p_2^3 - p_3^2}{p_1^2} = 9X^2U^2 - 18XYZU + 6XZ^3 - 3Y^2Z^2 + 8Y^3U.$$

Indeed, the image of the map

$$\rho = (p_1, \dots, p_4) : \mathbb{A}^4 \rightarrow \mathbb{A}^4$$

is contained in the hypersurface

$$F = \{X_1^2X_4 - X_2^3 + X_3^2 = 0\}$$

which is singular along the line $F_{\text{sing}} = \{X_1 = X_2 = X_3 = 0\}$. Being regular in codimension one, F is normal. We have

$$\bar{0} \in F_{\text{sing}} \quad \text{and} \quad \rho^{-1}(\bar{0}) = \rho^{-1}(F_{\text{sing}}) = \{Z = U = 0\} =: L \subseteq \mathbb{A}^4.$$

By the Weitzenböck Theorem (see e.g. [26]) there exists a quotient $E = \text{Spec}(\ker \partial_1)$. The inclusions

$$\mathbb{k}[p_1, p_2, p_3, p_4] \subseteq \ker \partial_1 \subseteq \mathbb{k}[X, Y, Z, T]$$

lead to morphisms

$$\mathbb{A}^4 \xrightarrow{\pi} E \xrightarrow{\mu} F, \quad \text{where} \quad \mu \circ \pi = \rho.$$

We claim that μ is an isomorphism. Since both E and F are normal affine threefolds, by the Hartogs Principle [6, Proposition 7.1] μ is an isomorphism if it is so in codimension one. In turn, it suffices to check that μ admits an inverse morphism defined on F_{reg} . The latter follows once we know that ρ separates the H_1 -orbits in \mathbb{A}^4 outside the plane L . Indeed, then π separates them as well, and μ induces a bijection between $\pi(\mathbb{A}^4 \setminus L) \subseteq E$ and $F_{\text{reg}} = \rho(\mathbb{A}^4 \setminus L)$.

The action of H_1 on \mathbb{A}^4 is given by

$$(7) \quad t \cdot \begin{pmatrix} x \\ y \\ z \\ u \end{pmatrix} = \begin{pmatrix} x + ty + \frac{t^2}{2}z + \frac{t^3}{6}u \\ y + tz + \frac{t^2}{2}u \\ z + tu \\ u \end{pmatrix}.$$

Let O be an H_1 -orbit contained in $\mathbb{A}^4 \setminus L$. Suppose first that $p_1|_O = U|_O = u \neq 0$. Letting $t = -z/u$ in (7) we get a point $A = (x, y, 0, u) \in O$. Since

$$p_2(A) = -2yu \quad \text{and} \quad p_3(A) = 3xu^2$$

we can recover the coordinates

$$(8) \quad y = -(p_2|_O)/2u \quad \text{and} \quad x = (p_3|_O)/3u^2.$$

Thus O is uniquely determined by the image $\rho(O) \in F$.

Suppose further that $p_1|O = U|O = 0$. Since $O \cap L = \emptyset$ then $Z|O = z \neq 0$. Taking $t = -y/z$ in (7) yields a point $A = (x, 0, z, 0) \in O$. Since

$$p_2(A) = z^2, p_3(A) = z^3, \quad \text{and} \quad p_4(A) = 6xz^3$$

we can recover the values¹⁴

$$(9) \quad z = (p_3|O)/(p_2|O) \quad \text{and} \quad x = (p_4|O)/6z^3.$$

Now both claims follow.

The generators p_1, \dots, p_4 of the algebra of H_1 -invariants vanish on the plane $L = \{Z = U = 0\}$ so that every H_1 -invariant is constant on L . Since ∂_2 is obtained from ∂_1 by interchanging X, Y , by symmetry also every H_2 -invariant is constant on L . Letting $G = \langle \text{Sat}(H_1, H_2) \rangle$ be the subgroup generated by H_1, H_2 , and all their replicas, it easily follows that G stabilizes L and that the G -action on L factors through the SL_2 -action. In particular, the action of G on its orbit $L \setminus \{0\}$ is not even 2-transitive, the linear dependence being an obstruction.

Observe finally that the three dimensional G -orbits in \mathbb{A}^4 are separated by the G -invariant function U .

2.4. G_Y -orbits. Let as before $G \subseteq \text{SAut}(X)$ be generated by a saturated set \mathcal{N} of locally nilpotent vector fields on an affine variety X . For a subvariety $Y \subseteq X$ we consider the subgroup $G_{\mathcal{N}, Y}$ of G as defined in 2.4.

Theorem 2.15. *If G acts on the affine variety $X = \text{Spec } A$ of dimension ≥ 2 with an open orbit O and $Y = X \setminus O$ is the complement, then $G_{\mathcal{N}, Y}$ acts transitively and hence infinitely transitively on O .*

Proof. Since $G_{\mathcal{N}, Y}$ is generated by a saturated set of locally nilpotent derivations, by Theorem 2.2 it suffices to show that $G_{\mathcal{N}, Y}$ acts transitively on O .

Using Corollary 1.20 we can choose $\partial_1, \dots, \partial_s \in \mathcal{N}$ spanning the tangent space $T_x X$ at each point $x \in O$. Letting I denote the ideal of Y in A , we claim that for every $\sigma = 1, \dots, s$ there is a nonzero function $f_\sigma \in I \cap \ker \partial_\sigma$. Indeed, Y being G -invariant, for every nonzero function $f \in I$ the span E of the orbit $H_\sigma.f$ in A , where $H_\sigma = \exp(\mathbb{k}\partial_\sigma) \subseteq G$, is an H_σ -invariant finite dimensional subspace contained in I . By the Lie-Kolchin Theorem there is a nonzero element $f_\sigma \in E$ fixed by H_σ . This proves the claim.

Let $p \in X$ be a general point so that $f_\sigma(p) \neq 0$ for $\sigma = 1, \dots, s$. We can normalize the invariants f_σ so that $f_\sigma(p) = 1$ and $f_\sigma|Y = 0$. The derivation $f_\sigma \partial_\sigma$ then vanishes on Y and so the replica $H_{\sigma, f_\sigma} = \exp(\mathbb{k}f_\sigma \partial_\sigma) \subseteq G_{\mathcal{N}, Y}$ of H_σ fixes Y pointwise while moving p in the direction of $\partial_\sigma(p)$. It follows that the $G_{\mathcal{N}, Y}$ -orbit of p is open.

Let now $q \in O$ be an arbitrary point. Choose $g \in G$ with $g.p = q$. Since g stabilizes Y the subgroup $H'_{\sigma, f_\sigma} = gH_{\sigma, f_\sigma}g^{-1} \subseteq G_{\mathcal{N}, Y}$ fixes Y pointwise and moves q into the direction of $dg(\partial_\sigma)(q)$. It follows that also the $G_{\mathcal{N}, Y}$ -orbit of q is open. Finally $G_{\mathcal{N}, Y}$ has O as an open orbit. \square

¹⁴Formulas (8) and (9) define sections of ρ in the open sets $U \neq 0$ and $Z \neq 0$, respectively. This shows that $\rho : \mathbb{A}^4 \setminus L \rightarrow F_{\text{reg}}$ is a principal \mathbb{A}^1 -bundle.

3. COLLECTIVE INFINITE TRANSITIVITY

3.1. Collective transitivity on G -varieties. By *collective infinite transitivity* we mean a possibility to transform simultaneously (that is, by the same automorphism) arbitrary finite sets of points from different orbits into a standard position. Applying the methods developed in Section 2 we can deduce the following generalization of Theorem 2.5. In the sequel X stands for an affine algebraic variety.

Theorem 3.1. *Let $G \subseteq \text{SAut}(X)$ be a subgroup generated by a saturated set \mathcal{N} of locally nilpotent vector fields, which has the orbit separation property on a G -invariant subset $\Omega \subseteq X$. Suppose that x_1, \dots, x_m and x'_1, \dots, x'_m are points in Ω with $x_i \neq x_j$ and $x'_i \neq x'_j$ for $i \neq j$ such that for each j the orbits $G.x_j$ and $G.x'_j$ are equal and of dimension ≥ 2 . Then there exists an element $g \in G$ such that $g.x_j = x'_j$ for $j = 1, \dots, m$.*

As in Section 2 this will be deduced from the following more technical result.

Theorem 3.2. *Let $G \subseteq \text{SAut}(X)$ be a subgroup generated by a saturated set \mathcal{N} of locally nilpotent vector fields. Suppose that \mathcal{N} has the orbit separation property on a G -invariant subset Ω . If $Z \subseteq \Omega$ is a finite subset and $O \subseteq \Omega$ is an orbit of dimension ≥ 2 , then the group $G_{\mathcal{N}, Z}$ acts transitively on $O \setminus Z$.*

Proof. With $Z_\mu = \{x_1, \dots, x_\mu\}$ let us show by induction on μ that $G_{\mathcal{N}, Z_\mu}$ acts transitively on $O \setminus Z_\mu$ for every G -orbit $O \subseteq \Omega$ of dimension ≥ 2 . For $\mu = 1$ this is just Corollary 2.12. Assuming for some $\mu < m$ that $G_{\mathcal{N}, Z_\mu}$ acts transitively on $O \setminus Z_\mu$, Corollary 2.12 also implies that $(G_{\mathcal{N}, Z_\mu})_{x_{\mu+1}} = G_{\mathcal{N}, Z_{\mu+1}}$ acts transitively on $O \setminus Z_{\mu+1}$. Note that by Lemma 2.8 at each step the set \mathcal{N}_{Z_μ} has again the orbit separation property on Ω so that Corollary 2.12 is indeed applicable. \square

Proof of Theorem 3.1. As in the proof Theorem 2.2 we proceed by induction on m , the case $m = 1$ being trivial. For the induction step suppose that there is already an automorphism $\alpha \in G$ with $\alpha.x_i = x'_i$ for $i = 1, \dots, m-1$. Applying Theorem 3.2 to $Z = \{x'_1, \dots, x'_{m-1}\}$ we can also find an automorphism $\beta \in G_{\mathcal{N}, Z}$ with $\beta(\alpha(x_m)) = x'_m$. Clearly then $g = \beta \circ \alpha$ has the required property. \square

3.2. Infinite transitivity on matrix varieties. In this subsection we apply our methods in a concrete setting where $X = \text{Mat}(n, m)$ is the set of all $n \times m$ matrices over \mathbb{k} endowed with the natural stratification by rank. We always assume that $mn \geq 2$. Let us precise the terminology. Let $X_r \subseteq X$ denote the subset of matrices of rank r . The product $\text{SL}_n \times \text{SL}_m$ acts naturally on X via the left-right multiplication preserving the strata X_r . For every $k \neq l$ we let $E_{kl} \in \mathfrak{sl}_n$ and $E^{kl} \in \mathfrak{sl}_m$ denote the nilpotent matrices with $x_{kl} = 1$ and the other entries equal zero. Let further $H_{kl} = I_n + \mathbb{k}E_{kl} \subseteq \text{SL}_n$ and $H^{kl} = I_m + \mathbb{k}E^{kl} \subseteq \text{SL}_m$ be the corresponding one-parameter unipotent subgroups acting on the stratification $X = \bigcup_r X_r$, and let δ_{kl} and δ^{kl} , respectively, be the corresponding locally nilpotent vector fields on X tangent to the strata.

We call *elementary* the one-parameter unipotent subgroups H_{kl} , H^{kl} , and all their replicas. In the following theorem we establish the collective infinite transitivity on the above stratification of the subgroup G of $\text{SAut}(X)$ generated by the two sides elementary subgroups (cf. [35]).

By a well known theorem of linear algebra, the subgroup $\mathrm{SL}_n \times \mathrm{SL}_m \subseteq G$ acts transitively on each stratum X_r (and so these strata are G -orbits) except for the open stratum X_n in the case where $m = n$. In the latter case the G -orbits contained in X_n are the level sets of the determinant.

Theorem 3.3. *Given two finite ordered collections \mathcal{B} and \mathcal{B}' of distinct matrices in $\mathrm{Mat}(n, m)$ of the same cardinality, with the same sequence of ranks¹⁵, and in the case where $m = n$ with the same sequence of determinants, we can simultaneously transform \mathcal{B} into \mathcal{B}' by means of an element $g \in G$, where $G \subseteq \mathrm{SAut}(\mathrm{Mat}(n, m))$ is the subgroup generated by all elementary one-parameter unipotent subgroups.*

Theorem 3.3 is an immediate consequence of Theorem 3.1 and Lemma 3.4 below. To formulate this Lemma, we let \mathcal{N} be the saturated set of locally nilpotent derivations on X generated by all locally nilpotent vector fields δ_{kl} and δ^{kl} ($k \neq l$) that is, the set of all conjugates of these derivations along with their replicas. The important observation is the following Lemma.

Lemma 3.4. *\mathcal{N} has the orbit separation property on $\Omega = X$.*

Proof. In view of Lemma 2.8 it suffices to show that the derivations δ_{kl} and δ^{kl} have the orbit separation property. Clearly it suffices to prove this for δ^{kl} . The action of the corresponding one-parameter subgroup $H^{kl} = \exp(\mathbb{k}\delta^{kl})$ on a matrix $B = (b_1, \dots, b_m) \in X$ with column vectors $b_1, \dots, b_m \in \mathbb{k}^n$ is explicitly given by

$$\exp(t\delta^{kl}).B = (b_1, \dots, b_l + tb_k, \dots, b_n),$$

where $b_l + tb_k$ is the l th column of the matrix on the right. Thus the H^{kl} -orbit of B has dimension one if and only if $b_k \neq 0$. The functions

$$B \mapsto b_{ij} \ (j \neq l) \quad \text{and} \quad B \mapsto \begin{vmatrix} b_{ik} & b_{il} \\ b_{jk} & b_{jl} \end{vmatrix} \ (i \neq j)$$

on $\mathrm{Mat}(n, m)$ are H^{kl} -invariants that obviously separate all H^{kl} -orbits of dimension one, as the reader may easily verify. \square

3.3. The case of symmetric and skew-symmetric matrices. We can apply the same reasoning to the varieties

$$X = \mathrm{Spec}(\mathbb{k}[T_{ij}]_{1 \leq i, j \leq n} / (T_{ij} - T_{ji})_{1 \leq i, j \leq n})$$

of symmetric $n \times n$ matrices over \mathbb{k} and to the variety

$$Y = \mathrm{Spec}(\mathbb{k}[T_{ij}]_{1 \leq i, j \leq n} / (T_{ij} + T_{ji})_{1 \leq i, j \leq n})$$

of skew symmetric matrices. The group SL_n acts on both varieties via

$$A.B = ABA^T, \text{ where } A \in \mathrm{SL}_n \text{ and } B \in X (\in Y, \text{ resp.}).$$

The subvariety X_r of symmetric matrices of rank r in X is stabilized by this action, and also the determinant of a matrix is preserved. In the skew symmetric case again the subvarieties Y_r of matrices of rank r are stabilized, and also the Pfaffian $\mathrm{Pf}(B)$ ¹⁶ of a matrix $B \in Y$ is preserved.

¹⁵In particular, we can choose \mathcal{B}' consisting of diagonal matrices.

¹⁶We keep the usual convention that the Pfaffian of a matrix of odd order equals zero.

By a well known theorem in linear algebra the orbits of the SL_n -action on X are the subsets X_r of matrices of rank r in X for $r < n$, whereas for $r = n$ the orbits are the level sets of the determinant. Similarly, the orbits of the SL_n -action on Y are the subsets Y_r for $r < n$, whereas for $r = n$ the orbits are the level sets of the Pfaffian.

As in subsection 3.2 the elementary matrix E_{ij} ($i \neq j$) generates a one-parameter subgroup $H_{ij} = I_n + \mathbb{k}E_{ij}$ that acts on X and on Y . The corresponding locally nilpotent vector field will be denoted by δ_{ij} .

Let $G_{\mathrm{sym}} \subseteq \mathrm{SAut}(X)$ and $G_{\mathrm{skew}} \subseteq \mathrm{SAut}(Y)$, respectively, be the subgroups generated by all H_{ij} along with their replicas. With these notations the following results hold.

Theorem 3.5. *Let M_1, \dots, M_k be a sequence of pairwise distinct symmetric matrices of order $n \geq 2$ over \mathbb{k} . Assume that M'_1, \dots, M'_k is another such sequence with*

$$\mathrm{rk}(M_i) = \mathrm{rk}(M'_i) \geq 2 \quad \text{and} \quad \det(M_i) = \det(M'_i) \quad \forall i = 1, \dots, k.$$

Then there exists an automorphism $g \in G_{\mathrm{sym}}$ with $g.M_i = M'_i$ for $i = 1, \dots, k$.

A similar result holds in the skew symmetric case.

Theorem 3.6. *Let M_1, \dots, M_k be a sequence of pairwise distinct skew-symmetric matrices of order $n \geq 2$ over \mathbb{k} . Assume that M'_1, \dots, M'_k is another such sequence with*

$$\mathrm{rk}(M_i) = \mathrm{rk}(M'_i) \quad \text{and} \quad \mathrm{Pf}(M_i) = \mathrm{Pf}(M'_i) \quad \forall i = 1, \dots, k.$$

Then there exists an automorphism $g \in G_{\mathrm{skew}}$ with $g.M_i = M'_i$ for $i = 1, \dots, k$.

We give a sketch of the proof in the symmetric case only and leave the skew-symmetric one to the reader. As in the case of generic matrices (see Theorem 3.3) Theorem 3.5 is an immediate consequence of Theorem 3.2 and Lemma 3.7 below. In this lemma we let \mathcal{N} be the saturated set of locally nilpotent derivations on X generated by all locally nilpotent vector fields δ_{kl} .

Lemma 3.7. *\mathcal{N} has the orbit separation property on $\Omega = X$.*

Proof. In view of Lemma 2.8 it suffices to show that the derivations δ_{kl} have the orbit separation property. We only treat the case $k < l$ the other one being similar. The action of the corresponding one-parameter subgroup $H_{kl} = \exp(\mathbb{k}\delta_{kl})$ on a matrix $B \in X$ with entries $b_{ij} = b_{ji}$ is explicitly given by $\exp(t\delta_{kl}).B = (b'_{ij})$, where

$$b'_{ij} = b_{ij} \text{ if } i, j \neq k, \quad b'_{ki} = b'_{ik} = b_{ik} + tb_{il} \text{ if } i \neq k, \quad \text{and} \quad b'_{kk} = b_{kk} + 2tb_{kl} + t^2b_{ll}.$$

Thus the H_{kl} -orbit of B has dimension 0 if and only if $b_{il} = 0 \forall i$. The functions

$$B \mapsto b_{ij} \ (i, j \neq k), \quad B \mapsto \begin{vmatrix} b_{ik} & b_{il} \\ b_{jk} & b_{jl} \end{vmatrix} \ (i, j \neq k), \quad \text{and} \quad B \mapsto \begin{vmatrix} b_{kk} & b_{kl} \\ b_{lk} & b_{ll} \end{vmatrix}$$

are H_{kl} -invariants that are easily seen to separate all H_{kl} -orbits of dimension one. \square

4. TANGENTIAL FLEXIBILITY, INTERPOLATION BY AUTOMORPHISMS, AND \mathbb{A}^1 -RICHNESS

4.1. Flexibility of the tangent bundle. We start with the following fact (see the Claim in the proof of Corollary 2.8 in [21]).

Lemma 4.1. *Let ∂ be a locally nilpotent vector field on the affine \mathbb{k} -scheme $X = \text{Spec } A$ and let $p \in X$ be a point. Assume that $f \in \ker \partial$ is an invariant of ∂ with $f(p) = 0$. If $\Phi = \exp(f\partial)$ is the automorphism associated with the locally nilpotent vector field $f\partial$, then*

$$(10) \quad d_p\Phi(w) = w + df(w)\partial(p) \quad \text{for all } w \in T_pX.$$

Proof. The tangent space T_pX is the space of all derivations $w : A \rightarrow \mathbb{k}$ centered at p . For such a tangent vector its image $d\Phi(w) \in T_pX$ is the derivation

$$\begin{aligned} A \ni g \mapsto w(\Phi(g)) &= w\left(\sum_{i \geq 0} \frac{f^i \partial^i(g)}{i!}\right) = \sum_{i \geq 0} \frac{1}{i!} w(f^i \partial^i(g)) \\ &= w(g) + w(f)\partial(g)(p), \end{aligned}$$

as $f(p) = 0$. Since by definition $w(f) = df(w)$, the result follows. \square

Now we can show the following result.

Theorem 4.2. *Let X be an affine algebraic variety and let $G \subseteq \text{Aut}(X)$ be a subgroup generated by a saturated set \mathcal{N} of locally nilpotent vector fields. Assume that \mathcal{N} satisfies the orbit separation property on a G -orbit O . Then for each point $p \in O$, associating to an automorphism $g \in G_{\mathcal{N},p}$ its tangent map $dg(p)$ yields a representation*

$$\tau : G_{\mathcal{N},p} \longrightarrow \text{GL}(T_pO) \quad \text{with} \quad \tau(G_{\mathcal{N},p}) = \text{SL}(T_pO).$$

Proof. The assertion is trivially true if $\dim O = 1$. Let us assume for the rest of the proof that $\dim O \geq 2$. For any one-parameter unipotent subgroup H in $G_{\mathcal{N},p}$ the image $\tau(H)$ is a subgroup of $\text{SL}(T_pO)$. Hence also $\tau(G_{\mathcal{N},p}) \subseteq \text{SL}(T_pO)$. Let us show the converse inclusion.

According to Proposition 1.7 there are locally nilpotent vector fields $\partial_1, \dots, \partial_s \in \mathcal{N}$ spanning T_xO at every point $x \in O$. Let $H_j = \exp(\mathbb{k}\partial_j)$ be the one-parameter subgroup associated with ∂_j . Using Remark 2.7(1) one can see that there are H_j -invariant open subsets $U(H_j) \subseteq O$ such that the geometric quotient $U(H_j)/H_j$ exists. Applying the orbit separation property, a suitable set of H_j -invariants from $\mathcal{O}_X(X)$ yields a generically injective map $\varrho_j : U(H_j)/H_j \rightarrow \mathbb{A}^N$ (see Remark 2.7). For a generic point $x \in \bigcap_{j=1}^s U(H_j)$ its image in $U(H_j)/H_j$ is a smooth point in which ϱ_j has maximal rank.

We may assume that $\partial_1(x), \dots, \partial_m(x)$, where $m = \dim O \leq s$, form a basis of T_xO . Hence for $j, \mu \in \{1, \dots, m\}$ with $\mu \neq j$ there exist ∂_j -invariant functions $f_{\mu j}$ on X such that $f_{\mu j}(x) = 0$ and $d_x f_{\mu j}(\partial_i(x)) = \delta_{\mu i}$. Consider the automorphism $\Phi_{\mu j}^t = \exp(t \cdot f_{\mu j} \partial_j) \in G_{\mathcal{N},x}$ for $t \in \mathbb{k}$. According to Lemma 4.1 its tangent map at x is

$$d_x \Phi_{\mu j}^t(\partial_i(x)) = \partial_i(x) + t \cdot d_x f_{\mu j}(\partial_i(x)) \cdot \partial_j(x) = \partial_i(x) + t \delta_{\mu i} \partial_j(x).$$

Thus representing the elements in $\text{GL}(T_xO)$ by matrices with respect to the basis $\partial_1(x), \dots, \partial_m(x)$, the elements $d_x \Phi_{\mu j}^t \in \text{GL}(T_xO)$, $t \in \mathbb{k}$, form just the one-parameter unipotent subgroup generated by the elementary matrix $E_{j\mu}$. Since such one-parameter subgroups generate $\text{SL}(T_xO)$, the image of $G_{\mathcal{N},x}$ in $\text{GL}(T_xO)$ contains $\text{SL}(T_xO)$ for a general point $x \in X$. Now the transitivity of G on O implies that the same is true for every point $p \in O$. \square

The following corollary is immediate.

Corollary 4.3. *Under the assumptions of Theorem 4.2 for each point $p \in O$ we have*

$$\mathcal{N}(p) := \{\partial(p) \in T_p X : \partial \in \mathcal{N}\} = T_p O.$$

In particular, the nilpotent cone $\text{LND}_p(G)$ coincides with the tangent space $T_p O$ for each $p \in O$.

Proof. Indeed, the group $G_{\mathcal{N},p}$ stabilizes $\mathcal{N}(p)$ and for $m = \dim O \geq 2$ the group $\text{SL}(T_p O)$ acts transitively on $T_p O \setminus \{0\}$. \square

Remark 4.4. The last assertion in Corollary 4.3 does not hold any more for a general special subgroup $G \subseteq \text{SAut}(X)$ which is not generated by a saturated set of locally nilpotent vector fields. For instance, if a semisimple algebraic group G acts on itself via left multiplications (i.e., $X = G$), then the cone $\text{LND}_e(G)$ is just the usual nilpotent cone in the Lie algebra $\text{Lie}(G) = T_e X$, which is a proper subcone.

To be more concrete, in the case $G = \text{SL}_2$ the nilcone $\text{LND}_e(G)$ is just the quadratic cone in $\mathfrak{sl}_2 \cong \mathbb{A}^3$ consisting of matrices with determinant 0.

We also have the following result on tangential flexibility.

Corollary 4.5. *Let X be a flexible affine variety and $G = \text{SAut}(X)$. If $\pi : E \rightarrow X$ is an irreducible and reduced G -homogeneous linear space¹⁷, which is over X_{reg} a vector bundle, then the total space E is also flexible. In particular, the tangent bundle TX and all its tensor bundles $E = (TX)^{\otimes a} \otimes (T^*X)^{\otimes b}$ are flexible.*

Proof. It suffices to check that the special automorphism group $G' = \text{SAut}(E)$ acts transitively on $E_{\text{reg}} = \pi^{-1}(X_{\text{reg}})$. Since E is G -homogeneous there is a natural inclusion $G \subseteq G'$. As X is flexible, $G \subseteq G'$ acts transitively on the zero section of E_{reg} . Moreover, X being affine for any point $e \in E_{\text{reg}}$ there is a section $V : X \rightarrow E$ with $V(\pi(e)) = e$. This section generates a \mathbb{G}_a -action $w \mapsto w + tV(\pi(w))$. With this action we can move e to the zero section of E , and the result follows. \square

Corollary 4.6. *Let X be a flexible affine variety of dimension ≥ 2 . Consider the special automorphism group $G = \text{SAut}(TX)$ of the tangent bundle TX , and let $Z \subseteq TX$ be the zero section. Then the group G_Z acts infinitely transitively on $TX_{\text{reg}} \setminus Z$.*

Proof. The special automorphism group $\text{SAut}(X)$ induces a special subgroup $\tilde{G} \subseteq G$ acting on TX_{reg} . Since X is flexible this action is transitive on the zero section, hence also on the set of fibers of $TX \rightarrow X$ over X_{reg} . On the other hand, by Theorem 4.2 the stationary subgroup \tilde{G}_p of a given point $p \in X_{\text{reg}}$ acts on $T_p X$ as $\text{SL}(T_p X)$. Since $\dim T_p X > 1$, it acts transitively off the origin. Finally the action of \tilde{G} on TX_{reg} is transitive off the zero section. Hence by Theorem 2.2 the group G_Z , being special and generated by a saturated set of locally nilpotent derivations, acts infinitely transitively on $TX_{\text{reg}} \setminus Z$. \square

4.7. For later use let us mention the following slightly more general version of Theorem 4.2. For a finite subset $Z \subseteq X$ and $p \in O$ we let $\mathcal{N}_{p,Z}^M \subseteq \mathcal{N}$ denote the set of all locally nilpotent vector fields $\partial \in \mathcal{N}$ such that ∂ has a zero at p and a zero of order $\geq M+1$ at

¹⁷in the sense of [17] Chap. II, 1.7.

all points of $Z \setminus \{p\}$. Let further $G_{p,Z}^M$ be the subgroup of G generated by all exponentials of elements in $\mathcal{N}_{p,Z}^M$. Replacing in Theorem 4.2 $G_{\mathcal{N},p}$ by $G_{p,Z}^M$ the following result holds.

Proposition 4.8. *If $\dim O \geq 2$ then the image of the group $G_{p,Z}^M$ in $\mathrm{GL}(T_p O)$ coincides with $\mathrm{SL}(T_p O)$.*

Proof. With the notation as in the proof of *loc.cit.*, by infinite transitivity (see Theorem 3.2) it suffices to show the assertion for the case that $x = p$ is general and Z consists of general points. Under this assumption we can find ∂_j -invariant functions h_j with $h_j(x) = 1$ which vanish in all points of $Z \setminus \{p\}$. Replacing in the proof of 4.2 f_{μ_j} by $h_j^{M+1} f_{\mu_j}$, the automorphisms $\Phi_{\mu_j}^t$ are the identity up to order M at the points of Z and remain unchanged at x . Now the same arguments as before give the conclusion. \square

Let further G_Z^M have the same meaning as $G_{p,Z}^M$ above, but without any constraint imposed at p . That is, G_Z^M is the subgroup of G generated by the saturated set \mathcal{N}_Z^M of locally nilpotent vector fields vanishing to order $M + 1 \geq 1$ in all points of Z . Then the same argument as before shows the following proposition.

Proposition 4.9. *Every point $p \in O \setminus Z$ is G_Z^M -flexible, hence $G_Z^M \cdot p = O \setminus Z$.*

4.2. Prescribed jets of automorphisms. Let us start with the following standard fact (see Proposition 6.4. in [20], cf. also Theorem 4.2). Recall that a *volume form* ω on a smooth algebraic variety X is a nowhere vanishing top-dimensional regular form on X ; it does exist if and only if $K_X = 0$ in $\mathrm{Pic}(X)$.

Lemma 4.10. *If X is an affine algebraic variety and $\omega \in \Omega_X^n$ a volume form on X_{reg} , then ω is preserved under every automorphism $g \in \mathrm{SAut}(X)$.*

Proof. It suffices to show that for every locally nilpotent vector field ∂ the form ω is invariant under an automorphism of $H = \exp(\mathbb{k}\partial)$. If $h_t = \exp(t\partial)$ then for every $x \in X_{\mathrm{reg}}$ the pullback $h_t^*(\omega)(x)$ is a multiple of $\omega(x)$, i.e. $h_t^*(\omega)(x) = f(x, t)\omega(x)$, where $f(x, t) \neq 0$ for all x, t . For a fixed x the function $f(x, t)$ is thus a polynomial in one variable without zero. Hence f is independent of t equal to $f(x, 1) = 1$. \square

4.11. We adopt the following notations and assumptions. If $\varphi : X \rightarrow X$ is a morphism then its m -jet $j_p^m \varphi$ at $p \in X$ can be regarded as a map of \mathbb{k} -algebras

$$j_p^m \varphi : \mathcal{O}_{X, \varphi(p)} / \mathfrak{m}_{\varphi(p)}^{m+1} \longrightarrow \mathcal{O}_{X, p} / \mathfrak{m}_p^{m+1},$$

where $\mathcal{O}_{X, x}$ denotes the local ring at a point $x \in X$ and \mathfrak{m}_x its maximal ideal.

We assume in the sequel that $p \in X_{\mathrm{reg}}$ is a regular point and $\varphi(p) = p$. Letting $A_m = \mathcal{O}_{X, p} / \mathfrak{m}_p^{m+1}$ the m -jet of φ yields a map of k -algebras

$$j^m \varphi = j_p^m \varphi : A_m \longrightarrow A_m,$$

which stabilizes the maximal ideal \mathfrak{m} of A_m and all of its powers \mathfrak{m}^k .

For $m \geq 1$ we let $\mathrm{Aut}_{m-1}(A_m)$ denote the set of \mathbb{k} -algebra isomorphisms $f : A_m \rightarrow A_m$ with $f \equiv \mathrm{id} \pmod{\mathfrak{m}^m}$. For every $f \in \mathrm{Aut}_{m-1}(A_m)$ the map $f - \mathrm{id}$ sends A_m into \mathfrak{m}^m and vanishes on the constants \mathbb{k} . As it vanishes as well on \mathfrak{m}^2 it induces a \mathbb{k} -linear map

$$\psi_f : \mathfrak{m} / \mathfrak{m}^2 \longrightarrow \mathfrak{m}^m = \mathfrak{m}_p^m / \mathfrak{m}_p^{m+1}.$$

Consider the \mathbb{k} -vector space $V = \mathfrak{m}/\mathfrak{m}^2$ so that \mathfrak{m}^m is the m th symmetric power $S^m V$. For every $m \geq 1$ our construction yields a map

$$(11) \quad \psi : \text{Aut}_{m-1}(A_m) \longrightarrow \text{Hom}_{\mathbb{k}}(\mathfrak{m}/\mathfrak{m}^2, \mathfrak{m}^m) \cong V^\vee \otimes S^m V,$$

where V^\vee stands for the dual module of V . For $m = 1$ this map associates to $f = j^1\varphi$ just the cotangent map $d\varphi(0)^\vee$.

In terms of local coordinates this construction can be interpreted as follows. The \mathbb{k} -algebra A_m is isomorphic to the quotient A/\mathfrak{m}_A^{m+1} of the formal power series ring $A = \mathbb{k}[[x_1, \dots, x_n]]$. Any map $f \in \text{Aut}_{m-1}(A_m)$ is represented by an m -jet of an n -tuple of power series $F = (F_1, \dots, F_n) \in A^n$ with $F_i \equiv x_i \pmod{\mathfrak{m}_A^m}$. Clearly for any $m \geq 1$ the m -form ψ_f corresponds to the m th order term of F .

With this notation we have the following lemma.

Lemma 4.12. (a) For every $m \geq 1$ the map ψ in (11) is bijective.

(b) If $m = 1$ then $\psi_{f \circ g} = \psi_f \circ \psi_g$ while for $m \geq 2$ we have $\psi_{f \circ g} = \psi_f + \psi_g$.

(c) If ∂ is a locally nilpotent vector field on X with a zero of order $m \geq 2$ at p then

$$\psi_{\exp(t\partial)} = t\psi_{\exp(\partial)}.$$

Proof. (a) is immediate using the coordinate description above.

(b) is easy and can be left to the reader. To deduce (c) we note that $\exp(t\partial) \in \text{SAut}(X)$ induces the map $\text{id} + t\hat{\partial} \in \text{Aut}_{m-1}(A_m)$, where $\hat{\partial}$ denotes the derivation on A_m induced by ∂ . Hence $\psi_{\exp(t\partial)} = t\hat{\partial}$, proving (c). \square

An n -tuple $F = (F_1, \dots, F_n) \in A^n$ as in 4.11 representing an m -jet $f = j^m F \in \text{Aut}_{m-1}(A_m)$ preserves a volume form ω on X_{reg} (or on (X, p)) if and only if the Jacobian determinant J_F of F is equal to 1. Modulo \mathfrak{m}^m this determinant depends only on f and not on the representative F of f . Hence we can set $J_f := J_F \pmod{\mathfrak{m}^m}$.¹⁸ We say in the sequel that an m -jet $f \in \text{Aut}_{m-1}(A_m)$ with $m \geq 1$ preserves a volume form if $J_f \equiv 1 \pmod{\mathfrak{m}^m}$. The latter condition can be detected in terms of ψ_f as follows.

Lemma 4.13. (a) If $m = 1$ then $f \in \text{Aut}(A_1)$ preserves a volume form if and only if $\psi_f \in \text{SL}(V)$.

In case $m \geq 2$ the map $f \in \text{Aut}_{m-1}(A_m)$ preserves a volume form if and only if ψ_f is in the kernel of the natural contraction map

$$\begin{aligned} \kappa_m : \text{Hom}_{\mathbb{k}}(V, S^m V) \cong V^\vee \otimes S^m V &\longrightarrow S^{m-1} V, \\ \lambda \otimes v_1 \cdots v_m &\longmapsto \sum_{\mu=1}^m \lambda(v_\mu) \cdot v_1 \cdots \hat{v}_\mu \cdots v_m. \end{aligned}$$

(b) $\ker \kappa_m$ is an irreducible $\text{SL}_n(V)$ -module for all $m \geq 1$.

Proof. In case $m = 1$ (a) is immediate. Suppose that $m \geq 2$. If $f = \text{id} + f_m \pmod{\mathfrak{m}^{m+1}}$ with an n -tuple of m -forms $f_m = (f_{m1}, \dots, f_{mn})$, then J_f is easily seen to be equal to

$$1 + \text{div } f_m = 1 + \frac{\partial f_{m1}}{\partial x_1} + \dots + \frac{\partial f_{mn}}{\partial x_n} \pmod{\mathfrak{m}^m},$$

¹⁸However J_f is not an element in A_m since it is not well defined modulo \mathfrak{m}^{m+1} .

where $\operatorname{div} f_m$ is the divergence of f_m . Thus $J_f \equiv 1 \pmod{\mathfrak{m}^m}$ if and only if $\operatorname{div} f_m = 0$. Writing $f_m \in V^\vee \otimes S^m V$ as $f_m = \sum_{i=1}^n \frac{\partial}{\partial x_i} \otimes f_{mi}$ the element $\operatorname{div} f_m$ in $S^{m-1} V$ corresponds just to the contraction $\kappa_m(f_m)$, proving (a).

(b) is a standard fact in representation theory, see e.g. [34, §IX.10.2]. \square

Now we can state our main result in this subsection.

Theorem 4.14. *Let X be an affine algebraic variety of dimension $n \geq 2$ equipped with an algebraic volume form ω defined on X_{reg} , and let $G \subseteq \operatorname{SAut}(X)$ be a subgroup generated by a saturated set \mathcal{N} of locally nilpotent derivations. If G acts on X with an open orbit O , then for every $m \geq 0$ and every finite subset $Z \subseteq O$ there exists an automorphism $g \in G$ with prescribed m -jets j_p^m at the points $p \in Z$, provided these jets preserve ω and inject Z into O .*

The proof will be reduced to the following lemma.

Lemma 4.15. *With the notation and assumptions of Theorem 4.2, suppose that j_p^m is an m -jet of an automorphism at a given point $p \in Z$, which is the identity up to order $m - 1 \geq 0$. Then for every $M > 0$ there is an automorphism $g \in G$ such that its m -jet at p is j_p^m while its M -jet at each other point $q \neq p$ of Z is the identity.*

Before proving Lemma 4.15 let us show how Theorem 4.5 follows.

Proof of Theorem 4.14. We proceed by induction on m . If $m = 0$ the assertion follows from the fact that G acts infinitely transitively on O . For the induction step suppose that we have an automorphism $g \in G$ with the prescribed jets up to order $m - 1 \geq 0$. Thus the m -jets $j_p^m = j_p^m \circ g^{-1}$ are up to order $m - 1$ the identity at every point $p \in Z$. If we find an automorphism $h \in G$ with m -jet equal to j_p^m for all $p \in Z$, then obviously the automorphism $h \circ g$ has the desired properties.

Thus replacing j_p^m by j_p^m we are reduced to show the assertion in the case that for all $p \in Z$ the m -jets j_p^m are the identity up to order $m - 1$, where $m \geq 1$.

Applying Lemma 4.15, for every point $p \in Z$ there is an automorphism $g_p \in G$ whose m -jet at p is the given one while its m -jets at all other points $q \in Z \setminus \{p\}$ are the identity. Obviously then the composition (in arbitrary order) $g = \prod_{p \in Z} g_p$ will have the required properties. \square

Proof of Lemma 4.15. In the case $m = 1$ the assertion follows from Theorem 4.2 and Proposition 4.8. So we may assume for the rest of the proof that $m \geq 2$.

Consider the set $\mathcal{N}_{mp,Z}^M$ of all locally nilpotent derivations in \mathcal{N} with a zero of order m at p and of order $M + 1$ at all other points $q \in Z \setminus \{p\}$. Let $G_{mp,Z}^M$ be the subgroup of G generated by the exponentials of elements in $\mathcal{N}_{mp,Z}^M$ so that an automorphism in $G_{mp,Z}^M$ is the identity up to order $(m - 1)$ at p and up to order M at all other points $q \in Z \setminus \{p\}$. With the notation as introduced in 4.11 let us consider the composed map

$$\Psi : G_{mp,Z}^M \longrightarrow \operatorname{Aut}_{m-1}(A_m) \xrightarrow{\psi} \operatorname{Hom}_{\mathbb{k}}(V, S^m V),$$

where ψ is as in (11) and the first arrow assigns to an automorphism its m -jet at p . Using Lemma 4.13(a) it suffices to show that Ψ maps $G_{mp,Z}^M$ surjectively onto the subspace $\ker \kappa_m$.

The group $G_{mp,Z}^M$ is generated by exponentials of vector fields in $\mathcal{N}_{mp,Z}^M$. Thus using Lemma 4.12(b), (c) the image $\text{im}(\Psi)$ of Ψ is a linear subspace of $\text{Hom}_{\mathbb{k}}(V, S^m V)$. We claim that this subspace is nonzero.

Indeed, consider a vector field $\partial \in \mathcal{N}$ with $\partial(p) \neq 0$ and the one-parameter subgroup $H = \exp(\mathbb{k}\partial)$. By the Rosenlicht theorem (see Remark 2.7(1)) on an open dense H -invariant subset $U(H) \subseteq O$ the group H admits a geometric quotient $U(H)/H$ defined by a finite collection of H -invariant regular functions on X . By infinite transitivity of the action of G on O we may assume that $Z \subseteq U(H)$ is such that the image of Z in the quotient $U(H)/H$ is contained in the regular part of $U(H)/H$, has the same cardinality as Z , and the projection $U(H) \rightarrow U(H)/H$ is smooth in the points of Z . Thus we can find a regular H -invariant function f on X with a simple zero at p , and another such function h with $h(p) = 1$ and $h(q) = 0$ for all $q \in Z \setminus \{p\}$. Replacing f by $h^{M+1}f$ we may assume that f has a zero of order $\geq M+1$ at all points of $Z \setminus \{p\}$ and a simple zero at p . Then $g = \exp(f^m \partial)$ is an automorphism in $G_{mp,Z}^M$ with $\Psi(g) = f^m \hat{\partial} \neq 0$, where $\hat{\partial}$ is the derivation of A_m induced by ∂ (cf. Lemma 4.12(c) and its proof). This proves the claim.

The group $G_{p,Z}^M$ acts on $G_{mp,Z}^M$ by conjugation $g.h = g \circ h \circ g^{-1}$, where $g \in G_{p,Z}^M$ and $h \in G_{mp,Z}^M$. If we write $h = \text{id} + h_m \pmod{\mathfrak{m}^{m+1}}$ with a map $h_m \in \text{Hom}_{\mathbb{k}}(V, S^m V)$ then $g.h = \text{id} + g \circ h_m \circ g^{-1} \pmod{\mathfrak{m}^{m+1}}$. The map g induces the cotangent map $(d_p g)^\vee$ on $V = (T_p X)^\vee$ and its m th symmetric power $S^m((d_p g)^\vee)$ on $S^m V$. Hence there is a commutative diagram

$$\begin{array}{ccc} G_{p,Z}^M \times G_{mp,Z}^M & \longrightarrow & G_{mp,Z}^M \\ d_p^\vee - \times \Psi \downarrow & & \downarrow \Psi \\ \text{SL}(V) \times \text{Hom}_{\mathbb{k}}(V, S^m V) & \longrightarrow & \text{Hom}_{\mathbb{k}}(V, S^m V), \end{array}$$

where the lower horizontal map is induced by the standard representation of $\text{SL}(V)$ on $S^m V$. Since the map $G_{p,Z}^M \rightarrow \text{SL}(V)$ is surjective (see Theorem 4.2 and Remark 4.7), the image $\text{im}(\Psi)$ of Ψ is a non-trivial $\text{SL}(V)$ -module. By Lemma 4.12 this representation is contained in the kernel of the contraction map κ_m . Since the latter kernel is irreducible (see Lemma 4.13(b)), it follows that $\text{im}(\Psi) = \ker \kappa_m$, as required. \square

Remark 4.16. If in the situation of Theorem 4.14 each of the jets j_p^m , $p \in Z$, fixes the point p and preserves a volume form,¹⁹ then the conclusion of Theorem 4.14 remains valid without the requirement that there is a global volume form on X_{reg} .

Remark 4.17. If X_{reg} does not admit a global volume form i.e., $K_{X_{\text{reg}}} \neq 0$, one can still formulate a necessary condition for interpolation of jets by an automorphism from a special group G , namely in terms of the ‘volume form monodromy’ of G . To define it we fix a volume form ω_x on the tangent space $T_x X$ at some point $x \in X_{\text{reg}}$, and consider the stabilizer subgroup $G_x \subseteq G$. Every element $g \in G_x$ transforms ω_x into $\chi_x(g) \cdot \omega_x$, where $\chi_x(g) \in \mathbb{G}_m = \mathbb{G}_m(\mathbb{k})$. The map

$$\chi_x : G_x \longrightarrow \mathbb{G}_m$$

¹⁹Note that this is a purely local condition, see the discussion before Lemma 4.13.

is then a character on G_x which equals 1 on $G_{N,x}$, see Theorem 4.2. If $y \in X$ is a second point and $h \in G$ is an automorphism with $h.x = y$ then $hG_xh^{-1} = G_y$ and $\chi_y(hgh^{-1}) = \chi_x(g)$ for all $g \in G_x$. In particular the image of χ_x forms a subgroup Γ of \mathbb{G}_m independent of $x \in O$, which is called the *volume form monodromy* of G .

The volume form monodromy can be a nontrivial discrete group as in the case of $X = \mathrm{SL}_2/N(\mathbb{T})$ and $G = \mathrm{SAut}(X)$, where $N(\mathbb{T}) \subseteq \mathrm{SL}_2$ is the normalizer of the maximal torus $\mathbb{T} \subseteq \mathrm{SL}_2$. Thus $N(\mathbb{T})$ is an extension of \mathbb{G}_m by $\mathbb{Z}/2\mathbb{Z}$ and $X = \mathbb{P}^2 \setminus C$, where $C \subseteq \mathbb{P}^2$ is a smooth conic; cf. [29, II]. Using technique from [20] one can show that here $\Gamma = \{\pm 1\}$.

4.3. \mathbb{A}^1 -richness. We remind the reader that an affine variety X is called \mathbb{A}^1 -rich if for every finite subset Z and every algebraic subset Y of codimension ≥ 2 there is a polynomial curve in X passing through Z and not meeting Y [23].

The following corollary is immediate from the Transversality Theorem 1.15. In the special case where $X = \mathbb{A}_{\mathbb{C}}^n$ this corollary yields the Gromov-Winkelmann theorem, see [40].

Corollary 4.18. *Let as before X be an affine variety and let $G \subseteq \mathrm{SAut}(X)$ be a subgroup generated by a saturated set \mathcal{N} of locally nilpotent derivations, which acts with an open orbit $O \subseteq X$. Then for any finite subset $Z \subseteq O$ and any closed subset $Y \subseteq X$ of codimension ≥ 2 with $Z \cap Y = \emptyset$ there is an orbit $C \cong \mathbb{A}^1$ of a \mathbb{G}_a -action on X which does not meet Y and passes through each point of Z having prescribed jets at these points.*

Proof. In the case $\dim X = 1$ this is trivially true. So assume that $\dim X \geq 2$. Let C be an orbit of a \mathbb{G}_a -action on O . Since G acts infinitely transitively on O we may assume that $Z \subseteq C$. By Theorem 4.14 and Remark 4.16, applying an appropriate automorphism $g' \in G$ we may suppose as well that C has prescribed m -jets at the points of Z . Indeed, the m -jets of automorphisms stabilizing a given point $p \in O$ and having at this point the jacobian determinant equal to 1 modulo \mathfrak{m}^m act transitively on the set of all m -jets of smooth curves at p .

By Proposition 4.9, using the notation as in 4.7, the special group G_Z^m acts transitively in $O \setminus Z$. Applying now the Transversality Theorem 1.15(b) to G_Z^m , $C \cap (O \setminus Z)$, and $Y \cap (O \setminus Z)$ we can find an element $g \in G_Z^m$ with $g.C \cap Y = \emptyset$. Thus the \mathbb{G}_a -orbit $g.C$ contains Z , has the prescribed jets at the points of Z , and does not meet Y . \square

We can deduce also the following fact.

Proposition 4.19. *Let $G \subseteq \mathrm{SAut}(X)$ be a subgroup generated by a saturated set \mathcal{N} of locally nilpotent derivations, which acts with an open orbit $O \subseteq X$. Then for any closed subset $Y \subseteq O$ of codimension ≥ 2 the group $G_{\mathcal{N},Y}$ acts with an open orbit.*

Proof. According to Proposition 1.7 there are locally nilpotent vector fields $\partial_1, \dots, \partial_s$ generating T_pX for all $p \in O$. Let $H_\sigma \subseteq G$ be the one-parameter subgroup associated to ∂_σ . By Rosenlicht's Theorem, for an open dense H_σ -invariant subsets $U(H_\sigma)$ in O there is a geometric quotient $U(H_\sigma)/H_\sigma$. Using the same reasoning as in the proof of Theorem 2.15 there is an H_σ -invariant function f_σ vanishing on $\overline{H_\sigma.Y}$ and equal to 1 at a given general point $p \in U(H_\sigma) \setminus \overline{H_\sigma.Y}$. Consequently $\exp(\mathbb{k}f_\sigma\partial_\sigma)$ stabilizes Y

and moves p in direction $\partial_\sigma(p)$. In other words, p is a $G_{\mathcal{N},Y}$ -flexible point. Applying Corollary 1.10(a) the result follows. \square

In contrast we have the following result.

Proposition 4.20. *Let G be a special subgroup of $\mathrm{SAut}(X)$ acting on X with an open orbit $O \subseteq X$. If the complement $X \setminus O$ contains a divisor D then $[D] \neq 0$ in $\mathrm{Cl}(X)_\mathbb{Q}$.*

Proof. Assume to the contrary that $[D] = 0$ in $\mathrm{Cl}(X)_\mathbb{Q}$. Then there is a function f on X with $D = \mathbb{V}(f)$ set theoretically. For every one dimensional unipotent subgroup H and $x \in O$ the function $f|_{H.x}$ is a polynomial on $H.x \cong \mathbb{k}$. As $H.x \subseteq O$ and so $D \cap H = \emptyset$, this polynomial has no zero and so is constant. Hence $H.x$ is contained in a level set of f . Since G is generated by such subgroups, the orbit O of G is contained in a level set of f and so it cannot be open, a contradiction. \square

Corollary 4.21. *Let G be a special subgroup of $\mathrm{SAut}(X)$. If X is \mathbb{Q} -factorial and a closed subset $Y \subseteq X$ contains a divisor, then the group $G_{\mathcal{N},Y}$ has no open orbit.*

Problems 4.22. 1. Assume as before that a group $G = \langle \mathcal{N} \rangle \subseteq \mathrm{SAut}(X)$, where $\mathcal{N} \subseteq \mathrm{LND}(G)$ is saturated, acts on a normal (or even smooth) affine variety X with an open orbit O . Is it true that the complement $Y = X \setminus O$ has codimension ≥ 2 in X ? This is true if $\dim X = 2$ by Gizatullin's Theorem [15]. For non-normal varieties there are counterexamples (see Example 5.10 below).

2. We do not know whether in the situation of Proposition 4.19 the group $G_{\mathcal{N},Y}$ acts transitively on $O \setminus Y$.

Remark 4.23. Every algebraic variety X contains a divisor Y such that the logarithmic Kodaira dimension $\bar{\kappa}(X \setminus Y)$ is ≥ 0 . In this case $X \setminus Y$ cannot carry a \mathbb{G}_a -action and so $G_Y = \{\mathrm{id}\}$ although X might be flexible. The simplest example of such a situation is given by the hypersurface $Y = \{X_1 \cdot \dots \cdot X_n = 0\}$ in $X = \mathbb{A}^n$, see also [18].

5. SOME APPLICATIONS

5.1. Unirationality, flexibility, and triviality of the Makar-Limanov invariant. Recall [12] that the *Makar-Limanov invariant* $\mathrm{ML}(X)$ of an affine variety X is the intersection of the kernels of all locally nilpotent derivations on X . In other words $\mathrm{ML}(X)$ is the subalgebra of the algebra $\mathcal{O}(X)$ consisting of all regular $\mathrm{SAut}(X)$ -invariants. Similarly [28] the *field Makar-Limanov invariant* $\mathrm{FML}(X)$ is defined as the subfield of $\mathbb{k}(X)$ of all rational $\mathrm{SAut}(X)$ -invariants. If it is trivial i.e., if $\mathrm{FML}(X) = \mathbb{k}$ then so is $\mathrm{ML}(X)$, while the converse is not true in general, see Example 5.3(1) below. The next proposition confirms, in particular, Conjecture 5.3 in [27].

Proposition 5.1. *An affine variety X possesses a flexible point if and only if the group $\mathrm{SAut}(X)$ acts on X with an open orbit, if and only if the field Makar-Limanov invariant $\mathrm{FML}(X)$ is trivial. In the latter case X is unirational.*

Proof. The first equivalence follows from Corollary 1.10(a) and the second one from Corollary 1.13. As for the last assertion, see Remark 5.2 below. \square

Remark 5.2. As follows from Proposition 1.2(b) for every G -orbit O of a special group $G \subseteq \mathrm{SAut}(X)$ there is a surjective map $\mathbb{A}^s \rightarrow O$. Hence any two points in O are

contained in the image of a morphism $\mathbb{A}^1 \rightarrow O$. In particular O is \mathbb{A}^1 -connected in the sense of [21, §6.2].

Let us mention some known counterexamples related to the problem of rationality of flexible varieties.

Examples 5.3. 1. Due to A. Liendo [27, §4.2] there are examples of non-unirational affine threefolds X such that $\text{ML}(X) = \mathbb{k}$. In these examples the variety X is birationally equivalent to $C \times \mathbb{A}^2$, where C is a curve of genus $g \geq 1$, and the general G -orbits are of dimension two. In particular, the invariant $\text{FML}(X)$ is non-trivial and there is no flexible point in X .

2. In turn, flexibility implies neither rationality nor stable rationality. Indeed, there exists a finite subgroup $F \subseteq \text{SL}(n, \mathbb{C})$, where $n \geq 4$, such that the smooth unirational affine variety $X = \text{SL}(n, \mathbb{C})/F$ is not stably rational, see [32, Example 1.22]. However, by Proposition 5.4 below X is flexible and the group $\text{SAut}(X)$ acts infinitely transitively on X .

5.2. Flexible quasihomogeneous varieties. An important class of flexible algebraic varieties consists of homogeneous spaces of semisimple algebraic groups. More generally, the following hold.

Proposition 5.4. *Let G be a connected affine algebraic group over \mathbb{k} without non-trivial characters²⁰, and let H be a closed subgroup of G . Then the homogeneous space G/H is flexible. In particular, if G/H is affine of dimension $n \geq 2$ then the group $\text{SAut}(G/H)$ acts infinitely transitively on G/H .*

Proof. The image of G is contained in $\text{SAut}(G/H)$. Thus the group $\text{SAut}(G/H)$ acts on the quotient G/H transitively and G/H is flexible; see Proposition 1.1 in [1]. The second assertion follows from the first one in view of Theorem 0.1 and Corollary 1.21. \square

The following problem arises.

Problem 5.5. *Characterize flexible varieties among affine varieties admitting an action of a semisimple algebraic group with an open orbit.*

For instance, if such a quasihomogeneous variety is smooth then in fact it is flexible. In the particular case $G = \text{SL}_2$ this was actually established in [29, III], where we borrowed the idea of the proof of the following theorem.

Theorem 5.6. *Suppose that a semisimple algebraic group G acts on a smooth affine variety $X = \text{Spec } A$ with an open orbit. Then X is homogeneous with respect to a connected affine algebraic group $\tilde{G} \supseteq G$ without non-trivial characters. In particular, X is flexible.*

Proof. Since by our assumption $A^G = \mathbb{k}$, due to Matsushima's Criterion and Luna's Étale Slice Theorem (see Theorems 4.17 and 6.7 in [33]), there is a unique closed G -orbit $O \subseteq X$, the stabilizer $H = \text{Stab}_G(x)$ of any point $x \in O$ is reductive, and

²⁰E.g., a semisimple algebraic group.

there is a linear representation W of H and a G -equivariant isomorphism with a (left) G -homogeneous fiber bundle

$$X \cong (G \times W)/H,$$

where H acts on $G \times W$ via

$$(12) \quad h.(g, w) = (gh^{-1}, h.w).$$

Comparing normal bundles we have necessarily $W = T_x X/T_x O$. According to [3] there exists a finite dimensional G -module V such that $V = W \oplus W'$, where $W' \subseteq V$ is a complementary H -submodule. Letting

$$\tilde{G} = G \rtimes V \quad \text{with} \quad (g_1, v_1).(g_2, v_2) = (g_1 g_2, g_2^{-1} v_1 + v_2),$$

$\tilde{H} = H \rtimes W'$, and $\tilde{H}_0 = \{e\} \rtimes W'$, we can identify \tilde{G}/\tilde{H}_0 and $G \times W$ as H -varieties, where H acts on $G \times W$ via (12). Since the subgroup $H \subseteq G$ normalizes \tilde{H}_0 in \tilde{G} it acts \tilde{G} -equivariantly on the right on \tilde{G}/\tilde{H}_0 . The latter fact can be used to deduce the isomorphisms of abstract varieties

$$\tilde{G}/\tilde{H} \cong (\tilde{G}/\tilde{H}_0)/H \cong (G \times W)/H \cong X.$$

By Proposition 5.4 X is flexible being a homogeneous variety of a connected affine algebraic group \tilde{G} without non-trivial characters (indeed, $\tilde{G} = (G, 0) \cdot (e, V)$, where both groups do not admit non-trivial characters). Now the proof is completed. \square

In the next theorem we provide a complete solution of Problem 5.5 for $G = \mathrm{SL}_2 := \mathrm{SL}_2(\mathbb{k})$ and X normal.

Theorem 5.7. *Every normal affine variety E admitting an SL_2 -action with an open orbit is flexible.*

For a homogeneous affine variety $E = \mathrm{SL}_2/H$ the result follows from Proposition 5.4. The proof in the general case given below is based on a description of normal SL_2 -varieties due to Popov [29, I] (see also [26, Chapter III, §4]) and a Cox ring SL_2 -construction due to Batyrev and Haddad [2]. Recall [29, I] that every non-homogeneous affine SL_2 -threefold with an open orbit is uniquely determined by a pair (h, m) , where m is the order of the generic isotropy group²¹ and $h = p/q \in (0, 1] \cap \mathbb{Q}$ is the height²² of the algebra of U -invariants, where $U \subseteq \mathrm{SL}_2$ is a maximal unipotent subgroup. Such an SL_2 -threefold with an invariant (h, m) is denoted by $E_{h,m}$. Notice that $E_{h,m}$ is smooth for $h = 1$ and singular for $h < 1$.

Assuming in the sequel that p and q are coprime positive integers we let

$$(13) \quad a = m/k \quad \text{and} \quad b = (q - p)/k, \quad \text{where} \quad k = \mathrm{gcd}(q - p, m).$$

Let $\mu_a = \langle \xi_a \rangle$ denote the cyclic group generated by a primitive root of unity $\xi_a \in \mathbb{G}_m = \mathbb{G}_m(\mathbb{k})$ of degree a . The SL_2 -variety $E_{h,m}$ is isomorphic to the categorical quotient of the hypersurface $D_b \subseteq \mathbb{A}^5$ with equation

$$(14) \quad Y^b = X_1 X_4 - X_2 X_3$$

²¹Which is a cyclic group.

²²Or rather a slope.

modulo the diagonal action of the group $\mathbb{G}_m \times \mu_a$ on $\mathbb{A}^5 = \text{Spec } \mathbb{k}[X_1, X_2, X_3, X_4, Y]$ via

$$\text{diag}(t^{-p}, t^{-p}, t^q, t^q, t^k) \times \text{diag}(\xi^{-1}, \xi^{-1}, \xi, \xi, 1), \quad t \in \mathbb{G}_m, \xi \in \mu_a.$$

Here the SL_2 -action on D_b is induced by the trivial action on the coordinate Y , while $\langle X_1, X_2 \rangle$ and $\langle X_3, X_4 \rangle$ are simple SL_2 -modules. This SL_2 -action on D_b commutes with the $(\mathbb{G}_m \times \mu_a)$ -action and so descends to the quotient. This gives a simple and uniform description of all non-homogeneous normal affine SL_2 -threefolds with an open orbit $E_{h,m}$ via the Cox realization as the quotient of the spectrum of the corresponding Cox ring by the action of the Neron-Severi quasitorus, see [2].

Proof of Theorem 5.7. Let E be a non-homogeneous SL_2 -variety with an open orbit. If $\dim E = 2$ then E is a toric surface, in fact a Veronese cone, and the group SL_2 is transitive off the vertex (see [29, II] or, alternatively, Theorem 0.2 in [1]). So the assertion follows.

Let further E be a non-homogeneous normal affine SL_2 -threefold. According to Popov's classification $E = E_{h,m}$ for some pair (h, m) .

In the case where $E = E_{h,m}$ is smooth that is, $h = 1$ the result follows from Theorem 5.6.

In the case where $E = E_{h,m}$ is singular i.e., $h = p/q < 1$, there is a unique singular point, say, $Q \in E$. The complement $E \setminus \{Q\}$ consists of two SL_2 -orbits O_1 and O_2 , where $O_1 \cong \text{SL}_2 / \mu_m$ while $O_2 \cong \text{SL}_2 / U_{a(p+q)}$ has the isotropy subgroup

$$U_{a(p+q)} = \left\{ \begin{pmatrix} \xi & \eta \\ 0 & \xi^{-1} \end{pmatrix} \mid \eta \in \mathbb{k}, \xi^{a(p+q)} = 1 \right\}.$$

Consider the hypersurface $D_b \subseteq \mathbb{A}^5$ as in (14). We can realize \mathbb{A}^5 as a matrix space:

$$\mathbb{A}^5 = \left\{ (X, Y) \mid X = \begin{pmatrix} X_1 & X_3 \\ X_2 & X_4 \end{pmatrix}, X_i, Y \in \mathbb{A}^1 \right\}.$$

Then according to [2] the 3-fold $E = E_{h,m}$ admits a realization as the categorical quotient of D_b by the action of the group $\mathbb{G}_m \times \mu_a$ via

$$(t, \xi).(X, Y) = \left(\begin{pmatrix} \xi^{-1}t^{-p}X_1 & \xi t^q X_3 \\ \xi^{-1}t^{-p}X_2 & \xi t^q X_4 \end{pmatrix}, t^k Y \right).$$

This action commutes with the natural SL_2 -action on D_b given by

$$A.(X, Y) = (AX, Y).$$

Hence the SL_2 -action on D_b descends to the quotient $E = E_{h,m}$. The hypersurface $Z = \{Y = 0\}$ in D_b is the inverse image of the unique two dimensional SL_2 -orbit closure in $E_{h,m}$. To show the transitivity (or the flexibility) of the group $\text{SAut}(X)$ in E_{reg} it suffices to find a locally nilpotent derivation ∂ of the algebra $\mathcal{O}(H_b)$ with $\partial(Y) \neq 0$ which preserves the $(\mathbb{Z} \times \mathbb{Z}_a)$ -bigrading on $\mathcal{O}(H_b)$ defined via

$$\deg X_1 = \deg X_2 = (-p, -\bar{1}), \quad \deg X_3 = \deg X_4 = (q, \bar{1}), \quad \text{and} \quad \deg Y = (k, \bar{0}).$$

Indeed, such a derivation induces a locally nilpotent derivation on $\mathcal{O}(E)$. Since $\partial(Y) \neq 0$ the restriction of the corresponding vector field to the image \bar{Z} of Z in E is nonzero and so the points of \bar{Z} with $\partial \neq 0$ are flexible. By transitivity, every point of $\bar{Z} \setminus \{Q\}$ is.

The variety D_b can be regarded as a suspension²³ over $\mathbb{A}^3 = \text{Spec } \mathbb{k}[X_2, X_3, Y]$. Namely,

$$D_b = \{X_1 X_4 = f(X_2, X_3, Y)\} \quad \text{where} \quad f = X_2 X_3 + Y^b.$$

According to [1] (see also Lemma 3.3 in [22, §5]) a desired bihomogeneous locally nilpotent derivation ∂ can be produced starting with a locally nilpotent derivation $\delta \in \text{Der } \mathbb{k}[X_2, X_3, Y]$. For instance, let δ be given by

$$\delta(X_2) = \delta(X_3) = 0, \quad \delta(Y) = X_2^c X_3^d.$$

Then ∂ can be defined via

$$(15) \quad \partial(X_1) = \partial(X_2) = \partial(X_3) = 0, \quad \partial(X_4) = \delta(f) = bX_2^c X_3^d Y^{b-1}, \quad \partial(Y) = X_1 X_2^c X_3^d$$

with a, b as in (13) and with appropriate values of the natural parameters c, d . Such a derivation ∂ preserves the $(\mathbb{Z} \times \mathbb{Z}_a)$ -bigrading²⁴ if and only if

$$\begin{aligned} -p - cp + dq &= k \\ k(b-1) - cp + dq &= q \\ -1 - c + d &\equiv 0 \pmod{a} \\ -c + d &\equiv 1 \pmod{a}. \end{aligned}$$

By virtue of (13) the second relation follows from the first one, while the last two are equivalent. Letting $c = s - 1$ we can rewrite the remaining relations as

$$(16) \quad \begin{aligned} dq - sp &= k \\ s &\equiv d \pmod{a}. \end{aligned}$$

Since $\gcd(p, q) = 1$ the first equation admits a solution (d_0, s_0) in natural numbers. For every $r \in \mathbb{N}$, the pair $(d_0 + rp, s_0 + rq)$ also represents such a solution. The second relation in (16) becomes

$$(17) \quad r(q - p) \equiv d_0 - s_0 \pmod{a}.$$

By (13) $k = \gcd(m, q - p)$, hence $\gcd(k, p) = 1$. The first equation in (16) written as

$$d_0(q - p) - p(s_0 - d_0) = k$$

implies that $k \mid (s_0 - d_0)$.

Let $l = \gcd(a, q - p) = \gcd(a, bk)$. Since $\gcd(a, b) = 1$ then $l \mid k$ and so (17) is equivalent to the congruence

$$r \cdot \frac{q - p}{l} \equiv \frac{d_0 - s_0}{l} \pmod{\frac{a}{l}}.$$

Since $\frac{q-p}{l}$ and $\frac{a}{l}$ are coprime the latter congruence admits a solution, say, r_0 . Letting finally

$$c = s_0 + r_0 q - 1, \quad d = d_0 + r_0 p$$

the locally nilpotent derivation ∂ as in (15) becomes homogeneous of bidegree $(0, \bar{0})$, as needed. Now the proof is completed. \square

²³See the definition of a suspension in the Introduction.

²⁴I.e. $\deg \partial(Y) = \deg Y$ and $\deg \partial(X_4) = \deg X_4$.

The question arises whether the smooth loci of singular affine SL_2 -threefolds are homogeneous as well, cf. Theorem 5.6. The answer is negative; the following proposition gives a more precise information.

Proposition 5.8. *Let $E = E_{h,m}$, where $h = p/q < 1$ with $\gcd(p, q) = 1$. The following conditions are equivalent:*

- (i) *The SL_2 -action on E extends to an action of a bigger affine algebraic group G on E which is transitive in E_{reg} ;*
- (ii) *The variety E is toric;*
- (iii) *$(q - p) \mid m$ or, equivalently, $b = 1$ in (13).*

Proof. Implication (i) \Rightarrow (ii) follows from Theorem 1 in [29, III]. According to this theorem, a normal affine threefold X with a unique singular point Q which admits an action of an affine algebraic group transitive on $X \setminus \{Q\}$, is toric.

The equivalence (ii) \Leftrightarrow (iii) follows from the results of [2] and [14]. Let us show the remaining implication (iii) \Rightarrow (i). If $b = 1$ in (13) then $D_b \cong \mathbb{A}^4 = \mathrm{Spec} \mathbb{k}[X_1, \dots, X_4]$. Hence the toric variety $E_{h,m}$ can be obtained as the quotient $\mathbb{A}^4 / (G_m \times \mu_a)$, where the group $G_m \times \mu_a$ with $a = m/(q - p)$ as in (13) acts diagonally on \mathbb{A}^4 via

$$(18) \quad (X_1, X_2, X_3, X_4) \longmapsto (\xi^{-1}t^{-p}X_1, \xi^{-1}t^{-p}X_2, \xi t^q X_3, \xi t^q X_4), \quad (t, \xi) \in G_m \times \mu_a.$$

Consider the action of the group $\mathrm{SL}_2 \times \mathrm{SL}_2$ on \mathbb{A}^4 via

$$(A_1, A_2).(X_1, X_2, X_3, X_4) = \left(A_1 \begin{pmatrix} X_1 \\ X_2 \end{pmatrix}, A_2 \begin{pmatrix} X_3 \\ X_4 \end{pmatrix} \right).$$

This action commutes with the $(G_m \times \mu_a)$ -action (18) and so descends to the quotient $E_{h,m}$. The induced $(\mathrm{SL}_2 \times \mathrm{SL}_2)$ -action on the quotient $E_{h,m}$ is transitive in the complement of the unique singular point Q . This yields (i). Now the proof is completed. \square

Corollary 5.9. *None of the non-toric affine threefolds $E = E_{h,m}$ with $h < 1$ admits an algebraic group action transitive in E_{reg} . However, the group $\mathrm{SAut}(E)$ acts infinitely transitively in E_{reg} .*

Let us finish this subsection with an example of a flexible non-normal affine variety with singular locus of codimension one.

Example 5.10. Consider the standard irreducible representation of the group SL_2 on the space of binary forms of degree three

$$V = \langle X^3, X^2Y, XY^2, Y^3 \rangle.$$

Restriction to the subvariety

$$E = \mathrm{SL}_2.X^2Y \cup \mathrm{SL}_2.X^3 \cup \{0\} \subseteq V$$

of forms with zero discriminant yields a non-normal SL_2 -embedding, see [26]. Since for a hypersurface in a smooth variety normality is equivalent to smoothness in codimension one, the divisor $D = \mathrm{SL}_2.X^3 \cup \{0\} \subseteq E$ coincides with the singular locus E_{sing} . The complement $E_{\mathrm{reg}} = \mathrm{SL}_2.X^2Y$ is the open SL_2 -orbit consisting of all flexible points of E . Hence E is flexible.

Observe that the normalization of E is isomorphic to $E_{\frac{1}{2},1}$. Indeed $m = 1$ because the stabilizer in SL_2 of a general point in E is trivial. On the other hand, the order of the stabilizer of the two dimensional orbit equals $p + q = 3$, hence $p = 1$ and $q = 2$.

6. APPENDIX: HOLOMORPHIC FLEXIBILITY

In this appendix we extend the notion of a flexible affine variety to the complex analytic setting (cf. [10]). We survey relations between holomorphic flexibility, Gromov's spray and the Andersen-Lempert theory. In particular, we show that every flexible variety admits a Gromov spray. This provides a new wide class of examples to which the Oka-Grauert-Gromov Principle can be applied. We refer the reader to the survey articles [11, §3] and [19] for a more thorough treatment and historical references.

6.1. Oka-Grauert-Gromov Principle for flexible varieties. The following notions were introduced in [16, §1.1.B].

Definition 6.1. (i) Let X be a complex manifold. A *dominating spray* on X is a holomorphic vector bundle $\rho : E \rightarrow X$ together with a holomorphic map $s : E \rightarrow X$, such that s restricts to the identity on the zero section Z while for each $x \in Z \cong X$ the tangent map $d_x s$ sends the fiber $E_x = \rho^{-1}(x)$ (viewed as a linear subspace of $T_x E$) surjectively onto $T_x X$.

(ii) Let $h : X \rightarrow B$ be a surjective submersion of complex manifolds. We say that it admits a *fiber dominating spray* if there is a holomorphic vector bundle E on X together with a holomorphic map $s : E \rightarrow X$ such that their restriction to each fiber $h^{-1}(b)$, $b \in B$, yields a spray on this fiber.

In these terms, the Oka-Grauert-Gromov Principle can be stated as follows.

Theorem 6.2. ([16, §4.5]) *Let $h : X \rightarrow B$ be a surjective submersion of Stein manifolds. If it admits a fiber dominating spray then the following hold.*

- (a) *Any continuous section of h is homotopic to a holomorphic one; and*
- (b) *any two holomorphic sections of h that are homotopic via continuous sections are also homotopic via holomorphic ones.*

Due to the following proposition, smooth affine algebraic G -fibrations with flexible fibers are appropriate for applying this principle (cf. [11, 3.4], [16]).

Proposition 6.3. (a) *Every flexible smooth affine algebraic variety X over \mathbb{C} admits a dominating spray.*

- (b) *Let $h : X \rightarrow B$ be a surjective submersion of smooth affine algebraic varieties over \mathbb{C} such that for some algebraically generated subgroup $G \subseteq \mathrm{Aut}(X)$ the orbits of G coincide with the fibers of h ²⁵. Then $X \rightarrow B$ admits a fiber dominating spray.*

Proof. It suffices to show (b). Indeed, due to Corollary 1.21, (a) is a particular case of (b).

By Proposition 1.7 there is a sequence of algebraic subgroups $\mathcal{H} = (H_1, \dots, H_s)$ of G such that the tangent space to the orbit $G.x$ at each point $x \in X$ is spanned by the tangent spaces at x to the orbits $H_i.x$, $i = 1, \dots, s$. Let $\exp : T_1(H_i) \rightarrow H_i$ be the

²⁵We say in this case that X is G -flexible over B .

exponential map. Letting $E = X \times \prod_{i=1}^s T_1(H_i)$ be the trivial vector bundle over X we consider the morphism

$$s : E \rightarrow X, \quad (x, (h_1, \dots, h_s)) \mapsto \Phi_{\mathcal{H},x}(\exp h_1, \dots, \exp h_s),$$

where $\Phi_{\mathcal{H},x}$ has the same meaning as in (1). This yields the desired dominating spray. \square

To extend Proposition 6.2 to the analytic setting we introduce below the notions of holomorphic flexibility. Recall that a holomorphic vector field on a complex manifold X is *completely integrable* if its phase flow defines a holomorphic action on X of the additive group $\mathbb{C}_+ = \mathbb{G}_a(\mathbb{C})$.

Definitions 6.4. (i) We say that a Stein space X is *holomorphically flexible* if the completely integrable holomorphic vector fields on X span the tangent space $T_x X$ at every smooth point of X .

(ii) Given a holomorphic submersion $h : X \rightarrow B$ of Stein manifolds, we say that X is *holomorphically flexible over B* if the completely integrable relative holomorphic vector fields on X span the relative tangent bundle of $X \rightarrow B$ at any point of X . In the latter case each fiber $h^{-1}(b)$, $b \in B$, is a holomorphically flexible Stein manifold.

Remarks 6.5. 1. The vector field $\delta = z \frac{d}{dz}$ on $X = \mathbb{C}^* = \mathbb{C} \setminus \{0\}$ is completely integrable. However, the derivation $\delta \in \text{Der}(\mathcal{O}(X))$ is not locally nilpotent. Hence $X = \mathbb{C}^*$ is not flexible in the sense used in this paper, while it is holomorphically flexible.

2. In the terminology of [38], a complex manifold X admits an *elliptic microspray* if the $\mathcal{O}_{\text{an}}(X)$ -module generated by all completely integrable holomorphic vector fields on X is dense in the $\mathcal{O}_{\text{an}}(X)$ -module of all holomorphic vector fields on X with respect to the compact-open topology.

We claim that a Stein manifold X admits an elliptic microspray if and only if X is holomorphically flexible. Indeed, admitting an elliptic microspray implies the holomorphic flexibility, because the holomorphic vector fields on a Stein manifold X span the tangent space at every point. As for the converse, we observe that on a holomorphically flexible manifold X the sheaf of germs of holomorphic vector fields is spanned by the sheaf of germs of holomorphic vector fields generated by completely integrable such fields. By Cartan's Theorem B, on a Stein manifold X the corresponding $\mathcal{O}_{\text{an}}(X)$ -modules coincide.

In the analytic setting, the following analog of Corollary 1.20 holds.

Lemma 6.6. *If a Stein manifold X is holomorphically flexible over a Stein manifold B then the relative tangent bundle of X over B is spanned a finite number of completely integrable relative holomorphic vector fields on X .*

Proof. In the absolute case i.e., B consists of a point, the assertion is just that of Lemma 4.1 in [19]. The proof of this lemma in [19] works without changes in the relative case as well. \square

With the same arguments as in the proof of Proposition 6.3 this implies that a Stein manifold X , which is holomorphically flexible over another Stein manifold B , admits a fiber dominating spray. Thus we obtain the following result.

Corollary 6.7. *Every Stein manifold X holomorphically flexible over another Stein manifold B admits a fiber dominating spray. Consequently, the Oka-Grauert-Gromov Principle is valid for $X \rightarrow B$.*

In particular, the Oka-Grauert principle holds for any holomorphically flexible Stein manifold X .

Comparing with the algebraic setting, in the analytic case we know little about invariants of completely integrable holomorphic vector fields. This leads to the following question.

Problem 6.8. *Does the group $\text{Aut}_{\text{an}}(X)$ of holomorphic automorphisms of a flexible connected Stein manifold X act infinitely transitively on X ?*

This group is transitive on X . Indeed, by the implicit function theorem every orbit of the group $\text{Aut}_{\text{an}}(X)$ is open. On the other hand, such an orbit is the complement of the union of all other orbits, thus it is closed. Hence there is only one orbit.

However, the infinite transitivity holds under a stronger assumption. We need the following notion from the Andersen-Lempert theory.

Definitions 6.9. (see [19], [39]) (i) We say that a complex manifold X has the *density property* if the Lie algebra generated by all completely integrable holomorphic vector fields on X is dense in the Lie algebra of all holomorphic vector fields on X in the compact-open topology.

(ii) Similarly, we say that an affine algebraic manifold X has the *algebraic density property* if the Lie algebra generated by all completely integrable algebraic vector fields on X coincides with the Lie algebra of all algebraic vector fields on X .

An analytic version of Theorem 0.1 can be stated as follows (cf. Theorem 5.5 in [11]).

Theorem 6.10. ([19, 2.13], [39]) *Let a Stein manifold X of dimension ≥ 2 possess the density property. Then the group $\text{Aut}_{\text{an}}(X)$ of holomorphic automorphisms of X acts infinitely transitively²⁶ on X . Moreover, for any discrete subset $Z \subseteq X$ and for any Stein space Y of positive dimension which admits a proper embedding into X , there is another proper embedding $\varphi : Y \hookrightarrow X$ which interpolates Z i.e., $Z \subseteq \varphi(Y)$.*

We refer the reader to [5] for a result on interpolation of a given discrete set of jets of automorphisms by an analytic automorphism of an affine space, similar to our Theorem 4.5.

6.2. Volume density property. As usual a holomorphic volume form ω on a complex manifold X is a nowhere vanishing top-dimensional holomorphic form on X . We need the following notions.

Definitions 6.11. (i) Given a submersion $X \rightarrow B$ of Stein manifolds and a volume form ω on X we say that X is *holomorphically volume flexible over B* , if Definition 6.4(ii) holds with all relative holomorphic vector fields considered there being ω -divergence-free. The latter means that the corresponding phase flow preserves ω .

²⁶By ‘infinite transitivity’ we mean, as before, m -transitivity for all $m \in \mathbb{N}$. Note however that transitivity for arbitrary discrete subsets does not hold already in $X = \mathbb{A}_{\mathbb{C}}^n$, as shows the famous example of Rosay and Rudin, see e.g., [11].

In the absolute case i.e., B is a point, we simply call the space X holomorphically volume flexible.

(ii) We say that X has the *volume density property* if Definition 6.9 holds with all fields in consideration being ω -divergence-free. The *algebraic volume density property* is defined likewise.

The holomorphic volume flexibility of a Stein manifold X is equivalent to the existence on X of an elliptic volume microspray as introduced in [38]. Lemma 6.6 and Corollary 6.7 admit analogs in this new context. However, the proofs become now more delicate. We address the interested reader to [19, 20].

The algebraic volume density property implies the usual volume density property [20]. However, we do not know whether a holomorphically volume flexible Stein manifold has automatically the volume density property (cf. [38]).

Concerning infinite transitivity, the following theorem is proven in [19, 2.1-2.2].

Theorem 6.12. *Let X be a Stein manifold of dimension ≥ 2 equipped with a holomorphic volume form. If X satisfies the holomorphic volume density property, then the conclusions of Theorem 6.10 hold, with volume preserving automorphisms.*

Given an algebraic volume form ω on a smooth affine algebraic variety X , every locally nilpotent vector field on X is automatically ω -divergence-free. Thus the usual flexibility implies the algebraic volume flexibility. Let us formulate the following related problem.

Problem 6.13. *Let X be a flexible smooth affine algebraic variety over \mathbb{C} equipped with an algebraic volume form. Does the algebraic volume density property hold for X ?*

We conclude with yet another problem.

Problem 6.14. Does there exist a flexible exotic algebraic structure on an affine space that is, a flexible smooth affine variety over \mathbb{C} diffeomorphic but not isomorphic to an affine space $\mathbb{A}_{\mathbb{C}}^n$?

Notice that for all exotic structures on $\mathbb{A}_{\mathbb{C}}^n$ known so far the Makar-Limanov invariant is non-trivial, whereas for a flexible such structure, by Proposition 5.1 even the field Makar-Limanov invariant must be trivial.

REFERENCES

- [1] I.V. Arzhantsev, K. Kuyumzhiyan, M. Zaidenberg: Flag varieties, toric varieties, and suspensions: three instances of infinite transitivity. arXiv:1003.3164, 25p.
- [2] V. Batyrev, F. Haddad: On the geometry of $SL(2)$ -equivariant flips. Mosc. Math. J. 8 (2008), 621–646.
- [3] A. Bialynicki-Birula, G. Hochschild, G. D. Mostow: Extensions of representations of algebraic linear groups. Amer. J. Math. 85 (1963), 131–144.
- [4] A. Borel: Les bouts des espaces homogènes de groupes de Lie. Ann. Math. (2) 58 (1953), 443–457.
- [5] G. T. Buzzard, F. Forstnerič: An interpolation theorem for holomorphic automorphisms of \mathbb{C}^n . J. Geom. Anal. 10 (2000), 101–108.
- [6] V. I. Danilov: Algebraic varieties and schemes. Algebraic geometry, I, 167–297, Encyclopaedia Math. Sci. 23, Springer, Berlin, 1994.

- [7] V. I. Danilov, M. H. Gizatullin: Examples of nonhomogeneous quasihomogeneous surfaces. *Math. USSR Izv.* 8 (1974), 43–60.
- [8] A. Dubouloz: Completions of normal affine surfaces with a trivial Makar-Limanov invariant. *Michigan Math. J.* 52 (2004), 289–308.
- [9] H. Flenner, S. Kaliman, and M. Zaidenberg: Smooth Affine Surfaces with Non-Unique \mathbb{C}^* -Actions. arXiv:0809.0651, 50p.; to appear in : *J. of Algebraic Geom.*
- [10] F. Forstnerič: Holomorphic flexibility properties of complex manifolds. *Amer. J. Math.* 128 (2006), 239–270.
- [11] F. Forstnerič: The homotopy principle in complex analysis: a survey. *Explorations in complex and Riemannian geometry*, 73–99, *Contemp. Math.*, 332, Amer. Math. Soc., Providence, RI, 2003.
- [12] G. Freudenburg: Algebraic Theory of Locally Nilpotent Derivations. *Encyclopaedia of Mathematical Sciences*, Vol. 136, Springer-Verlag, 2006.
- [13] J.-P. Furter, S. Lamy: Normal subgroup generated by a plane polynomial automorphism. E-print arXiv:0910.1616 (2010), 29 p.
- [14] S. A. Gaifullin: Affine toric $SL(2)$ -embeddings. *Sb. Math.* 199 (2008), 319–339.
- [15] M. H. Gizatullin: I. *Affine surfaces that are quasihomogeneous with respect to an algebraic group*, *Math. USSR Izv.* 5 (1971), 754–769; II. *Quasihomogeneous affine surfaces*, *ibid.* 1057–1081.
- [16] M. Gromov: Oka’s principle for holomorphic sections of elliptic bundles. *J. Amer. Math. Soc.* 2 (1989), 851–897.
- [17] A. Grothendieck: *Éléments de géométrie algébrique*. II. Étude globale élémentaire de quelques classes de morphismes. *Inst. Hautes Études Sci. Publ. Math.* 8 (1961), 222p.; IV. Étude locale des schémas et des morphismes de schémas III. *Inst. Hautes Études Sci. Publ. Math.* No. 28, 1966.
- [18] Z. Jelonek: A hypersurface which has the Abhyankar-Moh property. *Math. Ann.* 308 (1997), 73–84.
- [19] S. Kaliman, F. Kutzschebauch: On the present state of the Andersen-Lempert theory. arXiv:1003.3434 (2010), 42p. To appear in: Peter Russell’s Festschrift, Proceedings of the conference on Affine Algebraic Geometry held in Professor Russell’s honour, 1-5 June 2009, McGill Univ., Montreal.
- [20] S. Kaliman, F. Kutzschebauch: Algebraic volume density property of affine algebraic manifolds. *Invent. Math.* 181 (2010), 605–647.
- [21] S. Kaliman, F. Kutzschebauch: Criteria for the density property of complex manifolds. *Invent. Math.* 172 (2008), 71–87.
- [22] S. Kaliman, M. Zaidenberg: Affine modifications and affine hypersurfaces with a very transitive automorphism group. *Transform. Groups* 4 (1999), 53–95.
- [23] S. Kaliman, M. Zaidenberg: Miyanishi’s characterization of the affine 3-space does not hold in higher dimensions. *Ann. Inst. Fourier (Grenoble)* 50 (2000), 1649–1669 (2001).
- [24] S. L. Kleiman: The transversality of a general translate. *Compositio Math.* 28 (1974), 287–297.
- [25] F. Knop: Mehrfach transitive Operationen algebraischer Gruppen. *Arch. Math.* 41 (1983), 438–446.
- [26] H. Kraft: Geometrische Methoden in der Invariantentheorie. *Aspects of Mathematics*, D1. Friedr. Vieweg & Sohn, Braunschweig, 1984.
- [27] A. Liendo: Affine T -varieties of complexity one and locally nilpotent derivations. *Transform. Groups* 15 (2010), 389–425.
- [28] A. Liendo: \mathbb{G}_a -actions of fiber type on affine T -varieties. arXiv:0911.1110, 12p. (2009); *J. of Algebra* (to appear).
- [29] V. L. Popov: I. Quasihomogeneous affine algebraic varieties of the group $SL(2)$. *Math. USSR Izv.* 7 (1973), 793–831; II. Classification of affine algebraic surfaces that are quasihomogeneous with respect to an algebraic group. *ibid.* 7 (1973), 1039–1056; III. Classification

of three-dimensional affine algebraic varieties that are quasihomogeneous with respect to an algebraic group. *ibid.* 9 (1975), 535–576.

- [30] V. L. Popov: Generically multiple transitive algebraic group actions. Algebraic groups and homogeneous spaces, 481–523, Tata Inst. Fund. Res. Stud. Math., Mumbai, 2007.
- [31] V. L. Popov: Open Problems. In: Affine algebraic geometry. Papers from the Special Session at the 1st Joint AMS-RSME Meeting held in Seville, June 18–21, 2003, 12–16. J. Gutierrez, V. Shpilrain and Jie-Tai Yu (eds). Contemporary Mathematics, 369. American Mathematical Society, Providence, RI, 2005.
- [32] V. L. Popov: On the Makar-Limanov, Derksen invariants, and finite automorphism groups of algebraic varieties. arXiv:1001.1311 (2010), 22p.
- [33] V. L. Popov and E. B. Vinberg: Invariant Theory. In: Algebraic geometry IV, A. N. Parshin, I. R. Shafarevich (eds), Berlin, Heidelberg, New York: Springer-Verlag, 1994.
- [34] C. Procesi: Lie groups. An approach through invariants and representations. Universitext. Springer, New York, 2007.
- [35] Z. Reichstein: I. On automorphisms of matrix invariants. Trans. Amer. Math. Soc. 340 (1993), 353–371; II. On automorphisms of matrix invariants induced from the trace ring. Linear Algebra Appl. 193 (1993), 51–74.
- [36] R. Rentschler: Opérations du groupe additif sur le plane affine, C. R. Acad. Sci. **267** (1968), 384–387.
- [37] I. P. Shestakov, U. U. Umirbaev: The tame and the wild automorphisms of polynomial rings in three variables. J. Amer. Math. Soc. 17 (2004), 197–227.
- [38] D. Varolin: A general notion of shears, and applications. Michigan Math. J. 46 (1999), 533–553.
- [39] D. Varolin: The density property for complex manifolds and geometric structures. I. J. Geom. Anal. 11 (2001), 135–160. II. Internat. J. Math. 11 (2000), 837–847.
- [40] J. Winkelmann: On automorphisms of complements of analytic subsets in \mathbb{C}^n . Math. Z. 204 (1990), 117–127.

DEPARTMENT OF ALGEBRA, FACULTY OF MECHANICS AND MATHEMATICS, MOSCOW STATE UNIVERSITY, LENINSKIE GORY 1, GSP-1, MOSCOW, 119991, RUSSIA

E-mail address: arjantse@mcme.ru

FAKULTÄT FÜR MATHEMATIK, RUHR UNIVERSITÄT BOCHUM, GEB. NA 2/72, UNIVERSITÄTSSTR. 150, 44780 BOCHUM, GERMANY

E-mail address: Hubert.Flenner@rub.de

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MIAMI, CORAL GABLES, FL 33124, USA

E-mail address: kaliman@math.miami.edu

MATHEMATISCHES INSTITUT, UNIVERSITÄT BERN, SIDLERSTRASSE 5, CH-3012 BERN, SWITZERLAND,

E-mail address: frank.kutzschebauch@math.unibe.ch

UNIVERSIT GRENOBLE I, INSTITUT FOURIER, UMR 5582 CNRS-UJF, BP 74, 38402 ST. MARTIN D'HÈRES CÉDEX, FRANCE

E-mail address: Mikhail.Zaidenberg@ujf-grenoble.fr