

Roy–Steiner equations for πN scattering

Christoph Ditsche*^a, **Martin Hoferichter**^{ab}, **Bastian Kubis**^a and **Ulf-G. Meißner**^{ac}

^a*Helmholtz-Institut für Strahlen- und Kernphysik (Theorie) and Bethe Center for Theoretical Physics, Universität Bonn, D-53115 Bonn, Germany*

^b*Albert Einstein Center for Fundamental Physics, Institute for Theoretical Physics, Universität Bern, CH-3012 Bern, Switzerland*

^c*Institut für Kernphysik, Institute for Advanced Simulation, and Jülich Center for Hadron Physics, Forschungszentrum Jülich, D-52425 Jülich Germany*

E-mail: ditsche@hiskp.uni-bonn.de

Starting from hyperbolic dispersion relations for the invariant amplitudes of pion–nucleon scattering together with crossing symmetry and unitarity, one can derive a closed system of integral equations for the partial waves of both the s -channel ($\pi N \rightarrow \pi N$) and the t -channel ($\pi\pi \rightarrow \bar{N}N$) reaction, called Roy–Steiner equations. After giving a brief overview of the Roy–Steiner system for πN scattering, we demonstrate that the solution of the t -channel subsystem, which represents the first step in solving the full system, can be achieved by means of Muskhelishvili–Omnès techniques. In particular, we present results for the P -waves featuring in the dispersive analysis of the electromagnetic form factors of the nucleon.

*The 7th International Workshop on Chiral Dynamics,
August 6 -10, 2012
Jefferson Lab, Newport News, Virginia, USA*

*Speaker.

1. Introducing Roy–Steiner equations for πN scattering

Partial-wave dispersion relations (PWDRs) together with unitarity and crossing symmetry as well as isospin and chiral symmetry (i.e. all available symmetry constraints) have repeatedly proven to be a powerful tool for studying processes at low energies with high precision [1–4]. For πN scattering the (unsubtracted) hyperbolic dispersion relations (HDRs) for the usual Lorentz-invariant amplitudes read [5] (using the notation of [6], see [7] for more details)

$$\begin{aligned}
A^+(s, t) &= \frac{1}{\pi} \int_{s_+}^{\infty} ds' \left[\frac{1}{s' - s} + \frac{1}{s' - u} - \frac{1}{s' - a} \right] \text{Im} A^+(s', t') + \frac{1}{\pi} \int_{t_\pi}^{\infty} dt' \frac{\text{Im} A^+(s', t')}{t' - t}, \\
B^+(s, t) &= N^+(s, t) + \frac{1}{\pi} \int_{s_+}^{\infty} ds' \left[\frac{1}{s' - s} - \frac{1}{s' - u} \right] \text{Im} B^+(s', t') + \frac{1}{\pi} \int_{t_\pi}^{\infty} dt' \frac{s - u}{s' - u'} \frac{\text{Im} B^+(s', t')}{t' - t}, \\
N^+(s, t) &= g^2 \left[\frac{1}{m^2 - s} - \frac{1}{m^2 - u} \right], \quad (s - a)(u - a) = b = (s' - a)(u' - a), \quad (1.1)
\end{aligned}$$

and similarly for A^- , B^- , and N^- , where N^\pm are the nucleon pole terms and the “external” (unprimed) and “internal” (primed) kinematics are related by real hyperbola parameters a and b (as well as via $s + t + u = 2(m^2 + M_\pi^2) = s' + t' + u'$), so that HDRs allow for the combination of all physical regions, which is known to be crucial for a reliable continuation into the subthreshold region and hence for an accurate determination of the πN σ -term. Furthermore, the imaginary parts are only needed in regions where the corresponding partial-wave decompositions converge and the range of convergence can be maximized by tuning the free hyperbola parameter a . While the s -channel integrals start at the threshold $s_+ = W_+^2 = (m + M_\pi)^2$, the t -channel contributes already above the pseudothreshold $t_\pi = 4M_\pi^2$ far below the threshold $t_N = 4m^2$. Depending on the asymptotic behavior of the imaginary parts, in principle it could be necessary to subtract the HDRs to ensure the convergence of the integrals, thereby parameterizing high-energy information with polynomials containing a priori unknown subtraction constants. However, (additional) subtractions may also be introduced to lessen the dependence of the low-energy solution on high-energy input; the corresponding subtraction parameters then obey respective sum rules. For πN scattering it proves particularly useful to subtract at the subthreshold point ($s = u, t = 0$), as this preserves the $s \leftrightarrow u$ crossing symmetry (which can be made explicit in terms of the crossing variable $v = (s - u)/(4m)$ via $D^\pm(v, t) = A^\pm + vB^\pm = \pm D^\pm(-v, t)$). This is especially favorable for the t -channel subproblem and facilitates matching to chiral perturbation theory [8, 9] to determine the subtraction constants, which thus can be identified with the subthreshold expansion parameters.¹ In addition to the presentation in [7], we also introduce a (partial) third subtraction, which is related to the parameters a_{10}^+ and a_{01}^- of the subthreshold expansions (with $d_{0n}^+ = a_{0n}^+$ for all $n \geq 0$)

$$A^+(v, t) = \frac{g^2}{m} + d_{00}^+ + d_{01}^+ t + a_{10}^+ v^2 + \mathcal{O}(v^4, v^2 t, t^2), \quad A^-(v, t) = a_{00}^- v + a_{01}^- vt + \mathcal{O}(v^3, vt^2). \quad (1.2)$$

¹For the PWDRs of $\pi\pi$ scattering, called Roy equations [10], an analogous matching procedure for the $\pi\pi$ scattering lengths as pertinent subtraction parameters has been conducted in [11]. In contrast to $\pi\pi$ scattering, the πN scattering lengths can be extracted with high accuracy from hadronic-atom data [12, 13] and may thus serve as additional constraints on the subtraction constants in the Roy–Steiner system.

In order to derive the partial-wave HDRs, called Roy–Steiner (RS) equations, one needs to expand the s - and t -channel imaginary parts in (1.1) into the respective partial waves and subsequently project the full expanded equations onto either s - or t -channel partial waves; the resulting sets of integral equations together with the respective partial-wave unitarity relations then form the s - and t -channel RS subsystems. According to [5], the (unsubtracted) s -channel RS equations read (based on the MacDowell symmetry $f_{(\ell+1)-}^I(W) = -f_{\ell+}^I(-W)$ for all $\ell \geq 0$ [14])

$$f_{\ell+}^I(W) = N_{\ell+}^I(W) + \frac{1}{\pi} \int_{t_\pi}^{\infty} dt' \sum_J \left\{ G_{\ell J}(W, t') \operatorname{Im} f_+^J(t') + H_{\ell J}(W, t') \operatorname{Im} f_-^J(t') \right\} \\ + \frac{1}{\pi} \int_{W_+}^{\infty} dW' \sum_{\ell'=0}^{\infty} \left\{ K_{\ell\ell'}^I(W, W') \operatorname{Im} f_{\ell'+}^I(W') + K_{\ell\ell'}^I(W, -W') \operatorname{Im} f_{(\ell'+1)-}^I(W') \right\}, \quad (1.3)$$

where due to G -parity only even/odd J contribute for isospin $I = +/-$, respectively, and the partial-wave projections of the pole terms as well as the (lowest) kernels are analytically known, the latter including in particular the Cauchy kernel: $K_{\ell\ell'}^I(W, W') = \delta_{\ell\ell'} / (W' - W) + \dots$. The s -channel $I = \pm$ partial waves are intertwined by the usual unitarity relations, which are diagonal in the s -channel isospin basis $I_s \in \{1/2, 3/2\}$ only. Once the t -channel partial waves are known, the structure of the s -channel RS subsystem is therefore similar to the $\pi\pi$ Roy equations, cf. [1]. As shown in [7], the corresponding (unsubtracted) t -channel RS equations are given by

$$f_+^J(t) = \tilde{N}_+^J(t) + \frac{1}{\pi} \int_{W_+}^{\infty} dW' \sum_{\ell=0}^{\infty} \left\{ \tilde{G}_{J\ell}(t, W') \operatorname{Im} f_{\ell+}^J(W') + \tilde{G}_{J\ell}(t, -W') \operatorname{Im} f_{(\ell+1)-}^J(W') \right\} \\ + \frac{1}{\pi} \int_{t_\pi}^{\infty} dt' \sum_{J'} \left\{ \tilde{K}_{JJ'}^1(t, t') \operatorname{Im} f_+^{J'}(t') + \tilde{K}_{JJ'}^2(t, t') \operatorname{Im} f_-^{J'}(t') \right\} \quad (1.4)$$

and similarly for the f_-^J except for the fact that these do not receive contributions from the f_+^J . Here, only even or odd J' couple to even or odd J (corresponding to t -channel isospin $I_t = 0$ or $I_t = 1$), respectively, and $\tilde{K}_{JJ'}^1$ (as well as the analogous $\tilde{K}_{JJ'}^2$ for the f_-^J) contains the Cauchy kernel. Moreover, it turns out that only higher t -channel partial waves contribute to lower ones. Assuming Mandelstam analyticity, the equations (1.4) are valid for $\sqrt{t} \in [2M_\pi, 2.00 \text{ GeV}]$ using $a = -2.71M_\pi^2$, whereas (1.3) holds for $W \in [m + M_\pi, 1.38 \text{ GeV}]$ using $a = -23.19M_\pi^2$. The t -channel unitarity relations are diagonal in I_t and only linear in the f_\pm^J (below the first inelastic threshold t_{inel})

$$\operatorname{Im} f_\pm^J(t) = \sigma_t^\pi (t_J^I(t))^* f_\pm^J(t) \theta(t - t_\pi), \quad \sigma_t^\pi t_J^I(t) = \sin \delta_J^I(t) e^{i\delta_J^I(t)}, \quad \sigma_t^\pi(t) = \sqrt{1 - t_\pi/t},$$

from which one can infer Watson’s final state interaction theorem [15] stating that (in the “elastic” region) the phase of f_\pm^J is given by the phase δ_J^I of the respective $\pi\pi$ scattering partial wave t_J^I .

Due to the simpler recoupling scheme for the f_\pm^J , the t -channel RS subsystem can be recast as a (single-channel) Muskhelishvili–Omnès (MO) problem [16] with a finite matching point t_m [3] for f_+^0 , f_-^J , and the linear combinations $\Gamma^J(t) = m\sqrt{J/(J+1)} f_-^J(t) - f_+^J(t)$ with $\Gamma^J(t_m) = 0$ for all $J \geq 1$ of the generic form (the details are given in [7])

$$f(t) = \Delta(t) + \frac{1}{\pi} \int_{t_\pi}^{t_m} dt' \frac{\sin \delta(t') e^{-i\delta(t')} f(t')}{t' - t} + \frac{1}{\pi} \int_{t_m}^{\infty} dt' \frac{\operatorname{Im} f(t')}{t' - t} \equiv |f(t)| e^{i\delta(t)} \quad \text{for } t \leq t_m < t_{\text{inel}},$$




Figure 1: Flowchart of the solution strategy for the Roy–Steiner system for πN scattering. The third step consist in the self-consistent iteration (denoted by thick arrows) of the preceding steps until convergence.

where the inhomogeneities $\Delta(t)$ subsume the nucleon pole terms, all s -channel integrals, and the higher t -channel partial waves. For $t_\pi \leq t \leq t_m$, solving for $|f(t)|$ only according to Watson’s theorem requires $\delta(t)$ for $t_\pi \leq t \leq t_m$ and $\text{Im} f(t)$ for $t \geq t_m$. Introducing $n \geq 1$ subtractions does not change the general structure of the RS/MO system, e.g. the P -waves are given by

$$\Gamma^1(t) = \Delta_\Gamma^1(t) \Big|^{n\text{-sub}} + \frac{t^{n-1}(t-t_N)}{\pi} \int_{t_\pi}^{\infty} \frac{dt' \text{Im} \Gamma^1(t')}{t'^{n-1}(t'-t_N)(t'-t)}, \quad f_-^1(t) = \Delta_-^1(t) \Big|^{n\text{-sub}} + \frac{t^n}{\pi} \int_{t_\pi}^{\infty} \frac{dt' \text{Im} f_-^1(t')}{t'^n(t'-t)},$$

demonstrating that Γ^J and hence f_+^J is effectively subtracted by one power less than f_-^J , which motivates the additional (partial) third subtraction in A^\pm , cf. (1.2), that affects solely the f_+^J .

The solution strategy for the full RS system in the low-energy (or even subthreshold/pseudo-physical) regions, where only the lowest partial waves are relevant and inelastic contributions may be (approximately) neglected, is shown in Fig. 1; see [7] for more details.

2. The t -channel Muskhelishvili–Omnès problem: P -wave solutions

As the first step in the numerical solution of the full RS system, we check the consistency of our t -channel MO solutions with the results of the KH80 analysis [17], which are still used nowadays although no thorough error estimates are given (and despite the availability of more modern experimental data). Here, we present results for the P -waves in the (elastic) single-channel approximation of the MO problem, which is well justified for the P - and higher partial waves, whereas the S -wave requires a two-channel description including $\bar{K}K$ intermediate states as described in [7]. To produce the results (that will also serve as input for the solution of the s -channel RS subsystem, cf. Fig. 1) partly shown in Fig. 2, we have used as input $\pi\pi$ phase shifts from [18], s -channel partial waves ($l \leq 4$) from SAID [19] for $W \leq 2.5 \text{ GeV}$, and above the Regge model of [20]. To facilitate




Figure 2: n -subtracted MO solutions for the P -wave moduli.




Figure 3: Two-pion-continuum contribution to $\text{Im } G_E^V$ and $\text{Im } G_M^V$.

comparison with the results of KH80, we use the respective subthreshold parameter values and a πN coupling of $g^2/(4\pi) = 14.28$ [6, 17] (as starting point, the final values will result from the iteration procedure, cf. Fig. 1).² Moreover, KH80 uses different types of dispersion relations, in particular so-called fixed- t ones, which can be emulated (up to the t -channel contributions that are not present at all in the fixed- t case) by taking the “fixed- t limit” $|a| \rightarrow \infty$. As argued in [7], all t -channel input above $\sqrt{t_m} = 0.98 \text{ GeV}$ is set to zero, which forces the MO solutions to match zero at $t = t_m$. While Fig. 2 displays the results for $|a| \rightarrow \infty$, investigating the effect of using a different (i.e. higher) matching point leads to the same conclusion: with increasing number of subtractions, thus lowering the dependence on the high-energy input by introducing more subthreshold parameter contributions as subtraction polynomials, the solutions show a nice convergence pattern both in general (proving the internal consistency and numerical stability of our RS/MO framework) and in particular towards the KH80 results (being consistent with relying on KH80 values for g and the subtraction parameters). The P -waves feature prominently in the dispersive analysis of the nucleon electromagnetic form factors, see e.g. [21] and references therein, and in Fig. 3 we illustrate the effects on the spectral functions (by approximating the vector pion form factor F_π^V via a

²Modern analyses yield significant smaller values for the πN coupling, cf. e.g. $g^2/(4\pi) = 13.7 \pm 0.2$ of [13].

twice-subtracted Omnès representation, cf. [7])

$$\text{Im } G_E^v(t) = \frac{t(\sigma_t^\pi)^3}{8m} (F_\pi^V(t))^* f_+^1(t) \theta(t-t_\pi), \quad \text{Im } G_M^v(t) = \frac{t(\sigma_t^\pi)^3}{8\sqrt{2}} (F_\pi^V(t))^* f_-^1(t) \theta(t-t_\pi).$$

We are confident that a self-consistent iteration procedure between the solutions for the s - and t -channel eventually will yield a consistent and precise description (including error estimates) of the low-energy πN scattering amplitude in all kinematical channels.

References

- [1] B. Ananthanarayan, G. Colangelo, J. Gasser and H. Leutwyler, *Phys. Rept.* **353** (2001) 207 [hep-ph/0005297].
- [2] R. García-Martín *et al.*, *Phys. Rev. D* **83** (2011) 074004 [arXiv:1102.2183 [hep-ph]].
- [3] P. Büttiker, S. Descotes-Genon and B. Moussallam, *Eur. Phys. J. C* **33** (2004) 409 [hep-ph/0310283].
- [4] M. Hoferichter, D. R. Phillips and C. Schat, *Eur. Phys. J. C* **71** (2011) 1743 [arXiv:1106.4147 [hep-ph]].
- [5] G. E. Hite and F. Steiner, *Nuovo Cim. A* **18** (1973) 237.
- [6] G. Höhler, *Pion–Nukleon–Streuung: Methoden und Ergebnisse*, in Landolt-Börnstein, **9b2**, ed. H. Schopper, Springer Verlag, Berlin, 1983.
- [7] C. Ditsche, M. Hoferichter, B. Kubis and U.-G. Meißner, *JHEP* **1206** (2012) 043 [arXiv:1203.4758 [hep-ph]]; *JHEP* **1206** (2012) 063 [arXiv:1204.6251 [hep-ph]]; arXiv:1211.1485 [nucl-th].
- [8] P. Büttiker and U.-G. Meißner, *Nucl. Phys. A* **668** (2000) 97 [hep-ph/9908247].
- [9] T. Becher and H. Leutwyler, *JHEP* **0106** (2001) 017 [hep-ph/0103263].
- [10] S. M. Roy, *Phys. Lett. B* **36** (1971) 353.
- [11] G. Colangelo, J. Gasser and H. Leutwyler, *Phys. Lett. B* **488** (2000) 261 [hep-ph/0007112]; *Nucl. Phys. B* **603** (2001) 125 [hep-ph/0103088].
- [12] D. Gotta *et al.*, *Lect. Notes Phys.* **745** (2008) 165; T. Strauch *et al.*, *Eur. Phys. J. A* **47** (2011) 88 [arXiv:1011.2415 [nucl-ex]].
- [13] V. Baru *et al.*, *Phys. Lett. B* **694** (2011) 473 [arXiv:1003.4444 [nucl-th]]; *Nucl. Phys. A* **872** (2011) 69 [arXiv:1107.5509 [nucl-th]]; arXiv:1211.1145 [nucl-th].
- [14] S. W. MacDowell, *Phys. Rev.* **116** (1959) 774.
- [15] K. M. Watson, *Phys. Rev.* **95** (1954) 228.
- [16] N. I. Muskhelishvili, *Singular Integral Equations*, Wolters-Noordhoff Publishing, Groningen, 1953; R. Omnès, *Nuovo Cim.* **8** (1958) 316.
- [17] R. Koch and E. Pietarinen, *Nucl. Phys. A* **336** (1980) 331.
- [18] I. Caprini, G. Colangelo and H. Leutwyler, *Eur. Phys. J. C* **72** (2012) 1860 [arXiv:1111.7160 [hep-ph]]; in preparation.
- [19] SAID, <http://gwdac.phys.gwu.edu>; R. A. Arndt, W. J. Briscoe, I. I. Strakovsky and R. L. Workman, *Eur. Phys. J. A* **35** (2008) 311.
- [20] F. Huang *et al.*, *Eur. Phys. J. A* **44** (2010) 81 [arXiv:0910.4275 [nucl-th]].
- [21] I. T. Lorenz, H.-W. Hammer and U.-G. Meißner, *Eur. Phys. J. A* **48** (2012) 151 [arXiv:1205.6628 [hep-ph]].