

# A Genome-Wide Association Study to Detect QTL for Commercially Important Traits in Swiss Large White Boars

Doreen Becker<sup>1</sup>, Klaus Wimmers<sup>2</sup>, Henning Luther<sup>3</sup>, Andreas Hofer<sup>3</sup>, Tosso Leeb<sup>1\*</sup>

**1** Institute of Genetics, Vetsuisse Faculty, University of Bern, Bern, Switzerland, **2** Leibniz Institute for Farm Animal Biology, Dummerstorf, Germany, **3** SUISAG, Sempach, Switzerland

## Abstract

The improvement of meat quality and production traits has high priority in the pork industry. Many of these traits show a low to moderate heritability and are difficult and expensive to measure. Their improvement by targeted breeding programs is challenging and requires knowledge of the genetic and molecular background. For this study we genotyped 192 artificial insemination boars of a commercial line derived from the Swiss Large White breed using the PorcineSNP60 BeadChip with 62,163 evenly spaced SNPs across the pig genome. We obtained 26 estimated breeding values (EBVs) for various traits including exterior, meat quality, reproduction, and production. The subsequent genome-wide association analysis allowed us to identify four QTL with suggestive significance for three of these traits (p-values ranging from  $4.99 \times 10^{-6}$  to  $2.73 \times 10^{-5}$ ). Single QTL for the EBVs pH one hour post mortem (pH1) and carcass length were on pig chromosome (SSC) 14 and SSC 2, respectively. Two QTL for the EBV rear view hind legs were on SSC 10 and SSC 16.

**Citation:** Becker D, Wimmers K, Luther H, Hofer A, Leeb T (2013) A Genome-Wide Association Study to Detect QTL for Commercially Important Traits in Swiss Large White Boars. *PLoS ONE* 8(2): e55951. doi:10.1371/journal.pone.0055951

**Editor:** Stephen Moore, University of Queensland, Australia

**Received:** August 10, 2012; **Accepted:** January 4, 2013; **Published:** February 5, 2013

**Copyright:** © 2013 Becker et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This study was funded by a grant from the German Research Council (DFG LE 1032/23-1). The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** Henning Luther and Andreas Hofer are employed by SUISAG, the company who also provided tissue samples and estimated breeding values of the animals. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials.

\* E-mail: [tosso.leeb@vetsuisse.unibe.ch](mailto:tosso.leeb@vetsuisse.unibe.ch)

## Introduction

One of the most challenging tasks in pork production is the improvement of traits with low heritability [1–3]. Most commercially important traits are complex and influenced by multiple interacting factors including genetics and environment. The observed albeit low heritabilities suggest that these traits could be successfully improved by selection. However, the cost and difficulty to obtain direct measurements are limiting the improvements of these traits. In addition, the estimation of many breeding values relies only on the phenotypes of relatives, which limits their accuracy. Therefore, these traits are ideal candidates for the application of molecular genetic tools in future breeding programs. The identification of genes and polymorphisms associated with commercially important traits can provide useful markers for the selection of genetically superior animals. Knowledge of the genetic and molecular background is required to accelerate the genetic improvement.

Since the first QTL genome scan in pigs was reported [4] numerous QTL analyses have been conducted to identify QTL for various traits in pig production. For example Karlskov-Mortensen et al. reported QTL on chromosomes SSC 1, 4, 9, 10, 13 and 16 affecting fat deposition and lean meat content [5]. For various growth traits QTL were detected on SSC 1, 4, 7 and 8 by de Koning et al. [6]. Several QTL for pH at 45 minutes post mortem have been shown to be located on SSC3, 4, 5, 6, 8, 11, 13 and 17 [7]. Identification of multiple loci associated with one trait suggests an underlying complex genetic architecture. Originally, QTL

scans have been mainly performed on experimental crosses between a domestic breed and wild boar or Meishan [8–10]. Over the last decade, a number of QTL scans have been carried out on commercial pig line crosses, including Large White, Piétrain, Berkshire and Yorkshire [11,12].

So far, over 6,800 QTL for 585 different traits have been identified in pigs (PigQTLdb, <http://www.animalgenome.org/cgi-bin/QTLdb/SS/index>), more than in any other livestock species. The reported QTL were shown to influence meat quality, health, production, reproduction and exterior traits. However, only a few have been further investigated and led to discovery of associated or even causative mutations. These include a single base pair substitution in a non-coding region of *IGF2* on SSC 2 that explains variation in muscle mass and back fat thickness and a nonconservative substitution in the *PRKAG3* gene on SSC 15 having an effect on meat quality [13,14]. Since most loci explain just a small fraction of the phenotypic variation, the identification of the causative genetic variations underlying QTL remains challenging [15].

The recent development of the PorcineSNP60 BeadChip [16] facilitated by the efforts of the International Swine Genome Sequencing Consortium and Illumina now enables us to perform genome-wide association studies (GWAS) in pigs. Numerous GWAS for quantitative traits have been reported in humans [17,18] and other domestic animals [19,20]. However, due to a lack of SNP genotyping arrays, the first GWAS for quantitative traits in pigs have only recently been published. In the last few

months several GWAS investigating various traits of commercial or scientific interest in different pig populations were published [21–28]. The aim of this study was to identify QTL affecting economically important traits in a Swiss commercial boar line using the PorcineSNP60 BeadChip.

**Results**

We used 26 estimated breeding values (EBVs) rather than raw phenotypes for our GWAS. Breeding values have the advantage that they are free of systematic environmental effects on measured phenotypes, as these effects are considered in the statistical model used for the estimation of EBVs. Additionally, they reflect the genetic makeup more accurately because they do not solely rely on own records but include information from all measured relatives. The analyzed traits with available EBVs and key figures of their distribution are shown in Table S1. The EBVs are expressed as a deviation from the mean of a defined group of animals. Therefore the median is close to zero.

We genotyped 192 artificial insemination boars of a commercial line derived from the Swiss Large White breed for 62,163 SNP markers with an average distance of 49 kb. The average call rate per individual was 96.02%. We removed non-informative markers and markers with low call rate. After these quality control steps 186 individuals and 47,045 SNPs remained for the final analysis.


We calculated genome-wide pairwise identity-by-state distances and quantified the population stratification (Figure S1). Depending on the analyzed trait the calculated genomic inflation factor varied between 1.15 and 2 indicating that our material was highly stratified. Performing association studies with stratified samples can lead to false positive results, i.e. detected associations can be due to the underlying structure of the population instead of a biologically meaningful association with one or several genes. Therefore, we corrected for the population stratification in our association analysis using the egscore function implemented in GenABEL and by calculating stratified associations within clusters. These corrections in addition to genomic control reduced the genomic inflation factor to reasonable values between 1.00 and 1.01 in all analyses (Figure 1).

We set the p-value thresholds for moderately significant and highly significant associations at  $5 \times 10^{-5}$  and  $5 \times 10^{-7}$ , respectively. We did not detect any highly significant associations in our material. However, we detected four QTL with moderately significant associations (Table 1).

For the EBV pH1 we detected a QTL on SSC 14. We determined the positions of the associated markers in the Sscrofa 10.2 assembly of the pig genome. We observed two additional SNPs that are associated with EBV pH1, which are no longer contained in the Sscrofa 10.2 assembly. In the previous Sscrofa 9.2 assembly one of these two SNPs resided on SSC 11, while the other was on SSC 3, and may thus represent potential additional QTL for the EBV pH1.

For the EBV carcass length we observed two closely spaced associated SNPs on SSC 2 (Table 1). The highest significance in our study was observed for a QTL for the EBV rear view hind legs with a p-value after correction for population stratification of  $4.99 \times 10^{-6}$ . We observed two QTL supported by multiple closely spaced SNPs for this trait on SSC 10 and SSC 16, respectively (Table 1).

We grouped the animals according to genotype at the best-associated SNPs for the four detected QTL and analyzed their phenotype distribution (Figure 2). The QTL for EBV pH1 showed a largely additive effect. In contrast, the genotypes at the QTL for the EBV carcass length and the QTL on SSC 10 for EBV rear



**Figure 1. Manhattan plots of genome-wide association studies for EBVs of commercially important traits in pigs.** The red lines indicate the significance threshold for moderately significant associations ( $p = 5 \times 10^{-5}$ ). “Chromosome 0” harbors SNP markers that have not yet been mapped to the pig reference genome. The inserted quantile-quantile (QQ) plots show the observed versus expected log p-values. The straight line in the QQ plots indicates the distribution of SNP markers under the null hypothesis and the skew at the right edge indicate that these markers are stronger associated with the traits than it would be expected by chance. Please note that in Figure 1A there are appear to be associated SNPs on SSC 3, 11, and 14. The two associated SNPs on SSC 3 and 11 mapped to these chromosomes in the Sscrofa 9.2 assembly, but are no longer placed on any chromosome in the Sscrofa 10.2 assembly.  
doi:10.1371/journal.pone.0055951.g001

**Table 1.** Top allelic association hits in the GWAS for QTL affecting commercially important traits.

Trait	Marker	Chromosome	Position (Sscrofa build 10.2) <sup>b</sup>	Alleles	MAF	p <sub>raw</sub> <sup>a</sup>
pH1	ASGA0061594	14	14,730,418	C/T	0.49	1.57 × 10 <sup>-5</sup>
	H3GA0032045	n.d. <sup>b</sup>	n.d. <sup>b</sup>	A/C	0.35	2.59 × 10 <sup>-5</sup>
	ASGA0105130	n.d. <sup>b</sup>	n.d. <sup>b</sup>	C/T	0.49	2.74 × 10 <sup>-5</sup>
carcass length	ASGA0010032	2	42,938,876	A/G	0.38	2.73 × 10 <sup>-5</sup>
	H3GA0006598	2	42,886,909	C/T	0.23	4.76 × 10 <sup>-5</sup>
rear view hind legs	H3GA0045902	16	6,289,550	T/G	0.50	4.99 × 10 <sup>-6</sup>
	ASGA0072056	16	6,198,618	A/G	0.44	7.68 × 10 <sup>-6</sup>
	H3GA0045917	16	6,343,134	C/T	0.40	1.42 × 10 <sup>-5</sup>
	H3GA0045908	16	6,312,026	T/C	0.41	1.96 × 10 <sup>-5</sup>
	ALGA0058443	10	40,670,821	C/A	0.37	3.35 × 10 <sup>-5</sup>
	ALGA0058422	10	39,424,934	A/G	0.44	4.56 × 10 <sup>-5</sup>
	MARC0010334	10	39,538,944	T/C	0.44	4.56 × 10 <sup>-5</sup>
	ALGA0058431	10	39,626,717	C/T	0.44	4.56 × 10 <sup>-5</sup>
	DRGA0010453	10	39,667,084	C/T	0.44	4.56 × 10 <sup>-5</sup>

<sup>a</sup>p-values were calculated using  $\chi^2$  tests in an allelic association study.


<sup>b</sup>The positions of the associated SNPs on the latest version of the pig reference genome were determined by BLAST searches with the flanking sequences of the SNPs as provided by illumina with respect to the Sscrofa 10.2 assembly. Some of the flanking sequences did not give a significant BLAST hit with respect to this genome reference sequence.

doi:10.1371/journal.pone.0055951.t001

view hind legs indicated a recessive effect of the variant allele. Finally, at the QTL on SSC 16 for EBV rear view hind legs, heterozygous animals showed EBVs that are outside of the range of animals with the two alternative homozygous genotypes (overdominance).

**Discussion**

In this study we identified four QTL with moderate significance in a Swiss commercial pig population. We think that the relatively low number of detected QTL is primarily due to the small number of animals used in this study. The population substructure with different subclusters in our animal cohort (Figure S1) required



**Figure 2. Genotype-phenotype correlations.** The animals were grouped according to genotype at the best associated SNPs of each of the four detected QTL. Animal numbers are indicated at the bottom. The call rates for these SNPs were >98.9%. For each of the four QTL the homozygous alternative genotype classes have different phenotypic distributions ( $p < 0.05$ , Welch's t-test). (A) EBV pH1 distributions with respect to genotype at SNP ASGA0061594 on SSC 14. (B) EBV carcass length in boars with respect to genotype at SNP ASGA0010032 on SSC 2. (C) EBV rear view hind legs distributions in Swiss Large White boars with respect to genotype at SNP ALGA0058443 on SSC 10. (D) EBV rear view hind legs distributions in Swiss Large White boars with respect to genotype at SNP H3GA0045902 on SSC 16. doi:10.1371/journal.pone.0055951.g002

correction of the p-values for the effects of the population stratification, which also negatively affected the power of the GWAS. The limited power of our study design is reflected by the fact that we only found QTL with very even allele distributions. The best-associated SNPs at the four detected QTL had MAFs between 0.37 and 0.50. Considering all associated SNPs, the lowest MAF was 0.23 at marker H3GA0006598 on SSC 2. The detection of QTL caused by rare alleles of the same effect size would have required larger animal numbers.

For three of the four detected QTL, similar QTL have previously been identified at roughly the same locations in other pig populations. Lee et al. reported a QTL for carcass length in an experimental Meishan × wild boar family on SSC 2 at 20.8 cM [29]. Additionally, Evans et al. identified a QTL for carcass length on SSC 2 in the region of 0–10 cM in a Landrace population [30]. A QTL for “rear upright legs”, which might be a correlated trait to the rear view hind leg score of our analysis, was identified at 43 Mb on SSC 10 [31]. This QTL was identified by genome-wide association study in commercial pigs from Large White or Large White × Landrace crosses. Another study identified a QTL for rear leg score in a Japanese Landrace population at 82 cM on SSC 10 [32]. Finally, Lee et al. reported QTLs for the trait “back legs” in a Large White × Meishan crossbred population on SSC 10 and SSC 16 at 126 cM and 10 cM, respectively [33]. It is very difficult to evaluate whether these reported QTL are really the same QTL that we found as QTL from linkage analyses were typically mapped with very low resolution and have very large confidence intervals. If QTL are independently discovered in different populations, this suggests that they may indeed be due to a biologically relevant genetic variation rather than to confounding effects such as e.g. population stratification artifacts.

So far, we have no knowledge of a study concerning the trait pH one hour post mortem. There are several studies about the pH 45 minutes post mortem which closely correlates with the trait used in our analysis (PigQTLdb, <http://www.animalgenome.org/cgi-bin/QTLdb/SS/index>). However, a QTL for pH 45 minutes post mortem has not been reported on SSC 14. Given the moderate significance of our QTL, a replication study with independent animals would be desirable to confirm the results. Nonetheless, our findings offer a chance to unravel new QTL that contribute to the meat maturation.

The QTL for EBV pH1 detected in our study shows a largely additive effect (Fig. 2A) whereas the QTL for EBV carcass length and EBV rear view hind legs on SSC 10 indicate a recessive effect of the variant allele (Fig. 2B & C). This could be either due to coding variants or to regulatory variants that change the quantitative expression levels of the causative genes. On the other hand, the QTL for EBV rear view hind legs on SSC 16 does not show a simple additive effect (Fig. 2D). For this QTL, animals being heterozygous at the best associated SNP show more extreme phenotypic levels than animals with either homozygous genotype. It seems unlikely that such an overdominance effect can be caused by simple quantitative differences in mRNA expression of the underlying genes. One possible explanation would be provided by non-synonymous variants in genes encoding oligomeric proteins. In such a scenario, it is feasible that oligomers of truly identical proteins (in homozygous animals) have very different properties than oligomers of allelic variants (in heterozygous animals).

Our study provides another example of the usefulness of the PorcineSNP60 BeadChip for genome-wide association studies in pigs. This tool allows the detection of QTL for commercially important traits in pigs. However, it must also be noted that due to the imperfect pig reference genome assembly, the exact genome positions of many markers on this tool are not clear. We observed

very significant shifts of some associated markers between the Sscrofa9.2 and the Sscrofa10.2 assembly. Consequently, the marker spacing may also be expected to be somewhat irregular. Thus, it is quite possible that some real QTL might have been missed due to insufficient marker coverage of the PorcineSNP60 BeadChip. With rapidly increasing genomic resources for the pig, it may be expected that improved genotyping tools will soon become available.

In conclusion, we have mapped four QTL by genome-wide association mapping in Swiss commercial pigs. Three of these QTL coincide with previously detected QTL for similar traits in other independent pig populations while the QTL for EBV pH1 on SSC 14 is described for the first time.

## Materials and Methods

### Animals and phenotypic data

We obtained previously archived tissue samples of 192 artificial insemination boars from the commercial Premo<sup>®</sup> line, which is derived from the Swiss Large White breed. The animals were from the breeding company SUISAG ([www.suisag.ch](http://www.suisag.ch)) and born between 2004 and 2009. All animals underwent a performance and progeny test according to the test scheme of SUISAG [34]. We obtained 26 corresponding estimated breeding values (EBV) for different pig production traits and used them as phenotypes in our analysis (Table S1). The EBVs were routinely estimated by SUISAG in 3 separate analyses using multiple trait animal models and BLUP [35]. For exterior traits linear description scores by trained technicians on animals tested in the central testing station Sempach and on-farm tested selection candidates were considered. The trait rear view hind legs describes the hind legs from an extreme X- (score 1) to an extreme O- form (score 7). Carcass length was measured in centimeters from the cranial edge of the first cervical vertebra to the cranial edge of the pelvic bone 24 h post mortem in all station tested pigs. The estimation for production traits of field tested crossbred progeny of artificial insemination sires were included in addition to the animals scored for exterior traits. pH1 was measured 1 hour post mortem at the *musculus longissimus dorsi* in all station tested pigs. Mating and litter records obtained from herdbook farms were used for the estimation of breeding values for reproduction traits. There were less animals with EBVs on reproduction traits, as artificial insemination sires need litter records of daughters to reach an acceptable accuracy.

### SNP array genotyping

We isolated genomic DNA from tissue samples with the Nucleon Bacc2 kit (GE Healthcare) according to the manufacturer's protocol. DNA samples with a ratio of A260/280 higher than 1.8 and a concentration of approximately 50 ng/μl were genotyped at the Leibniz Institute for Farm Animal Biology, Dummerstorf, Germany using the Illumina Porcine60SNP BeadChip containing 62,163 markers. We used BLASTN to determine the positions of the markers in the Sscrofa 10.2 genome reference assembly of the pig.

### Quality control and genome-wide association analyses

We analyzed the data with the GenABEL package [36] in the R environment. Initially, we removed all individuals with a call rate <95% and checked the dataset for replicates and gender mismatch. We excluded markers strongly deviating from Hardy-Weinberg equilibrium ( $p < 0.0001$ ), markers having a call rate <95%, and markers with a minor allele frequency of <5%. After

these quality control steps 186 individuals and 47,045 SNPs remained for the analysis.

We calculated genome-wide pairwise identity-by-state (IBS) distances to measure population stratification as we used samples of animals with expected diverse genetics. Based on genetic distances between individuals projected into two-dimensional space using multidimensional scaling we grouped the animals into three subpopulations. Additionally, the procedure eggscore which uses principal component analysis [37] was used to correct for population stratification in the dataset. We performed allelic genome-wide association analyses for all 26 EBVs. We considered  $p$ -values  $< 5 \times 10^{-7}$  as indicative for strong evidence of association and  $p$ -values between  $5 \times 10^{-5}$  and  $5 \times 10^{-7}$  as indicative for moderate evidence of association according to the recommendation of the Wellcome Trust Case Control Consortium [38].

## Supporting Information

**Figure S1 Multidimensional scaling (MDS) plot showing the genomic kinship between the analyzed animals.** This plot visualizes the overall genetic distances between the boars

## References

- Sonesson AK, de Greef KH, Meuwissen THE (1998) Genetic parameters and trends of meat quality, carcass composition and performance traits in two selected lines of large white pigs. *Livest Prod Sci* 57: 23–32.
- van Wijk HJ, Arts DJ, Matthews JO, Webster M, Ducro BJ, et al. (2005) Genetic parameters for carcass composition and pork quality estimated in a commercial production chain. *J Anim Sci* 83: 324–333.
- Hermesch S, Luxford BG, Graser HU (2000) Genetic Parameters for lean meat yield, meat quality, reproduction and feed efficiency traits for Australian pigs I. Description of traits and heritability estimates. *Livest Prod Sci* 65: 239–248.
- Andersson L, Haley CS, Ellegren H, Knott SA, Johansson M, et al. (1994) Genetic-mapping of quantitative trait loci for growth and fatness in pigs. *Science* 263: 1771–1774.
- Karlskov-Mortensen P, Bruun CS, Braunschweig MH, Sawera M, Markljung E, et al. (2006) Genome-wide identification of quantitative trait loci in a cross between Hampshire and Landrace I: carcass traits. *Anim Genet* 37: 156–162.
- de Koning DJ, Rattink AP, Harlizius B, Groenen MAM, Brascamp EW, et al. (2001) Detection and characterization of quantitative trait loci for growth and reproduction traits in pigs. *Livest Prod Sci* 72: 185–198.
- Rohrer GA, Keele JW (1998) Identification of quantitative trait loci affecting carcass composition in swine: II. Muscling and wholesale product yield traits. *J Anim Sci* 76: 2255–2262.
- Andersson-Eklund L, Marklund L, Lundström K, Haley CS, Andersson K, et al. (1998) Mapping Quantitative Trait Loci for carcass and meat quality traits in a Wild Boar × Large White intercross. *J Anim Sci* 76: 694–700.
- Geldermann H, Müller E, Moser G, Reiner G, Bartenschlager H, et al. (2003) Genome-wide linkage and QTL mapping in porcine F2 families generated from Pietrain, Meishan and Wild boar crosses. *J Anim Breed Genet* 120: 363–393.
- Knott SA, Marklund L, Haley CS, Andersson K, Davies W, et al. (1998) Multiple marker mapping of quantitative trait loci in a cross between outbred wild boar and large white pigs. *Genetics* 149: 1069–1080.
- Nezer C, Moreau L, Wagenaar D, Georges M (2002) Results of a whole genome scan targeting QTL for growth and carcass traits in a Pietrain × Large White intercross. *Genet Sel Evol* 34: 371–387.
- Malek M, Dekkers JCM, Lee HK, Baas TJ, Rothschild MF (2001) A molecular genome scan analysis to identify chromosomal regions influencing economic traits in the pig. I. Growth and body composition. *Mamm Genome* 12: 630–636.
- van Laere AS, Nguyen M, Braunschweig M, Nezer C, Colette C, et al. (2003) A regulatory mutation in IGF2 causes a major QTL effect on muscle growth in the pig. *Nature* 425: 832–836.
- Milan D, Jeon JT, Looft C, Amarger V, Robic A, et al. (2000) A mutation in PRKAG3 associated with excess glycogen content in pig skeletal muscle. *Science* 288: 1248–1251.
- Glazier AM, Nadeau JH, Aitman TJ (2002) Finding genes that underlie complex traits. *Science* 298: 2345–2349.
- Ramos AM, Crooijmans RPMA, Affara NA, Amaral AJ, Archibald AL, et al. (2009) Design of a High Density SNP Genotyping Assay in the Pig Using SNPs Identified and Characterized by Next Generation Sequencing Technology. *PLoS ONE* 4: e6524.
- Potkin SG, Guffanti G, Lakatos A, Turner JA, Kruggel F, et al. (2009) Hippocampal Atrophy as a Quantitative Trait in a Genome-Wide Association Study Identifying Novel Susceptibility Genes for Alzheimer's Disease. *PLoS ONE* 4: e6501.
- Estrada K, Krawczak M, Schreiber S, van Duijn K, Stolk L, et al. (2009) A genome-wide association study of northwestern Europeans involves the C-type natriuretic peptide signaling pathway in the etiology of human height variation. *Hum Mol Genet* 18: 3516–3524.
- Sutter NB, Bustamante CD, Chase K, Gray MM, Zhao K, et al. (2007) A single IGF1 allele is a major determinant of small size in dogs. *Science* 316: 112–115.
- Bolormaa S, Porto Neto LR, Zhang YD, Bunch RJ, Harrison BE, et al. (2011) A genome-wide association study of meat and carcass traits in Australian cattle. *J Anim Sci* 89: 2297–2309.
- Onteru SK, Fan B, Du Z-Q, Garrick DJ, Stalder KJ, et al. (2011) A whole-genome association study for pig reproductive traits. *Anim Genet* 43: 18–26.
- Duijvesteijn N, Knol EF, Merks JW, Crooijmans RP, Groenen MA, et al. (2010) A genome-wide association study on androstene levels in pigs reveals a cluster of candidate genes on chromosome 6. *BMC Genet* 11: 42.
- Uimari P, Sironen A, Sevón-Aimonen M-L (2011) Whole-genome SNP association analysis of reproduction traits in the Finnish Landrace pig breed. *Genet Sel Evol* 43: 42.
- Gregersen VR, Conley LN, Sorensen KK, Gulbrandsen B, Velandier IH, et al. (2012) Genome-wide association scan and phased haplotype construction for quantitative trait loci affecting boar taint in three pig breeds. *BMC Genomics* 13: 22.
- Wang JY, Luo YR, Fu WX, Lu X, Zhou JP, et al. (2012) Genome-wide association studies for hematological traits in swine. *Anim Genet* doi: 10.1111/j.1365-2052.2012.02366.x.
- Schneider JF, Rempel LA, Snelling WM, Wiedmann RT, Nonneman DJ, et al. (2012) Genome-wide association study of swine farrowing traits. Part II: Bayesian analysis of marker data. *J Anim Sci* 90: 3360–3367.
- Wilson K, Zanella R, Ventura C, Johansen HL, Framstad T, et al. (2012) Identification of chromosomal locations associated with tail biting and being a victim of tail-biting behaviour in the domestic pig (*Sus scrofa domestica*). *J Appl Genet* 53: 449–456.
- Fontanesi L, Schiavo G, Galimberti G, Calò DG, Scotti E, et al. (2012) A genome wide association study for backfat thickness in Italian Large White pigs highlights new regions affecting fat deposition including neuronal genes. *BMC Genomics* 13: 583.
- Lee SS, Chen Y, Moran C, Cepica S, Reiner G, et al. (2003) Linkage and QTL mapping for *Sus scrofa* chromosome 2. *J Anim Breed Genet* 120: 11–19.
- Evans GJ, Giuffra E, Sanchez A, Kerje S, Davalos G, et al. (2003) Identification of Quantitative Trait Loci for Production Traits in Commercial Pig Populations. *Genetics* 164: 621–627.
- Fan B, Onteru SK, Du Z-Q, Garrick DJ, Stalder KJ, et al. (2011) Genome-wide association study identifies loci for body composition and structural soundness traits in pigs. *PLoS One* 6: e14726.
- Uemoto Y, Sato S, Ohnishi C, Hirose K, Kameyama K, et al. (2010) Quantitative trait loci for leg weakness traits in a Landrace purebred population. *Anim Sci J* 81: 28–33.
- Lee GJ, Archibald AL, Garth GB, Law AS, Nicholson D, et al. (2003) Detection of quantitative trait loci for locomotion and osteochondrosis-related traits in Large White × Meishan pigs. *Anim Sci* 76: 155–165.
- Homepage SUIISAG (2012) Available: <http://www.suisag.ch/Zucht/Leistungspr%C3%BCfung/Endproduktepr%C3%BCfung/tabid/109/Default.aspx>. Accessed 18 July 2012.

35. Henderson CR (1975) Best linear unbiased Estimation and Prediction under a selection model. *Biometrics* 31: 423–447.
36. Aulchenko YS, Ripke S, Isaacs A, van Duijn CM (2007) GenABEL: an R library for genome-wide association analysis. *Bioinformatics* 23: 1294–1296.
37. Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, et al. (2006) Principal component analysis corrects for stratification in genome-wide association studies. *Nat Genet* 38: 904–909.
38. The Wellcome Trust Case Control Consortium (2007) Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* 447: 661–678.