

University of Nebraska - Lincoln
DigitalCommons@University of Nebraska - Lincoln

Dissertations, Theses, and Student Research Papers
in Mathematics

Mathematics, Department of

8-2015

Systems of parameters and the Cohen-Macaulay property

Katharine Shultis

University of Nebraska-Lincoln, s-kshulti1@math.unl.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/mathstudent>



Part of the [Algebra Commons](#)

Shultis, Katharine, "Systems of parameters and the Cohen-Macaulay property" (2015). *Dissertations, Theses, and Student Research Papers in Mathematics*. 66.

<http://digitalcommons.unl.edu/mathstudent/66>

This Article is brought to you for free and open access by the Mathematics, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Dissertations, Theses, and Student Research Papers in Mathematics by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

SYSTEMS OF PARAMETERS AND THE COHEN-MACAULAY PROPERTY

by

Katharine Shultis

A DISSERTATION

Presented to the Faculty of

The Graduate College at the University of Nebraska

In Partial Fulfilment of Requirements

For the Degree of Doctor of Philosophy

Major: Mathematics

Under the Supervision of Srikanth B. Iyengar and Roger Wiegand

Lincoln, Nebraska

August, 2015

SYSTEMS OF PARAMETERS AND THE COHEN-MACAULAY PROPERTY

Katharine Shultis, Ph.D.

University of Nebraska, 2015

Advisers: Srikanth B. Iyengar and Roger Wiegand

Let R be a commutative, Noetherian, local ring and M a finitely generated R -module. Consider the module of homomorphisms $\text{Hom}_R(R/\mathfrak{a}, M/\mathfrak{b}M)$ where $\mathfrak{b} \subseteq \mathfrak{a}$ are parameter ideals of M . When $M = R$ and R is Cohen-Macaulay, Rees showed that this module of homomorphisms is isomorphic to R/\mathfrak{a} , and in particular, a free module over R/\mathfrak{a} of rank one. In this work, we study the structure of such modules of homomorphisms for a not necessarily Cohen-Macaulay R -module M .

ACKNOWLEDGMENTS

The completion of this dissertation has been a long time coming, and would not have been possible without the support of my mentors, family, and friends. I would like to begin with my advisors: Srikanth Iyengar and Roger Wiegand. When I decided to come to Nebraska to study commutative algebra, I knew that I would find a good advisor who would help me muddle my way through my Ph.D. Instead, I am thrilled that I found two fantastic advisors who have helped me not only finish, but thrive along the way.

Sri has encouraged me both when I feel frustrated and when I feel excited by my research, and I cannot determine which encouragement has been more important. His patience has helped me through when I feel stuck and helpless, and his brilliance has pushed me to be a better mathematician. One of my favorite things about Sri is that when he agreed to be my advisor, the first thing he wanted to know was what my goals were. Thank you, Sri, for letting me know early on that your goal was to help me find my dream job.

When Roger told me he would be “honored” to be my second advisor, I felt like he had it backwards: I was honored that he would work with me. Having Roger as a second advisor has helped my research to be stronger. Knowing that he will regularly ask me if I have “found any new eigenvalues” pushes me to work harder. In addition to helping me with mathematics, Roger has also helped me with my writing. Thank you, Roger, for reading my work carefully and teaching me to be a better writer.

Thank you to the rest of my committee, Brian Harbourne, Wenliang Zheng, and Byrav Ramamurthy, and to my teaching mentor, Jim Lewis, for all of your help, advice, and support. In particular, I would like to thank Brian for his careful reading of this dissertation and his willingness to step into a larger advising role when Sri left

UNL. I also want to particularly thank Jim for laughing in just the right way when I was feeling insecure about my teaching ability and wanted to know if he was going to write me a good letter or if I should ask someone else.

My mentors before I arrived in Nebraska helped pave the way for me to get to this point. In particular, I appreciate the advice of Lance Small who told me that I should “move to Nebraska, buy a winter coat, and learn that all rings are commutative” when I suddenly discovered that I needed a new plan for graduate school. I can no longer tell you if these are his exact words, but this was exactly the piece of advice I needed to help me move forward when it seemed like I would fail.

Christopher Towse should probably be credited with my choice to study mathematics. As a senior in high school I visited Scripps College as part of a scholarship interviewing process and met with Chris. His enthusiasm for mathematics was contagious and he explained to me that unique factorization is not something that should be taken for granted. Learning that 36 factors into irreducible elements in $2\mathbb{Z}$ as both $6 \cdot 6$ and as $2 \cdot 18$ has stuck with me for the last twelve years and that conversation with Chris may have been the moment I should have realized I was a mathematician.

My family has always been my strongest cheering squad. If I need a place to vent, cry, or even gloat, my parents are always just a phone call away. They listen carefully to my woes, are excited for me when I succeed, and provide not only a great sounding board but also suggestions when I need help figuring out what to do. They “walk me home” when it is dark, cold, or even just lonely. My brother is proud of me when I succeed and has wanted to celebrate my successes even more than his own birthday! A girl couldn’t ask for a better family.

My friends have succeeded at the sometimes incredibly hard job of keeping my spirits and hope up over the past eight years of graduate school. After I pushed my way into their office for our second year, my officemates Sarah Behrens and Caitlyn

Parmelee have somehow managed to not only put up with my particular brand of crazy but also to learn to love it. I appreciate all the email reviewing, snowflake cutting, “family dinners”, and crazy outings. Most importantly, “I like it when we’re friends!”

Thank you to Lauren Keough for the many margarita nights and for inviting me to trivia when I was certain I had failed my comp. Weekly trivia outings have kept me sane by providing a fun mid-week break for the past several years. Thank you to Marilyn Johnson for not only helping me navigate all the paperwork involved in having three advisors, none of whom are in the state of Nebraska, but also for the advice and support over the past five years.

Despite never fully understanding my love of mathematics or my desire to earn a Ph.D., my non-mathematician friends have been encouraging and supportive of this graduate school adventure. Kristen Wright and Karen Liberty have been important to me for over half my life. I am trying to decide if that means I should refer to them as staples or fixtures in my life, but both sound equally silly. My friendships with my Tinarina, Carrie Barrie, and Kristen Poo (Christina Lindal, Carrie Tanner, and Kristen Kurimoto if you prefer their formal names) have stayed strong despite the fact that after graduating from Scripps, we haven’t lived in the same building and are now several states away. I am super excited to be living close to you three again!

Before I stop naming people, I want to thank two more members of my mathematical family by name. Ryan Karr, Roger’s eleventh student, overheard me asking Roger a question at Bottle Smashing last year, and promptly took it upon himself to come up with a counterexample to my claim. For pointing out that the ring in Example 2.10 has positive depth and a system of parameters consisting only of zerodivisors, I am extremely grateful to Ryan. I had spent too many hours trying to prove that a ring with a system of parameters consisting only of zerodivisors must

have zero depth.

Learning that Frank Moore, Sri's younger mathematical brother, knows how to work magic with Macaulay2, I promptly asked him if he knew how to use the software program to determine whether or not a module is decomposable. After sending him many harassing emails, he finally agreed to work with me on this problem. I showed Frank my painstaking computations and he played the game "name that tune in five lines or less." While it is more than five lines of code, the result of that lunchtime conversation is Remark 3.5 and I am very grateful to Frank for its existence.

I would also like to thank all of the faculty and graduate students who overlapped with me at UNL. Shortly after moving to Nebraska, I told people that "despite the weather, I'm a thousand times happier in Nebraska than I was in San Diego." This was, and is, because of the strong community of support in the UNL mathematics department. To all of my other mentors, mathematical family, biological family, and friends: I apologize that your names do not appear here. Please know that I appreciate your help, advice and support and that I would write you a story telling you what I appreciate most about you, if only I had infinite time and space. And you, dear reader of these obscenely long acknowledgements, thank you for your support!

GRANT INFORMATION

This research was partially supported by National Science Foundation (NSF) grants DMS-1201889 and DMS-1503044 and U.S. Department of Education grant P200A120068 (GAANN).

Contents

Contents	viii
1 Introduction	1
1.1 Dimension and Systems of Parameters	2
1.2 Depth	6
1.3 The I -torsion Functor	7
1.4 Cohen-Macaulay Modules	9
1.5 Free Modules	10
1.6 Rees' Theorem	10
1.7 Main Results	12
2 Proof of Results	15
2.1 Preliminary Results	15
2.2 Dimension One	24
2.3 Higher Dimensions	29
3 Examples	33
Bibliography	46

Chapter 1

Introduction

This thesis makes a contribution to the study of systems of parameters of local rings and modules over local rings. The module of study is well understood in the Cohen-Macaulay case, and so our focus is on the non-Cohen-Macaulay case. In this thesis *rings* are assumed to be commutative. Throughout, a *local ring* is a Noetherian ring with a unique maximal ideal. The notation (R, \mathfrak{m}) indicates that R is a local ring whose maximal ideal is \mathfrak{m} .

In this chapter, we provide background material. The notion of a Cohen-Macaulay module centers on two key invariants of the module: the dimension and the depth. In Section 1.1 we will focus on the dimension of a module and its systems of parameters. In Section 1.2 we will define the depth of a module and related terms. In Section 1.3 we will define the I -torsion functor and state some basic properties of the functor. In Section 1.4 we will define Cohen-Macaulay modules and indicate some relationships among the notions defined in the first two sections when a module is Cohen-Macaulay. In Section 1.5 we discuss the basics of free modules. In Section 1.6 we will discuss the historical background necessary for understanding where the main results fit into the larger picture. Finally, in Section 1.7 we will state the main results.

1.1 Dimension and Systems of Parameters

A *prime ideal* of a ring R is a proper ideal \mathfrak{p} such that R/\mathfrak{p} is an integral domain. An equivalent characterization is that \mathfrak{p} is prime if whenever $a, b \in R$ with $ab \in \mathfrak{p}$, then $a \in \mathfrak{p}$ or $b \in \mathfrak{p}$. A chain of prime ideals

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n$$

is said to have *length* n . The supremum of the lengths of chains of primes in a ring R is called the *dimension of R* , written $\dim R$.

Example 1.1. Consider the ring $R = k[[x, y]]/(x^2, xy^m)$ for $m \geq 1$. Since $R/(x) \cong k[[y]]$ and $R/(x, y) \cong k$ are integral domains and $(x) \subsetneq (x, y)$, we know that the dimension of R is at least one. To show that the dimension is at most one, we will use the notion of a system of parameters.

For a ring R , recall that the *radical* of an ideal $I \subseteq R$ is the ideal

$$\sqrt{I} := \{a \in R \mid a^n \in I \text{ for some } n \in \mathbb{N}\}.$$

If $\sqrt{I} = \mathfrak{m}$ is a maximal ideal, we say that I is *\mathfrak{m} -primary*.

Let (R, \mathfrak{m}) be a local ring of dimension d . There exist sets of d elements which generate \mathfrak{m} -primary ideals, but no ideal generated by fewer than d elements is \mathfrak{m} -primary. For a proof of this fact, see [7, Theorem 13.4]. By Krull's Height Theorem [7, Theorem 13.5], the height of an ideal generated by r elements is no more than r , so no ideal generated by fewer than d elements is \mathfrak{m} -primary. A set of d elements generating an \mathfrak{m} -primary ideal is called a *system of parameters*. An ideal generated by a system of parameters is called a *parameter ideal*. We sometimes refer to an element of a

system of parameters as a *parameter*.

Example 1.2. With $R = k[[x, y]]/(x^2, xy^m)$ as in Example 1.1, consider the ideal (y) . Since $x^2 = 0 \in (y)$, both x and y are in $\sqrt{(y)}$ so that $\mathfrak{m} = (x, y) \subseteq \sqrt{(y)}$. Thus $\mathfrak{m} = \sqrt{(y)}$, and (y) is an \mathfrak{m} -primary ideal. In Example 1.1 we saw $\dim R \geq 1$. Thus R has dimension one and y is a parameter of R .

We now extend the notions of dimension and system of parameters to modules. Let R be a ring and M an R -module. The *dimension* of M , denoted $\dim_R M$, can be defined to be the dimension of the ring $R/\text{ann}_R(M)$ where

$$\text{ann}_R(M) := \{r \in R \mid rM = 0\}$$

is the *annihilator* of M . Since $\text{ann}_R(R) = (0)$ we recover the original definition when the module is the ring itself. More generally, the dimension of the R -module R/I is equal to the dimension of R/I as a ring.

Example 1.3. With $R = k[[x, y]]/(x^2, xy^m)$ as in Example 1.1 and Example 1.2, consider the R -module $M = y^m R$. For $r \in R$, we know $ry^m = 0$ only if $r \in (x)$. As $y^m \in M$ this tells us that $\text{ann}_R(M) \subseteq (x)$. Since $xM = xy^m R = 0$, we see that $\text{ann}_R(M) = \{r \in R \mid r(y^m R) = 0\} = (x)$. We can now compute the dimension of M . Since $R/(x) \cong k[[y]]$, we obtain

$$\dim_R M = \dim(R/\text{ann}_R(M)) = \dim(R/(x)) = \dim(k[[y]]) = 1.$$

Next we introduce systems of parameters for modules. Let R be a local ring and M an R -module of dimension d . There exist sets of d elements $a_1, \dots, a_d \in R$ such that $M/(a_1, \dots, a_d)M$ has finite length, that is, such that $M/(a_1, \dots, a_d)M$ is

both an Artinian and a Noetherian R -module. However for any $a_1, \dots, a_n \in R$ with $0 \leq n < d$, the module $M/(a_1, \dots, a_n)M$ has infinite length. For a proof of this fact, see [7, Theorem 13.4].

A set of d elements $a_1, \dots, a_d \in R$ with the property that $M/(a_1, \dots, a_d)M$ has finite length is called a *system of parameters of M* . As in the ring case, an ideal generated by a system of parameters is called a *parameter ideal of M* , and we sometimes refer to an element of a system of parameters as a *parameter*. The next proposition says that this definition agrees with the earlier one when $M = R$.

Proposition 1.4. *Let (R, \mathfrak{m}) be a local ring and I any ideal of R . The R -module R/I has finite length if and only if the ideal I is \mathfrak{m} -primary.*

Proof. Recall [1, Proposition 1.14]: The radical of an ideal is the intersection of the prime ideals that contain it. Also, if $A \rightarrow B$ is a homomorphism of rings and L is a B -module, then the length of L as an A -module is the same as its length as a B -module. Indeed, the A -submodules of L are precisely the B -submodules of L .

As R is Noetherian, so is R/I . Thus R/I has finite length if and only if it is Artinian. Since R is Noetherian, we know R/I is Artinian if and only if R/I has dimension zero [1, Theorem 8.5]. Moreover, the ring R/I has dimension zero if and only if the maximal ideal of R is the only prime ideal which contains I , which is equivalent to $\mathfrak{m} = \sqrt{I}$ by [1, Proposition 1.14]. \square

Example 1.5. With $R = k[[x, y]]/(x^2, xy^m)$ and $M = y^m R$ as in Example 1.3, we claim that $\{y\}$ is a system of parameters for M , and moreover, $M/yM = (y^m)/(y^{m+1})$ has length one. Indeed, the map

$$R/(x, y) \rightarrow (y^m)/(y^{m+1})$$

given by $1 + (x, y) \mapsto y^m + (y^{m+1})$ is R -linear and bijective. Thus M/yM is a simple R -module, M/yM has length one, and y is a parameter of M .

The next proposition is useful for the induction technique used in the proofs of the main results.

Proposition 1.6. *Let R be a commutative, local, Noetherian ring and M an R -module. If $\{a_1, \dots, a_i\}$ is part of a system of parameters of M , then*

$$\dim_R(M/(a_1, \dots, a_i)M) = \dim_R(M) - i.$$

Proof. Set $\overline{M} = M/a_1M$, $d = \dim_R(M)$, and $\delta = \dim_R(\overline{M})$. We proceed by induction on i . Since a_1 is a parameter of M , we can find elements $x_2, \dots, x_d \in R$ such that $M/(a_1, x_2, \dots, x_d)M$ has finite length. Note that

$$\overline{M}/(x_2, \dots, x_d)\overline{M} \cong M/(a_1, x_2, \dots, x_d)M,$$

which has finite length. Hence $\dim_R(\overline{M}) \leq \dim_R(M) - 1$. For the other inequality, let b_2, \dots, b_δ be a system of parameters of \overline{M} . Then

$$M/(a_1, b_2, \dots, b_\delta)M \cong \overline{M}/(b_2, \dots, b_\delta)\overline{M}$$

has finite length, and it follows that $\dim_R(M) \leq \dim_R(\overline{M}) + 1$. This completes the proof for the case $i = 1$.

For $i > 1$, note that $\{a_2, \dots, a_i\}$ is part of a system of parameters of \overline{M} . The

induction hypothesis then gives

$$\begin{aligned} \dim_R(\overline{M}/(a_2, \dots, a_i)\overline{M}) &= \dim_R(\overline{M}) - (i - 1) \\ &= \dim_R(M) - 1 - (i - 1) \\ &= \dim_R(M) - i. \end{aligned}$$

This is the desired result, since $\overline{M}/(a_2, \dots, a_i)\overline{M} \cong M/(a_1, \dots, a_i)M$. \square

1.2 Depth

Depth is another important invariant of a ring and its modules. Roughly speaking, depth measures how many independent elements of the ring behave like indeterminates. An element of a ring R is called a *regular element* on a module M if it is not a zero-divisor on M . That is, $r \in R$ is a regular element on M if $rm \neq 0$ for any nonzero $m \in M$.

A sequence $r_1, \dots, r_n \in R$ is called a *weak regular sequence on M* if r_i is a regular element on $M/(r_1, \dots, r_{i-1})M$ for all $i = 1, \dots, n$. We take the ideal generated by the empty set to be the zero ideal, so the condition for $i = 1$ is that r_1 is a regular element on $M/(0)M = M$. A sequence $r_1, \dots, r_n \in R$ is a *regular sequence on M* if it is a weak regular sequence on M and, in addition, $M/(r_1, \dots, r_n)M \neq 0$. A *maximal regular sequence* is a regular sequence that cannot be extended to a longer one.

All maximal regular sequences which are contained in a fixed ideal have the same length. See [7, Theorem 16.7] for a proof of this fact. For a local ring (R, \mathfrak{m}) , the *depth* of an R -module M is defined to be the length of any maximal regular sequence contained in \mathfrak{m} .

Example 1.7. With $R = k[[x, y]]/(x^2, xy^m)$ and $M = y^m R$, as in Example 1.3 and Example 1.5, we claim that R has depth zero and M has depth one.

First suppose $p \in R$ is a non-unit. Then $p = ax + by$ for some $a, b \in R$ and $pxy^{m-1} = ax^2y^{m-1} + bxy^m = 0$ in R . Thus the only regular elements of R are units; however, if $p \in R$ is a unit, then $R/(p) = (0)$. Hence, there are no regular sequences on R of length one and R has depth zero.

Next note that the element $y \in R$ is a regular element on M . Moreover, $M/yM = (y^m)/(y^{m+1}) \neq 0$, and so y is a regular sequence on M . We claim that y is a maximal regular sequence. To see this directly, let $r \in R$. If r is a non-unit we may write $r = ax + by$ for some $a, b \in R$. Then $ry^m = axy^m + by^{m+1} \in (y^{m+1}) = yM$. Thus r is not a regular element on $(y^m)/(y^{m+1}) = M/yM$. If $r \in R$ is a unit, then $(r, y) = R$, so $M/(r, y)M = (0)$. Thus y is a maximal length regular sequence on M and M has depth one.

1.3 The I -torsion Functor

Let R be a ring, $I \subseteq R$ an ideal, M an R -module, and $N \subseteq M$ a submodule. We define $(N :_M I)$ to be the set

$$(N :_M I) := \{m \in M \mid mI \subseteq N\}.$$

This is a submodule of M . When M is clear from context, we sometimes write $(N : I)$ instead of $(N :_M I)$. When $I = (a)$ is a principal ideal, we typically write $(N :_M a)$ instead of $(N :_M (a))$.

Let R be a commutative Noetherian ring, $I \subseteq R$ an ideal, and M an R -module.

The submodule $\Gamma_I(M)$ of M is defined as

$$\Gamma_I(M) := \bigcup_{i=1}^{\infty} (0 :_M I^i).$$

For any R -linear map $f : M \rightarrow N$, we define $\Gamma_I(f) : \Gamma_I(M) \rightarrow \Gamma_I(N)$ to be the map induced by f . This makes $\Gamma_I(-)$ a covariant functor from the category of R -modules to itself.

Example 1.8. Consider the ring $R = k[[x, y]]/(x^2, xy^m)$ and module $M = y^m R$ from Example 1.7. The ring R is local with maximal ideal $\mathfrak{m} = (x, y)$. For $i \geq m$ it is clear that $\mathfrak{m}^i = (y^i)$ since all other degree i monomials in x and y are zero. It is also easy to see that $(0 :_R \mathfrak{m}^i) = (0 :_R y^i) = (x)$ for $i \geq m$ and $(0 :_M \mathfrak{m}^i) = (0 :_{y^m R} y^i) = (0)$ for $i \geq m$. Thus $\Gamma_{\mathfrak{m}}(R) = (x)$ and $\Gamma_{\mathfrak{m}}(M) = (0)$.

It is straightforward to verify the following properties of the I -torsion functor.

Proposition 1.9. *Let R be a Noetherian ring, $I \subseteq R$ an ideal, and M a finitely generated R -module.*

- (a) *The I -torsion functor, $\Gamma_I(-)$, is left exact; [5, Exercise 7.2].*
- (b) *If J is an ideal of R with $\sqrt{J} = \sqrt{I}$, then $\Gamma_J(M) \cong \Gamma_I(M)$; [5, Proposition 7.3].*

Now suppose R is a local ring with maximal ideal \mathfrak{m} .

- (c) *M has depth zero if and only if $\Gamma_{\mathfrak{m}}(M) \neq 0$; [5, Remark 9.4].*
- (d) *$\Gamma_{\mathfrak{m}}(M)$ is Artinian; [5, Exercise 7.7].*

1.4 Cohen-Macaulay Modules

Let R be a ring. If M is a nonzero R -module, then there is an inequality:

$$\text{depth}_R(M) \leq \dim_R(M);$$

see [3, Proposition 1.2.12]. In the extremal case when the depth and dimension are equal, or when $M = 0$, we say that M is *Cohen-Macaulay*. We say a local ring R is *Cohen-Macaulay* if it is Cohen-Macaulay as a module over itself.

Example 1.10. In Example 1.3, Example 1.5 and Example 1.7, we saw that $R = k[[x, y]]/(x^2, xy^m)$ has dimension one but depth zero and $M = y^m R$ has dimension and depth both equal to one. Thus R is not a Cohen-Macaulay ring, but M is a Cohen-Macaulay module.

Example 1.11. Consider the ring $R = k[[x, y]]/(x^2)$. It is easy to see that $(x) \subsetneq (x, y)$ is a chain of prime ideals of R and $\sqrt{(y)} = (x, y) = \mathfrak{m}$. Thus R has dimension one and $\{y\}$ is a system of parameters of R . To see that R has depth one, we need only produce a regular element of R which is not a unit, since the depth cannot exceed the dimension. It is clear that y is such a regular element and R is a Cohen-Macaulay ring.

In the ring $R = k[[x, y]]/(x^2, xy^m)$, the parameter y is not a regular element as R has depth zero. However, in the ring $R = k[[x, y]]/(x^2)$, the parameter y forms a regular sequence on R . More generally, for any Cohen-Macaulay module M , the elements $a_1, \dots, a_i \in R$ are part of a system of parameters of M if and only if a_1, \dots, a_i is a regular sequence on M [3, Theorem 2.1.2(d)].

1.5 Free Modules

Let R be a ring. An R -module F is said to be *free* if there is a linearly independent set of generators of F . That is, if there is a set of generators, $\{x_\alpha\}_{\alpha \in A}$, that satisfies the property: $\sum_{\alpha \in A} r_\alpha x_\alpha = 0$ if and only if $r_\alpha = 0$ for all $\alpha \in A$. We say a set of linearly independent generators is a *basis* of the free module F . If F is a nonzero free R -module, then $\text{ann}_R(F) = (0)$. Indeed, let $\{x_\alpha\}_{\alpha \in A}$ be a basis of the free module F . If $r \in R$ and $rF = 0$, then in particular $rx_\alpha = 0$ for all $\alpha \in A$. However, by linear independence of the basis, we know that $rx_\alpha = 0$ implies that $r = 0$.

Consider a commutative ring R , an ideal I , and an R/I -module M . Note that M is also an R -module via the action $r \cdot m := (r+I)m$ for any $r \in R$ and $m \in M$. Clearly $I \subseteq \text{ann}_R(M)$. If M is R/I -free then $a \in \text{ann}_R(M)$ also implies $a + I \in \text{ann}_{R/I}(M)$. Equivalently, if M is a free R/I -module and $a \in \text{ann}_R M$, then $a + I = 0$ in R/I , i.e. $a \in I$. Thus for any R/I -module M we know $\text{ann}_R(M) \supseteq I$, and we have equality when M is free.

However, there are non-free R/I -modules M such that $\text{ann}_R(M) = I$. For example, for any ring R and proper ideals $I \subsetneq J$ of R . The R/I -module $M = R/I \oplus R/J$ is not free, but $\text{ann}_R(M) = I$.

1.6 Rees' Theorem

The purpose of this section is to provide historical context in order to see how the main results fit into the larger picture. We begin by recalling Rees' Theorem [9, Theorem 2.1], as reformulated in the book by Bruns and Herzog [3, Lemma 1.2.4].

Theorem 1.12. *Let R be a ring, M, N be R -modules, and x_1, \dots, x_n elements of*

$\text{ann}_R(N)$ which form a weak regular sequence on M . Then

$$\text{Ext}_R^n(N, M) \cong \text{Hom}_R(N, M/(x_1, \dots, x_n)M).$$

The following proposition is a direct consequence of Rees' Theorem.

Proposition 1.13. *Let R be a local ring, M a Cohen-Macaulay R -module of dimension d and $\mathfrak{b} \subseteq \mathfrak{a}$ parameter ideals of M . Then*

$$\text{Hom}_R(R/\mathfrak{a}, M/\mathfrak{b}M) \cong M/\mathfrak{a}M.$$

Proof. Let \mathfrak{a} and \mathfrak{b} be generated by the systems of parameters $\{a_1, \dots, a_d\}$ and $\{b_1, \dots, b_d\}$ respectively. Since M is Cohen-Macaulay, both a_1, \dots, a_d and b_1, \dots, b_d are regular sequences on M [3, Theorem 2.1.2(d)]. Applying Theorem 1.12 to b_1, \dots, b_d and then to a_1, \dots, a_d yields the following isomorphisms:

$$\begin{aligned} \text{Hom}_R(R/\mathfrak{a}, M/\mathfrak{b}M) &\cong \text{Ext}_R^d(R/\mathfrak{a}, M) \\ &\cong \text{Hom}_R(R/\mathfrak{a}, M/\mathfrak{a}M) \\ &\cong \text{Hom}_{R/\mathfrak{a}}(R/\mathfrak{a}, M/\mathfrak{a}M) \\ &\cong M/\mathfrak{a}M. \end{aligned}$$

This is the desired result. □

When $M = R$ is Cohen-Macaulay and $\mathfrak{b} \subseteq \mathfrak{a}$ are parameter ideals of R , Proposition 1.13 says that we have the following isomorphism of R/\mathfrak{a} -modules:

$$\text{Hom}_R(R/\mathfrak{a}, R/\mathfrak{b}) \cong R/\mathfrak{a}.$$

That is, the module of homomorphisms is a free R/\mathfrak{a} -module of rank one, and hence is indecomposable. Recently, K. Bahmanpour and R. Naghipour [2, Theorem 2.4] proved the following converse of this statement.

Proposition 1.14. *If R is not Cohen-Macaulay there exist parameter ideals $\mathfrak{b} \subseteq \mathfrak{a}$ such that the R/\mathfrak{a} -modules $\text{Hom}_R(R/\mathfrak{a}, R/\mathfrak{b})$ and R/\mathfrak{a} are not isomorphic.*

The focus of this work is to study the structure of the module of homomorphisms $\text{Hom}_R(R/\mathfrak{a}, M/\mathfrak{b}M)$ when M is not Cohen-Macaulay and $\mathfrak{b} \subseteq \mathfrak{a}$ are parameter ideals. We focus on showing conditions under which $\text{Hom}_R(R/\mathfrak{a}, M/\mathfrak{b}M)$ is decomposable and conditions that imply $\text{Hom}_R(R/\mathfrak{a}, R/\mathfrak{b})$ is not a free R/\mathfrak{a} -module.

1.7 Main Results

In this section we state the main results and compare them to each other. This first result is only for modules of dimension one and depth zero. We show that the module $\text{Hom}_R(R/aR, M/bM)$ is decomposable if the parameter b is chosen to be a multiple of a sufficiently high power of the parameter a .

Theorem 1.15. *[10, Theorem 3.1] Let (R, \mathfrak{m}) be a local ring, and M a nonzero finitely generated R -module of dimension one and depth zero. Choose an integer n such that $\mathfrak{m}^n M \cap \Gamma_{\mathfrak{m}}(M) = (0)$. For any parameter a of M , and any parameter b of M with $b \in (a^{n+1})$, the following R -module is decomposable:*

$$\text{Hom}_R(R/aR, M/bM).$$

For higher dimensional modules, we also have a theorem to show that the module $\text{Hom}_R(R/\mathfrak{a}, M/\mathfrak{b}M)$ is decomposable; however, this result is weaker since it is not as

explicit as the result for modules of dimension one.

Theorem 1.16. [10, Theorem 4.1] *Let R be a local ring and M a finitely generated R -module of dimension d . If M is not Cohen-Macaulay, then, for any system of parameters $\mathbf{a} = a_1, \dots, a_d$ of M , there exist positive integers n_1, \dots, n_d such that the following R -module is decomposable:*

$$\mathrm{Hom}_R(R/(\mathbf{a}), M/(a_1^{n_1}, \dots, a_d^{n_d})M).$$

For a dimension one ring, we show that $\mathrm{Hom}_R(R/(a), R/(b))$ is decomposable and not a free $R/(a)$ -module if the parameter a is chosen to be in a sufficiently high power of the maximal ideal and the parameter b is a multiple of a^2 .

Theorem 1.17. [10, Theorem 3.3] *Let (R, \mathfrak{m}) be a local ring of dimension one and depth zero, and n an integer such that $\mathfrak{m}^n \cap \Gamma_{\mathfrak{m}}(R) = (0)$. For any parameter a in \mathfrak{m}^n and any parameter b in (a^2) , the $R/(a)$ -module*

$$\mathrm{Hom}_R(R/(a), R/(b))$$

is decomposable and has a non-free summand.

For higher dimensional rings, we also have a theorem to show that $\mathrm{Hom}_R(R/\mathbf{a}, R/\mathbf{b})$ is decomposable and is not R/\mathbf{a} -free; however, as was the case when comparing Theorem 1.15 to Theorem 1.16, this result is weaker as it is less explicit than the result for one dimensional rings.

Theorem 1.18. [10, Theorem 4.2] *Let R be a local ring of dimension d . If R is not Cohen-Macaulay, then for any system of parameters a_1, \dots, a_d of R , there exist integers $n_1, \dots, n_d, N_1, \dots, N_d \in \mathbb{N}$ with $N_i \geq n_i$ for $i = 1, \dots, d$ such that the*

$R/(a_1^{n_1}, \dots, a_d^{n_d})$ -module

$$\mathrm{Hom}_R(R/(a_1^{n_1}, \dots, a_d^{n_d}), R/(a_1^{N_1}, \dots, a_d^{N_d}))$$

is decomposable and has a non-free summand.

Theorem 1.15 and Theorem 1.17 are for one dimensional rings and modules respectively and are explicit in indicating when $\mathrm{Hom}_R(R/\mathfrak{a}, M/\mathfrak{b}M)$ is decomposable and when it has a non-free summand. Theorem 1.16 and Theorem 1.18 are similar, but for higher dimensional rings and modules.

In Chapter 2 we will state some preliminary results and prove the main results stated in this section. In Chapter 3 we will provide examples in order to explore some questions about the powers appearing in the main results.

Chapter 2

Proof of Results

In this chapter we present proofs of our main results. In Section 2.1 we present some preliminary results. Section 2.2 focuses on our results in dimension one. These results are stronger and more explicit than those in Section 2.3, which focuses on our results in higher dimensions.

2.1 Preliminary Results

In this section we include proofs of many of the results due to lack of adequate references. Recall the *support* of an R -module, M , is defined to be

$$\text{Supp}_R(M) := \{\mathfrak{p} \in \text{Spec } R \mid M_{\mathfrak{p}} \neq 0\}.$$

An R -module M is said to be *faithful* if $\text{ann}_R(M) = 0$. If M is a faithful R -module, then $\dim_R M = \dim R$. Indeed, by definition of faithful, $\text{ann}_R(M) = 0$. The next two results are well-known.

Lemma 2.1. *Let R be a ring, $I \subseteq R$ an ideal of R , and M a finitely generated, faithful R -module. Then $\text{Supp}_R(R/I) = \text{Supp}_R(M/IM)$.*

Proof. If \mathfrak{p} is a prime ideal of R such that $\mathfrak{p} \notin \text{Supp}_R(R/I)$, then $I_{\mathfrak{p}} = R_{\mathfrak{p}}$. Thus $I_{\mathfrak{p}}M_{\mathfrak{p}} = R_{\mathfrak{p}}M_{\mathfrak{p}} = M_{\mathfrak{p}}$ so that $\mathfrak{p} \notin \text{Supp}_R(M/IM)$.

For the other inclusion, suppose \mathfrak{p} is a prime ideal of R such that $\mathfrak{p} \notin \text{Supp}_R(M/IM)$. Then $M_{\mathfrak{p}} = I_{\mathfrak{p}}M_{\mathfrak{p}}$. Nakayama's Lemma [7, Theorem 2.2] gives the existence of an element $x \in R_{\mathfrak{p}}$ such that $xM_{\mathfrak{p}} = 0$ and $x-1 \in I_{\mathfrak{p}}$. Say $x-1 = \frac{a}{t}$ with $a \in I$ and $t \in R \setminus \mathfrak{p}$. Rewriting this gives $x = 1 + \frac{a}{t}$ or equivalently $xt = t + \frac{a}{1}$. As $xtM_{\mathfrak{p}} = t(xM_{\mathfrak{p}}) = 0$ then $(t + \frac{a}{1})M_{\mathfrak{p}} = 0$. As M is finitely generated, this means that there exists some $s \in R \setminus \mathfrak{p}$ such that $s(t+a)M = 0$. As M is a faithful R -module, $s(t+a) = 0 \in \mathfrak{p}$. Since $s \notin \mathfrak{p}$ this implies that $t+a \in \mathfrak{p}$. Hence $a \notin \mathfrak{p}$, since otherwise we would also have $t \in \mathfrak{p}$, a contradiction. We have thus found an element $a \in I \setminus \mathfrak{p}$ and so $(R/I)_{\mathfrak{p}} = 0$ and hence $\mathfrak{p} \notin \text{Supp}_R(R/I)$. \square

Lemma 2.2. *Let (R, \mathfrak{m}) be a local ring and M a nonzero finitely generated R -module. Then $\text{Supp}_R(M) = \{\mathfrak{m}\}$ if and only if M has finite length.*

Proof. First suppose that M has finite length. We'll show $\text{Supp}_R(M) = \{\mathfrak{m}\}$ by induction on the length of M . When M has length one, $0 \subsetneq M$ is a composition series. In other words, M is a simple R -module, so is isomorphic to R/\mathfrak{m} . Since $\text{Ass}_R(R/\mathfrak{m}) = \{\mathfrak{m}\}$, we know that $\text{Ass}_R(M) = \{\mathfrak{m}\}$. Since $\text{Ass}_R(M) \subseteq \text{Supp}_R(M)$ and their minimal elements coincide [7, Theorem 6.5], we have that $\text{Supp}_R(M) = \{\mathfrak{m}\}$ as well. Now suppose that M has length t and that every nonzero R -module of length less than t has \mathfrak{m} as the only element of its support. Let

$$0 = M_0 \subsetneq M_1 \subsetneq \cdots \subsetneq M_{t-1} \subsetneq M_t = M$$

be a composition series of M . As \mathfrak{m} is the only maximal ideal of R , we know that $M_i/M_{i-1} \cong R/\mathfrak{m}$ for each $i = 1, \dots, t$. Consider the short exact sequence

$$0 \rightarrow M_{t-1} \rightarrow M \rightarrow M/M_{t-1} \rightarrow 0.$$

We know that M_{t-1} has length $t - 1$ and so $\text{Supp}_R(M_{t-1}) = \{\mathfrak{m}\}$. Also, $M/M_{t-1} \cong R/\mathfrak{m}$ and so $\text{Supp}_R(M/M_{t-1}) = \{\mathfrak{m}\}$ as well. If $\mathfrak{p} \in \text{Supp}_R(M)$, either M_{t-1} or M/M_{t-1} is nonzero upon localization at \mathfrak{p} since localization is exact. Since this only holds for $\mathfrak{p} = \mathfrak{m}$, we know that $\text{Supp}_R(M) = \{\mathfrak{m}\}$ as claimed.

Now suppose that $\text{Supp}_R(M) = \{\mathfrak{m}\}$. This implies $\text{Ass}_R(M) = \{\mathfrak{m}\}$. We wish to show that M has finite length. Since $\mathfrak{m} \in \text{Ass}_R M$, we may choose a nonzero submodule M_1 of M with $M_1 \cong R/\mathfrak{m}$. Thus M has finite length submodules, and since M is Noetherian, there exists a submodule M' maximal with respect to having finite length. We wish to show $M' = M$. If not, then $M/M' \neq 0$ so that $\text{Supp}_R(M/M') = \{\mathfrak{m}\}$. As this implies $\mathfrak{m} \in \text{Ass}_R(M/M')$, there is an injection $R/\mathfrak{m} \hookrightarrow M/M'$. We thus have a submodule $M'' \subseteq M$ with $M'' \supsetneq M'$ and $M''/M' \cong R/\mathfrak{m}$. As the length of M'' is thus one more than the length of M' , we have arrived at a contradiction. Hence $M' = M$ and M has finite length. \square

The next result is also well known. We provide a proof for completeness.

Proposition 2.3. *Let a_1, \dots, a_d be elements of R and $\bar{a}_1, \dots, \bar{a}_d$ their images in $R/\text{ann}_R(M)$. Then $\{a_1, \dots, a_d\}$ is a system of parameters of M if and only if $\{\bar{a}_1, \dots, \bar{a}_d\}$ is a system of parameters of the ring $R/\text{ann}_R(M)$.*

Proof. Note $\{a_1, \dots, a_d\}$ is a system of parameters of M as an R -module if and only if $\{\bar{a}_1, \dots, \bar{a}_d\}$ is a system of parameters of M as an $R/\text{ann}_R(M)$ -module. We may

thus assume $\text{ann}_R(M) = 0$. The desired result is thus that $\{a_1, \dots, a_d\}$ is a system of parameters of M if and only if $\{a_1, \dots, a_d\}$ is a system of parameters of R .

Suppose $\{a_1, \dots, a_d\}$ is a system of parameters of M and write $\mathfrak{a} = (a_1, \dots, a_d)$, so $M/\mathfrak{a}M$ has finite length as an R -module. By Lemma 2.2 we know $\text{Supp}_R(M/\mathfrak{a}M) = \{\mathfrak{m}\}$. Lemma 2.1 then tells us that $\text{Supp}_R(R/\mathfrak{a}) = \{\mathfrak{m}\}$. By Lemma 2.2 we have that R/\mathfrak{a} has finite length and $\mathfrak{a} = (a_1, \dots, a_d)$ is a parameter ideal of R .

For the other direction, suppose $\{a_1, \dots, a_d\}$ is a system of parameters of R and write $\mathfrak{a} = (a_1, \dots, a_d)$ so that R/\mathfrak{a} has finite length. As the length of R/\mathfrak{a} as an R -module is the same as its length as an R/\mathfrak{a} -module, we know that R/\mathfrak{a} is both an Artinian and Noetherian ring. (Of course, all Artinian rings are Noetherian [1, Theorem 8.5].) As $M/\mathfrak{a}M$ is a finitely generated R/\mathfrak{a} -module, it must be both Artinian and Noetherian as well, and hence have finite length as an R/\mathfrak{a} -module. The length of $M/\mathfrak{a}M$ as an R -module is the same as its length as an R/\mathfrak{a} -module, and hence $M/\mathfrak{a}M$ has finite length as an R -module as well. Thus $\mathfrak{a} = (a_1, \dots, a_d)$ is a parameter ideal of M . \square

The integer n appearing in the next result plays a key role in the main results.

Lemma 2.4. *Let (R, \mathfrak{m}) be a local ring and M a finitely generated R -module. There exists an integer n such that $\mathfrak{m}^n M \cap \Gamma_{\mathfrak{m}}(M) = (0)$.*

Proof. Since $\Gamma_{\mathfrak{m}}(M)$ is Artinian (see Proposition 1.9), the descending chain of submodules

$$(\mathfrak{m}M \cap \Gamma_{\mathfrak{m}}(M)) \supseteq (\mathfrak{m}^2M \cap \Gamma_{\mathfrak{m}}(M)) \supseteq \dots$$

must stabilize. That is, there is some $n \in \mathbb{N}$ such that

$$\mathfrak{m}^{n+i}M \cap \Gamma_{\mathfrak{m}}(M) = \mathfrak{m}^nM \cap \Gamma_{\mathfrak{m}}(M)$$

for all integers $i \geq 0$. Thus

$$\begin{aligned} \mathfrak{m}^n M \cap \Gamma_{\mathfrak{m}}(M) &= \bigcap_{i \geq n} (\mathfrak{m}^i M \cap \Gamma_{\mathfrak{m}}(M)) \\ &\subseteq \bigcap_{i \geq 1} \mathfrak{m}^i M \\ &= (0). \end{aligned}$$

The last equality is by the Krull Intersection Theorem [7, Theorem 8.10]. \square

Remark 2.5. In fact, for any finite-length submodule $L \subseteq M$, we have $L \subseteq \Gamma_{\mathfrak{m}}(M)$, and hence $\mathfrak{m}^n M \cap L = (0)$ where n is the integer of Lemma 2.4.

The next result is in the spirit of [8, Prop 4.7.13]. We include a proof in order to obtain specific bounds on the powers of a in this special case.

Proposition 2.6. *Let R be any commutative ring, M an R -module, and $a, b \in R$. Then, for arbitrary positive integers $p \leq q \leq r$, we have an equality*

$$(ba^r M : a^p) = a^{r-q}(ba^q M : a^p) + (0 :_M a^p).$$

Proof. First let $x \in (ba^r M : a^p)$. Then $a^p x = ba^r y$ for some y in M . Now

$$a^p(x - ba^{r-p}y) = 0,$$

so that $x - ba^{r-p}y \in (0 :_M a^p)$. Additionally,

$$ba^{r-p}y = a^{r-q} \cdot ba^{q-p}y \in a^{r-q}(ba^q M : a^p).$$

We now have

$$x = ba^{r-p}y + (x - ba^{r-p}y) \in a^{r-q}(ba^qM : a^p) + (0 :_M a^p).$$

For the other inclusion, it is clear that $(0 :_M a^p) \subseteq (ba^rM : a^p)$, so it suffices to prove $a^{r-q}(ba^qM : a^p) \subseteq (ba^rM : a^p)$. To that end, let $a^{r-q}y \in a^{r-q}(ba^qM : a^p)$ with $y \in (ba^qM : a^p)$. We can write $a^p y = ba^q w$ for some w in M . Thus we may rewrite $a^p \cdot a^{r-q}y$ as follows:

$$\begin{aligned} a^p a^{r-q}y &= a^{r-q} a^p y \\ &= a^{r-q} ba^q w \\ &= ba^r w, \end{aligned}$$

which is in ba^rM . Thus $a^{r-q}(ba^qM : a^p) + (0 :_M a^p) \subseteq (ba^rM : a^p)$ as desired. \square

The next result will be applied in Section 2.3 in the situation where $I = (a_1, \dots, a_d)$ is a parameter ideal and J is of the form $(a_1^{n_1}, a_2, \dots, a_d)$ for a positive integer n_1 .

Lemma 2.7. *Let R be a Noetherian ring, $J \subseteq I$ proper ideals of R with $\sqrt{I} = \sqrt{J}$, and N an R -module. If $\text{Hom}_R(R/J, N)$ is decomposable, then so is $\text{Hom}_R(R/I, N)$.*

Proof. Suppose that $\text{Hom}_R(R/J, N) = X \oplus Y$ where X and Y are nonzero R -modules. Since Hom and \otimes are adjoint functors, there are isomorphisms

$$\begin{aligned} \text{Hom}_R(R/I, N) &\cong \text{Hom}_R((R/I) \otimes_R (R/J), N) \\ &\cong \text{Hom}_R(R/I, \text{Hom}_R(R/J, N)) \\ &\cong \text{Hom}_R(R/I, X \oplus Y) \\ &\cong \text{Hom}_R(R/I, X) \oplus \text{Hom}_R(R/I, Y). \end{aligned}$$

By symmetry, it suffices to show that $\text{Hom}_R(R/I, X) \neq 0$. It is clear that $JX = (0)$ since $X \subseteq \text{Hom}_R(R/J, N)$. Choose $\mathfrak{p} \in \text{Ass}_R X$ and note that $J \subseteq \mathfrak{p}$. Indeed, we know $JX = (0)$ and any $\mathfrak{p} \in \text{Ass}_R(X)$ has the form $\mathfrak{p} = \text{ann}(x)$ for some $x \in X$. Since $Jx = (0)$, we obtain that $J \subseteq \mathfrak{p}$. Since $\sqrt{J} = \sqrt{I}$, one has $I \subseteq \mathfrak{p}$, so there are maps

$$R/I \twoheadrightarrow R/\mathfrak{p} \hookrightarrow X.$$

The composition of these maps is nonzero, and so $\text{Hom}_R(R/I, X) \neq (0)$ as desired. \square

Remarks 2.8. 1. The hypothesis that $J \subseteq I$ is necessary. For any pair of ideals I, J , if we let $N = R/I$, then

$$\text{Hom}_R(R/I, N) = \text{Hom}_R(R/I, R/I) \cong R/I$$

is indecomposable. However, it is possible that $\text{Hom}_R(R/J, N)$ is decomposable. For instance, let k be a field and consider the ring $R = k[[x, y]]/(x^2, xy)$ along with the ideals $I = (y^2)$, $J = (y)$ of R . Here $\sqrt{I} = \sqrt{J} = (x, y)$ but $J \not\subseteq I$. In Example 3.2 we will show

$$\text{Hom}_R(R/J, R/I) = \text{Hom}_R(R/(y), R/(y^2)) \cong \frac{(y)}{(y^2)} \oplus \frac{(x, y^2)}{(y^2)}.$$

2. The hypothesis that $\sqrt{I} = \sqrt{J}$ is also necessary. For example, let k be a field, $R = k[[x, y, z]]/(x^2, xyz)$, $N = R/(y^2)$, $I = (y, z)$, and $J = (y)$. We have $J \subseteq I$, but $\sqrt{J} = (x, y) \subsetneq (x, y, z) = \sqrt{I}$. Using this notation we obtain

$$\text{Hom}_R(R/J, N) \cong \frac{(y^2) : y}{(y^2)} = \frac{(y)}{(y^2)} \oplus \frac{(xz, y^2)}{(y^2)}$$

decomposes but

$$\mathrm{Hom}_R(R/I, N) \cong \frac{(y^2) : (y, z)}{(y^2)} = \frac{(xy, y^2)}{(y^2)}$$

is cyclic, and hence indecomposable.

This next result will be used in Section 2.3 in an induction argument.

Lemma 2.9. *Let R be a local ring and M an R -module of dimension $d \geq 2$. If M is not Cohen-Macaulay, then for any system of parameters a_1, \dots, a_d of M , there exist positive integers i and s such that $M/a_i^s M$ is not Cohen-Macaulay.*

Proof. If some a_i is M -regular, then $M/a_i M$ is not Cohen-Macaulay, so we may assume that each a_i is a zero-divisor on M . Suppose, by way of contradiction, that $M/a_1^s M$ is Cohen-Macaulay for each $s \geq 1$. Then a_2, \dots, a_d is a regular sequence on $M/a_1^s M$ for all integers $s \geq 1$. In particular a_2 is $M/a_1^s M$ -regular for all integers $s \geq 1$. We claim this implies a_2 is M -regular. Indeed, suppose $a_2 m = 0$ for some $m \in M$. Then $a_2 \bar{m} = 0$ in $M/a_1^s M$ for all integers $s \geq 1$, so that $m \in a_1^s M$ for all integers $s \geq 1$. By the Krull Intersection Theorem [7, Theorem 8.10], we have $m = 0$ and hence a_2 is M -regular which gives the desired contradiction. \square

The next example, noticed by Ryan Karr, shows that even when every element in a of the parameters is a zero-divisor, M may have positive depth.

Example 2.10. Consider the ring $R = k[[x, y, z]]/(x^2, xyz)$. It is clear that the rings $R/(x, y, z) \cong k$, $R/(x, y) \cong k[[z]]$, and $R/(x) \cong k[[y, z]]$ are all domains and so the dimension of R is at least 2. Since x is nilpotent, we have $x \in \sqrt{(y, z)}$ so that $\sqrt{(y, z)} = (x, y, z) = \mathfrak{m}$. Hence the dimension of R is 2 and $\{y, z\}$ is a system of parameters of R . Both y and z are zero-divisors in R since $xz \cdot y = xy \cdot z = 0$ and $xz, xy \neq 0$ in R .

To see that R has positive depth, note that $\text{Ass}_R R = \{(x), (x, y), (x, z)\}$. As $y - z \notin (x) \cup (x, y) \cup (x, z)$ we know that it is a regular element on R [7, Theorem 6.1(ii)]. We saw in Example 1.7 that $R/(y - z) \cong k[[x, y]]/(x^2, xy^2)$ has depth zero. Hence $y - z$ is a maximal length regular sequence on R and R has depth one. Thus we have found a ring of positive depth where the system of parameters consists only of zero-divisors.

Note that both $R/(y)$ and $R/(z)$ are Cohen-Macaulay rings of dimension one in this example, but that $R/(y^n)$ and $R/(z^n)$ have dimension one and depth zero for all integers $n \geq 1$.

The next example shows that for an arbitrary parameter a of a non-Cohen-Macaulay module, M , it is possible that $M/a^s M$ is Cohen-Macaulay for all s . Hence, we cannot strengthen Lemma 2.9 to say that for any parameter a of a non-Cohen-Macaulay module M , there is an integer s such that $M/a^s M$ is non-Cohen-Macaulay.

Example 2.11. Consider the ring $R = k[[x, y, z]]/(x^2, xy)$. It is clear that the rings $R/(x, y, z) \cong k$, $R/(x, y) \cong k[[z]]$, and $R/(x) \cong k[[y, z]]$ are all domains and so the dimension of R is at least two. Since x is nilpotent, $x \in \sqrt{I}$ for every ideal I of R . Thus, $\sqrt{(y, z)} = (x, y, z)$ is the maximal ideal of R . Hence R has dimension two and $\{y, z\}$ is a system of parameters of R . As z is a regular element of R which is in the maximal ideal, R must have at least depth one. However, $R/(z) \cong k[[x, y]]/(x^2, xy)$ has depth zero; see Example 1.7. Thus z is a maximal length regular sequence and R has depth one.

Next, we consider the quotients $S_n := R/(y^n) = k[[x, y, z]]/(x^2, xy, y^n)$ for $n \in \mathbb{N}$. We claim these quotients are all Cohen-Macaulay rings of dimension one. Indeed, $S_n/(x, y, z)S_n \cong k$ and $S_n/(x, y)S_n \cong k[[z]]$ are both domains, so S_n has dimension at least one for all n . Since $\sqrt{(y)S_n} = (x, y, z)S_n$ for all $n \in \mathbb{N}$ we have that $\{y\}$ is a

system of parameters for S_n . Thus $\dim S_n = 1$ for all n . To see that S_n has depth one, we simply need to note that z is a regular element in the maximal ideal of S_n .

2.2 Dimension One

We start the proofs of the main results with those for modules of dimension one and depth zero since we are able to obtain stronger bounds in this case. We show $\mathrm{Hom}_R(R/aR, M/bM)$ decomposes if the parameter b is chosen to be in a sufficiently high power of the ideal generated by an arbitrary parameter a . Recall:

Theorem 1.15. *[10, Theorem 3.1] Let (R, \mathfrak{m}) be a local ring, and M a nonzero finitely generated R -module of dimension one and depth zero. Choose an integer n such that $\mathfrak{m}^n M \cap \Gamma_{\mathfrak{m}}(M) = (0)$. For any parameter a of M , and any parameter b of M with $b \in (a^{n+1})$, the following R -module is decomposable:*

$$\mathrm{Hom}_R(R/aR, M/bM).$$

Remark 2.12. The integer n in the statement exists by Lemma 2.4. Note that $n \geq 1$ because $\Gamma_{\mathfrak{m}}(M) \neq (0)$; see Proposition 1.9.

Proof of Theorem 1.15. Set $S := R/\mathrm{ann}_R(M)$, and let $(-)$ denote the image in S . Then \bar{a} and \bar{b} are parameters of M as an S -module. Moreover there is an R -module isomorphism

$$\mathrm{Hom}_S(S/\bar{a}S, M/\bar{b}M) \cong \mathrm{Hom}_R(R/aR, M/bM).$$

By replacing R with S , we may thus assume that M is a faithful R -module.

Write $b = ca^{n+1}$. Since M is faithful, we have $\sqrt{(a)} = \mathfrak{m}$ (see Proposition 2.3)

and so

$$(0 :_M a) \subseteq \Gamma_{(a)}(M) = \Gamma_{\mathfrak{m}}(M).$$

Thus we know

$$(0 :_M a) \cap ca^n M \subseteq \Gamma_{\mathfrak{m}}(M) \cap \mathfrak{m}^n M = (0). \quad (2.1)$$

By Proposition 2.6, with $p = q = 1$ and $r = n + 1$, we have

$$(ca^{n+1}M : a) = a^n(caM : a) + (0 :_M a). \quad (2.2)$$

We now claim that

$$a^n(caM : a) = ca^n M.$$

Indeed, it is clear that $ca^n M \subseteq a^n(caM : a)$. For the reverse inclusion, let x be in $a^n(caM : a)$ and write $x = a^n m$ for some $m \in (caM : a)$. We have $am = cam'$ for some $m' \in M$. Then

$$x = a^n m = a^{n-1} \cdot am = ca^n m' \in ca^n M.$$

Equation (2.2) is thus equivalent to

$$(bM : a) = ca^n M + (0 :_M a). \quad (2.3)$$

Next we claim that

$$ca^{n+1}M = ca^n M \cap [(0 :_M a) + ca^{n+1}M]. \quad (2.4)$$

It is clear that $ca^{n+1}M \subseteq ca^n M \cap [(0 :_M a) + ca^{n+1}M]$. For the other inclusion, let x

be in $ca^n M \cap [(0 :_M a) + ca^{n+1}M]$, and write

$$x = ca^n y = z + ca^{n+1}w$$

for some $y, w \in M$ and $z \in (0 :_M a)$. Then

$$\begin{aligned} z &= ca^n y - ca^{n+1}w \\ &\in (0 :_M a) \cap ca^n M = (0) \end{aligned} \quad \text{by (2.1).}$$

Equation (2.4) follows. Now there are isomorphisms

$$\begin{aligned} \text{Hom}_R(R/aR, M/bM) &\cong \frac{(bM : a)}{bM} \\ &\cong \frac{ca^n M + (0 :_M a)}{ca^{n+1}M} && \text{by (2.3)} \\ &\cong \frac{ca^n M}{ca^{n+1}M} \oplus \frac{(0 :_M a) + ca^{n+1}M}{ca^{n+1}M} && \text{by (2.4)}. \end{aligned}$$

All that remains to prove is that both summands are nonzero.

If the summand on the left were zero, then $ca^n M = (a) \cdot ca^n M$ so that $ca^n M = (0)$ by Nakayama's Lemma. This would be a contradiction as $ca^{n+1} = b$ is a parameter of M .

If the summand on the right were zero, then

$$(0 :_M a) \subseteq ca^{n+1}M.$$

By Equation (2.1) we would then have

$$(0 :_M a) = (0 :_M a) \cap ca^{n+1}M = (0).$$

This would also give a contradiction as $\text{depth}_R M = 0$. Thus $\text{Hom}_R(R/aR, M/bM)$ is decomposable, as desired. \square

When R is a Cohen-Macaulay ring, we know from Proposition 1.13 that the R/\mathfrak{a} -module $\text{Hom}_R(R/\mathfrak{a}, R/\mathfrak{b}) \cong R/\mathfrak{a}$ is not only indecomposable, but also free. When R is one-dimensional and not Cohen-Macaulay, we can prove that in addition to being decomposable, this module will be non-free if the parameters are chosen to be in sufficiently high powers of the maximal ideal. In comparing this to Theorem 1.15 we see that the requirements for showing $\text{Hom}_R(R/\mathfrak{a}, R/\mathfrak{b})$ is not a free R/\mathfrak{a} -module are greater than those required to show that it is decomposable since the integer n in the two theorems is the same when $M = R$. Recall:

Theorem 1.17. [10, Theorem 3.3] *Let (R, \mathfrak{m}) be a local ring of dimension one and depth zero, and n an integer such that $\mathfrak{m}^n \cap \Gamma_{\mathfrak{m}}(R) = (0)$. For any parameter a in \mathfrak{m}^n and any parameter b in (a^2) , the $R/(a)$ -module*

$$\text{Hom}_R(R/(a), R/(b))$$

is decomposable and has a non-free summand.

Remark 2.13. Again, the integer n in the statement exists by Lemma 2.4 and must be positive since $\Gamma_{\mathfrak{m}}(R) \neq (0)$; see Proposition 1.9.

Proof of Theorem 1.17. We will first prove that the module decomposes. Both the proof of this fact and the decomposition obtained are similar to those found in the proof of Theorem 1.15. Write $I = \Gamma_{\mathfrak{m}}(R)$. For any $x \in \mathfrak{m}^n$, we know $(x) \cap I = (0)$ and hence $xI = (0)$. If $x \in \mathfrak{m}^n$ is also a parameter, then we know $\sqrt{(x)} = \mathfrak{m}$ and so $\Gamma_{(x)}(R) = I$, and $(0 : x) = I$ as well. To see that $(0 : x) = I$, note that $\sqrt{(x)} = \mathfrak{m}$

and $xI = 0$ imply

$$I \subseteq (0 : x) \subseteq \Gamma_{(x)}(R) = I,$$

whence $(0 : x) = I$.

Let $a \in \mathfrak{m}^n$ and $b \in (a^2)$ be parameters and write $b = ca^2$. Applying Proposition 2.6 with $p = q = 1$ and $r = 2$, we obtain the equality

$$((ca^2) : a) = a((ca) : a) + (0 : a). \quad (2.5)$$

We now note that $a((ca) : a) = (ca)$. We may thus rewrite Equation (2.5) as

$$((b) : a) = (ca) + I. \quad (2.6)$$

Next we wish to show that

$$(ca^2) = (ca) \cap [I + (ca^2)]. \quad (2.7)$$

The inclusion \subseteq is clear. For the other inclusion, let $x \in (ca) \cap [I + (ca^2)]$ and write $x = rca = s + r'ca^2$ for some $r, r' \in R$ and $s \in I$. Then

$$\begin{aligned} s &= rca - r'ca^2 \\ &\in (a) \cap I \\ &\subseteq \mathfrak{m}^n \cap I = (0). \end{aligned}$$

Thus $s = 0$ and $x = r'ca^2 \in (ca^2)$. This gives the existence of the following isomor-

phisms of $R/(a)$ -modules:

$$\begin{aligned} \mathrm{Hom}_R(R/(a), R/(b)) &\cong \frac{((b) : a)}{(b)} \\ &\cong \frac{(ca) + I}{(ca^2)} && \text{by (2.6)} \\ &\cong \frac{(ca)}{(ca^2)} \oplus \frac{I + (ca^2)}{(ca^2)} && \text{by (2.7).} \end{aligned}$$

Next we show that both summands are nonzero.

If the summand on the left were zero, then Nakayama's Lemma would imply that $ca = 0$, a contradiction as $ca^2 = b$ is a parameter and hence nonzero.

If the summand on the right were zero, then $I \subseteq (ca^2)$ so that

$$\Gamma_{\mathfrak{m}}(R) = I = I \cap (ca^2) \subseteq I \cap (a) = (0),$$

a contradiction as the depth of R is zero (see Proposition 1.9).

We now show that the summand on the left, that is, $(ca)/(ca^2)$, is not a free $R/(a)$ -module. To that end, recall that $I \cap (a) = (0)$, but $I \neq (0)$, so we can choose an element $y \in I \setminus (a)$. We know $ya = 0$ since $aI = (0)$. In particular, $yca \in (ca^2)$. Thus $y + (a)$ is a nonzero element of

$$\mathrm{ann}_{R/(a)} \left(\frac{(ca)}{(ca^2)} \right),$$

and hence $(ca)/(ca^2)$ is not free as an $R/(a)$ -module. □

2.3 Higher Dimensions

In higher dimensions, we can also prove a decomposition theorem. Recall:

Theorem 1.16. [10, Theorem 4.1] *Let R be a local ring and M a finitely generated R -module of dimension d . If M is not Cohen-Macaulay, then, for any system of parameters $\mathbf{a} = a_1, \dots, a_d$ of M , there exist positive integers n_1, \dots, n_d such that the following R -module is decomposable:*

$$\mathrm{Hom}_R(R/(\mathbf{a}), M/(a_1^{n_1}, \dots, a_d^{n_d})M).$$

Proof. As in the proof of Theorem 1.15, we may reduce to the case that M is a faithful module. We proceed by induction on d , the case $d = 1$ being covered by Theorem 1.15.

Assume, now, that $d \geq 2$. By Lemma 2.9, we can find some positive integer $i \leq d$ and a positive integer n_i such that $M/a_i^{n_i}M$ is not Cohen-Macaulay. We may harmlessly assume $i = 1$. Set

$$\bar{R} := R/(a_1^{n_1}), \quad \bar{M} := M/a_1^{n_1}M, \quad \text{and} \quad \bar{\mathbf{a}} := (\bar{a}_2, \dots, \bar{a}_d).$$

Then $\bar{\mathbf{a}}$ is a parameter ideal of \bar{M} . Since \bar{M} has dimension $d - 1$ and is not Cohen-Macaulay, by induction there are positive integers n_2, \dots, n_d such that the R -module

$$U := \mathrm{Hom}_{\bar{R}}(\bar{R}/\bar{\mathbf{a}}, \bar{M}/(\bar{a}_2^{n_2}, \dots, \bar{a}_d^{n_d})\bar{M})$$

is decomposable. Since there is an isomorphism

$$U \cong \mathrm{Hom}_R(R/(a_1^{n_1}, a_2, \dots, a_d), M/(a_1^{n_1}, a_2^{n_2}, \dots, a_d^{n_d})M),$$

we apply Lemma 2.7, with $N = M/(a_1^{n_1}, \dots, a_d^{n_d})M$, $I = (a_1, \dots, a_d)$, and $J = (a_1^{n_1}, a_2, \dots, a_d)$, to obtain the existence of the desired decomposition. \square

Recall the result below, which is a version of Theorem 1.17 for rings of arbitrary dimension.

Theorem 1.18. [10, Theorem 4.2] *Let R be a local ring of dimension d . If R is not Cohen-Macaulay, then for any system of parameters a_1, \dots, a_d of R , there exist integers $n_1, \dots, n_d, N_1, \dots, N_d \in \mathbb{N}$ with $N_i \geq n_i$ for $i = 1, \dots, d$ such that the $R/(a_1^{n_1}, \dots, a_d^{n_d})$ -module*

$$\mathrm{Hom}_R(R/(a_1^{n_1}, \dots, a_d^{n_d}), R/(a_1^{N_1}, \dots, a_d^{N_d}))$$

is decomposable and has a non-free summand.

Proof. We proceed by induction on d .

For $d = 1$, we choose n such that $\mathfrak{m}^n \cap \Gamma_{\mathfrak{m}}(R) = (0)$ and set n_1 and N_1 to be n and $2n$, respectively. Theorem 1.17 then gives the desired decomposition and existence of a non-free summand.

Now suppose that $d \geq 2$. By Lemma 2.9, we can find integers i and n_i such that $R/(a_i^{n_i})$ is not Cohen-Macaulay. We may harmlessly assume $i = 1$. Set $S := R/(a_1^{n_1})$ let $(\bar{})$ denote the image in S . Then $\bar{a}_2, \dots, \bar{a}_d$ is a system of parameters of S and, by induction, there exist integers $n_2, \dots, n_d, N_2, \dots, N_d$ such that the $S/(\bar{a}_2^{n_2}, \dots, \bar{a}_d^{n_d})$ -module

$$U := \mathrm{Hom}_S(S/(\bar{a}_2^{n_2}, \dots, \bar{a}_d^{n_d}), S/(\bar{a}_2^{N_2}, \dots, \bar{a}_d^{N_d}))$$

is decomposable and has a non-free summand. Note that

$$S/(\bar{a}_2^{n_2}, \dots, \bar{a}_d^{n_d}) \cong R/(a_1^{n_1}, \dots, a_d^{n_d})$$

and

$$S/(\overline{a_1}^{N_2}, \dots, \overline{a_d}^{N_d}) \cong R/(a_1^{n_1}, a_2^{N_2}, \dots, a_d^{N_d}).$$

Setting $N_1 = n_1$ we then have

$$U \cong \text{Hom}_R(R/(a_1^{n_1}, \dots, a_d^{n_d}), R/(a_1^{N_1}, \dots, a_d^{N_d})),$$

and this is decomposable and has a non-free summand. □

Chapter 3

Examples

Choosing $b = a^t$ in Theorem 1.15 gives $\text{Hom}_R(R/aR, M/a^tM)$ is decomposable for all integers $t \geq n + 1$. Similarly, if a is a parameter of R , Theorem 1.17 says that $\text{Hom}_R(R/(a^t), R/(a^T))$ is decomposable and has a non-free summand for $t \geq n$ and $T \geq 2t$. Thus, these theorems provide lower bounds sufficient to show that the module is decomposable and not free. One natural question to ask is whether or not these bounds are optimal, that is, are there smaller values of t and T that will also cause these modules to be decomposable and have non-free summands.

Theorem 1.16 and Theorem 1.18, which are for higher dimensional rings and modules, say there exist integers η_i , n_i , and N_i such that both

$$A := \text{Hom}_R(R/\mathfrak{a}, M/(a_1^{\eta_1}, \dots, a_d^{\eta_d})M)$$

and

$$B := \text{Hom}_R(R/(a_1^{n_1}, \dots, a_d^{n_d}), R/(a_1^{N_1}, \dots, a_d^{N_d}))$$

are decomposable and B has a non-free summand. These statements are not explicit about the powers which will work; however, the proofs indicate a method for itera-

tively choosing the integers to be sufficiently large so that the modules will decompose and B will have a non-free summand. For $d \geq 2$, examples seem to indicate that

$$\mathrm{Hom}_R(R/\mathfrak{a}, R/(a_1^{n_1}, \dots, a_d^{n_d}))$$

decomposes for all $n_i \geq N$ with N chosen to be sufficiently large. However, Example 3.6 shows that Theorem 1.15 is not strong enough to use the induction technique in Theorem 1.16 to prove the existence of such an integer N . So the question remains: Can we find bounds on the powers of the parameters that guarantee the modules A and B above are decomposable and B has a non-free summand?

The purpose of this chapter is to explore this question by way of examples. In particular, we focus on the structure of the R/\mathfrak{a} -module $\mathrm{Hom}_R(R/\mathfrak{a}, R/\mathfrak{b})$ for concrete examples of R , \mathfrak{a} , and \mathfrak{b} .

Let us take $M = R$ in Theorem 1.16. Our first example shows that sometimes

$$\mathrm{Hom}_R(R/(a_1, \dots, a_d), R/(a_1^{n_1}, \dots, a_d^{n_d})) \cong R/(a_1, \dots, a_d)$$

even when R is not Cohen-Macaulay and at least one of the n_i 's is greater than one.

Example 3.1. Let $R = k[[x, y]]/(x^2, xy^2)$. This ring has dimension one and depth zero as was shown in Examples 1.2 and 1.7. Consider the parameter y of R . Let $r \in R$ and write r using coset notation:

$$r = r_0 + r_1x + r_2y + r_3xy + r_4y^2 + r_5y^3 + \dots + (x^2, xy^2)$$

with $r_i \in k$. Looking at

$$ry = r_0y + r_1xy + r_2y^2 + r_3xy^2 + r_4y^3 + r_5y^4 + \dots + (x^2, xy^2)$$

we see that $ry \in (y^2) \subset R$ if and only if $r_0 = r_1 = 0$. This happens exactly when $r \in (y) \subset R$. Thus

$$\mathrm{Hom}_R(R/(y), R/(y^2)) \cong \frac{(y^2) :_R y}{(y^2)} = \frac{(y)}{(y^2)} \cong R/(y).$$

The second isomorphism is given by $ry + (y^2) \leftrightarrow r + (y)$.

The next example shows that the bound in Theorem 1.15 is close to optimal.

Example 3.2. Let $R = k[[x, y]]/(x^2, xy^m)$. This ring has dimension one and depth zero as was shown in Example 1.2 and Example 1.7. Consider the parameter y of $R = k[[x, y]]/(x^2, xy^m)$ and set

$$U_t := \mathrm{Hom}_R(R/(y), R/(y^t)) \cong \frac{(y^t) : y}{(y^t)}.$$

We will show U_t is cyclic (and hence indecomposable) for $t \leq m$ and decomposable for $t > m$. For any $r \in R$ we may write r in coset notation:

$$r = \sum_{i=0}^{\infty} r_i y^i + \sum_{i=0}^{m-1} r'_i x y^i + (x^2, xy^m)$$

with $r_i, r'_i \in k$. In this notation

$$ry = \sum_{i=0}^{\infty} r_i y^{i+1} + \sum_{i=0}^{m-2} r'_i x y^{i+1} + (x^2, xy^m). \quad (3.1)$$

Let $t \leq m$. We know that $ry \in (y^t) \subset R$ if the only nonzero terms of the sums in (3.1) are those with $i+1 < t$. So $ry \in (y^t) \subset R$ if and only if $r_i = r'_i = 0$ for $i < t-1$.

This gives that $r \in ((y^t) :_R y)$ if and only if

$$r = \sum_{i=t-1}^{\infty} r_i y^i + \sum_{i=t-1}^{m-1} r'_i x y^i + (x^2, x y^m).$$

That is, $r \in ((y^t) :_R y)$ if and only if $r \in (y^{t-1}) \subset R$. Hence $U_t \cong (y^{t-1})/(y^t)$ is cyclic.

Now let $t > m$. Since $x y^m = 0$ we now have that $r y \in (y^t)$ if and only if the only nonzero terms in (3.1) are those with in the first sum with $i + 1 < t$. This gives that $r y \in (y^t)$ if and only if

$$r = \sum_{i=t-1}^{\infty} r_i y^i + r'_{m-1} x y^{m-1} + (x^2, x y^m).$$

That is, $r \in ((y^t) :_R y)$ if and only if $r \in (y^{t-1}, x y^{m-1})$. Since $t > m$ we know $x y^{t-1} = 0$ in R . As $k[[x, y]]$ has unique factorization, we have $(y^{t-1}) \cap (x y^{m-1}, y^t) = (y^t)$ in R .

Thus

$$U_t \cong \frac{(y^{t-1}, x y^{m-1})}{(y^t)} \cong \frac{(y^{t-1})}{(y^t)} \oplus \frac{(x y^{m-1}, y^t)}{(y^t)}$$

is decomposable for all $t > m$.

Note that $\mathfrak{m}^i = (y^i)$ for $i \geq m + 1$. For $i \geq m + 1$ and $r \in R$ we have that $r y^i = 0$ if and only if $r \in (x)$ and so

$$\Gamma_{\mathfrak{m}}(R) = \bigcup_{i=1}^{\infty} (0 : \mathfrak{m}^i) = \bigcup_{i=m+1}^{\infty} (x) = (x).$$

Moreover $\mathfrak{m}^i \cap \Gamma_{\mathfrak{m}}(R) = (0)$ precisely when $i \geq m + 1$ and so Theorem 1.15 gives only that U_t decomposes for $t \geq m + 2$. Since our computations showed that U_t actually decomposes for $t \geq m + 1$, the bound obtained in Theorem 1.15 is, at worst, one away from a tight bound.

Before we present the next example, we recall the following well-known result:

Proposition 3.3. [6, Proposition 3.1, Theorem 3.7] *Let R be a ring and M an R -module of finite length. The non-units of the non-commutative ring $\text{End}_R(M)$ form a two-sided ideal if and only if M is indecomposable.*

In particular, this means that for parameter ideals $\mathfrak{a}, \mathfrak{b}$ of an R -module, M , the module $U := \text{Hom}_R(R/\mathfrak{a}, M/\mathfrak{b}M)$ is indecomposable if and only if the non-units of $\text{End}_R(U)$ form a two-sided ideal.

The next example shows that it is possible that the module $\text{Hom}_R(R/\mathfrak{a}, R/\mathfrak{b})$ is neither cyclic nor decomposable. It also shows that the bound in Theorem 1.15 may be quite far from optimal.

Example 3.4. Consider the parameter y^2 of $R = k[[x, y]]/(x^2, xy^m)$ for $m \geq 3$. Set

$$U_t := \text{Hom}_R(R/(y^2), R/(y^t)).$$

We claim that U_t is

$$\left\{ \begin{array}{ll} \text{cyclic,} & \text{if } t < m + 1, \\ \text{indecomposable, but not cyclic,} & \text{if } t = m + 1 \\ \text{decomposable,} & \text{if } t > m + 1. \end{array} \right. \quad (3.2)$$

For small values of t , this is easily seen to be cyclic (and hence indecomposable).

If $t = 0$, then

$$U_t = \text{Hom}_R(R/(y^2), R) \cong (0) : y^2 = (xy^{m-2})$$

is cyclic. When $t = 1$, we have that

$$U_t = \text{Hom}_R(R/(y^2), R/(y)) \cong \frac{(y) : y^2}{(y)} = R/(y)$$

is also cyclic. When $t = 2$, then

$$U_t = \text{Hom}_R(R/(y^2), R/(y^2)) \cong \frac{(y^2) : y^2}{(y^2)} = R/(y^2)$$

is also cyclic. For what follows, we'll focus on the case that $t \geq 3$ and the representation of U_t as $\frac{(y^t) : y^2}{(y^t)}$. We will also use the following notation. Let $r \in R$ and write r in coset notation:

$$r = \sum_{i=0}^{\infty} a_i y^i + \sum_{i=0}^{m-1} b_i x y^i + (x^2, x y^m). \quad (3.3)$$

In this notation

$$r y^2 = \sum_{i=0}^{\infty} a_i y^{i+2} + \sum_{i=0}^{m-3} b_i x y^{i+2} + (x^2, x y^m). \quad (3.4)$$

Suppose that $3 \leq t < m + 1$. We show that U_t is cyclic for all such values of t . We have that $r y^2 \in (y^t)$ if and only if the only nonzero terms of (3.4) are those in either sum with $i + 2 \geq t$. This means that $a_i = b_i = 0$ for $i = 0, 1, \dots, t - 1$. As $t - 2 \leq m - 2$ this gives that $r y^2 \in (y^t)$ if and only if

$$r = \sum_{i=t-2}^{\infty} a_i y^i + \sum_{i=t-2}^{m-2} b_i x y^i + (x^2, x y^m).$$

That is, $r \in ((y^t) : y^2)$ if and only if $r \in (y^{t-2})$. Thus, whenever $3 \leq t < m + 1$, the module $U_t \cong (y^{t-2})/(y^t)$ is cyclic.

Next consider the case that $t > m + 1$. We again have that $r y^2 \in (y^t)$ if and only if the only nonzero terms of (3.4) are those in either sum with $i + 2 \geq t$. Since $t - 2 > m - 2$ we know that the only nonzero summands appear in the first sum. This means that $a_i = 0$ for $i = 0, 1, \dots, t - 1$ and $b_i = 0$ for $i = 0, \dots, m - 3$. Thus

$ry^2 \in (y^t)$ if and only if

$$r = \sum_{i=t-2}^{\infty} a_i y^i + \sum_{i=m-2}^{m-1} b_i x y^i + (x^2, x y^m).$$

That is, $r \in ((y^t) : y^2)$ if and only if $r \in (y^{t-2}, x y^{m-2})$. Moreover, we claim that

$$(y^{t-2}) \cap (x y^{m-2}, y^t) = (y^t).$$

Indeed, we have that $t \geq m + 2$ and so writing $f \in (y^{t-2}) \cap (x y^{m-2}, y^t)$ as

$$f = a y^{t-2} = b x y^{m-2} + c y^t$$

with $a, b, c \in R$ we see that $b x y^{m-2}$ must be of the form $b' x y^{t-2} = 0$, since R is a quotient of the unique factorization domain $k[[x, y]]$. Hence $f \in (y^t)$. The other inclusion is clear. This gives the desired decomposition for $t > m + 1$:

$$U_t \cong \frac{(y^t) : y^2}{(y^t)} = \frac{(x y^{m-2}, y^{t-2})}{(y^t)} \cong \frac{(x y^{m-2}, y^t)}{(y^t)} \oplus \frac{(y^{t-2})}{(y^t)}.$$

We finally consider the case where $t = m + 1$. For this case we have that ry^2 is in $(y^t) = (y^{m+1})$ if and only if the only nonzero terms of (3.4) are those with $i + 2 \geq m + 1$, or equivalently $i \geq m - 1$. Since $m - 1 > m - 3$ all of these nonzero terms are in the first sum of (3.4). Thus $ry^2 \in (y^t)$ if and only if $a_i = 0$ for $i < m - 1$ and $b_i = 0$ for $i < m - 2$. This gives $r \in ((y^t) : y^2)$ if and only if $r \in (x y^{m-2}, y^{m-1})$ so that

$$U_t = \text{Hom}_R(R/(y^2), R/(y^t)) \cong \frac{(y^t) : y^2}{(y^t)} = \frac{(x y^{m-2}, y^{m-1})}{(y^t)}.$$

We now claim that this module is indecomposable. To that end, note that the

set $\{y^{m-1}, y^m, xy^{m-2}, xy^{m-1}\}$ forms a k -basis for U_t . We will show the non-units of $\text{End}_R(U_t)$ form a two-sided ideal. Let $\phi \in \text{End}_R(U_t)$ and write

$$\phi(y^{m-1}) = ay^{m-1} + by^m + cxy^{m-2} + dxy^{m-1}$$

and

$$\phi(xy^{m-2}) = ey^{m-1} + fy^m + gxy^{m-2} + hxy^{m-1}$$

where $a, b, c, d, e, f, g, h \in k$. Since ϕ is R -linear, we know that $y\phi(xy^{m-2}) = \phi(xy^{m-1}) = x\phi(y^{m-1})$. Hence,

$$ey^m + gxy^{m-1} = y\phi(xy^{m-2}) = x\phi(y^{m-1}) = axy^{m-1}$$

which implies $e = 0$ and $g = a$. In turn, this gives

$$\phi(y^m) = y\phi(y^{m-1}) = ay^m + cxy^{m-1}.$$

We may thus represent ϕ as a matrix in $M_4(k)$ using $y^{m-1}, y^m, xy^{m-2}, xy^{m-1}$ as the ordering of the basis:

$$\phi = \begin{pmatrix} a & 0 & 0 & 0 \\ b & a & f & 0 \\ c & 0 & a & 0 \\ d & c & h & a \end{pmatrix}.$$

Hence $\text{End}_R(U_t)$ is the set of all matrices of the form of ϕ . Note that the determinant of such a matrix is a^4 and so $\phi \in \text{End}_R(U_t)$ is a unit if and only if the diagonal entry in the matrix representation of ϕ is nonzero. Let $I \subseteq \text{End}_R(U_t)$ be the set of all

non-units of $\text{End}_R(U_t)$. That is, I is the set of matrices in $M_4(k)$ of the form

$$\begin{pmatrix} 0 & 0 & 0 & 0 \\ b & 0 & f & 0 \\ c & 0 & 0 & 0 \\ d & c & h & 0 \end{pmatrix}$$

where $b, c, d, f, h \in k$. The sum of any two matrices in I is clearly in I . Also, letting $X \in I$ and $Y \in \text{End}_R(U_t)$ we have

$$XY = \begin{pmatrix} 0 & 0 & 0 & 0 \\ b & 0 & f & 0 \\ c & 0 & 0 & 0 \\ d & c & h & 0 \end{pmatrix} \cdot \begin{pmatrix} a' & 0 & 0 & 0 \\ b' & a' & f' & 0 \\ c' & 0 & a' & 0 \\ d' & c' & h' & a' \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ a'b + fc' & 0 & a'f & 0 \\ a'c & 0 & 0 & 0 \\ a'd + b'c + c'h & a'c & f'c + a'h & 0 \end{pmatrix}$$

and

$$YX = \begin{pmatrix} a' & 0 & 0 & 0 \\ b' & a' & f' & 0 \\ c' & 0 & a' & 0 \\ d' & c' & h' & a' \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 & 0 & 0 \\ b & 0 & f & 0 \\ c & 0 & 0 & 0 \\ d & c & h & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ a'b + f'c & 0 & a'f & 0 \\ a'c & 0 & 0 & 0 \\ bc' + ch' + a'd & a'c & f'c + a'h & 0 \end{pmatrix}$$

which are both elements of I . Thus, I is a two sided ideal of $\text{End}_R(U_t)$ so that U_t is an indecomposable R -module by Proposition 3.3.

This completes the proof of the assertions in (3.2). We showed in Example 3.2 that $\mathfrak{m}^i \cap \Gamma_{\mathfrak{m}}(R) = (0)$ if and only if $i \geq m + 1$. Theorem 1.15 thus only predicts that U_t decomposes for $t \geq 2m + 4$ and so in this case the bound obtained in Theorem 1.15 is especially poor for large values of m .

Remark 3.5. A more efficient way to compute $\text{End}_R(U_{m+1})$ for a fixed value of m and field k in Example 3.4 is using Macaulay2 [4]. The following code, generated with the help of Frank Moore, determines the form of an arbitrary element of $\text{End}_R(U_t)$ given a module U_t which is an ideal in a quotient of a polynomial ring R :

```
linearMaps = method()
linearMaps (Module, Symbol) := (Ut, T) -> (
  --- We first set up the objects we need to work with.
  R := ring Ut;
  gensUt := flatten entries gens Ut;
  dimenUt := numgens source basis Ut;
  numgensUt := #gensUt;
  varList := toList(T_(1,1)..T_(numgensUt,dimnUt));
  B := QQ[varList,MonomialOrder=>Lex];
  rGens := gens R;
  idealR := ideal R;
  Q := B[rGens];
  S := Q/sub(idealR,Q);
  --- This builds the k-matrix representing an R-linear map.
  basisUt := sub((gens Ut) * (matrix basis Ut),S);
  M1 := transpose sub(genericMatrix(B,dimnUt,numgensUt),S);
  M2 := (transpose basisUt);
  genMatr := M1 * M2;
  M3 := transpose genMatr * sub(matrix basis Ut,S);
  Temp1 := coefficients(M3, Monomials=>flatten entries basisUt);
  myMatrix := last Temp1;
```

```

--- Now we find the relationships between the coefficients
--- coming from the fact that some elements of the basis
--- can be obtained in more than one way.
syzMatr := transpose genMatr * syz sub(matrix {gensUt},S);
Temp2 := flatten entries basisUt;
Temp3 := coefficients(syzMatr, Monomials => Temp2);
myEquations := select(flatten entries last Temp3, f -> f != 0);
--- Finally, we apply the relations to the k-matrix
--- to obtain an arbitrary element of End_R(Ut).
C := B/apply(myEquations,f -> sub(f,B));
QQQ := C[rGens];
SS := QQQ/sub(idealR, QQQ);
sub(myMatrix,C)
)

```

With this algorithm in Macaulay2, we are prepared to compute an arbitrary element of $\text{End}_R(U_t)$. In order to do this for the example $R = \mathbb{Q}[[x, y]]/(x^2, xy^m)$ and $U_t = \text{Hom}_R(R/(y^2), R/(y^t))$ with $m = 3$ and $t = 4$ we use the following input. The first two input lines define the ring R and module U_t .

```

i1 : R = QQ[x,y]/ideal(x^2,x*y^3);
i2 : Ut = Hom(coker matrix{{y^2}},coker matrix{{y^4}});
i3 : M = linearMaps(Ut,T)

```

The output for the third line is

```

o3 = | T_(2,3) 0      0      0      |
      | T_(1,2) T_(2,3) T_(2,2) 0    |

```

$$\begin{array}{|cccc|} \hline T_-(1,3) & 0 & T_-(2,3) & 0 \\ \hline T_-(1,4) & T_-(1,3) & T_-(2,4) & T_-(2,3) \\ \hline \end{array}$$

which is an arbitrary element of $\text{End}_R(U_t)$. By making the following identification of the coefficients used in Example 3.4 and by Macaulay2:

$$\begin{array}{ll} T_-(1,1) = \mathbf{a} = T_-(2,3) & T_-(2,1) = \mathbf{e} = 0 \\ T_-(1,2) = \mathbf{b} & T_-(2,2) = \mathbf{f} \\ T_-(1,3) = \mathbf{c} & T_-(2,4) = \mathbf{h} \\ T_-(1,4) = \mathbf{d} & \end{array}$$

we see that this is the same as the matrix as the one painstakingly computed by hand in Example 3.4.

Theorem 1.15 and Theorem 1.17, which give bounds on the powers needed to make the module $\text{Hom}_R(R/\mathfrak{a}, M/\mathfrak{b}M)$ decompose and be non-free, apply only in dimension one. However, examples seem to indicate that the R/\mathfrak{a} -module

$$\text{Hom}_R(R/\mathfrak{a}, R/(a_1^{n_1}, \dots, a_d^{n_d}))$$

is neither free nor indecomposable if the n_i are large enough. One such example is explained below.

Example 3.6. Consider the ring $R = k[[x, y, z]]/(x^2, xyz)$. We saw in Example 2.10 that this ring has dimension two and depth one. Note that $\{y, z\}$ is a system of parameters of R . If $n_1 \geq 2$, then $S_{n_1} := R/(y^{n_1})$ is not Cohen-Macaulay. Indeed, as y^{n_1} is a parameter, S_{n_1} has dimension one. The depth of S_{n_1} is zero since the nonzero element xy^{n_1-1} is in the socle. Letting \mathfrak{m} be the maximal ideal of S_{n_1} we have $\mathfrak{m}^i \cap \Gamma_{\mathfrak{m}}(S_{n_1}) = 0$ if and only if $i \geq n_1 + 2$. By symmetry, the same holds for the ring

$T_{n_2} := R/(z^{n_2})$. Thus Theorem 1.15 gives that $U_{n_1, n_2} := \text{Hom}_R(R/(y, z), R/(y^{n_1}, z^{n_2}))$ decomposes for all $n_1, n_2 \geq 2$ with $|n_1 - n_2| > 2$. However, direct computation shows that U_{n_1, n_2} actually decomposes as

$$U_{n_1, n_2} \cong \frac{(xy^{n_1-1}, y^{n_1}, z^{n_2})}{(y^{n_1}, z^{n_2})} \oplus \frac{(xz^{n_2-1}, y^{n_1}, z^{n_2})}{(y^{n_1}, z^{n_2})} \oplus \frac{(y^{n_1-1}z^{n_2-1}, y^{n_1}, z^{n_2})}{(y^{n_1}, z^{n_2})}$$

for all $n_1, n_2 \geq 2$. See Figure 3.1 for a visual representation of this.




Figure 3.1: In this figure, a lattice point (n_1, n_2) corresponds to the module $\text{Hom}_R(R/(y, z), R/(y^{n_1}, z^{n_2}))$ from Example 3.6. The modules corresponding to lattice points in the light grey regions are known to decompose due to Theorem 1.15. The modules corresponding to lattice points in the middle dark grey region are known to decompose by direct computation. The modules corresponding to lattice points where $n_1 = 1$ or $n_2 = 1$ are indecomposable since $R/(y)$ and $R/(z)$ are both Cohen-Macaulay rings.

Bibliography

- [1] M. F. Atiyah and I. G. Macdonald. *Introduction to commutative algebra*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969.
- [2] Kamal Bahmanpour and Reza Naghipour. A new characterization of Cohen-Macaulay rings. *J. Algebra Appl.*, 13(8):1450064, 7, 2014.
- [3] Winfried Bruns and Jürgen Herzog. *Cohen-Macaulay rings*, volume 39 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1993.
- [4] Daniel R. Grayson and Michael E. Stillman. Macaulay2, a software system for research in algebraic geometry. Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [5] Srikanth B. Iyengar, Graham J. Leuschke, Anton Leykin, Claudia Miller, Ezra Miller, Anurag K. Singh, and Uli Walther. *Twenty-four hours of local cohomology*, volume 87 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2007.
- [6] Nathan Jacobson. *Basic algebra. II*. W. H. Freeman and Company, New York, second edition, 1989.

- [7] Hideyuki Matsumura. *Commutative ring theory*, volume 8 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 1989. Translated from the Japanese by M. Reid.
- [8] D. G. Northcott. *Lessons on rings, modules and multiplicities*. Cambridge University Press, London, 1968.
- [9] D. Rees. A theorem of homological algebra. *Proc. Cambridge Philos. Soc.*, 52:605–610, 1956.
- [10] Katharine Shultis. Systems of parameters and the Cohen-Macaulay property. Preprint, 2014. arXiv:1412.5912.