

University of Nebraska - Lincoln
DigitalCommons@University of Nebraska - Lincoln

Nebraska Game and Parks Commission -- Staff
Research Publications

Nebraska Game and Parks Commission

2005

Metazoan Parasites of Young-of-the-Year Paddlefish from Lewis and Clark Lake, Nebraska, U.S.A.

Brenda M. Pracheil

Michigan State University, pracheilbm@ornl.gov

Gerald E. Mestl

Nebraska Game and Parks Commission

Patrick M. Muzzall

Michigan State University

Follow this and additional works at: <https://digitalcommons.unl.edu/nebgamestaff>

Pracheil, Brenda M.; Mestl, Gerald E.; and Muzzall, Patrick M., "Metazoan Parasites of Young-of-the-Year Paddlefish from Lewis and Clark Lake, Nebraska, U.S.A." (2005). *Nebraska Game and Parks Commission -- Staff Research Publications*. 65.
<https://digitalcommons.unl.edu/nebgamestaff/65>

This Article is brought to you for free and open access by the Nebraska Game and Parks Commission at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Nebraska Game and Parks Commission -- Staff Research Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Research Note

Metazoan Parasites of Young-of-the-Year Paddlefish from Lewis and Clark Lake, Nebraska, U.S.A.

BRENDA M. PRACHEIL,^{1,3} GERALD E. MESTL,² AND PATRICK M. MUZZALL¹

¹ Department of Zoology, Michigan State University, East Lansing, Michigan 48824, U.S.A. (e-mail: pracheil@msu.edu) and

² Nebraska Game and Parks Commission, 2200 N 33rd Street, Lincoln, Nebraska 68503, U.S.A.

ABSTRACT: Young-of-the-year paddlefish, *Polyodon spathula* (Polyodontidae), from Lewis and Clark Lake, an impoundment of the Missouri River in Nebraska, U.S.A., were surveyed for parasites. In 2001 and 2002, 28 and 48 fish were examined for parasites, respectively. Only the nematode *Rhabdochona decaturensis* infected fish collected in 2001, but 8 parasite taxa (*R. decaturensis*, *Spinitectus* sp., *Camallanus* sp., *Contracaecum* sp., *Marsipometra* sp., *Diclybothrium hamulatum*, *Ergasilus elongatus*, and 1 unidentified leech) infected fish collected in 2002. *Rhabdochona decaturensis* was the most common parasite, occurring in 21.4% of fish in 2001 and 79.2% of fish in 2002. Prevalence of other parasite species infecting fish from 2002 was 33% or less. The helminth community of young-of-the-year paddlefish in both years was dominated by nematodes. This study is the first report on parasites of young-of-the-year paddlefish and documents a new host record for *R. decaturensis*.

KEY WORDS: *Polyodon spathula*, paddlefish, parasites, *Camallanus*, *Contracaecum*, *Diclybothrium hamulatum*, *Ergasilus elongatus*, Hirudinea, *Marsipometra*, Piscicolidae, *Rhabdochona decaturensis*, *Spinitectus*, Lewis and Clark Lake, Missouri River, Nebraska, United States.

The paddlefish, *Polyodon spathula* (Walbaum, 1792), is among the most ancient species of freshwater bony fishes in the United States and occurs in large rivers of the Mississippi River drainage (Bemis et al., 1997). Although adult paddlefish have been the subject of considerable parasite survey (Linton, 1898; Cooper, 1918; Pearse, 1924; Simer, 1929, 1930; Meyer, 1940; Bangham and Venard, 1942; Wilson, 1956; Causey, 1957; Meyer, 1960: unpublished thesis, Iowa State University, Ames, Iowa; Huggins, 1972; Schmidt et al., 1974; Lockard and Parsons, 1975; Suppes and Meyer, 1975; Raikova et al., 1979; Robinson and Jahn, 1980; Miyazaki et al., 1988; Holloway et al., 1991), there is no published survey of parasites of young-of-the-year paddlefish. This study reports the parasites of young-of-the-year paddlefish

from Lewis and Clark Lake, Nebraska, U.S.A., and briefly characterizes their helminth community.

Lewis and Clark Lake is an impoundment of the Missouri River on the border of Nebraska and South Dakota, U.S.A., measuring ca. 40 by 4 km with a maximum depth of ca. 17 m. Young-of-the-year paddlefish were collected from this lake, using a 7.9-m (headrope) semiballoon otter trawl fished on the bottom of the old river channel (42°50'5"N; 97°34'2"W). Collections were made weekly 5–19 July 2001 and 26 June–24 July 2002. (In 2002, only 5 fish were collected in June: 48 were collected in July.) Fish were fixed in 10% formalin for 2 mo and stored in 70% alcohol.

On necropsy, the entire fish was examined for parasites. Monogenean and copepod prevalences reported in this study may differ from those found on fish examined at fresh necropsy but still represent minimum prevalence values for monogenean and copepod infections on young-of-the-year paddlefish in Lewis and Clark Lake. Parasites were processed using conventional parasitological techniques. Voucher specimens have been deposited in the United States National Parasite Collection (USNPC), Beltsville, Maryland as follows: *Rhabdochona decaturensis* Gustafson, 1949 (USNPC 096440); *Diclybothrium hamulatum* Simer, 1929 (Price, 1942) (USNPC 096441); and *Ergasilus elongatus* Wilson, 1916 (USNPC 096439). Specimens of the remaining species are retained by the senior author (B.M.P.).

Use of prevalence, mean intensity, and mean abundance is consistent with that recommended by Bush et al. (1997). Species richness is the number of parasite species in an examined fish. Values for Brillouin's index for use in diversity and evenness (Pielou, 1975; Magurran, 1988) and Simpson's dominance were calculated using common logarithms for all parasites irrespective of their sites of infection. Only prevalence was recorded for monogeneans and copepods, and they are not included in diversity, evenness, and dominance values. Values are reported as mean \pm SD

³ Corresponding author.

Table 1. Prevalence, mean abundance, and mean intensity of parasites in young-of-the-year paddlefish, *Polyodon spathula*, from Lewis and Clark Lake, Nebraska, U.S.A.*

Parasite	Prevalence (%)	Mean abundance \pm SD	Mean intensity \pm SD
<i>Dicelybothrium hamulatum</i> †	12.5	0.3 \pm 0.7	2.0 \pm 1.1
<i>Marsipometra</i> sp.†	6.3	0.1 \pm 0.2	1.0 \pm 0.0
<i>Camallanus</i> sp.‡	8.3	0.1 \pm 0.3	1.0 \pm 0.0
<i>Contracaecum</i> sp.‡	14.6	0.2 \pm 0.4	1.1 \pm 0.4
<i>Rhabdochona decaturensis</i> *‡	21.4	0.6 \pm 1.6	2.7 \pm 2.7
<i>Rhabdochona decaturensis</i> †	79.2	10.0 \pm 13.7	12.6 \pm 14.2
<i>Spinitectus</i> sp.‡	33.3	0.5 \pm 0.9	1.6 \pm 0.9
<i>Ergasilus elongatus</i> †	8.3	0.1 \pm 9.9	18.0 \pm 33.3
Pisicolididae gen. sp.†	2.1	<0.1 \pm 0.2	2.0 \pm 0

* Values from fish collected in 2001. All other values from fish collected in 2002.

† Gravid adults.

‡ Immatures.

followed parenthetically by range values where appropriate.

Paddlefish length measurements are reported as mean eye-to-fork length (Ruelle and Hudson, 1977). In 2001, the mean length of 28 fish sampled was 67.9 mm \pm 24.5 (35–108 mm), and in 2002, the mean length of 48 fish sampled was 93.5 mm \pm 27.7 (28–135 mm). Paddlefish in the 2002 sample were significantly larger than those in the 2001 sample (Student's *t*-test, $t = -4.3$, $P < 0.001$, 74 df). Because fish length between years was significantly different, species richness, prevalence, mean intensity, and mean abundance of *R. decaturensis* were compared between years in fish of similar length (75–108 mm) ranging from the minimum length of fish from either year infected with a parasite to the maximum fish length recorded in 2001. Gonads were not well developed, and sex could not be determined for most fish.

Paddlefish examined in 2001 were infected only with *R. decaturensis*, whereas fish from 2002 were infected with 8 parasite taxa including 1 monogenean, *D. hamulatum*; 1 cestode, *Marsipometra* sp.; 4 nematodes, *R. decaturensis*, *Camallanus* sp., *Contracaecum* sp., and *Spinitectus* sp.; 1 species of pisicolid leech; and 1 copepod, *E. elongatus* (Table 1). No gravid parasites were found in the paddlefish collected in 2001, but gravid individuals of *D. hamulatum*, *Marsipometra* sp., *R. decaturensis*, and *E. elongatus* occurred in paddlefish collected in 2002.

Mean parasite species richness for 2001 and 2002 was 0.2 \pm 0.4 and 1.7 \pm 1.3, respectively. Mean

species richness values were significantly different between years overall (Student's *t*-test on natural log-transformed data, $P < 0.001$, 74 df) and between years for fish of similar length (Student's *t*-test on square-root transformed data, $P < 0.001$, 49 df). Prevalence ($\chi^2 = 22.3$, $P < 0.001$, 1 df), natural log-transformed mean abundance (Student's *t*-test, $P < 0.001$, 74 df), and natural log-transformed intensity (Student's *t*-test, $P = 0.012$, 42 df) of *R. decaturensis* were significantly higher in 2002 than in 2001 overall, but in fish of similar size, there was no significant difference in prevalence (χ^2 , $P > 0.05$, 1 df), natural log-transformed mean abundance (Student's *t*-test, $P > 0.05$, 25 df), or natural log-transformed mean intensity (Student's *t*-test, $P > 0.05$, 32 df) between years. Intensity of *R. decaturensis* and fish length were significantly correlated in 2002 (Spearman's rank correlation, $r_s = 0.53$, $P < 0.05$) but not in 2001 (Spearman's rank correlation, $P > 0.05$).

With the exception of *Rhabdochona*, all parasite genera reported in this study are previously known from adult paddlefish. The paddlefish is a new host record for *R. decaturensis*. *Spinitectus carolini* was tentatively identified in cultured paddlefish by an employee with the Missouri Department of Conservation (Hoffman, 1999), but there are no other reports on *Spinitectus* sp. infecting paddlefish in the literature. Individuals of *Spinitectus* sp. found in this study were immature females.

Several factors including diet and fish age and size may be contributing to the between year differences in prevalence, intensity, and abundance of *R. decaturensis* and also to the between-year difference in parasite species richness values. For the first months of life, young-of-the-year paddlefish prey on a variety of macroinvertebrates such as *Daphnia* and *Hexagenia* spp. (Michaletz et al., 1982). As fish become older, they eat a wider variety of food items and filter feed on planktonic crustaceans such as copepods (Meyer, 1960: unpublished thesis, Iowa State University, Ames, Iowa; Ruelle and Hudson, 1977). Although the precise age of each fish was not determined in this study, it follows that the overall larger fish collected in 2002 were older, had a more diverse diet, and ate greater quantities of food than the overall smaller fish collected in 2001. Thus, the fish sampled in 2002 were probably exposed to both a greater diversity and absolute number of potential intermediate hosts.

Overall prevalence of young-of-the-year paddlefish from Lewis and Clark Lake in 2001 and 2002 was 21.4% and 85.4%, respectively. The internal helminth community in both years was dominated by nema-

todes. Among fish collected in 2002, there was a significant correlation between parasite species richness and fish length (Spearman's rank correlation, $r_s = 0.66$, $P < 0.05$). The mean diversity, evenness, dominance, and species richness values for the internal parasites of infected paddlefish collected in 2002 were 0.075 ± 0.129 , 0.202 ± 0.260 , 0.846 ± 0.253 , and 1.7 ± 1.3 , respectively. High relative intensities of *R. decaturensis* produced these high dominance, low diversity, and low evenness values.

We thank Kirk Steffensen, Jason Skold, David Tsoodle, Steve Freeling, Ken Hatten, Clint Williams, Austin Budden, and Jake Charvat, Nebraska Game and Parks Commission, for their assistance in the field.

LITERATURE CITED

- Bangham, R. V., and C. E. Venard.** 1942. Studies on parasites of Reelfoot Lake fish. IV. Distribution and checklist of parasites. *Journal of the Tennessee Academy of Sciences* 54:22–38.
- Bemis, W. E., E. K. Findeis, and L. Grande.** 1997. An overview of Acipenseriformes. *Environmental Biology of Fishes* 48:25–71.
- Bush, A. O., K. D. Lafferty, J. M. Lotz, and A. W. Shostak.** 1997. Parasitology meets ecology on its own terms: Margolis et al. revisited. *Journal of Parasitology* 83:575–583.
- Causey, D.** 1957. Parasitic copepoda from Louisiana fresh water fish. *American Midland Naturalist* 58:378–382.
- Cooper, A. R.** 1918. North American pseudophyllidean cestodes from fishes. *Illinois Biological Monographs* 4:1–243.
- Hoffman, G. L.** 1999. Parasites of North American Freshwater Fishes. Cornell University Press, Ithaca, New York. 539 pp.
- Holloway, H. L., T. A. Dick, and C. A. Ottinger.** 1991. *Polypodium hydriforme* (Coelenterata) in paddlefish from the upper Missouri River drainage. *Journal of Aquatic Animal Health* 3:210–212.
- Huggins, E. J.** 1972. Parasites of fishes in South Dakota. *South Dakota Experiment Station Bulletin* 484:1–73.
- Linton, E.** 1898. Notes on the cestode parasites of fishes. *Proceedings of the United States National Museum* 20:423–456.
- Lockard, L. L., and R. R. Parsons.** 1975. Some parasites of paddlefish (*Polyodon spathula*) from the Yellowstone River, Montana. *Great Basin Naturalist* 35:425–426.
- Magurran, A. E.** 1988. Ecological Diversity and its Measurement. Princeton University Press, Princeton, New Jersey. 179 pp.
- Meyer, M. C.** 1940. A revision of the leeches (Piscicolidae) living on fresh-water fishes of North America. *Transactions of the American Microscopical Society* 59:354–376.
- Michaletz, P. H., C. F. Rabeni, W. W. Taylor, and T. R. Russell.** 1982. Feeding ecology and growth of young of the year paddlefish in hatchery ponds. *Transactions of the American Fisheries Society* 111:700–709.
- Miyazaki, T., W. A. Rogers, and K. J. Semmens.** 1988. Gastro-intestinal histopathology of paddlefish, *Polyodon spathula* (Walbaum), infected with larval *Hysterothylacium dollfusi* Schmidt, Leiby and Kritsky, 1974. *Journal of Fish Diseases* 11:245–250.
- Pearse, A. S.** 1924. The parasites of lake fishes. *Wisconsin Academy of Sciences, Arts and Letters* 21:161–194.
- Pielou, E. C.** 1975. Ecological Diversity. Wiley-Interscience, New York, New York. 165 pp.
- Raikova, E. V., V. C. Suppes, and G. L. Hoffman.** 1979. The parasitic coelenterate, *Polypodium hydriforme*, Ussov, from the eggs of the American acipenseriform *Polyodon spathula*. *Journal of Parasitology* 65:804–810.
- Robinson, G. L., and L. A. Jahn.** 1980. Some observations of fish parasites in Pool 20, Mississippi River. *Transactions of the American Microscopical Society* 99:206–212.
- Ruelle, R., and P. L. Hudson.** 1977. Paddlefish (*Polyodon spathula*): growth and food of young-of-the-year and a suggested technique for measuring length. *Transactions of the American Fisheries Society* 106:609–613.
- Schmidt, G. D., P. D. Leiby, and D. C. Kritsky.** 1974. Studies on helminths of North Dakota. VIII. Nematodes from the paddlefish *Polyodon spathula* (Walbaum), including *Thynnascaris dollfusi* sp. nov. (Nematoda: Stomachidae). *Canadian Journal of Zoology* 52:261–263.
- Simer, P. H.** 1929. Fish trematodes from the lower Tallahatchie River. *American Midland Naturalist* 11:563–588.
- Simer, P. H.** 1930. A preliminary study of the cestodes of the spoonbill *Polyodon spathula*. *Transactions of the Illinois State Academy of Science* 22:139–145.
- Suppes, V. C., and F. P. Meyer.** 1975. *Polypodium* sp. (Coelenterata) infection of paddlefish (*Polyodon spathula*) eggs. *Journal of Parasitology* 61:772–774.
- Wilson, W. D.** 1956. Notes on the cestodes in paddlefish, *Polyodon spathula* (Walbaum), from the Missouri River. *Kansas Academy of Sciences* 59:459–460.