

**Serveur Académique Lausannois SERVAL [serval.unil.ch](https://serval.unil.ch)**

## **Author Manuscript**

### **Faculty of Biology and Medicine Publication**

**This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.**

Published in final edited form as:

**Title:** Meta-analysis identifies 13 new loci associated with waist-hip ratio and reveals sexual dimorphism in the genetic basis of fat distribution.

**Authors:** Heid IM, Jackson AU, Randall JC, Winkler TW, Qi L, Steinhorsdottir V, Thorleifsson G, Zillikens MC, Speliotes EK, Mägi R, Workalemahu T, White CC, Bouatia-Naji N, Harris TB, Berndt SI, Ingelsson E, Willer CJ, Weedon MN, Luan J, Vedantam S, Esko T, Kilpeläinen TO, Kutalik Z, Li S, Monda KL, Dixon AL, Holmes CC, Kaplan LM, Liang L, Min JL, Moffatt MF, Molony C, Nicholson G, Schadt EE, Zondervan KT, Feitosa MF, Ferreira T, Lango Allen H, Weyant RJ, Wheeler E, Wood AR, MAGIC., Estrada K, Goddard ME, Lettre G, Mangino M, Nyholt DR, Purcell S, Smith AV, Visscher PM, Yang J, McCarroll SA, Nemes J, Voight BF, Absher D, Amin N, Aspelund T, Coin L, Glazer NL, Hayward C, Heard-Costa NL, Hottenga JJ, Johansson A, Johnson T, Kaakinen M, Kapur K, Ketkar S, Knowles JW, Kraft P,

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.



Published in final edited form as:

Nat Genet. 2010 November ; 42(11): 949–960. doi:10.1038/ng.685.

## Meta-analysis identifies 13 new loci associated with waist-hip ratio and reveals sexual dimorphism in the genetic basis of fat distribution

A full list of authors and affiliations appears at the end of the article.

### Abstract

Users may view, print, copy, download and text and data- mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use: [http://www.nature.com/authors/editorial\\_policies/license.html#terms](http://www.nature.com/authors/editorial_policies/license.html#terms)

Corresponding Authors: Cecilia M. Lindgren, Wellcome Trust Centre for Human Genetics, University of Oxford, Roosevelt Drive, Oxford, OX3 7BN, UK, [celi@well.ox.ac.uk](mailto:celi@well.ox.ac.uk), Karen L. Mohlke, University of North Carolina, Chapel Hill, NC 27599-7264, [mohlke@med.unc.edu](mailto:mohlke@med.unc.edu), Caroline S. Fox, NHLBI's Framingham Heart Study, 73 Mt Wayte Ave Suite #2, Framingham MA 01702, [foxca@nhlbi.nih.gov](mailto:foxca@nhlbi.nih.gov), Mark I. McCarthy, Oxford Centre for Diabetes, Endocrinology and Metabolism, University of Oxford, Churchill Hospital, Old Road, Headington, Oxford OX3 7LJ, United Kingdom, [Mark.mccarthy@drl.ox.ac.uk](mailto:Mark.mccarthy@drl.ox.ac.uk), Iris M. Heid, Department of Epidemiology and Preventive Medicine, Regensburg University Medical Center, Franz-Josef-Strauss Allee 11, 93055 Regensburg, Germany, [iris.heid@klinik.uni-regensburg.de](mailto:iris.heid@klinik.uni-regensburg.de).

\*These authors contributed jointly.

\*\*These senior authors jointly directed this work.

#### AUTHOR CONTRIBUTION

**Writing group:** IB, CSF, IMH (lead), CML (lead), MIM, KLMoh, LQ, VSt, GT, MCZ

**Waist phenotype working group:** TLA, NB, IB, LAC, CMD, CSF, TBH, IMH, AUJ, CML (lead), RJFL, RM, MIM, KLMoh, LQ, JCR, EKS, VSt, KSt, GT, UT, CCW, TW, TWW, HEW, MCZ

**Data cleaning & analysis:** SIB, IMH (lead), EI, AUJ, HL, CML (lead), RJFL (lead), JL, RM, LQ, JCR, EKS, GT, SV, MNW, EW, CJW, TWW, TW

**Sex-specific analyses:** SIB, TE, IMH, AUJ, TOK, ZK, SL, CML, RJFL, RM, KLMon, KEN, LQ, JCR (lead), VSt, GT, TWW (lead)

**eQTL and expression analyses:** SIB, ALD, CCH, JNH, FK, LMK, CML, LL, RJFL, JL, MFM, JLM, CM, GN, EES, EKS, VSt, GT, KTZ

**Pathway and CNV analyses:** CML, SAM, MIM, JN, VSt, GT, BFV

**Secondary analyses:** SIB, IBB, NC, KE, TMF, MFF, TF, MEG, JNH, EI, GL, CML, HL, RM, MMas, MIM, KLMon, DRN, JRO, SP, JRBP, JCR, AVS, EKS, PMV, MNW, CJW, RJW, EW, ARW, JY

**Study-specific analyses:** GRA, DA, NA, TA, TLA, NB, CC, PSC, LC, LAC, DIC, MNC, CMD, TE, KE, EF, MFF, TF, APG, NLG, MEG, CHay, NLH, IMH, JJH, AUJ, ÅJ, TJoh, JOJ, JRK, MKaa, KKap, SKet, JWK, PKra, ATK, ZK, JKet, CLam, RJFL, CLec, HL, MFL, CML, JL, RWL, RM, MMas, BM, KLMon, APM, NN, KEN, DRN, JRO, KKO, CO, MJP, OPol, IPro, NP, MP, LQ, JCR, NWR, SR, FR, NRR, CS, LJS, Ksil, EKS, KSt, SS, AVS, NS, US, VSt, DPS, IS, MLT, TMT, NJT, AT, GT, AU, SV, VVat, LV, PMV, RMW, RW, RJW, SW, MNW, CCW, CJW, TWW, ARW, JY, JHZ, MCZ

**Study-specific genotyping:** DA, TLA, LDA, NB, IB, AJB, EB, LLB, IBB, HC, DIC, INMD, MDci, MRE, PE, KE, NBF, MF, APG, HG, CG, EJCG, CJG, THan, ALH, NH, CHay, AAH, JJH, FBH, DJH, JH, WI, MRJ, ÅJ, JOJ, JWK, PKov, ATK, HKK, JKet, PKra, RNL, CML, RJFL, JL, MLL, MAM, MMas, WLM, MIM, JBIM, MJN, MN, DRN, KKO, CO, OPed, LP, MJP, GP, ANP, NP, LQ, NWR, FR, NRR, CS, AJS, NS, ACS, MT, BT, AU, GU, VVat, PMV, HW, PZ

**Study-specific phenotyping:** HA, PA, DA, AMA, TLA, BB, SRB, RB, EB, IBB, JPB, MDör, CMD, PE, MFF, CSF, TMF, MF, SG, JG, LCG, THan, ASH, CHen, ALH, ATH, KHH, AHof, FBH, DJH, BI, TI, TJor, PJ, MRJ, ÅJ, AJ, ALJ, JOJ, FK, LK, JKuu, KKva, RK, SKet, JWK, IK, SKos, VK, MKäh, PKov, OL, RNL, BL, JL, GML, RJFL, TL, MMas, MIM, CO, BMP, OPed, CGPP, JFP, IPic, KP, OPol, AP, LQ, MR, IR, OR, VSa, JSar, PEHS, Ksil, NJS, JHS, TDS, DPS, RS, HMS, JSin, TT, AT, MU, PV, CBV, LV, JV, DRW, GBW, SHW, GW, JCW, AFW, LZ, PZ

**Study-specific management:** GRA, AMA, BB, YBS, RNB, HB, JSB, SB, MB, EB, DIB, IBB, JPB, MJC, FSC, LAC, GD, CMD, SE, GE, PF, CSF, TMF, LCG, VG, UG, MEG, THan, CHen, KH, AHam, TBH, ATH, AHof, FBH, DJH, BI, TI, CI, TJor, MRJ, ALJ, FK, KTK, WHLK, RK, JKap, MKäh, ML, DAL, LJJ, CML, RJFL, TL, MMar, TM, AMET, KM, MIM, KLMoh, PBM, KEN, MSN, DRN, BO, CO, OPed, LP, BWP, PPP, BMP, LJP, TQ, AR, IR, OR, PMR, VSa, PS, DS, ARS, NS, TDS, KSt, DPS, ACS, MS, TT, JT, UT, AT, MU, AU, TTV, PV, HV, JV, PMV, NJW, HEW, JFW, JCW, AFW

**Steering committee:** GRA, TLA, IB, SIB, MB, IBB, PD, CMD, CSF, TMF, LCG, THar, JNH (Chair), DJH, EI, RK, RJFL, MIM, KLMoh, KEN, JRO, LP, DS, DPS, UT, HEW

#### FINANCIAL DISCLOSURES

The authors declare competing financial interests. Details can be found in the Supplementary Note.

Waist-hip ratio (WHR) is a measure of body fat distribution and a predictor of metabolic consequences independent of overall adiposity. WHR is heritable, but few genetic variants influencing this trait have been identified. We conducted a meta-analysis of 32 genome-wide association studies for WHR adjusted for body-mass-index (up to 77,167 participants), following up 16 loci in an additional 29 studies (up to 113,636 subjects). We identified 13 novel loci in or near *RSPO3*, *VEGFA*, *TBX15-WARS2*, *NFE2L3*, *GRB14*, *DNM3-PIGC*, *ITPR2-SSPN*, *LY86*, *HOXC13*, *ADAMTS9*, *ZNRF3-KREMEN1*, *NISCH-STAB1*, and *CPEB4* ( $P$   $1.9 \times 10^{-9}$  to  $1.8 \times 10^{-40}$ ), and the known signal at *LYPLAL1*. Seven of these loci exhibited marked sexual dimorphism, all with a stronger effect on WHR in women than men ( $P$  for sex-difference  $1.9 \times 10^{-3}$  to  $1.2 \times 10^{-13}$ ). These findings provide evidence for multiple loci that modulate body fat distribution, independent of overall adiposity, and reveal powerful gene-by-sex interactions.

## Keywords

genome-wide association; waist-hip-ratio; body fat distribution; central obesity; meta-analysis; genetics; visceral adipose tissue; metabolism; body composition; Expression Quantitative Trait Loci; sex difference

---

Central obesity and body fat distribution, as measured by waist circumference (WC) and waist-hip-ratio (WHR), are associated with individual risk of type 2 diabetes (T2D)<sup>1,2</sup> and coronary heart disease<sup>3</sup>, and with all-cause mortality<sup>4</sup>. These effects are independent of overall adiposity as measured by body mass index (BMI). WHR is of particular interest as a measure of body fat distribution, since it integrates the adverse metabolic risk associated with increasing WC with the more protective role of gluteal fat deposition with respect to diabetes, hypertension, and dyslipidemia<sup>5,6</sup>.

There is abundant evidence that body fat distribution is influenced by genetic loci distinct from those regulating BMI and overall adiposity. First, even after accounting for BMI, individual variation in WHR is heritable<sup>7,8</sup>, with estimates ranging from 22–61%<sup>7–10</sup>. Second, the striking abnormalities of regional fat deposition associated with lipodystrophic syndromes demonstrate that genetic variation can have dramatic effects on the development and maintenance of specific fat depots<sup>11,12</sup>. Third, in a previous genome-wide association analysis, we identified a locus near *LYPLAL1* strongly associated with WHR independent of any effects on BMI<sup>13</sup>, providing proof-of-principle for the genetic control of body fat distribution, distinct from that of overall adiposity.

Within the GIANT (Genetic Investigation of Anthropometric Traits) consortium, we performed a large-scale meta-analysis of genome-wide association (GWA) studies informative for WHR, using adjustment for BMI to focus discovery towards genetic loci associated with body fat distribution rather than overall adiposity<sup>14–16</sup>.

## RESULTS

### Genome-wide significant association of WHR with 14 SNPs

We conducted a two-stage study among individuals of European descent (Supplementary Table 1 and Online Methods). In the discovery stage, up to 2,850,269 imputed and

genotyped single nucleotide polymorphisms (SNPs) were examined in 32 GWA studies comprising up to 77,167 participants informative for anthropometric measures of body fat distribution. We performed a fixed-effects meta-analysis of WHR, employing study-specific linear regression adjusted for BMI and age, stratified by gender, and using an additive genetic model. After genomic control adjustment per study and in the meta-analysis, these analyses revealed a substantial excess of low p-values (Figure 1 a, b).

We selected SNPs representing the top 16 independent (> 1 Mb distance) regions of association (discovery  $P < 1.4 \times 10^{-6}$ , Table 1) and evaluated them in 29 additional, independent studies (up to 113,636 individuals) using a mixture of *in silico* data and *de novo* genotyping. In these follow-up studies, 14 of the 16 showed strong directionally-consistent evidence for replication ( $P < 1.0 \times 10^{-3}$ ) and ten reached genome-wide significance ( $P < 5.0 \times 10^{-8}$ ). Joint analysis of the discovery and follow-up results revealed genome-wide significant associations for 14 signals ( $P$  between  $1.9 \times 10^{-9}$  and  $1.8 \times 10^{-40}$ , Table 1).

Between-study heterogeneity was low ( $I^2 < 30\%$ ) for all but two signals (*GRB14* and *LYPLALI*, see Supplementary Note) and all 14 associations remained genome-wide significant in a random-effects meta-analysis (Supplementary Table 2).

One of these SNPs, rs4846567, is in linkage disequilibrium (LD,  $r^2 = 0.64$ ,  $D' = 0.84$ ; HapMap CEU) with the previously reported WHR-associated variant (rs2605100) near the *LYPLALI* gene13. The remaining 13 loci were in or near genes not previously associated with WHR or other measures of adiposity: *RSPO3*, *VEGFA*, *TBX15-WARS2*, *NFE2L3*, *GRB14*, *DNM3-PIGC*, *ITPR2-SSPN*, *LY86*, *HOXC13*, *ADAMTS9*, *ZNRF3-KREMEN1*, *NISCH-STAB1*, and *CPEB4* (Figure 2). These 14 loci explain 1.03% of the variance in WHR (after adjustment for BMI, age, and sex), with each locus contributing from 0.02% (*ZNRF3-KREMEN1*) to 0.14% (*RSPO3*) based on effect estimates in the follow-up stage.

### Sexual dimorphism at several of the WHR loci

Given the known sexual dimorphism of WHR and evidence from variance decomposition studies that this reflects sex-specific genetic effects<sup>17</sup>, we performed sex-specific meta-analyses for the 14 WHR associated SNPs. These analyses included up to 108,979 women (42,735 discovery, 66,244 follow-up) or 82,483 men (34,601 discovery, 47,882 follow-up). In joint analysis of discovery and follow-up data, 12 of the 14 SNPs reached genome-wide significance in women, but only 3 in men (Table 2). At all but one locus (*TBX15-WARS2*), effect-size estimates were numerically greater in women. At seven of the loci (those near *RSPO3*, *VEGFA*, *GRB14*, *LYPLALI*, *HOXC13*, *ITPR2-SSPN* and *ADAMTS9*), there were marked differences in sex-specific beta-coefficients ( $P$  ranging from  $1.9 \times 10^{-3}$  to  $1.2 \times 10^{-13}$ ). All loci displayed consistent patterns of sex-specific differences in both discovery and follow-up studies (Table 2). These 14 loci explain 1.34% of the variance in WHR (after adjustment for BMI and age) in women, but only 0.46% in men.

### Association with other anthropometric measures

By focusing on WHR after adjustment for BMI, our goal was to detect effects on body fat distribution independent of those influencing overall adiposity. As expected, we found very

little evidence that known BMI-associated variants were detected in our WHR analysis. Of the 10 loci identified shown to be associated with BMI in previous genome-wide association studies 14,15,18, only two showed nominally significant ( $P < 0.05$ ) associations for BMI-adjusted WHR in the discovery analysis (*FTO*: rs805013614,  $P = 0.03$ ,  $N = 77,074$ ; *TMEM18*: rs654823815,  $P = 3.0 \times 10^{-3}$ ,  $N = 77,016$ ).

We also tested the 14 WHR-associated SNPs for their effect on BMI using data from up to 242,530 participants available from the GIANT consortium (including most of the studies available for WHR association). Of the 14 WHR loci, four (near *TBX15-WARS2*, *CPEB4*, *LYPLAL1* and *GRB14*) also showed evidence of association with BMI ( $4.1 \times 10^{-3}$   $P$   $3.2 \times 10^{-6}$ ) with the WHR-increasing allele associated with decreased BMI (Supplementary Table 3). When adding an interaction term of SNP and BMI into the model, we observed that BMI modified the WHR association at the *LY86* locus ( $P$  for interaction =  $9.5 \times 10^{-5}$ ) with a larger WHR effect among the obese compared to the non-obese (see Supplementary Note).

To determine whether the WHR-associated signals exert their effects primarily through an effect on waist (WC) or hip circumference (HIP), we performed meta-analyses for these specific phenotypes in the discovery and follow-up studies (Supplementary Table 1 and 3). Overall, we observed stronger associations for HIP than for WC. Effect-size estimates were numerically greater for HIP than for WC at eleven of the 14 loci, and there were nominal associations ( $P < 0.05$ ) with HIP for twelve of the WHR-associated loci but only four associations with WC. In both sexes, the WHR-associated loci displaying nominal association with HIP always featured the WHR-increasing allele associated with reduced HIP. In contrast, we observed sexual dimorphism in the pattern of WC associations. In women, the WHR-increasing allele at all 14 loci was associated with increased WC, whereas this was only true for 6 of these loci in men (Figure 3). At *GRB14*, for example, the WHR-increasing allele was associated with increased WC in women ( $P = 3.6 \times 10^{-4}$ ) but decreased WC in men ( $P = 6.8 \times 10^{-3}$ ). These differences in the relationships between WC, HIP and WHR underlie some of the sexual dimorphism in the patterns of WHR association.

### Enrichment of association with metabolic traits

We evaluated the 14 WHR-associated loci for their relationships with related metabolic traits using GWA data provided by trait-specific consortia<sup>19–21</sup> as well as our *de novo* genotyped follow-up studies. As expected, given the sample overlap between this GWA data with our WHR GWA data as well as known trait correlations (Supplementary Table 4), we observed directionally consistent enrichment of associations ( $P < 0.05$ ) between the 14 WHR-associated alleles and increased triglycerides, LDL-cholesterol, fasting insulin, and HOMA-derived measures of insulin resistance (binomial  $P$  from  $3.2 \times 10^{-4}$  to  $1.8 \times 10^{-8}$ ; Table 3 and Supplementary Table 5a). For example, the WHR-increasing allele at *GRB14* shows strong associations with increased triglycerides ( $P = 7.4 \times 10^{-9}$ ), fasting insulin levels ( $P = 5.0 \times 10^{-6}$ ) and insulin resistance ( $P = 1.9 \times 10^{-6}$ ). Eleven of 14 WHR-associated loci showed directionally consistent associations with T2D, three of these (*ADAMTS9*, *NISCH-STAB1*, and *ITPR2-SSPN*) reaching nominal significance ( $P < 0.05$ ) (Table 3 and Supplementary Table 5a,b). Because the association signals for correlated traits in this

analysis are vulnerable to overestimation given the overlap in GWAS samples examined, we repeated these analyses restricted to our *de novo* genotyped follow-up studies. Although this also resulted in lower sample size, similar patterns of enrichment were still observed (Supplementary Table 5c).

### Pathway analysis of the WHR-associated loci and potential biological role

To identify potential functional connections and pathway relationships between genes mapping at the WHR-associated loci, we focused on the 95 genes located in a 2 Mb interval centered around each of the 48 independent SNPs that attained a  $P < 1.0 \times 10^{-5}$  in the WHR discovery studies.

First, we performed a survey of the published literature using GRAIL22, to search for connectivity between the genes and specific keywords that describe these functional connections (see Online Methods). Although there was no evidence, after correcting for multiple testing, that the connectivity between these genes was greater than chance, we identified 8 genes with nominal significance ( $P < 0.05$ ) for potential functional connectivity (*PLXND*, *HOXC10*, *TBX15*, *RSPO3*, *HOXC4*, *HOXC6*, *KREMEN1* and *HOXC11*). The keywords associated with these connections included “vegf”, “homeobox”, “patterning”, “mesenchyme”, “embryonic”, “development” and “angiogenesis”.

Additionally, we performed pathway analyses using the PANTHER database<sup>23</sup> based on the same set of 95 genes (Online methods and Supplementary Note). This analysis generated some evidence for over-representation of “developmental processes” ( $P = 5.8 \times 10^{-8}$ ) and “mRNA transcription regulation” ( $P = 2.7 \times 10^{-6}$ ), but neither retained nominal significance after adjustment for bias (e.g. due to non-random SNP coverage in relation to genes) and the number of biological processes tested (Supplementary Note, Supplementary Table 6).

Finally, we examined the described functional roles of some of the most compelling candidates based on either proximity to the signal or other analyses described in this paper. These uncovered possible roles in adipocyte development (*TBX15*), pattern formation during embryonic development (*HOXC13*), angiogenesis (*VEGFA*, *RSPO3*, *STAB1*), Wnt/beta-catenin signaling (*RSPO3*, *KREMEN1*), insulin signaling (*ADAMTS9*, *GRB14*, *NISCH*), lipase activity (*LYPLAL1*), lipid biosynthesis (*PIGC*) and intracellular calcium signaling (*ITPR2*) (see Supplementary Note for details).

### Evaluation of copy number variants (CNVs) and nonsynonymous changes

Both common and rare CNVs have been reported to be associated with overall adiposity<sup>14,15,24,25</sup>, but the impact of CNVs on fat distribution has not been evaluated previously. To examine the potential contribution of common CNVs to variation in WHR, we looked for evidence of association in our GWA discovery meta-analysis, using a set of 6,018 CNV-tagging SNPs, which collectively capture >40% of common CNVs >1 kb<sup>26,27</sup> (Online Methods, Supplementary Note).

One CNV-tagging SNP (rs1294421, *LY86*) was observed amongst our 14 WHR-associated loci. This SNP is in strong LD ( $r^2 = 0.98$ ) with a 2,832 bp duplication variant (CNVR2760.1)<sup>27</sup>, located 12 kb from an expressed sequence tag (BC039678) and 87 kb

from *LY86*, such that the duplication allele is associated with reduced WHR. The duplicated region consists entirely of noncoding sequence but includes part of a predicted enhancer sequence (E.5552.1)28.

To identify other putatively causal variants in our associated regions, we searched for non-synonymous coding SNPs in strong LD ( $r^2 > 0.7$ ) with the most strongly associated SNPs at each locus using data from the HapMap (Build 21) and 1000 Genomes Project (April and August 2009 releases). In this search, one lead SNP (rs6784615, at the *NISCH-STAB1* locus) was correlated with non-synonymous changes in two nearby genes, *DNAH1* (Val441Leu, Arg1285Trp and Arg3809Cys) and *GLYCTK* (Leu170Val). Fine-mapping and functional studies will be required to determine whether the *DNAH1* or *GLYCTK* SNPs or the *LY86* CNV are causal for the WHR-associations at these loci.

### Evaluation of effect of the WHR associations on expression in relevant tissues

Expression-QTL (eQTL) data can implicate regional transcripts that mediate trait-associations, and we therefore examined the 14 WHR-associated loci using eQTL data from human subcutaneous adipose tissue (SAT)29 (two separate sample sets, N=610 and N=603), omental fat30 (N=740), liver30 (N=518), blood29 (N=745), and lymphocytes31 (N=830) (Online methods, Supplementary Note).

At six of the loci, the WHR-associated SNP was either the strongest SNP associated with significant ( $P < 1.0 \times 10^{-5}$ ) expression of a local (within 1 Mb) gene transcript or explained the majority of the association between the most significant eQTL SNP and the gene transcript in conditional analyses (adjusted  $P > 0.05$ ; Table 4). For example, the WHR-associated SNP rs1011731 (near *DNM3-PIGC*) was strongly-associated with expression of *PIGC* in lymphocytes ( $P = 5.9 \times 10^{-10}$ ); furthermore, rs1011731 is in high LD ( $r^2 = 1.00$ ,  $D' = 1.00$ , HapMap CEU) with the SNP with the strongest effect on *PIGC* expression (rs991790), and this *cis*-eQTL association is abolished by conditioning on rs1011731. These analyses therefore indicate that these two signals are coincident and that *PIGC* is a strong candidate for mediating the WHR-association at rs1011731. We found similar evidence for coincidence of the WHR signal with expression for rs984222 (*TBX15* in omental fat), rs1055144 (expressed sequence tag AA553656 in SAT), rs10195252 (*GRB14* in SAT), rs4823006 (*ZNRF3* in SAT and omental fat), and rs6784615 (*STAB1* in blood)(Table 4). Taken together, the overlap between trait association and gene expression at these loci suggests that the WHR associations are driven through altered expression of *PIGC*, *TBX15*, AA553656, *GRB1*, *ZNRF3* and *STAB1*.

### Differential RNA expression of gluteal compared to abdominal fat tissue

To determine whether genes within the WHR-associated loci showed evidence of differential transcription in distinct fat-depots, we compared expression levels in gluteal or abdominal SAT in 49 individuals. We focused on the 15 genes with the strongest credentials for causal involvement (on the basis of proximity to the lead SNP and/or other biological or functional data: Table 1) for which expression data were available. Five of these (*RSPO3*, *TBX15*, *ITPR2*, *WARS2* and *STAB1*) were differentially expressed between the two tissues (F-test, corrected for false discovery rate across the 15 expressed genes,  $P < 0.05$ ;

Supplementary Table 7). This supported the hypothesis that, at some loci at least, the association with WHR reflects depot-specific differences in expression patterns.

## DISCUSSION

Overall, our findings demonstrate that the genetic regulation of body fat distribution involves loci and processes that are largely distinct from those that influence BMI and risk of obesity. This finding is consistent with the evidence that WHR displays substantial heritability even after adjustment for BMI. The loci that emerge from this study display no overlap with those shown to be associated with BMI, either in previous reports<sup>14,15,16</sup> or in the expanded meta-analysis recently completed by the GIANT consortium<sup>32</sup>.

Another point of distinction between our findings and those for BMI relates to the evidence for sexual dimorphism that we observed at several of the WHR-associated loci. Sex differences in the regulation of body fat distribution have long been acknowledged without a clear understanding of the underlying molecular mechanisms. These differences become apparent during puberty and are generally attributed to the influence of sex hormones<sup>33</sup>. Consistent with our findings, variance decomposition studies have shown that the genetic contribution to the overall variance in WHR, waist or hip circumference is greater in women<sup>17</sup>. While there is some evidence for loci with differential sex effects influencing lipids<sup>34</sup>, uric acid levels<sup>35</sup> and risk of schizophrenia<sup>36</sup>, we are unaware of prior reports indicating such strong enrichment of female-specific associations for any other phenotype, including BMI<sup>32</sup>.

The primary objective of genetic discovery efforts is to characterize the specific mechanisms involved in the regulation of the trait of interest. Despite the considerable challenges associated with moving from common variant association signals to definition of the causal alleles and pathways, we have identified strong candidates at several of the loci. For example, the *cis*-eQTL data implicate *GRB14* as a compelling candidate for the WHR-association on chromosome 2, and we were able to show that the same *GRB14* variants are also associated with triglyceride and insulin levels, consistent with previous association of this locus with HDL-cholesterol<sup>37</sup>. These inferences about the role of *GRB14* are supported by evidence that *Grb14*-deficient mice exhibit improved glucose homeostasis despite lower circulating insulin levels, and enhanced insulin signaling in liver and skeletal muscle<sup>38</sup>. The signal near *ADAMTS9* overlaps a previously-reported T2D locus<sup>39</sup>, and the lead SNP for WHR in our study is identical to the SNP displaying the strongest T2D association in an expanded T2D meta-analysis<sup>40</sup>. Given evidence that *ADAMTS9* T2D-risk alleles are associated with insulin resistance in peripheral tissues<sup>41</sup>, these findings are consistent with a primary effect of *ADAMTS9* variants on body fat distribution. At the chromosome 6 locus, *VEGFA* is the most apparent biological candidate, given the presumed role of *VEGFA* as a mediator of adipogenesis<sup>42</sup> and evidence that serum levels of VEGFA are correlated with obesity<sup>43,44</sup>. Finally, at the *TBX15-WARS2* locus, *TBX15* emerges as the strongest candidate based on the *cis*-eQTL data in omental fat, marked depot-specific differences in adipose tissue expression in mice and humans, and associations between *TBX15* expression in visceral fat and WHR<sup>45,46</sup>.



Our efforts to use pathway- and literature-mining approaches to look for functional enrichment of the genes mapping to associated regions met with only limited success, but did provide some support for overrepresentation of developmental processes. Developmental genes have been implicated in fat accumulation and distribution<sup>45,46</sup>, and recent evidence supports a link between developmental genes and body fat distribution, including HOXC13<sup>47</sup> and TBX15<sup>45,48</sup>. Developmental genes may in part determine the adipocyte-specific expression patterns that have been observed in different fat depots<sup>45</sup>. Taken together, our findings point to a set of genes influencing body fat distribution that have their principal effects in adipose tissue. This is in contrast to the predominantly central (hypothalamic) processes that are involved in the regulation of body mass index and overall adiposity<sup>49</sup>.

By providing novel insights into the regulation of body fat distribution, the present study raises a number of issues for future investigation. From the genetic perspective, re-sequencing, dense-array genotyping and fine-mapping approaches will be required to characterize causal variants at the loci we have identified, and to support further discoveries that may account for the substantial proportion of genetic variance unexplained by our findings. From the clinical perspective, it will be important to explore the relationship of these variants to more refined measures of body fat distribution derived from detailed imaging studies, to use the variants identified to characterise the causal relationships between body fat distribution and related metabolic and cardiovascular traits, and to explore ethnic differences in patterns of body fat distribution. Efforts to tackle overall obesity through therapeutic or lifestyle-based modulation of overall energy balance have proved extremely challenging to implement, and the manipulation of processes associated with more beneficial patterns of fat distribution offers an alternative perspective for future drug discovery.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Authors

Iris M. Heid<sup>1,2,\*</sup>, Anne U. Jackson<sup>3,\*</sup>, Joshua C. Randall<sup>4,\*</sup>, Thomas W. Winkler<sup>1,\*</sup>, Lu Qi<sup>5,6,\*</sup>, Valgerdur Steinthorsdottir<sup>7,\*</sup>, Gudmar Thorleifsson<sup>7,\*</sup>, M. Carola Zillikens<sup>8,9</sup>, Elizabeth K. Speliotes<sup>10,11</sup>, Reedik Mägi<sup>4</sup>, Tsegaselassie Workalemahu<sup>5</sup>, Charles C. White<sup>12</sup>, Nabila Bouatia-Naji<sup>13,14</sup>, Tamara B. Harris<sup>15</sup>, Sonja I. Berndt<sup>16</sup>, Erik Ingelsson<sup>17</sup>, Cristen J. Willer<sup>3</sup>, Michael N. Weedon<sup>18</sup>, Jian'an Luan<sup>19</sup>, Sailaja Vedantam<sup>10,20</sup>, Tõnu Esko<sup>21,22,23</sup>, Tuomas O. Kilpeläinen<sup>19</sup>, Zoltán Kutalik<sup>24,25</sup>, Shengxu Li<sup>19</sup>, Keri L. Monda<sup>26</sup>, Anna L. Dixon<sup>27</sup>, Christopher C. Holmes<sup>28,29</sup>, Lee M. Kaplan<sup>11,30,31</sup>, Liming Liang<sup>32,33</sup>, Josine L. Min<sup>34</sup>, Miriam F. Moffatt<sup>35</sup>, Cliona Molony<sup>36</sup>, George Nicholson<sup>29</sup>, Eric E. Schadt<sup>37,38</sup>, Krina T. Zondervan<sup>39</sup>, Mary F. Feitosa<sup>40</sup>, Teresa Ferreira<sup>4</sup>, Hana Lango Allen<sup>18</sup>, Robert J. Weyant<sup>3</sup>, Eleanor Wheeler<sup>41</sup>, Andrew R. Wood<sup>18</sup>, MAGIC<sup>42</sup>, Karol Estrada<sup>8,9,43</sup>, Michael E. Goddard<sup>44,45</sup>, Guillaume Lettre<sup>46,47</sup>, Massimo Mangino<sup>48</sup>, Dale R. Nyholt<sup>49</sup>, Shaun Purcell<sup>50,51,52</sup>, Albert Vernon Smith<sup>53,54</sup>, Peter M. Visscher<sup>55</sup>, Jian

Yang<sup>55</sup>, Steven A. McCaroll<sup>50,56,57</sup>, James Nemes<sup>56</sup>, Benjamin F. Voight<sup>50,56,57</sup>, Devin Absher<sup>58</sup>, Najaf Amin<sup>43</sup>, Thor Aspelund<sup>53,54</sup>, Lachlan Coin<sup>59</sup>, Nicole L. Glazer<sup>60,61</sup>, Caroline Hayward<sup>62</sup>, Nancy L. Heard-Costa<sup>63</sup>, Jouke-Jan Hottenga<sup>64</sup>, Åsa Johansson<sup>65,66</sup>, Toby Johnson<sup>24,25,67,68</sup>, Marika Kaakinen<sup>69,70</sup>, Karen Kapur<sup>24,25</sup>, Shamika Ketkar<sup>40</sup>, Joshua W. Knowles<sup>71</sup>, Peter Kraft<sup>32,33</sup>, Aldi T. Kraja<sup>40</sup>, Claudia Lamina<sup>2,72</sup>, Michael F. Leitzmann<sup>1</sup>, Barbara McKnight<sup>73</sup>, Andrew P. Morris<sup>4</sup>, Ken K. Ong<sup>19</sup>, John R.B. Perry<sup>18</sup>, Marjolein J. Peters<sup>8,9</sup>, Ozren Polasek<sup>74,75</sup>, Inga Prokopenko<sup>4,76</sup>, Nigel W. Rayner<sup>4,76</sup>, Samuli Ripatti<sup>77,78</sup>, Fernando Rivadeneira<sup>8,9,43</sup>, Neil R. Robertson<sup>4,76</sup>, Serena Sanna<sup>79</sup>, Ulla Sovio<sup>59</sup>, Ida Surakka<sup>77,78</sup>, Alexander Teumer<sup>80</sup>, Sophie van Wingerden<sup>43</sup>, Veronique Vitart<sup>62</sup>, Jing Hua Zhao<sup>19</sup>, Christine Cavalcanti-Proença<sup>13,14</sup>, Peter S. Chines<sup>81</sup>, Eva Fisher<sup>82</sup>, Jennifer R. Kulzer<sup>83</sup>, Cecile Lecoeur<sup>13,14</sup>, Narisu Narisu<sup>81</sup>, Camilla Sandholt<sup>84</sup>, Laura J. Scott<sup>3</sup>, Kaisa Silander<sup>77,78</sup>, Klaus Stark<sup>85</sup>, Mari-Liis Tammesoo<sup>21</sup>, Tanya M. Teslovich<sup>3</sup>, Nicholas John Timpson<sup>86</sup>, Richard M. Watanabe<sup>87,88</sup>, Ryan Welch<sup>3</sup>, Daniel I. Chasman<sup>30,89</sup>, Matthew N. Cooper<sup>90</sup>, John-Olov Jansson<sup>91</sup>, Johannes Kettunen<sup>77,78</sup>, Robert W. Lawrence<sup>90</sup>, Niina Pellikka<sup>77,78</sup>, Markus Perola<sup>77,78</sup>, Liesbeth Vandenput<sup>92</sup>, Helene Alavere<sup>21</sup>, Peter Almgren<sup>93</sup>, Larry D. Atwood<sup>63</sup>, Amanda J. Bennett<sup>76</sup>, Reiner Biffar<sup>94</sup>, Lori L. Bonnycastle<sup>81</sup>, Stefan R. Bornstein<sup>95</sup>, Thomas A. Buchanan<sup>87,96</sup>, Harry Campbell<sup>97</sup>, Ian N.M. Day<sup>86</sup>, Mariano Dei<sup>79</sup>, Marcus Dörr<sup>98</sup>, Paul Elliott<sup>59,99</sup>, Michael R. Erdos<sup>81</sup>, Johan G. Eriksson<sup>100,101,102,103,104</sup>, Nelson B. Freimer<sup>105</sup>, Mao Fu<sup>106</sup>, Stefan Gaget<sup>13,14</sup>, Eco J.C. Geus<sup>64</sup>, Anette P. Gjesing<sup>84</sup>, Harald Grallert<sup>2</sup>, Jürgen Gräßler<sup>107</sup>, Christopher J. Groves<sup>76</sup>, Candace Guiducci<sup>10</sup>, Anna-Liisa Hartikainen<sup>108</sup>, Neelam Hassanali<sup>76</sup>, Aki S. Havulinna<sup>109</sup>, Karl-Heinz Herzig<sup>70,110,111</sup>, Andrew A. Hicks<sup>112</sup>, Jennie Hui<sup>90,113,114</sup>, Wilmar Igl<sup>65</sup>, Pekka Jousilahti<sup>109</sup>, Antti Jula<sup>115</sup>, Eero Kajantie<sup>101,116</sup>, Leena Kinnunen<sup>117</sup>, Ivana Kolcic<sup>74</sup>, Seppo Koskinen<sup>109</sup>, Peter Kovacs<sup>118</sup>, Heyo K. Kroemer<sup>119</sup>, Vjekoslav Krzelj<sup>120</sup>, Johanna Kuusisto<sup>121</sup>, Kirsti Kvaloy<sup>122</sup>, Jaana Laitinen<sup>123</sup>, Olivier Lantieri<sup>124</sup>, G. Mark Lathrop<sup>125</sup>, Marja-Liisa Lokki<sup>126</sup>, Robert N. Luben<sup>127</sup>, Barbara Ludwig<sup>95</sup>, Wendy L. McArdle<sup>128</sup>, Anne McCarthy<sup>129</sup>, Mario A. Morcken<sup>81</sup>, Mari Nelis<sup>21,22,23</sup>, Matt J. Neville<sup>76</sup>, Guillaume Paré<sup>130</sup>, Alex N. Parker<sup>131</sup>, John F. Peden<sup>4,132</sup>, Irene Pichler<sup>112</sup>, Kirsi H. Pietiläinen<sup>133,134</sup>, Carl G.P. Platou<sup>122,135</sup>, Anneli Pouta<sup>108,136</sup>, Martin Ridderstråle<sup>137</sup>, Nilesh J. Samani<sup>138,139</sup>, Jouko Saramies<sup>140</sup>, Juha Sinisalo<sup>141</sup>, Jan H. Smit<sup>142</sup>, Rona J. Strawbridge<sup>143</sup>, Heather M. Stringham<sup>3</sup>, Amy J. Swift<sup>81</sup>, Maris Teder-Laving<sup>22,23</sup>, Brian Thomson<sup>10</sup>, Gianluca Usala<sup>79</sup>, Joyce B.J. van Meurs<sup>8,9,43</sup>, Gert-Jan van Ommen<sup>144,145</sup>, Vincent Vatin<sup>13,14</sup>, Claudia B. Volpato<sup>112</sup>, Henri Wallaschofski<sup>146</sup>, G. Bragi Walters<sup>7</sup>, Elisabeth Widen, Sarah H. Wild<sup>97</sup>, Gonneke Willemsen<sup>64</sup>, Daniel R. Witte<sup>147</sup>, Lina Zgaga<sup>74</sup>, Paavo Zitting<sup>148</sup>, John P. Beilby<sup>113,114,149</sup>, Alan L. James<sup>114,150</sup>, Mika Kähönen<sup>151</sup>, Terho Lehtimäki<sup>152</sup>, Markku S. Nieminen<sup>141,141</sup>, Claes Ohlsson<sup>92</sup>, Lyle J. Palmer<sup>90,114</sup>, Olli Raitakari<sup>153,154</sup>, Paul M. Ridker<sup>30,89</sup>, Michael Stumvoll<sup>155,156</sup>, Anke Tönjes<sup>155,157</sup>, Jorma Viikari<sup>158</sup>, Beverley Balkau<sup>159,160</sup>, Yoav Ben-Shlomo<sup>161</sup>, Richard N. Bergman<sup>87</sup>, Heiner Boeing<sup>82</sup>, George Davey Smith<sup>86</sup>, Shah Ebrahim<sup>162,163</sup>, Philippe Froguel<sup>13,14,164</sup>, Torben Hansen<sup>84,165</sup>, Christian Hengstenberg<sup>166,167</sup>, Kristian Hveem<sup>122</sup>, Bo Isomaa<sup>103,168</sup>, Torben

Jørgensen<sup>169,170</sup>, Fredrik Karpe<sup>76,171</sup>, Kay-Tee Khaw<sup>127</sup>, Markku Laakso<sup>121</sup>, Debbie A. Lawlor, Michel Marre<sup>172,173</sup>, Thomas Meitinger<sup>174,175</sup>, Andres Metspalu<sup>21,22,23</sup>, Kristian Midthjell<sup>122</sup>, Oluf Pedersen<sup>84,176,177</sup>, Veikko Salomaa<sup>109</sup>, Peter E.H. Schwarz<sup>178</sup>, Tiinamaija Tuomi<sup>103,179,180</sup>, Jaakko Tuomilehto<sup>117,181,182</sup>, Timo T. Valle<sup>117</sup>, Nicholas J. Wareham<sup>19</sup>, Alice M. Arnold<sup>73,183</sup>, Jacques S. Beckmann<sup>24,184</sup>, Sven Bergmann<sup>24,25</sup>, Eric Boerwinkle<sup>185</sup>, Dorret I. Boomsma<sup>64</sup>, Mark J. Caulfield<sup>68</sup>, Francis S. Collins<sup>81</sup>, Gudny Eiriksdottir<sup>53</sup>, Vilmundur Gudnason<sup>53,54</sup>, Ulf Gyllensten<sup>65</sup>, Anders Hamsten<sup>143</sup>, Andrew T. Hattersley<sup>18</sup>, Albert Hofman<sup>9,43</sup>, Frank B. Hu<sup>5,6,32</sup>, Thomas Illig<sup>2</sup>, Carlos Iribarren<sup>186,187</sup>, Marjo-Riitta Jarvelin<sup>59,69,70,136</sup>, W.H. Linda Kao<sup>188</sup>, Jaakko Kaprio<sup>77,133,189</sup>, Lenore J. Launer<sup>15</sup>, Patricia B. Munroe<sup>68</sup>, Ben Oostra<sup>190</sup>, Brenda W. Penninx<sup>142,191,192</sup>, Peter P. Pramstaller<sup>112,193,194</sup>, Bruce M. Psaty<sup>195,196</sup>, Thomas Quertermous<sup>71</sup>, Aila Rissanen<sup>134</sup>, Igor Rudan<sup>97,120</sup>, Alan R. Shuldiner<sup>106,197</sup>, Nicole Soranzo<sup>41,48</sup>, Timothy D. Spector<sup>48</sup>, Ann-Christine Syvanen<sup>198</sup>, Manuela Uda<sup>79</sup>, André Uitterlinden<sup>8,9,43</sup>, Henry Völzke<sup>199</sup>, Peter Vollenweider<sup>200</sup>, James F. Wilson<sup>97</sup>, Jacqueline C. Witteman<sup>9,43</sup>, Alan F. Wright<sup>62</sup>, Gonçalo R. Abecasis<sup>3</sup>, Michael Boehnke<sup>3</sup>, Ingrid B. Borecki<sup>40,201</sup>, Panos Deloukas<sup>41</sup>, Timothy M. Frayling<sup>18</sup>, Leif C. Groop<sup>93</sup>, Talin Haritunians<sup>202</sup>, David J. Hunter<sup>5,6,32</sup>, Robert C. Kaplan<sup>203</sup>, Kari E. North<sup>26,204</sup>, Jeffrey R. O'Connell<sup>106</sup>, Leena Peltonen, David Schlessinger<sup>206</sup>, David P. Strachan<sup>207</sup>, Joel N. Hirschhorn<sup>10,20,208</sup>, Themistocles L. Assimes<sup>71</sup>, H.-Erich Wichmann<sup>2,209,210</sup>, Unnur Thorsteinsdottir<sup>7,211</sup>, Cornelia M. van Duijn<sup>9,43</sup>, Kari Stefansson<sup>7,211,\*\*</sup>, L. Adrienne Cupples<sup>12,\*\*</sup>, Ruth J.F. Loos<sup>19,\*\*</sup>, Inês Barroso<sup>41,212,\*\*</sup>, Mark I. McCarthy<sup>4,76,171,\*\*</sup>, Caroline S. Fox<sup>213,\*\*</sup>, Karen L. Mohlke<sup>83,\*\*</sup>, and Cecilia M. Lindgren<sup>4,76,\*\*</sup>

## Affiliations

<sup>1</sup>Regensburg University Medical Center, Department of Epidemiology and Preventive Medicine, 93053 Regensburg, Germany <sup>2</sup>Institute of Epidemiology, Helmholtz Zentrum München - German Research Center for Environmental Health, 85764 Neuherberg, Germany <sup>3</sup>Department of Biostatistics, Center for Statistical Genetics, University of Michigan, Ann Arbor, Michigan 48109, USA <sup>4</sup>Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, OX3 7BN, UK <sup>5</sup>Department of Nutrition, Harvard School of Public Health, Boston, Massachusetts 02115, USA <sup>6</sup>Channing Laboratory, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts 02115, USA <sup>7</sup>deCODE Genetics, 101 Reykjavik, Iceland <sup>8</sup>Department of Internal Medicine, Erasmus MC, Rotterdam, 3015GE, The Netherlands <sup>9</sup>Netherlands Genomics Initiative (NGI)-sponsored Netherlands Consortium for Healthy Aging (NCHA) <sup>10</sup>Metabolism Initiative and Program in Medical and Population Genetics, Broad Institute, Cambridge, Massachusetts 02142, USA <sup>11</sup>Division of Gastroenterology, Massachusetts General Hospital, Boston, Massachusetts 02114, USA <sup>12</sup>Department of Biostatistics, Boston University School of Public Health, Boston, Massachusetts 02118, USA <sup>13</sup>CNRS UMR8199-IBL-Institut Pasteur de Lille, F-59019 Lille, France <sup>14</sup>University Lille Nord de France, 59000 Lille, France <sup>15</sup>Laboratory of Epidemiology, Demography, Biometry, National Institute on Aging,

National Institutes of Health, Bethesda, Maryland 20892, USA <sup>16</sup>Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Department of Health and Human Services, Bethesda, Maryland 20892, USA <sup>17</sup>Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, 17177 Stockholm, Sweden <sup>18</sup>Genetics of Complex Traits, Peninsula College of Medicine and Dentistry, University of Exeter, Exeter, EX1 2LU, UK <sup>19</sup>MRC Epidemiology Unit, Institute of Metabolic Science, Addenbrooke's Hospital, Cambridge, CB2 0QQ, UK <sup>20</sup>Divisions of Genetics and Endocrinology and Program in Genomics, Children's Hospital, Boston, Massachusetts 02115, USA <sup>21</sup>Estonian Genome Center, University of Tartu, Tartu 50410, Estonia <sup>22</sup>Estonian Biocenter, Tartu 51010, Estonia <sup>23</sup>Institute of Molecular and Cell Biology, University of Tartu, Tartu 51010, Estonia <sup>24</sup>Department of Medical Genetics, University of Lausanne, 1005 Lausanne, Switzerland <sup>25</sup>Swiss Institute of Bioinformatics, 1015 Lausanne, Switzerland <sup>26</sup>Department of Epidemiology, School of Public Health, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27514, USA <sup>27</sup>Department of Pharmacy and Pharmacology, University of Bath, Bath, BA1 1RL, UK <sup>28</sup>MRC Harwell, Harwell Science and Innovation Campus, Oxfordshire, OX11 0RD, UK <sup>29</sup>Department of Statistics, University of Oxford, Oxford OX1 3TG, UK <sup>30</sup>Harvard Medical School, Boston, Massachusetts 02115, USA <sup>31</sup>MGH Weight Center, Massachusetts General Hospital, Boston, Massachusetts 02114, USA <sup>32</sup>Department of Epidemiology, Harvard School of Public Health, Boston, Massachusetts 02115, USA <sup>33</sup>Department of Biostatistics, Harvard School of Public Health, Boston, Massachusetts 02115, USA <sup>34</sup>Human Genetics, Leiden University Medical Center, Leiden 2333, The Netherlands <sup>35</sup>National Heart and Lung Institute, Imperial College London, London SW3 6LY, UK <sup>36</sup>Merck Research Laboratories, Merck & Co., Inc., Boston, Massachusetts 02115, USA <sup>37</sup>Pacific Biosciences, Menlo Park, California 94025, USA <sup>38</sup>Sage Bionetworks, Seattle, Washington 98109, USA <sup>39</sup>Genetic and Genomic Epidemiology Unit, Wellcome Trust Centre for Human Genetics, OX3 7BN, Oxford <sup>40</sup>Department of Genetics, Washington University School of Medicine, St Louis, Missouri 63110, USA <sup>41</sup>Wellcome Trust Sanger Institute, Hinxton, Cambridge, CB10 1SA, UK <sup>42</sup>On behalf of the MAGIC (Meta-Analyses of Glucose and Insulin-related traits Consortium) investigators <sup>43</sup>Department of Epidemiology, Erasmus MC, Rotterdam, 3015GE, The Netherlands <sup>44</sup>University of Melbourne, Parkville 3010, Australia <sup>45</sup>Department of Primary Industries, Melbourne, Victoria 3001, Australia <sup>46</sup>Montreal Heart Institute, Montreal, Quebec, H1T 1C8, Canada <sup>47</sup>Department of Medicine, Université de Montréal, Montreal, Quebec, H3T 1J4, Canada <sup>48</sup>Department of Twin Research and Genetic Epidemiology, King's College London, London, SE1 7EH, UK <sup>49</sup>Neurogenetics Laboratory, Queensland Institute of Medical Research, Queensland 4006, Australia <sup>50</sup>Center for Human Genetic Research, Massachusetts General Hospital, Boston, Massachusetts 02114, USA. <sup>51</sup>The Broad Institute of Harvard and MIT, Cambridge, Massachusetts 02142, USA <sup>52</sup>Department of Psychiatry, Harvard Medical School, Boston, Massachusetts 02115, USA <sup>53</sup>Icelandic Heart Association, Kopavogur, Iceland <sup>54</sup>University of Iceland, Reykjavik, Iceland <sup>55</sup>Queensland Statistical Genetics

Laboratory, Queensland Institute of Medical Research, Queensland 4006, Australia  
<sup>56</sup>Program in Medical and Population Genetics, Broad Institute of Harvard and  
 Massachusetts Institute of Technology, Cambridge, Massachusetts 02142, USA  
<sup>57</sup>Department of Molecular Biology, Massachusetts General Hospital, Boston,  
 Massachusetts 02114, USA <sup>58</sup>Hudson Alpha Institute for Biotechnology, Huntsville,  
 Alabama 35806, USA <sup>59</sup>Department of Epidemiology and Biostatistics, School of  
 Public Health, Faculty of Medicine, Imperial College London, London, W2 1PG, UK  
<sup>60</sup>Department of Medicine, University of Washington, Seattle, Washington 98101,  
 USA <sup>61</sup>Cardiovascular Health Research Unit, University of Washington, Seattle,  
 Washington 98101, USA <sup>62</sup>MRC Human Genetics Unit, Institute for Genetics and  
 Molecular Medicine, Western General Hospital, Edinburgh, EH4 2XU, Scotland, UK  
<sup>63</sup>Department of Neurology, Boston University School of Medicine, Boston,  
 Massachusetts 02118, USA <sup>64</sup>Department of Biological Psychology, VU University  
 Amsterdam, 1081 BT Amsterdam, The Netherlands <sup>65</sup>Department of Genetics and  
 Pathology, Rudbeck Laboratory, University of Uppsala, SE-75185 Uppsala, Sweden  
<sup>66</sup>Department of Cancer Research and Molecular Medicine, Faculty of Medicine,  
 Norwegian University of Science and Technology (NTNU), Trondheim, N-7489,  
 Norway <sup>67</sup>Clinical Pharmacology, William Harvey Research Institute, Barts and The  
 London School of Medicine and Dentistry, Queen Mary, University of London,  
 London, UK <sup>68</sup>Clinical Pharmacology and Barts and The London Genome Centre,  
 William Harvey Research Institute, Barts and The London School of Medicine and  
 Dentistry, Queen Mary University of London, Charterhouse Square, London EC1M  
 6BQ, UK <sup>69</sup>Institute of Health Sciences, University of Oulu, 90014 Oulu, Finland  
<sup>70</sup>Biocenter Oulu, University of Oulu, 90014 Oulu, Finland <sup>71</sup>Department of  
 Medicine, Stanford University School of Medicine, Stanford, California 94305, USA  
<sup>72</sup>Division of Genetic Epidemiology, Department of Medical Genetics, Molecular and  
 Clinical Pharmacology, Innsbruck Medical University, 6020 Innsbruck, Austria  
<sup>73</sup>Departments of Biostatistics, University of Washington, Seattle, Washington  
 98195, USA <sup>74</sup>Andrija Stampar School of Public Health, Medical School, University  
 of Zagreb, 10000 Zagreb, Croatia <sup>75</sup>Gen-Info Ltd, 10000 Zagreb, Croatia <sup>76</sup>Oxford  
 Centre for Diabetes, Endocrinology and Metabolism, University of Oxford, Oxford,  
 OX3 7LJ, UK <sup>77</sup>Institute for Molecular Medicine Finland (FIMM), University of  
 Helsinki, 00014, Helsinki, Finland <sup>78</sup>National Institute for Health and Welfare,  
 Department of Chronic Disease Prevention, Unit of Public Health Genomics, 00014,  
 Helsinki, Finland <sup>79</sup>Istituto di Neurogenetica e Neurofarmacologia del CNR,  
 Monserrato, 09042, Cagliari, Italy <sup>80</sup>Interfaculty Institute for Genetics and Functional  
 Genomics, Ernst-Moritz-Arndt-University Greifswald, 17487 Greifswald, Germany  
<sup>81</sup>National Human Genome Research Institute, National Institutes of Health,  
 Bethesda, Maryland 20892, USA <sup>82</sup>Department of Epidemiology, German Institute  
 of Human Nutrition Potsdam-Rehbruecke, 14558 Nuthetal, Germany <sup>83</sup>Department  
 of Genetics, University of North Carolina, Chapel Hill, North Carolina 27599, USA  
<sup>84</sup>Hagedorn Research Institute, 2820 Gentofte, Denmark <sup>85</sup>Regensburg University  
 Medical Center, Clinic and Policlinic for Internal Medicine II, 93053 Regensburg,  
 Germany <sup>86</sup>MRC Centre for Causal Analyses in Translational Epidemiology,

Department of Social Medicine, Oakfield House, Bristol, BS8 2BN, UK  
<sup>87</sup>Department of Physiology and Biophysics, Keck School of Medicine, University of Southern California, Los Angeles, California 90033, USA <sup>88</sup>Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, California 90089, USA <sup>89</sup>Division of Preventive Medicine, Brigham and Women's Hospital, Boston, Massachusetts 02215, USA <sup>90</sup>Centre for Genetic Epidemiology and Biostatistics, University of Western Australia, Crawley, Western Australia 6009, Australia <sup>91</sup>Department of Physiology, Institute of Neuroscience and Physiology, Sahlgrenska Academy, University of Gothenburg, 405 30 Gothenburg, Sweden <sup>92</sup>Department of Internal Medicine, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, 413 45 Gothenburg, Sweden <sup>93</sup>Lund University Diabetes Centre, Department of Clinical Sciences, Lund University, 20502 Malmö, Sweden <sup>94</sup>Zentrum für Zahn-, Mund- und Kieferheilkunde, 17489 Greifswald, Germany <sup>95</sup>Department of Medicine III, University of Dresden, 01307 Dresden, Germany <sup>96</sup>Division of Endocrinology, Keck School of Medicine, University of Southern California, Los Angeles, California 90033, USA <sup>97</sup>Centre for Population Health Sciences, University of Edinburgh, Teviot Place, Edinburgh, EH8 9AG, Scotland <sup>98</sup>Department of Internal Medicine B, Ernst-Moritz-Arndt University, 17475 Greifswald, Germany <sup>99</sup>MRC-HPA Centre for Environment and Health, London W2 1PG, UK <sup>100</sup>Department of General Practice and Primary health Care, University of Helsinki, Helsinki, Finland <sup>101</sup>National Institute for Health and Welfare, 00271 Helsinki, Finland <sup>102</sup>Helsinki University Central Hospital, Unit of General Practice, 00280 Helsinki, Finland <sup>103</sup>Folkhalsan Research Centre, 00250 Helsinki, Finland <sup>104</sup>Vasa Central Hospital, 65130 Vasa, Finland <sup>105</sup>Center for Neurobehavioral Genetics, University of California, Los Angeles, California 90095, USA <sup>106</sup>Department of Medicine, University of Maryland School of Medicine, Baltimore, Maryland 21201, USA <sup>107</sup>Department of Medicine III, Pathobiochemistry, University of Dresden, 01307 Dresden, Germany <sup>108</sup>Department of Clinical Sciences/Obstetrics and Gynecology, University of Oulu, 90014 Oulu, Finland <sup>109</sup>National Institute for Health and Welfare, Department of Chronic Disease Prevention, Chronic Disease Epidemiology and Prevention Unit, 00014, Helsinki, Finland <sup>110</sup>Institute of Biomedicine, Department of Physiology, University of Oulu, 90014 Oulu, Finland <sup>111</sup>Department of Psychiatry, Kuopio University Hospital and University of Kuopio, 70210 Kuopio, Finland <sup>112</sup>Institute of Genetic Medicine, European Academy Bozen/Bolzano (EURAC), Bolzano/Bozen, 39100, Italy. Affiliated Institute of the University of Lübeck, Lübeck, Germany <sup>113</sup>PathWest Laboratory of Western Australia, Department of Molecular Genetics, J Block, QEII Medical Centre, Nedlands, Western Australia 6009, Australia <sup>114</sup>Busselton Population Medical Research Foundation Inc., Sir Charles Gairdner Hospital, Nedlands, Western Australia 6009, Australia <sup>115</sup>National Institute for Health and Welfare, Department of Chronic Disease Prevention, Population Studies Unit, 20720 Turku, Finland <sup>116</sup>Hospital for Children and Adolescents, Helsinki University Central Hospital and University of Helsinki, 00029 HUS, Finland <sup>117</sup>National Institute for Health and Welfare, Diabetes Prevention Unit, 00271 Helsinki, Finland

<sup>118</sup>Interdisciplinary Centre for Clinical Research, University of Leipzig, 04103 Leipzig, Germany <sup>119</sup>Institut für Pharmakologie, Universität Greifswald, 17487 Greifswald, Germany <sup>120</sup>Croatian Centre for Global Health, School of Medicine, University of Split, Split 21000, Croatia <sup>121</sup>Department of Medicine, University of Kuopio and Kuopio University Hospital, 70210 Kuopio, Finland <sup>122</sup>HUNT Research Centre, Department of Public Health and General Practice, Norwegian University of Science and Technology, 7600 Levanger, Norway <sup>123</sup>Finnish Institute of Occupational Health, 90220 Oulu, Finland <sup>124</sup>Institut inter-regional pour la sante (IRSA), F-37521 La Riche, France. <sup>125</sup>Centre National de Genotypage, Evry, Paris 91057, France <sup>126</sup>Transplantation Laboratory, Haartman Institute, University of Helsinki, 00014, Helsinki, Finland <sup>127</sup>Department of Public Health and Primary Care, Institute of Public Health, University of Cambridge, Cambridge CB2 2SR, UK <sup>128</sup>Avon Longitudinal Study of Parents and Children (ALSPAC) Laboratory, Department of Social Medicine, University of Bristol, Bristol, BS8 2BN, UK <sup>129</sup>Division of Health, Research Board, An Bord Taighde Sláinte, Dublin, 2, Ireland <sup>130</sup>Department of Pathology and Molecular Medicine, McMaster University, Hamilton, Ontario L8N3Z5, Canada <sup>131</sup>Amgen, Cambridge, Massachusetts 02139, USA <sup>132</sup>Department of Cardiovascular Medicine, University of Oxford, Level 6 West Wing, John Radcliffe Hospital, Headley Way, Headington, Oxford, OX3 9DU <sup>133</sup>Finnish Twin Cohort Study, Department of Public Health, University of Helsinki, 00014, Helsinki, Finland <sup>134</sup>Obesity Research unit, Department of Psychiatry, Helsinki University Central Hospital, Helsinki, Finland <sup>135</sup>Department of Medicine, Levanger Hospital, The Nord-Trøndelag Health Trust, 7600 Levanger, Norway <sup>136</sup>National Institute for Health and Welfare, 90101 Oulu, Finland <sup>137</sup>Department of Clinical Sciences, Lund University, 20502 Malmö, Sweden <sup>138</sup>Department of Cardiovascular Sciences, University of Leicester, Glenfield Hospital, Leicester, LE3 9QP, UK <sup>139</sup>Leicester NIHR Biomedical Research Unit in Cardiovascular Disease, Glenfield Hospital, Leicester, LE3 9QP, UK <sup>140</sup>South Karelia Central Hospital, 53130 Lappeenranta, Finland <sup>141</sup>Division of Cardiology, Cardiovascular Laboratory, Helsinki University Central Hospital, 00029 Helsinki, Finland <sup>142</sup>Department of Psychiatry/EMGO Institute, VU University Medical Center, 1081 BT Amsterdam, The Netherlands <sup>143</sup>Atherosclerosis Research Unit, Department of Medicine, Solna, Karolinska Institutet, Karolinska University Hospital, 171 76 Stockholm, Sweden <sup>144</sup>Department of Human Genetics, Leiden University Medical Center, 2333 ZC Leiden, the Netherlands <sup>145</sup>Center of Medical Systems Biology, Leiden University Medical Center, 2333 ZC Leiden, the Netherlands <sup>146</sup>Institut für Klinische Chemie und Laboratoriumsmedizin, Universität Greifswald, 17475 Greifswald, Germany <sup>147</sup>Steno Diabetes Center, 2820 Gentofte, Denmark <sup>148</sup>Department of Physiatrics, Lapland Central Hospital, 96101 Rovaniemi, Finland <sup>149</sup>School of Pathology and Laboratory Medicine, University of Western Australia, Nedlands, Western Australia 6009, Australia <sup>150</sup>School of Medicine and Pharmacology, University of Western Australia, Perth, Western Australia 6009, Australia <sup>151</sup>Department of Clinical Physiology, University of Tampere and Tampere University Hospital, 33520 Tampere, Finland <sup>152</sup>Department of Clinical Chemistry,

University of Tampere and Tampere University Hospital, 33520 Tampere, Finland  
<sup>153</sup>Research Centre of Applied and Preventive Cardiovascular Medicine, University of Turku, 20520 Turku, Finland <sup>154</sup>The Department of Clinical Physiology, Turku University Hospital, 20520 Turku, Finland <sup>155</sup>Department of Medicine, University of Leipzig, 04103 Leipzig, Germany <sup>156</sup>LIFE Study Centre, University of Leipzig, Leipzig, Germany <sup>157</sup>Coordination Centre for Clinical Trials, University of Leipzig, Härtelstr. 16-18, 04103 Leipzig, Germany <sup>158</sup>Department of Medicine, University of Turku and Turku University Hospital, 20520 Turku, Finland <sup>159</sup>INSERM CESP Centre for Research in Epidemiology and Public Health U1018, Epidemiology of diabetes, obesity and chronic kidney disease over the lifecourse, 94807 Villejuif, France <sup>160</sup>University Paris Sud 11, UMRS 1018, 94807 Villejuif, France  
<sup>161</sup>Department of Social Medicine, University of Bristol, Bristol, BS8 2PS, UK <sup>162</sup>The London School of Hygiene and Tropical Medicine, London, WC1E 7HT, UK  
<sup>163</sup>South Asia Network for Chronic Disease <sup>164</sup>Department of Genomics of Common Disease, School of Public Health, Imperial College London, W12 0NN, London, UK  
<sup>165</sup>Faculty of Health Science, University of Southern Denmark, 5000 Odense, Denmark <sup>166</sup>Klinik und Poliklinik für Innere Medizin II, Universität Regensburg, 93053 Regensburg, Germany <sup>167</sup>Regensburg University Medical Center, Innere Medizin II, 93053 Regensburg, Germany <sup>168</sup>Department of Social Services and Health Care, 68601 Jakobstad, Finland <sup>169</sup>Research Centre for Prevention and Health, Glostrup University Hospital, 2600 Glostrup, Denmark <sup>170</sup>Faculty of Health Science, University of Copenhagen, 2100 Copenhagen, Denmark <sup>171</sup>NIHR Oxford Biomedical Research Centre, Churchill Hospital, Oxford, OX3 7LJ, UK  
<sup>172</sup>Department of Endocrinology, Diabetology and Nutrition, Bichat-Claude Bernard University Hospital, Assistance Publique des Hôpitaux de Paris, F-75018 Paris, France <sup>173</sup>Cardiovascular Genetics Research Unit, Université Henri Poincaré-Nancy 1, 54000, Nancy, France <sup>174</sup>Institute of Human Genetics, Klinikum rechts der Isar der Technischen Universität München, 81675 Munich, Germany <sup>175</sup>Institute of Human Genetics, Helmholtz Zentrum München - German Research Center for Environmental Health, 85764 Neuherberg, Germany <sup>176</sup>Institute of Biomedical Sciences, University of Copenhagen, 2200 Copenhagen, Denmark <sup>177</sup>Faculty of Health Science, University of Aarhus, 8000 Aarhus, Denmark <sup>178</sup>Department of Medicine III, Prevention and Care of Diabetes, University of Dresden, 01307 Dresden, Germany <sup>179</sup>Department of Medicine, Helsinki University Central Hospital, 00290 Helsinki, Finland <sup>180</sup>Research Program of Molecular Medicine, University of Helsinki, 00014 Helsinki, Finland <sup>181</sup>Hjelt Institute, Department of Public Health, University of Helsinki, 00014 Helsinki, Finland <sup>182</sup>South Ostrobothnia Central Hospital, 60220 Seinäjoki, Finland <sup>183</sup>Collaborative Health Studies Coordinating Center, Seattle, Washington 98115, USA <sup>184</sup>Service of Medical Genetics, Centre Hospitalier Universitaire Vaudois (CHUV) University Hospital, 1011 Lausanne, Switzerland <sup>185</sup>Human Genetics Center and Institute of Molecular Medicine, University of Texas Health Science Center, Houston, Texas 77030, USA <sup>186</sup>Division of Research, Kaiser Permanente Northern California, Oakland, California 94612, USA <sup>187</sup>Department of Epidemiology and Biostatistics, University of California, San



Francisco, San Francisco, California 94107, USA <sup>188</sup>Department of Epidemiology and Medicine, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland 21205, USA <sup>189</sup>National Institute for Health and Welfare, Department of Mental Health and Substance Abuse Services, Unit for Child and Adolescent Mental Health, 00271 Helsinki, Finland <sup>190</sup>Department of Clinical Genetics, Erasmus MC, Rotterdam, 3015GE, The Netherlands <sup>191</sup>Department of Psychiatry, Leiden University Medical Centre, 2300 RC Leiden, The Netherlands <sup>192</sup>Department of Psychiatry, University Medical Centre Groningen, 9713 GZ Groningen, The Netherlands <sup>193</sup>Department of Neurology, General Central Hospital, Bolzano, Italy <sup>194</sup>Department of Neurology, University of Lübeck, Lübeck, Germany. <sup>195</sup>Departments of Epidemiology, Medicine and Health Services, University of Washington, Seattle, Washington 98195, USA <sup>196</sup>Group Health Research Institute, Group Health, Seattle, Washington 98101, USA <sup>197</sup>Geriatrics Research and Education Clinical Center, Baltimore Veterans Administration Medical Center, Baltimore, Maryland 21201, USA <sup>198</sup>Uppsala University / Dept. of Medical Sciences, Molecular Medicine, 751 85 Uppsala, Sweden <sup>199</sup>Institut für Community Medicine, 17489 Greifswald, Germany <sup>200</sup>Department of Internal Medicine, Centre Hospitalier Universitaire Vaudois (CHUV) University Hospital, 1011 Lausanne, Switzerland <sup>201</sup>Division of Biostatistics, Washington University School of Medicine, St.Louis, Missouri 63110, USA <sup>202</sup>Medical Genetics Institute, Cedars-Sinai Medical Center, Los Angeles, California 90048, USA <sup>203</sup>Department of Epidemiology and Population Health, Albert Einstein College of Medicine, Bronx, New York 10461, USA <sup>204</sup>Carolina Center for Genome Sciences, School of Public Health, University of North Carolina Chapel Hill, Chapel Hill, North Carolina 27514, USA <sup>205</sup>Department of Medical Genetics, University of Helsinki, 00014 Helsinki, Finland <sup>206</sup>Laboratory of Genetics, National Institute on Aging, Baltimore, Maryland 21224, USA <sup>207</sup>Division of Community Health Sciences, St George's, University of London, London, SW17 0RE, UK <sup>208</sup>Department of Genetics, Harvard Medical School, Boston, Massachusetts 02115, USA <sup>209</sup>Klinikum Grosshadern, 81377 Munich, Germany <sup>210</sup>Ludwig-Maximilians-Universität, Institute of Medical Informatics, Biometry and Epidemiology, Chair of Epidemiology, 81377 Munich, Germany <sup>211</sup>Faculty of Medicine, University of Iceland, 101 Reykjavík, Iceland <sup>212</sup>University of Cambridge Metabolic Research Labs, Institute of Metabolic Science Addenbrooke's Hospital, CB2 0QQ, Cambridge, UK <sup>213</sup>Division of Intramural Research, National Heart, Lung and Blood Institute, Framingham Heart Study, Framingham, Massachusetts 01702, USA

## ACKNOWLEDGEMENTS

Academy of Finland (104781, 120315, 129269, 117797, 121584, 126925, 129418, 129568, 77299, 124243, 213506, 129680, 129494, 10404, 213506, 129680, 114382, 126775, 127437, 129255, 129306, 130326, 209072, 210595, 213225, 216374); ADA Mentor-Based Postdoctoral Fellowship grant; Affymetrix, Inc for genotyping services (N02-HL-6-4278); ALF/LUA Gothenburg; Althingi (the Icelandic Parliament); Amgen; AstraZeneca AB; Augustinus Foundation; Becket Foundation; Biocentrum Helsinki; Biomedicum Helsinki Foundation; Boston Obesity Nutrition Research Center (DK46200); British Diabetes Association (1192); British Diabetic Association Research; British Heart Foundation (97020, PG/02/128); Busselton Population Medical Research Foundation; Cambridge NIHR Comprehensive Biomedical Research Centre; CamStrad; Chief Scientist Office of the Scottish

Government; Contrat Plan Etat Région de France; Danish Centre for Health Technology Assessment; Danish Diabetes Association; Danish Ministry of Internal Affairs and Health; Danish Heart Foundation; Danish Pharmaceutical Association; Danish Research Council; DIAB Core (German Network of Diabetes); Diabetes UK; Donald W. Reynolds Foundation; Dr Knut Krohn, Microarray Core Facility of the Interdisciplinary Centre for Clinical Research (IZKF), University of Leipzig, Germany; Dresden University of Technology Funding Grant, Med Drive; EMGO+ institute; Emil and Vera Cornell Foundation; Erasmus Medical Center and Erasmus University, Rotterdam; Estonian Government SF0180142s08; European Commission (2004310, 212111, 205419, 245536, DG XII, HEALTH-F4-2007-201413, FP7/2007-2013, QLG1-CT-2000-01643, QLG2-CT-2002-01254, LSHG-CT-2006-018947, LSHG-CT-2006-01947, LSHG-CT-2004-512066, LSHM-CT-2007-037273, EU/WLRT-2001-01254, LSHG-CT-2004-518153, SOC 95201408 05F02, Marie Curie Intra-European Fellowship); Federal Ministry of Education and Research, Germany (01ZZ9603, 01ZZ0103, 01ZZ0403, 03ZIK012, 01 EA 9401); Federal State of Mecklenburg- West Pomerania; Finnish Diabetes Research Foundation; Finnish Diabetes Research Society; Finnish Foundation for Pediatric Research; Finnish Foundation of Cardiovascular Research; Finnish Medical Society; Finska Läkaresällskapet; Finnish Ministry of Education; Folkhälsan Research Foundation; Fond Européen pour le Développement Régional; Fondation LeDucq; Foundation for Life and Health in Finland; GEN-AU "GOLD" from Austria; German Bundesministerium fuer Forschung und Technology (#01 AK 803 A-H, # 01 IG 07015 G); German National Genome Research Net NGFN2 and NGFNplus (01GS0823, FKZ 01GS0823); German Research Council (KFO-152); GlaxoSmithKline; Göteborg Medical Society; Gyllenberg Foundation; Health Care Centers in Vasa, Närpes and Korsholm; Healthway, Western Australia; Helmholtz Center Munich; Helsinki University Central Hospital; Hjartavernd (the Icelandic Heart Association); Ib Henriksen Foundation; IZKF (B27); Jalmari and Rauha Ahokas Foundation; Juho Vainio Foundation; Juvenile Diabetes Research Foundation International (JDRF); Karolinska Institute and the Stockholm County Council (560183); Knut and Alice Wallenberg Foundation; Lundbeck Foundation Centre of Applied Medical Genomics for Personalized Disease Prediction, Prevention and Care; Lundberg Foundation; MC Health; Ministry of Cultural Affairs of the Federal State of Mecklenburg-West Pomerania, Germany; South Tyrol Ministry of Health; Ministry of Science, Education and Sport of the Republic of Croatia (216-1080315-0302); Medical Research Council UK (G0000649, G0601261, G9521010D, G0000934, G0500539, G0600331, PrevMetSyn); Montreal Heart Institute Foundation; MRC Centre for Obesity related Metabolic disease; Municipal Health Care Center and Hospital in Jakobstad; Municipality of Rotterdam; Närpes Health Care Foundation; National Health and Medical Research Council of Australia and the Great Wine Estates Auctions; Netherlands Centre for Medical Systems Biology (SPI 56-464-1419); Netherlands Ministry for Health, Welfare and Sports; Netherlands Ministry of Education, Culture and Science; Netherlands Genomics Initiative; Netherlands Consortium for Healthy Aging (050-060-810); Netherlands Organisation of Scientific Research NWO Investments (175.010.2005.011, 911-03-012, 904-61-090,904-61-193,480-04-004,400-05-717); National Institute on Aging Intramural Research Program; National Institutes of Health (CA047988, CA65725, CA87969, CA49449, CA67262, CA50385, , DK075787, DK062370, DK58845, DK072193, K23-DK080145, K99HL094535, N01-HC85079 through N01-HC85086, N01-HG-65403; N01-AG-12100, N01-HC-25195, N01-HC35129, N01-HC15103, N01-HC55222, N01-HC75150, N01-HC45133, N01-HC55015, N01-HC55016, N01-HC-55018, N01-HC-55019, N01-HC-55020, N01-HC-55021, N01-HC-55022, N01-AG-1-2109, HL71981, HG005581, HG002651, HL084729, HL043851, HHSN268200625226C, K23-DK080145; MH084698, P30-DK072488, R01-DK075787, R01 HL087652, R01-HL087641, R01-HL59367, R01-HL086694, R01-HL087647, R01-HL087679, R01-HL087700, R01-AG031890, R01-HL088119, R01-DK068336, R01-DK075681, R01-DK-073490, R01-DK075787, R01-MH63706; U01-HL72515, U01-GM074518, U01-HL084756, U01-HG004399, UO1-CA098233, UL1-RR025005, UL1-RR025005, U01-HG004402, U01-DK062418; U01 HL080295, T32-HG00040, 263-MA-410953; 1RL1-MH083268-01, intramural project 1Z01-HG000024); National Institute for Health Research (NIHR); Neuroscience Campus Amsterdam; Novo Nordisk Foundation; Novo Nordisk Inc., Research Foundation of Copenhagen County; Ollqvist Foundation; Paavo Nurmi Foundation; Päivikki and Sakari Sohlberg Foundation; Pew Scholarship for the Biomedical Sciences); Perklén Foundation; Petrus and Augusta Hedlunds Foundation; Research Institute for Diseases in the Elderly (014-93-015, RIDE, RIDE2); Sahlgrenska Center for Cardiovascular and Metabolic Research (CMR, A305:188); Siemens Healthcare, Erlangen, Germany; Signe and Ane Gyllenberg Foundation; Sigrid Juselius Foundation; Social Insurance Institution of Finland; Social Ministry of the Federal State of Mecklenburg-West Pomerania, Germany; South Tyrolean Sparkasse Foundation; State of Bavaria, Germany; Support for Science Funding programme; Swedish Cultural Foundation in Finland; Swedish Foundation for Strategic Research (SSF); Swedish Heart-Lung Foundation; Swedish Medical Research Council (8691, K2007-66X-20270-01-3, K2010-54X-09894-19-3); Swedish Society of Medicine; Swiss National Science Foundation (33CSO-122661); the Royal Society; the Royal Swedish Academy of Science; Torsten and Ragnar Söderberg's Foundation; Turku University Hospitals; UK Department of Health Policy Research Programme; University and Research of the Autonomous Province of Bolzano; University Hospital Medical funds to Tampere; University Hospital Oulu, Biocenter, University of Oulu, Finland (75617); Västra Götaland Foundation; Wellcome Trust (077016/Z/05/Z, 068545/Z/02, 072960, 076113, 083270, 085301, 079557, 081682, 075491, 076113/B/04/Z, 091746/Z/10/Z, 079895, WT086596/Z/08/Z, WT Research Career Development Fellowship; WT Career Development Award); Western Australian Genetic Epidemiology Resource and the Western Australian DNA Bank (both National Health and Medical Research Council of Australia Enabling Facilities); Yrjö Jahnsson Foundation.

## Abbreviations


<b>BMI</b>	Body-mass-index
<b>WC</b>	waist circumference
<b>WHR</b>	waist-hip ratio
<b>HIP</b>	hip circumference
<b>GWA</b>	genome-wide association
<b>SNP</b>	single-nucleotide polymorphism
<b>LD</b>	linkage disequilibrium
<b>eQTL</b>	Expression Quantitative Trait Loci

## REFERENCES

- Carey VJ, et al. Body fat distribution and risk of non-insulin-dependent diabetes mellitus in women. The Nurses' Health Study. *Am J Epidemiol.* 1997; 145:614–619. [PubMed: 9098178]
- Wang Y, Rimm EB, Stampfer MJ, Willett WC, Hu FB. Comparison of abdominal adiposity and overall obesity in predicting risk of type 2 diabetes among men. *Am J Clin Nutr.* 2005; 81:555–563. [PubMed: 15755822]
- Canoy D. Distribution of body fat and risk of coronary heart disease in men and women. *Curr Opin Cardiol.* 2008; 23:591–598. [PubMed: 18830075]
- Pischon T, et al. General and abdominal adiposity and risk of death in Europe. *N Engl J Med.* 2008; 359:2105–2120. [PubMed: 19005195]
- Snijder MB, et al. Associations of hip and thigh circumferences independent of waist circumference with the incidence of type 2 diabetes: the Hoorn Study. *Am J Clin Nutr.* 2003; 77:1192–1197. [PubMed: 12716671]
- Snijder MB, et al. Trunk fat and leg fat have independent and opposite associations with fasting and postload glucose levels: the Hoorn study. *Diabetes Care.* 2004; 27:372–377. [PubMed: 14747216]
- Mills GW, et al. Heritability estimates for beta cell function and features of the insulin resistance syndrome in UK families with an increased susceptibility to type 2 diabetes. *Diabetologia.* 2004; 47:732–738. [PubMed: 15298351]
- Souren NY, et al. Anthropometry, carbohydrate and lipid metabolism in the East Flanders Prospective Twin Survey: heritabilities. *Diabetologia.* 2007; 50:2107–2116. [PubMed: 17694296]
- Rose KM, Newman B, Mayer-Davis EJ, Selby JV. Genetic and behavioral determinants of waist-hip ratio and waist circumference in women twins. *Obes Res.* 1998; 6:383–392. [PubMed: 9845227]
- Selby JV, et al. Genetic and behavioral influences on body fat distribution. *Int J Obes.* 1990; 14:593–602. [PubMed: 2228394]
- Agarwal AK, Garg A. Genetic disorders of adipose tissue development, differentiation, and death. *Annu Rev Genomics Hum Genet.* 2006; 7:175–199. [PubMed: 16722806]
- Garg A. Acquired and inherited lipodystrophies. *N Engl J Med.* 2004; 350:1220–1234. [PubMed: 15028826]
- Lindgren CM, et al. Genome-wide association scan meta-analysis identifies three Loci influencing adiposity and fat distribution. *PLoS Genet.* 2009; 5 e1000508.
- Thorleifsson G, et al. Genome-wide association yields new sequence variants at seven loci that associate with measures of obesity. *Nat Genet.* 2009; 41:18–24. [PubMed: 19079260]
- Willer CJ, et al. Six new loci associated with body mass index highlight a neuronal influence on body weight regulation. *Nat Genet.* 2009; 41:25–34. [PubMed: 19079261]
- Loos RJ, et al. Common variants near MC4R are associated with fat mass, weight and risk of obesity. *Nat Genet.* 2008; 40:768–775. [PubMed: 18454148]

17. Zillikens MC, et al. Sex-specific genetic effects influence variation in body composition. *Diabetologia*. 2008; 51:2233–2241. [PubMed: 18839131]
18. Frayling TM, et al. A common variant in the FTO gene is associated with body mass index and predisposes to childhood and adult obesity. *Science*. 2007; 316:889–894. [PubMed: 17434869]
19. Kathiresan S, et al. Common variants at 30 loci contribute to polygenic dyslipidemia. *Nat Genet*. 2009; 41:56–65. [PubMed: 19060906]
20. Dupuis J, et al. New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nat Genet*. 42:105–116. [PubMed: 20081858]
21. Saxena R, et al. Genetic variation in GIPR influences the glucose and insulin responses to an oral glucose challenge. *Nat Genet*. 42:142–148. [PubMed: 20081857]
22. Raychaudhuri S, et al. Identifying relationships among genomic disease regions: predicting genes at pathogenic SNP associations and rare deletions. *PLoS Genet*. 2009; 5 e1000534.
23. Thomas PD, et al. PANTHER: a library of protein families and subfamilies indexed by function. *Genome Res*. 2003; 13:2129–2141. [PubMed: 12952881]
24. Bochukova EG, et al. Large, rare chromosomal deletions associated with severe early-onset obesity. *Nature*. 2009
25. Walters RG, et al. A new highly penetrant form of obesity due to deletions on chromosome 16p11.2. *Nature*. 463:671–675. [PubMed: 20130649]
26. Conrad DF, et al. Origins and functional impact of copy number variation in the human genome. *Nature*. 2009
27. Genome-wide association study of CNVs in 16,000 cases of eight common diseases and 3,000 shared controls. *Nature*. 464:713–7420. [PubMed: 20360734]
28. Pennacchio LA, Loots GG, Nobrega MA, Ovcharenko I. Predicting tissue-specific enhancers in the human genome. *Genome Res*. 2007; 17:201–211. [PubMed: 17210927]
29. Emilsson V, et al. Genetics of gene expression and its effect on disease. *Nature*. 2008; 452:423–428. [PubMed: 18344981]
30. Schadt EE, et al. Mapping the genetic architecture of gene expression in human liver. *PLoS Biol*. 2008; 6 e107.
31. Dixon AL, et al. A genome-wide association study of global gene expression. *Nat Genet*. 2007; 39:1202–1207. [PubMed: 17873877]
32. Speliotes EK, et al. Association analyses of 249,796 individuals reveal eighteen new loci associated with body mass index. *Nat Genet*. (in press).
33. Wells JC. Sexual dimorphism of body composition. *Best Pract Res Clin Endocrinol Metab*. 2007; 21:415–430. [PubMed: 17875489]
34. Aulchenko YS, et al. Loci influencing lipid levels and coronary heart disease risk in 16 European population cohorts. *Nat Genet*. 2009; 41:47–55. [PubMed: 19060911]
35. Doring A, et al. SLC2A9 influences uric acid concentrations with pronounced sex-specific effects. *Nat Genet*. 2008; 40:430–436. [PubMed: 18327256]
36. Shifman S, et al. Genome-wide association identifies a common variant in the reelin gene that increases the risk of schizophrenia only in women. *PLoS Genet*. 2008; 4:e28. [PubMed: 18282107]
37. Ridker PM, et al. Polymorphism in the CETP gene region, HDL cholesterol, and risk of future myocardial infarction: Genomewide analysis among 18 245 initially healthy women from the Women's Genome Health Study. *Circ Cardiovasc Genet*. 2009; 2:26–33. [PubMed: 20031564]
38. Cooney GJ, et al. Improved glucose homeostasis and enhanced insulin signalling in Grb14-deficient mice. *EMBO J*. 2004; 23:582–593. [PubMed: 14749734]
39. Zeggini E, et al. Meta-analysis of genome-wide association data and large-scale replication identifies additional susceptibility loci for type 2 diabetes. *Nat Genet*. 2008; 40:638–645. [PubMed: 18372903]
40. Voight BF, et al. Twelve type 2 diabetes susceptibility loci identified through large-scale association analysis. *Nature Genetics*. accepted.


41. Boesgaard TW, et al. Variant near ADAMTS9 known to associate with type 2 diabetes is related to insulin resistance in offspring of type 2 diabetes patients--EUGENE2 study. *PLoS One*. 2009; 4:e7236.
42. Nishimura S, et al. Adipogenesis in obesity requires close interplay between differentiating adipocytes, stromal cells, and blood vessels. *Diabetes*. 2007; 56:1517–1526. [PubMed: 17389330]
43. Silha JV, Krsek M, Sucharda P, Murphy LJ. Angiogenic factors are elevated in overweight and obese individuals. *Int J Obes (Lond)*. 2005; 29:1308–1314. [PubMed: 15953938]
44. Garcia de la Torre N, et al. Effects of weight loss after bariatric surgery for morbid obesity on vascular endothelial growth factor-A, adipocytokines, and insulin. *J Clin Endocrinol Metab*. 2008; 93:4276–4281. [PubMed: 18713823]
45. Gesta S, et al. Evidence for a role of developmental genes in the origin of obesity and body fat distribution. *Proc Natl Acad Sci U S A*. 2006; 103:6676–6681. [PubMed: 16617105]
46. Gesta S, Tseng YH, Kahn CR. Developmental origin of fat: tracking obesity to its source. *Cell*. 2007; 131:242–256. [PubMed: 17956727]
47. Lanctot C, Kaspar C, Cremer T. Positioning of the mouse Hox gene clusters in the nuclei of developing embryos and differentiating embryoid bodies. *Exp Cell Res*. 2007; 313:1449–1459. [PubMed: 17346703]
48. Candille SI, et al. Dorsoventral patterning of the mouse coat by Tbx15. *PLoS Biol*. 2004; 2:E3.
49. O'Rahilly S. Human genetics illuminates the paths to metabolic disease. *Nature*. 2009; 462:307–314. [PubMed: 19924209]
50. Krotkiewski M, Bjorntorp P, Sjostrom L, Smith U. Impact of obesity on metabolism in men and women. Importance of regional adipose tissue distribution. *J Clin Invest*. 1983; 72:1150–1162. [PubMed: 6350364]
51. Fox CS, et al. Abdominal visceral and subcutaneous adipose tissue compartments: association with metabolic risk factors in the Framingham Heart Study. *Circulation*. 2007; 116:39–48. [PubMed: 17576866]
52. Scherzer R, et al. Simple anthropometric measures correlate with metabolic risk indicators as strongly as magnetic resonance imaging-measured adipose tissue depots in both HIV-infected and control subjects. *Am J Clin Nutr*. 2008; 87:1809–1817. [PubMed: 18541572]
53. Vega GL, et al. Influence of body fat content and distribution on variation in metabolic risk. *J Clin Endocrinol Metab*. 2006; 91:4459–4466. [PubMed: 16926254]
54. Li Y, Willer C, Sanna S, Abecasis G. Genotype imputation. *Annu Rev Genomics Hum Genet*. 2009; 10:387–406. [PubMed: 19715440]
55. Marchini J, Howie B, Myers S, McVean G, Donnelly P. A new multipoint method for genome-wide association studies by imputation of genotypes. *Nat Genet*. 2007; 39:906–913. [PubMed: 17572673]
56. Servin B, Stephens M. Imputation-based analysis of association studies: candidate regions and quantitative traits. *PLoS Genet*. 2007; 3:e114. [PubMed: 17676998]
57. Cox, DR.; Hinkley, DV. *Theoretical Statistics*. London: Chapman and Hall; 1979.
58. Devlin B, Roeder K. Genomic control for association studies. *Biometrics*. 1999; 55:997–1004. [PubMed: 11315092]
59. Stouffer, SA.; Suchman, EA.; De Vinney, LC.; Star, SA.; Williams, RM. *Adjustment during army life*. Princeton, NJ: Princeton University Press; 1949.
60. Whitlock MC. Combining probability from independent tests: the weighted Z-method is superior to Fisher's approach. *Journal of Evolutionary Biology*. 2005; 18:1368–1373. [PubMed: 16135132]
61. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate - a Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B-Methodological*. 1995; 57:289–300.



**Figure 1. Genome-wide association analyses for WHR in discovery studies**


A. Manhattan plot shows results of the WHR association meta-analysis in discovery studies ( $P$  on the y-axis and SNP genomic position on the x-axis). Colored genomic loci indicate significant association ( $P < 5 \times 10^{-8}$ ) detected previously (blue)<sup>13</sup>, in our GWA stage (red), and after the meta-analysis combining GWA and follow-up studies (orange). Two loci tested in the follow-up stage did not achieve genome-wide significance (green).

B. Quantile-quantile (QQ) plot of SNPs for the discovery meta-analysis of WHR (black) and after removing SNPs within 1 Mb of either the recently reported *LYPLAL1* signal (blue) or the 14 significant associations (green). The grey area represents the 95% confidence interval around the test statistic under the null distribution.



**Figure 2. Regional plots of 14 loci with genome-wide significant association**

SNP association with WHR in meta-analysis of discovery studies for 14 loci ( $-\log_{10} P$  on the y-axis and SNP genomic position on the x-axis). In each panel, an index SNP is denoted with a purple diamond and plotted using the  $P$  attained across discovery and follow-up data (Table 1). Estimated recombination rates are plotted in blue. SNPs are colored to reflect LD with the index SNP (pair-wise  $r^2$  values from HapMap CEU). Gene and microRNA annotations are from the UCSC genome browser.




**Figure 3. Association of the 14 WHR loci with waist and hip circumference**


Beta-coefficients for waist circumference (WC, x-axis) and hip circumference (HIP, y-axis) in women and men derived from the joint discovery and follow-up analysis. *P* for WC and HIP are represented by color. In men, grey gene labels refer to those SNPs that were not significant in the male-specific WHR analysis. More details can be found in Supplementary Table 3.



**Women**



**Men**



**Figure 4.**

Table 1

Fourteen SNPs associated with WHR at genome-wide significance levels

SNP	Chr	Position (b36)	Nearby genes	EA <sup>a</sup>	EAF <sup>b</sup>	Discovery			Follow-up			Combined		
						P	Beta	N	P	Beta	N	P	Beta	N
<b>SNPs evaluated in follow-up achieving genome-wide significance</b>														
rs9491696	6	127,494,332	<i>RSPO3</i>	G	0.520	2.10E-14	0.037	77,164	3.27E-28	0.045	113,582	1.84E-40	0.042	
rs6905288	6	43,866,851	<i>VEGFA</i>	A	0.562	4.72E-10	0.033	77,129	1.18E-16	0.039	95,430	5.88E-25	0.036	
rs984222	1	119,305,366	<i>TBX15-WARS2</i>	G	0.365	3.81E-14	0.037	77,167	1.56E-12	0.031	109,623	8.69E-25	0.034	
rs1055144	7	25,837,634	<i>NFE2L3</i>	T	0.210	1.49E-08	0.034	77,145	3.26E-18	0.043	113,636	9.97E-25	0.040	
rs10195252	2	165,221,337	<i>GRB14</i>	T	0.599	3.23E-10	0.031	77,119	3.18E-16	0.036	102,449	2.09E-24	0.033	
rs4846567	1	217,817,340	<i>LYPAL1</i>	G	0.283	2.37E-12	0.037	77,167	3.15E-10	0.032	91,820	6.89E-21	0.034	
rs1011731	1	170,613,171	<i>DNM3-PTG</i>	G	0.572	1.72E-10	0.031	77,094	7.47E-09	0.026	92,018	9.51E-18	0.028	
rs718314	12	26,344,550	<i>ITPR2-SSPN</i>	G	0.741	2.41E-08	0.031	77,167	1.49E-10	0.030	107,503	1.14E-17	0.030	
rs1294421	6	6,688,148	<i>LY86</i>	G	0.387	6.31E-09	0.029	77,154	2.69E-10	0.028	102,189	1.75E-17	0.028	
rs1443512	12	52,628,951	<i>HOXC13</i>	A	0.239	3.33E-08	0.031	77,165	2.92E-10	0.030	112,353	6.38E-17	0.031	
rs6795735	3	64,680,405	<i>ADAMTS9</i>	C	0.406	2.47E-07	0.025	77,162	6.75E-08	0.026	84,480	9.79E-14	0.025	
rs4823006	22	27,781,671	<i>ZNRF3-KREMEN1</i>	A	0.569	4.47E-08	0.027	77,086	2.41E-05	0.019	93,911	1.10E-11	0.023	
rs6784615	3	52,481,466	<i>NISCH-STAB1</i>	T	0.941	3.18E-07	0.052	76,859	1.56E-04	0.036	109,028	3.84E-10	0.043	
rs6861681	5	173,295,064	<i>CPEB4</i>	A	0.340	1.40E-06	0.026	77,164	2.13E-04	0.019	85,722	1.91E-09	0.022	
<b>Further SNPs evaluated in follow-up but not achieving genome-wide significant in the combined analysis</b>														
rs2076529	6	32,471,933	<i>BTNL2</i>	C	0.570	2.22E-08	0.041	34,532	0.012	0.011	92,778	3.71E-07	0.020	
rs7081678	10	32,030,629	<i>ZEB1</i>	A	0.085	5.76E-07	0.045	76,270	0.094	0.013	100,527	5.57E-06	0.027	

P-values and beta-coefficients (per change of WHR increasing allele) for the association with WHR on the inverse normal transformed ranked scale in the meta-analyses of discovery studies (up to 77,167), follow-up studies (up to 113,636 subjects), and both combined (up to 190,781 subjects). Fourteen of the 16 SNPs put forward to follow-up show genome-wide significant results ( $P < 5 \times 10^{-8}$ ) in the combined analysis. P-values in the discovery were genomic control corrected per study and in the meta-analysis. Details on between-study heterogeneity in Supplementary Table 1c.

<sup>a</sup>Effect Allele: WHR increasing allele on the forward strand;

<sup>b</sup>Effect allele frequency

**Table 2**  
Evidence of sex-differences in the WHR association at seven of the 14 associated loci

SNP	Nearby Genes	Men						Women						Sex difference	
		Discovery		Follow-up		Combined		Discovery		Follow-up		Combined		Beta	P
		P	Beta	P	Beta	P	Beta	P	Beta	P	Beta	P	Beta		
rs9491696	<i>RSP03</i>	1.68E-04	0.026	6.97E-09	0.036	1.05E-11	0.031	1.62E-12	0.047	8.84E-22	0.053	1.93E-32	0.050	1.94E-03	
rs6905288	<i>VEGFA</i>	0.066	0.013	2.09E-04	0.025	7.38E-05	0.020	7.72E-13	0.052	3.14E-15	0.051	2.27E-26	0.052	5.20E-06	
rs984222	<i>TBX15-WARS2</i>	3.32E-09	0.041	2.43E-05	0.029	9.41E-13	0.035	1.21E-07	0.036	1.33E-08	0.033	1.02E-14	0.034	0.951	
rs1055144	<i>NFE2L3</i>	6.00E-04	0.029	5.67E-08	0.040	2.52E-10	0.035	2.34E-06	0.040	7.13E-12	0.046	1.41E-16	0.044	0.270	
rs10195252	<i>GRB14</i>	0.201	0.009	0.114	0.011	0.043	0.010	6.33E-15	0.053	4.95E-21	0.054	3.84E-34	0.054	1.41E-11	
rs4846567	<i>LYPLALI</i>	0.191	0.010	0.982	0.000	0.358	0.005	4.84E-18	0.064	8.12E-17	0.055	4.95E-33	0.059	1.18E-13	
rs1011731	<i>DNM3-PIGC</i>	4.88E-07	0.034	1.95E-03	0.022	7.81E-09	0.028	2.13E-05	0.028	7.03E-07	0.030	6.90E-11	0.029	0.855	
rs718314	<i>ITPR2-SSPN</i>	0.177	0.010	2.02E-03	0.022	1.41E-03	0.017	8.29E-10	0.047	4.21E-09	0.038	2.41E-17	0.042	4.67E-04	
rs1294421	<i>LY86</i>	4.18E-03	0.020	7.00E-06	0.030	1.63E-07	0.025	3.44E-08	0.038	7.32E-06	0.026	2.40E-12	0.031	0.357	
rs1443512	<i>HOXC13</i>	0.184	0.011	9.74E-04	0.024	9.45E-04	0.018	1.43E-09	0.048	3.09E-08	0.035	6.38E-16	0.040	2.23E-03	
rs6795735	<i>ADAMTS9</i>	0.011	0.017	0.614	0.004	0.027	0.011	7.85E-07	0.033	2.95E-11	0.042	1.92E-16	0.038	8.50E-05	
rs4823006	<i>ZNRF3-KREMEN1</i>	6.87E-03	0.019	0.094	0.012	1.94E-03	0.015	6.86E-08	0.037	3.81E-05	0.024	3.24E-11	0.030	0.032	
rs6784615	<i>NISCH-STAB1</i>	1.51E-03	0.045	0.033	0.032	1.68E-04	0.039	6.23E-05	0.057	1.72E-03	0.039	6.01E-07	0.047	0.574	
rs6861681	<i>CPEB4</i>	1.88E-03	0.023	0.045	0.015	3.03E-04	0.019	2.14E-04	0.027	1.58E-03	0.021	1.55E-06	0.024	0.555	

*P*-values and beta-coefficients (per change of WHR increasing allele in the sex-combined analysis as in Table 1) for the WHR association are given for the discovery (up to 34,601 men, 42,735 women), the follow-up (up to 47,882 men, 65,780 women) and the combined meta-analysis (up to 81,301 men, 107,429 women). Also given are the *P*-values for testing for difference between sex-specific beta-coefficients in the combined meta-analysis; SNPs with *P* for sex difference < 3.6E-03 (=0.05/14) were considered to show significant sex difference.

**Table 3**  
WHR signals show enrichment of association with other traits related to metabolic disorders

Trait	Sample size <sup>a</sup>	SNPs in concordant direction <sup>b</sup>	SNPs in concordant direction with $P < .05$ <sup>c</sup>	$P$	
Triglycerides	43,826	14	6,10E-5	7	1.79E-8
HDL-C	45,561	13	9,16E-4	4	3.20E-4
LDL-C	43,889	10	0.090	1	0.298
Fasting glucose	63,849	10	0.090	1	0.298
Fasting insulin	54,883	13	9,16E-4	5	1.62E-5
HOMA-IR	53,625	13	9,16E-4	6	6.17E-7
2-hr glucose	27,011	7	0.605	0	1.000
Type 2 diabetes	10,128 <sup>f</sup>	11	0.029	3	4.62E-3

The 14 WHR SNPs were tested for association with other traits by meta-analysis of GWA data from previous reports 19–21, 39 together with our non-overlapping *de novo* genotyped follow-up studies. HDL-C, high density lipoprotein cholesterol; LDL-C, low density lipoprotein cholesterol; HOMA-IR, index of insulin resistance; 2-hr glucose, glucose levels 2 h after an oral glucose challenge.

<sup>a</sup>Maximum number of subjects available for any of the 14 SNPs

<sup>b</sup>Number of the 14 SNPs for which the WHR increasing allele is associated with the trait in the concordant direction (i.e. increased levels except for HDL-C) and corresponding binomial  $P$ -value to test whether this number is greater than by chance not accounting for the correlation between the traits.

<sup>c</sup>Number of SNPs in concordant direction that show  $P < 0.05$  for the association with the trait and corresponding binomial  $P$ -value as in <sup>b</sup>.

<sup>f</sup>4,549 cases, 5579 controls

Table 4

Expression quantitative trait locus analysis for 11 out of the 14 WHR signals.

WHR SNP	Tissue	Gene	Effect <sup>a</sup>	WHR SNP association with transcript (P)		Transcript peak SNP <sup>b</sup>	r <sup>2c</sup>	Peak SNP association with transcript (P)	
				Unadj.	Adj. for peak SNP			Unadj.	Adj. for WHR SNP
rs9491696	SAT-D	RSPO3	+	1.10E-07	0.03	rs1936795	0.26	2.20E-13	7.40E-08
<b>rs984222</b>	<b>Omental</b>	<b>TBX15</b>	+	<b>7.90E-10</b>	<b>1.00</b>	<b>rs984222</b>	<b>1.00</b>	<b>7.90E-10</b>	<b>1.00</b>
	Omental	WARS2	+	5.11E-36	0.03	rs10802075	0.27	1.31E-163	1.33E-88
	Subcutaneous fat	WARS2	+	1.67E-25	0.01	rs10802075	0.22	3.88E-110	1.01E-63
	Lymph.	WARS2	-	4.30E-18	5.47E-05	rs2645305	0.27	5.57E-40	6.88E-26
	Liver	WARS2	+	2.57E-17	0.07	rs1057990	0.26	6.69E-59	1.97E-32
	SAT-D	WARS2	+	1.10E-18	0.51	rs1057990	0.26	5.80E-130	5.80E-100
	Blood	WARS2	+	6.10E-17	0.11	rs1057990	0.26	6.30E-75	1.10E-54
<b>rs1055144</b>	<b>SAT-D</b>	<b>AA553656<sup>d</sup></b>	-	<b>1.20E-11</b>	<b>0.96</b>	<b>rs7798002</b>	<b>0.95</b>	<b>7.20E-12</b>	<b>0.32</b>
	SAT-M	<b>AA553656<sup>d</sup></b>	-	<b>2.46E-07</b>	<b>0.65</b>	<b>rs1451385</b>	<b>0.77</b>	<b>5.93E-08</b>	<b>0.38</b>
<b>rs10195252</b>	<b>SAT-D</b>	<b>GRB14</b>	+	<b>4.40E-11</b>	<b>1.00</b>	<b>rs10195252</b>	<b>1.00</b>	<b>4.40E-11</b>	<b>1.00</b>
	SAT-M	<b>GRB14</b>	+	<b>5.51E-06</b>	<b>1.00</b>	<b>rs10184004</b>	<b>1.00</b>	<b>5.51E-06</b>	<b>1.00</b>
	Omental	<b>GRB14</b>	+	<b>1.02E-13</b>	<b>1.00</b>	<b>rs10195252</b>	<b>1.00</b>	<b>1.02E-13</b>	<b>1.00</b>
	SAT-M	<i>SLC38A11</i>	-	3.93E-06	0.66	rs10184126	0.18	7.76E-44	8.57E-34
	SAT-D	<i>SLC38A11</i>	-	3.70E-09	0.35	rs10184126	0.18	2.40E-94	7.40E-82
<b>rs1011731</b>	Blood	<i>C1orf105</i>	+	3.80E-16	0.20	rs2157451	0.28	1.30E-33	8.20E-18
	<b>Lymph.</b>	<i>PIGC</i>	-	<b>5.87E-10</b>	<b>1.00</b>	<b>rs991790</b>	<b>1.00</b>	<b>5.65E-10</b>	<b>1.00</b>
rs718314	Lymph.	<i>ITPR2</i>	+	1.79E-09	0.98	rs7976877	0.45	2.21E-18	1.91E-06
	Blood	<i>ITPR2</i>	-	2.40E-09	0.20	rs2570	0.41	2.40E-37	1.80E-28
rs1294421	SAT-M	<i>BC039678</i>	-	2.43E-07	0.38	rs1294404	0.64	1.89E-16	3.42E-04
	Omental	<i>BC039678</i>	-	1.09E-06	0.33	rs912056	0.71	8.28E-17	4.26E-05
rs6795735	SAT-D	<i>ADAMTS9</i>	-	1.50E-06	0.04	rs7372321	0.11	1.10E-09	2.30E-05
	Omental	<i>AK022320</i>	-	7.99E-15	0.64	rs4521216	0.02	5.15E-42	1.49E-19
	SAT-D	<i>AK022320</i>	-	2.24E-10	0.98	rs4521216	0.02	9.62E-37	7.58E-19
<b>rs4823006</b>	<b>SAT-D</b>	<b>ZNRF3</b>	-	<b>2.40E-08</b>	<b>0.63</b>	<b>rs3178915</b>	<b>0.81</b>	<b>6.70E-11</b>	<b>8.90E-04</b>

WHR SNP	Tissue	Gene	Effect <sup>a</sup>	WHR SNP association with transcript (P)		Transcript peak SNP <sup>b</sup>	r <sup>2c</sup>	Peak SNP association with transcript (P)	
				Unadj.	Adj. for peak SNP			Unadj.	Adj. for WHR SNP
	SAT-M	<i>ZNRF3</i>	-	1.08E-18	0.93	rs6005975	0.79	1.59E-19	0.50
	Omental	<i>ZNRF3</i>	-	9.13E-18	0.98	rs6005975	0.79	6.07E-21	0.27
<b>rs6784615</b>	Blood	<i>STAB1</i>	+	2.80E-09	0.32	rs9846089	0.83	9.40E-10	0.08
rs6861681	Lymph.	<i>CPEB4</i>	+	3.79E-22	0.89	rs7705502	0.87	4.95E-29	2.00E-03
	Blood	<i>HMP19</i>	+	1.60E-16	0.97	rs10516107	0.83	1.10E-21	4.30E-06

Association between the 14 WHR SNPs and expression of transcripts located within 1 Mb of the WHR SNP in two sets of abdominal subcutaneous adipose tissue (SAT-D from deCODE and SAT-M from Massachusetts General Hospital), omental fat, liver, lymphocytes, and blood (see Supplementary Note); results are given if the unadjusted WHR SNP association showed  $P < 1.00E-05$ . Findings are highlighted in bold font, where the WHR SNP was the transcript peak SNP or where the WHR signal and the *cis*-eQTL signal were considered coincident (i.e. the transcript peak SNP was highly correlated with the WHR SNP,  $r^2 > 0.7$ , and the transcript peak association disappeared by adjusting on the WHR SNP,  $P > 0.05$ ); see also Online Methods

Unadj. = unadjusted. Adj. = adjusted.

<sup>a</sup>Effect direction for the WHR increasing allele.

<sup>b</sup>SNP with the strongest association with the transcript in the region (transcript peak SNP).

<sup>c</sup>Correlation (Hapmap CEU, B36) between the WHR SNP and transcript peak SNP.

<sup>d</sup>The transcript labeled AA553656 was detected as Contig27623\_RC and corresponds to chr7: 25,854,143-25,854,203 (B36)