

NERIDRONATE TREATMENT IN CHILDREN WITH OSTEOGENESIS IMPERFECTA (OI): THE EFFECTS ON BONE MINERAL DENSITY (BMD) AND VERTEBRAL DEFORMITIES

M. Celli¹, D. Diacinti², A. Zambrano¹, P. D'Eufemia¹, M. Tetti¹, R. Del Fiacco², V. Baldini², E. D'Erasmo²

¹ Department of Pediatrics, University of Rome "La Sapienza", Rome, Italy

² Department of Clinical Sciences, University of Rome "La Sapienza", Rome, Italy

To evaluate the effects of neridronate treatment on bone mass and vertebral deformities in children with OI.

In 21 children (8 M, 13 F; age 1-12 aa), with OI (types I, III e IV) in therapy with neridronate (2 mg/kg i.v. every 3 months), were evaluated lumbar bone mass (BMD e BMAD) and vertebral deformities by morphometric X-ray absorptiometry (MXA) using QDR4500A (Hologic). By MXA were calculated the wedging index [(1-ah/ph)x100%] and concavity index [(1-mh/ph)x100%].

A significant increase of BMD was observed after 12 months, from 0.08 ± 0.01 to 0.09 ± 0.01 (+17.1%; $p < 0.01$); the indexes of vertebral deformity were reduced (wedging: 4.64% n.s.; biconcavity: -7.5% n.s.) and inversely correlated to the BMD ($r = -0.372$) and to the BMAD ($r = -0.376$).

The semiquantitative (SQ) assessment of the radiographs and the MXA revealed the same number and grade of vertebral fractures in 13/20 children.

Our data confirm the effectiveness of the therapy with neridronate increasing the bone density; the relationship between the morphometric indexes and the BMD suggest the possibility to use the MXA (low-dose method), in the follow-up for the children with IO for the identification of vertebral fractures.