

# Conformal symmetry of the Lange-Neubert evolution equation

V.M. Braun<sup>1</sup> and A.N. Manashov<sup>1,2</sup>

<sup>1</sup> *Institut für Theoretische Physik, Universität Regensburg, D-93040 Regensburg, Germany*

<sup>2</sup> *Department of Theoretical Physics, St.-Petersburg University, 199034, St.-Petersburg, Russia*

## Abstract

The Lange-Neubert evolution equation describes the scale dependence of the wave function of a meson built of an infinitely heavy quark and light antiquark at light-like separations, which is the hydrogen atom problem of QCD. It has numerous applications to the studies of  $B$ -meson decays. We show that the kernel of this equation can be written in a remarkably compact form, as a logarithm of the generator of special conformal transformation in the light-ray direction. This representation allows one to study solutions of this equation in a very simple and mathematically consistent manner. Generalizing this result, we show that all heavy-light evolution kernels that appear in the renormalization of higher-twist  $B$ -meson distribution amplitudes can be written in the same form.

**1.** Studies of heavy meson weak decays have been instrumental to uncover the flavor sector of the Standard model and can be a gate to new physics at TeV scales, if it exists. Considerable effort has been invested to understand the QCD dynamics of heavy meson decays in the heavy quark limit. The  $B$ -meson distribution amplitude (DA), first introduced in [1], provides the key nonperturbative input in the QCD factorization approach [2] for weak decays involving light hadrons in the final state.

Following an established convention we define the  $B$ -meson DA as the renormalized matrix element of the bilocal operator built of an effective heavy quark field  $h_v(0)$  and a light antiquark  $\bar{q}(zn)$  at a light-like separation:

$$\langle 0 | \bar{q}(zn) \not{n} [zn, 0] \Gamma h_v(0) | \bar{B}(v) \rangle = -\frac{i}{2} F(\mu) \text{Tr} [\gamma_5 \not{n} \Gamma P_+] \Phi_+(z, \mu) \quad (1)$$

with

$$[zn, 0] \equiv \text{Pexp} \left[ ig \int_0^1 d\alpha n_\mu A^\mu(\alpha zn) \right]. \quad (2)$$

Here  $v_\mu$  is the heavy quark velocity,  $n_\mu$  is the light-like vector,  $n^2 = 0$ , such that  $n \cdot v = 1$ ,  $P_+ = \frac{1}{2}(1 + \not{n})$  is the projector on upper components of the heavy quark spinor,  $\Gamma$  stands for an arbitrary Dirac structure,  $|\bar{B}(v)\rangle$  is the  $\bar{B}$ -meson state in the heavy quark effective theory

(HQET) and  $F(\mu)$  is the decay constant in HQET, which is used for normalization. The effective heavy quark can be related to the Wilson line through the following equation [3]:

$$\langle 0|h_v(0)|h,v\rangle = [0, -v\infty] = \text{Pexp} \left[ ig \int_{-\infty}^0 d\alpha v_\mu A^\mu(\alpha v) \right], \quad (3)$$

so that the operator in Eq. (1) can be viewed as a single light antiquark attached to the Wilson line with a cusp containing one lightlike and one timelike segment.

The invariant function  $\Phi_+(z, \mu)$  where  $z$  is a real number defines what is usually called the leading twist B-meson DA in position space. Its Fourier transform is

$$\begin{aligned} \phi_+(k, \mu) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} dz e^{ikz} \Phi_+(z - i0, \mu), \\ \Phi_+(z, \mu) &= \int_0^{\infty} dk e^{-ikz} \phi_+(k, \mu), \end{aligned} \quad (4)$$

where in the first equation the integration contour goes below the singularities of  $\Phi_+(z, \mu)$  that are located in the upper-half plane. The parameter  $\mu$  is the renormalization (factorization) scale. We tacitly imply using dimensional regularization with modified minimum subtraction.

The scale dependence of the DA is driven by the renormalization of the corresponding nonlocal operator

$$O_+(z) = \bar{q}(zn) \not{n} [zn, 0] \Gamma h_v(0).$$

The corresponding one-loop  $Z$ -factor was computed by Lange and Neubert (LN) [4], giving rise to an evolution equation which is convenient to write, for our purposes, as a renormalization group equation for the operator  $O_+(z)$  [5, 6]:

$$\left( \mu \frac{\partial}{\partial \mu} + \beta(g) \frac{\partial}{\partial g} + \frac{\alpha_s C_F}{\pi} \mathcal{H} \right) O_+(z, \mu) = 0, \quad (5)$$

where

$$[\mathcal{H}f](z) = \int_0^1 \frac{d\alpha}{\alpha} \left( f(z) - \bar{\alpha} f(\bar{\alpha} z) \right) + \ln(i\mu z) f(z) - \frac{5}{4} f(z), \quad \bar{\alpha} \equiv 1 - \alpha. \quad (6)$$

This equation thus governs the scale dependence of the  $B$ -meson DA in position space,  $\Phi_+(z, \mu)$ . It is fully equivalent to the original LN equation for the DA in momentum space,  $\phi_+(k, \mu)$ , as it is easy to show by Fourier transformation.

**2.** We will demonstrate that the LN kernel (6) can be written in terms of the generators of collinear conformal transformations

$$S_+ = z^2 \partial_z + 2jz, \quad S_0 = z \partial_z + j, \quad S_- = -\partial_z, \quad (7)$$

where  $j = 1$  is the conformal spin of the light quark. They satisfy the standard  $SL(2)$  commutation relations

$$[S_+, S_-] = 2S_0, \quad [S_0, S_\pm] = \pm S_\pm. \quad (8)$$

The starting observation is that the integral operator  $\mathcal{H}$  (LN kernel) can be written in a somewhat different form by studying its action on the test functions  $f(z) = z^p$ ,  $z\partial_z f(z) = pf(z)$ . Here and below  $\partial_z = \partial/\partial z$ . In this way one obtains

$$[\mathcal{H}f](z) = \left[ \psi(z\partial_z + 2) - \psi(1) + \ln(i\mu z) - \frac{5}{4} \right] f(z). \quad (9)$$

Next, we use the identity for a fractional derivative  $(i\partial_z)^a$  defined as the multiplication operator  $k^a$  in momentum representation [7]:

$$(i\partial_z)^a = (iz)^{-a} \frac{\Gamma(a - z\partial_z)}{\Gamma(-z\partial_z)}. \quad (10)$$

It holds for the functions  $f(z)$  that are holomorphic in the lower complex half-plane  $\Im m z < 0$ ,  $z \in \mathbb{C}_-$ , and vanish at infinity. Fourier transform for such functions goes over positive momenta  $f(z) = \int_0^\infty dk e^{-ikz} \tilde{f}(k)$ ,  $(i\partial_z)^a f(z) = \int_0^\infty dk e^{-ikz} k^a \tilde{f}(k)$ , corresponding in our case to positive values of the light-quark energy  $\omega = k/2$  in the B-meson rest frame, cf. Eq. (4). Expanding this identity around  $a = 0$  one gets

$$\ln(i\partial_z) = \psi(-z\partial_z) - \ln(iz) \quad (11)$$

and making an inversion  $z \rightarrow -1/z$

$$\ln(iz^2\partial_z) = \psi(z\partial_z) + \ln(iz). \quad (12)$$

Finally, since for any function  $f(z\partial_z)z = zf(z\partial_z + 1)$ , we can write this identity as

$$z^{-2} \ln(iz^2\partial_z) z^2 = \ln[i(z^2\partial_z + 2z)] = \ln(iS^+) = \psi(z\partial_z + 2) + \ln(iz). \quad (13)$$

Comparing with Eq. (6) we see that

$$\mathcal{H} = \ln(i\mu S^+) - \psi(1) - \frac{5}{4} \quad (14)$$

which is our main result. Note that the scale  $\mu$  under the logarithm is necessary simply because  $S_+$  has dimension  $[\text{mass}]^{-1}$ .

Alternatively, the same expression can be derived starting from the commutation relations for the LN kernel obtained in Ref. [6]:

$$[S_+, \mathcal{H}] = 0, \quad [S_0, \mathcal{H}] = 1. \quad (15)$$

Since the problem has one degree of freedom — the light-cone coordinate of the light quark — it follows from  $[S_+, \mathcal{H}] = 0$  that the operator  $\mathcal{H}$  must be a *function* of  $S_+$ ,  $\mathcal{H} = h(S_+)$ . This function can be found using the second commutation relation. Let  $S = S_0 + 1$ . Then  $S_+ = zS$  and the relation  $[S_0, h(S_+)] = 1$  can be written equivalently as  $[S, h(zS)] = 1$ . Taking into account that  $[S, zS] = zS$  one obtains an equation on the function  $h(s)$

$$s h'(s) = 1 \implies h(s) = \ln s + \text{constant}, \quad (16)$$

reproducing the result in Eq. (14) up to a (scheme-dependent) constant.

**3.** The main advantage of Eq. (14) is that diagonalization of the kernel  $\mathcal{H}$  can be traded for a much simpler task of diagonalization of the first-order differential operator  $S_+$  (7). Eigenfunctions of  $S_+$  take a simple form<sup>1</sup>

$$Q_s(z) = -\frac{1}{z^2} e^{is/z}, \quad iS_+ Q_s(z) = s Q_s(z), \quad (17)$$

so that

$$\mathcal{H} Q_s(z) = \left[ \ln(\mu s) - \psi(1) - \frac{5}{4} \right] Q_s(z). \quad (18)$$

A further advantage is that one can use  $SL(2)$  representation theory methods to work with these solutions, see e.g. Ref. [11] for a short discussion of this technique. In particular one can make use of the standard  $SL(2)$  invariant scalar product [12] (for spin  $j = 1$ )

$$\langle \Phi | \Psi \rangle = \frac{1}{\pi} \int_{\mathbb{C}_-} d^2 z \overline{\Phi(z)} \Psi(z), \quad (19)$$

where the (two-dimensional) integration goes over the lower half-plane  $\mathbb{C}_-$ ,  $\Im m z < 0$ . The generator  $iS^+$  is self-adjoint w.r.t. this scalar product. The eigenfunctions (17) are orthogonal to each other and form a complete set

$$\langle Q_{s'} | Q_s \rangle = \frac{1}{s} \delta(s - s'), \quad \int_0^\infty ds s Q_s(z) \overline{Q_s(z')} = \frac{e^{-i\pi}}{(z - \bar{z}')^2}. \quad (20)$$

The function on the r.h.s. of the completeness relation is called reproducing kernel [13]. It acts as a unit operator so that for any function holomorphic in the lower half plane

$$\Psi(z) = \frac{1}{\pi} \int_{\mathbb{C}_-} d^2 z' \frac{e^{-i\pi}}{(z - \bar{z}')^2} \Psi(z'). \quad (21)$$

Hence the  $B$ -meson DA (1) can be expanded as

$$\Phi_+(z, \mu) = \int_0^\infty ds s \eta(s, \mu) Q_s(z) = -\frac{1}{z^2} \int_0^\infty ds s e^{is/z} \eta(s, \mu), \quad \eta(s, \mu) = \langle Q_s | \Phi \rangle. \quad (22)$$

The integration goes over all possible eigenvalues of the step-up generator  $S_+$  that corresponds to special conformal transformations along the light-ray  $n^\mu$ . This representation is very similar to the one suggested in Ref. [8].

The scale-dependence of the coefficients  $\eta(s, \mu)$  is governed by the renormalization-group equation

$$\left( \mu \frac{\partial}{\partial \mu} + \beta(\alpha_s) \frac{\partial}{\partial \alpha_s} + \Gamma_{cusp}(\alpha_s) \ln(\mu s/s_0) \right) F(\mu) \eta(s, \mu) = 0, \quad (23)$$

---

<sup>1</sup>The sign is chosen such that  $Q_s(z)$  are real and positive for  $z = -i\tau$ ,  $\tau > 0$ .

where  $s_0 = e^{5/4-\gamma_E}$  and  $\Gamma_{cusp}(\alpha_s) = \frac{\alpha_s}{\pi} C_F + \dots$  is the cusp anomalous dimension [9, 10].

The solution of this equation takes the form

$$\begin{aligned} F(\mu) \eta(s, \mu) &= F(\mu_0) \eta(\xi, \mu_0) \times \exp \left\{ - \int_{\mu_0}^{\mu} \frac{d\tau}{\tau} \Gamma_{cusp}(\alpha_s(\tau)) \ln(\tau s/s_0) \right\} \\ &= F(\mu_0) \eta(\xi, \mu_0) \left( \frac{\mu_0 s}{s_0} \right)^{r(\mu)} B(\mu), \end{aligned} \quad (24)$$

where

$$\begin{aligned} r(\mu) &= - \int_{\alpha(\mu_0)}^{\alpha(\mu)} \frac{d\alpha}{\beta(\alpha)} \Gamma_{cusp}(\alpha_s) = 2C_F/\beta_0 \ln \left( \frac{\alpha(\mu)}{\alpha(\mu_0)} \right) + \dots, \\ B(\mu) &= \exp \left\{ - \int_{\alpha(\mu_0)}^{\alpha(\mu)} \frac{d\alpha}{\beta(\alpha)} \Gamma_{cusp}(\alpha) \int_{\alpha(\mu_0)}^{\alpha} \frac{d\alpha'}{\beta(\alpha')} \right\}. \end{aligned} \quad (25)$$

In practical applications the momentum (energy) representation for the  $B$ -meson DA  $\phi_+(k, \mu)$  as defined in (4) is more convenient. This can be derived easily by observing that exponential functions  $e^{-ipz}$ ,  $p > 0$  are mutually orthogonal and form a complete set w.r.t. the same scalar product

$$\langle e^{-ipz} | e^{-ip'z} \rangle = \frac{1}{p} \delta(p - p'). \quad (26)$$

Hence

$$\Phi_+(z, \mu) = \int_0^\infty dp p e^{-ipz} \langle e^{-ipz} | \Phi_+(z, \mu) \rangle = \int_0^\infty dp p e^{-ipz} \int_0^\infty ds s \eta(s, \mu) \langle e^{-ipz} | Q_s(z) \rangle \quad (27)$$

and therefore

$$\phi_+(k, \mu) = \frac{1}{2\pi} \int_{-\infty}^\infty dz e^{ikz} \Phi_+(z - i0, \mu) = k \int_0^\infty ds s \eta(s, \mu) \langle e^{-ikz} | Q_s(z) \rangle. \quad (28)$$

Using

$$\langle e^{-ikz} | Q_s(z) \rangle = \frac{1}{\sqrt{ks}} J_1(2\sqrt{ks}) \quad (29)$$

we finally obtain

$$\phi_+(k, \mu) = \int_0^\infty ds \sqrt{ks} J_1(2\sqrt{ks}) \eta(s, \mu), \quad (30)$$

where  $J_1(x)$  is the Bessel function. The representation in Eq. (30) is equivalent to the one suggested by Bell, Feldmann, Wang and Yip in Ref. [8], who noticed that the evolution equation is significantly simplified in this manner. In their notation, cf. second line in Eq. (2.17),  $s \eta(s, \mu) \equiv \rho_+(1/s, \mu)$ .

The orthogonality relation (26) combined with the projection (29) leads to a familiar relation for the Bessel functions

$$\int_0^\infty ds J_1(2\sqrt{ps}) J_1(2\sqrt{p's}) = \delta(p - p'), \quad (31)$$

which can be used to invert Eq. (30) and express  $\eta(s, \mu)$  in terms of  $\phi_+(k, \mu)$ .

Note that the representation in (14) is valid for the evolution kernel in momentum space as well, but the generator  $S_+$  has to be taken in the adjoint representation

$$\mathcal{S}_+ = i[k\partial_k^2 + 2j\partial_k], \quad j = 1. \quad (32)$$

The Bessel functions appearing in (29), (30) are eigenfunctions of  $\mathcal{S}_+$ , indeed:

$$s\langle e^{-ikz}|Q_s(z)\rangle = \langle e^{-ikz}|iS_+Q_s(z)\rangle = \langle iS_+e^{-ikz}|Q_s(z)\rangle = i\mathcal{S}_+\langle e^{-ikz}|Q_s(z)\rangle. \quad (33)$$

Of particular interest for the QCD description of  $B$ -decays is the value of the first negative moment

$$\lambda_B^{-1}(\mu) = \int_0^\infty \frac{dk}{k} \phi_+(k, \mu) = \int_0^\infty d\tau \Phi_+(-i\tau, \mu) = \int_0^\infty ds \eta(s, \mu). \quad (34)$$

As demonstrated in [8], QCD factorization expressions for  $B$  decay amplitudes can conveniently be written in terms of  $\eta(s, \mu)$  as well, so that we do not dwell on this topic here.

**5.** The same representation can be derived for arbitrary two-particle heavy-light one-loop kernels that contribute to the evolution equations for higher-twist  $B$ -meson DAs [6]. The difference to the leading twist is that the two-particle evolution equations are not closed: The two-particle,  $2 \rightarrow 2$ , kernels appear as parts of larger mixing matrices involving  $2 \rightarrow 3$  parton transitions, however,  $3 \rightarrow 2$  transitions do not occur at the one-loop level.

Explicit expressions for all  $2 \rightarrow 2$  heavy-light kernels have been derived in Ref. [6], see Sec. 3.2. They can be written in terms of an integral operator

$$[\mathcal{H}_j f](z) = \int_0^1 \frac{d\alpha}{\alpha} \left[ f(z) - \bar{\alpha}^{2j-1} f(\bar{\alpha}z) \right] + \ln(i\mu z) f(z) - [\sigma_h + \sigma_\ell] f(z), \quad (35)$$

where  $j$  is the conformal spin of the light parton  $\ell$  (quark or gluon) and the constants  $\sigma_h = 1/2$ ,  $\sigma_{\text{quark}} = 3/4$ ,  $\sigma_{\text{gluon}} = \beta_0/4N_c$  ( $\beta_0 = 11/3N_c - 2/3n_f$ ) are related to the anomalous dimensions of the fields. Conformal spin of a parton is defined as  $j = (d + s)/2$  where  $d$  is canonical dimension and  $s$  is spin projection on the light cone, see [14]. For a quark  $j = 1$  for the “plus” projection that contributes to the leading-twist  $B$ -meson DA (1), in which case (35) reproduces (6), and  $j = 1/2$  for the “minus” projection that is relevant for the DA  $\Phi_-(z, \mu)$ , cf. [2]. In turn, for a gluon  $j = 3/2$  for the leading-twist projection and  $j = 1$  for the higher-twist.

Following the above derivation for  $j = 1$  we obtain the following representation for the kernel in the general case:

$$\mathcal{H}_j = \ln(i\mu S_+^{(j)}) - \psi(1) - \sigma_h - \sigma_\ell, \quad (36)$$

where the generator of special conformal transformations  $S_+^{(j)}$  for spin  $j$  is defined in Eq. (7). The eigenfunctions of  $S_+^{(j)}$  have the form

$$Q_s^{(j)}(z) = \frac{e^{-i\pi j}}{z^{2j}} e^{is/z}, \quad iS_+^{(j)} Q_s^{(j)}(z) = s Q_s^{(j)}(z). \quad (37)$$

They are orthogonal and form a complete set with respect to the  $SL(2)$  scalar product [13]

$$\langle \Phi | \Psi \rangle_j = \frac{2j-1}{\pi} \int_{\mathbb{C}_-} \mathcal{D}_j z \overline{\Phi(z)} \Psi(z), \quad (38)$$

where  $\mathcal{D}_j z = d^2 z [i(z - \bar{z})]^{2j-2}$ . One obtains

$$\langle Q_s^{(j)} | Q_{s'}^{(j)} \rangle_j = \frac{\Gamma(2j)}{s^{2j-1}} \delta(s - s'), \quad \frac{1}{\Gamma(2j)} \int_0^\infty ds s^{2j-1} Q_s^{(j)}(z) \overline{Q_{s'}^{(j)}(z')} = \frac{e^{-i\pi j}}{(z - \bar{z}')^{2j}}. \quad (39)$$

The expression on the r.h.s. of the second integral defines the reproducing kernel for arbitrary spin  $j$  [13], i.e. for arbitrary function (holomorphic in the lower plane)

$$\Psi(z) = \frac{2j-1}{\pi} \int_{\mathbb{C}_-} \mathcal{D}_j z \frac{e^{-i\pi j}}{(z - \bar{z}')^{2j}} \Psi(z). \quad (40)$$

The functions  $Q_s^j(z)$  diagonalize the renormalization group kernel

$$\mathcal{H}_j Q_s^j(z) = [\ln(\mu s) - \psi(1) - \sigma_h - \sigma_\ell] Q_s^j(z) \quad (41)$$

so that it is natural to write matrix elements of generic heavy-light operators as an expansion

$$\Phi_j(z, \mu) = \int_0^\infty ds s^{2j-1} \eta_j(s, \mu) Q_s^{(j)}(z), \quad (42)$$

where  $\Phi_j(z, \mu)$  is analogue of  $\Phi_+(z, \mu)$  (1).

The expansion coefficients  $\phi_j(k, \mu)$  appearing in the Fourier transform

$$\Phi_j(z, \mu) = \int_0^\infty dk e^{-ikz} \phi_j(k, \mu) \quad (43)$$

can be found making use of the following relations:

$$\begin{aligned} \langle e^{-ikz} | e^{-ik'z} \rangle_j &= \Gamma(2j) k^{1-2j} \delta(k - k'), \\ \langle e^{-ikz} | Q_s^{(j)} \rangle_j &= \Gamma(2j) (ks)^{1/2-j} J_{2j-1}(2\sqrt{ks}). \end{aligned} \quad (44)$$

In this way one obtains

$$\phi_j(p, \mu) = \int_0^\infty ds \eta_j(s, \mu) (sp)^{j-1/2} J_{2j-1}(2\sqrt{ps}). \quad (45)$$

In particular for  $j = 1/2$  corresponding to the  $B$ -meson DA  $\phi_-(k, \mu)$  [2] the conformal expansion goes over Bessel functions  $J_0(2\sqrt{ks})$  as compared to  $J_1(2\sqrt{ks})$  for the leading twist, cf. [8].

**6.** To summarize, we have constructed a conformal expansion of the distribution amplitudes of heavy-light mesons in terms of eigenfunctions of the generator of special conformal transformations. This construction is similar in spirit to the well-known expansion of DAs of light mesons in Gegenbauer polynomials which are eigenfunctions of two-particle  $SL(2)$  Casimir operators, see e.g. [5]. Similar to the latter case, this expansion can serve as a basis for the construction of approximations of phenomenological relevance.

As we have shown, this expansion is a consequence of the commutation relations (15) and it would be very interesting to find out whether these relations hold true to all orders in perturbation theory for a conformal theory like  $N = 4$  SYM. The consequences of our results for the DAs of baryons made of one heavy and two light quarks should be studied as well.

## 7. Acknowledgements

This work was supported by the DFG, grant BR2021/5-2.

## References

- [1] A. Szczepaniak, E. M. Henley and S. J. Brodsky, Phys. Lett. B **243** (1990) 287.
- [2] M. Beneke, G. Buchalla, M. Neubert and C. T. Sachrajda, Phys. Rev. Lett. **83** (1999) 1914; Nucl. Phys. B **591** (2000) 313.
- [3] G. P. Korchemsky and A. V. Radyushkin, Phys. Lett. B **279** (1992) 359.
- [4] B. O. Lange and M. Neubert, Phys. Rev. Lett. **91** (2003) 102001.
- [5] V. M. Braun, D. Yu. Ivanov and G. P. Korchemsky, Phys. Rev. D **69** (2004) 034014.
- [6] M. Knödlseeder and N. Offen, JHEP **1110** (2011) 069.
- [7] A. V. Belitsky, S. E. Derkachov and A. N. Manashov, arXiv:1401.7307 [hep-th].
- [8] G. Bell, T. Feldmann, Y.-M. Wang and M. W. Y. Yip, JHEP **1311** (2013) 191.
- [9] A. M. Polyakov, Nucl. Phys. B **164** (1980) 171.
- [10] G. P. Korchemsky and A. V. Radyushkin, Nucl. Phys. B **283** (1987) 342.
- [11] S. E. Derkachov, G. P. Korchemsky and A. N. Manashov, JHEP **0307** (2003) 047.
- [12] I.M. Gelfand, M.I. Graev, N.Ya. Vilenkin, Generalized functions. Vol. 5: Integral geometry and representation theory,” Academic Press (New York, 1966).
- [13] B.C. Hall. Holomorphic Methods in Mathematical Physics - Contemporary Mathematics, Volume 260, pp. 1-59. quant-ph/9912054
- [14] V. M. Braun, G. P. Korchemsky and D. Mueller, Prog. Part. Nucl. Phys. **51** (2003) 311.