

ON SQUARES OF IRREDUCIBLE CHARACTERS

EMMANUEL ZHMUD

Kharkiv State University, Ukraine

ABSTRACT. We study the finite groups G with a faithful irreducible character whose square is a linear combination of algebraically conjugate irreducible characters of G . In conclusion, we offer another proof of one theorem of Isaacs-Zisser.

There are a few papers treating the finite groups possessing an irreducible character whose powers are linear combinations of appropriate irreducible characters, for example, [BC] and [IZ]. Our note is inspired by these two papers, especially, the second one.

In what follows, G is a finite group. We use standard notation of finite group theory (see [BZ]). Recall that if $\chi \in \text{Irr}(G)$, then the generalized character $\chi^{(2)}$ (see [BZ, Chapter 4]) is defined as follows:

$$\chi^{(2)}(g) = \chi(g^2) \quad (g \in G).$$

Next, $\text{Char}(G)$ denotes the set of characters of a group G and, if θ is a generalized character of G , then

$$\text{Irr}(\theta) = \{\chi \in \text{Irr}(G) \mid \langle \theta, \chi \rangle \neq 0\}.$$

The quasikernel $Z(\chi)$ of $\chi \in \text{Char}(G)$ is defined as follows:

$$Z(\chi) = \{g \in G \mid |\chi(g)| = \chi(1)\}.$$

It is known that $Z(\chi)$ is a normal subgroup of G containing $\ker(\chi)$ and $Z(G/\ker(\chi)) \leq Z(\chi)/\ker(\chi)$ with equality if, in addition, $\chi \in \text{Irr}(G)$. In what follows, we use freely results stated in this paragraph.

2010 *Mathematics Subject Classification.* 20C15.

Key words and phrases. Generalized character, algebraic conjugate, irreducible characters, quasikernel, exterior power.

Professor Zhmud (1918–2007) died in 29 December 2007. This note was prepared by Y. Berkovich, based on a letter dated March 14, 2000.

Let $\chi \in \text{Irr}(G)$. It is known that

$$(1) \quad \text{Irr}(\chi^{(2)}) \subseteq \text{Irr}(\chi^2).$$

Indeed, $\chi^{(2)} = \chi^2 - 2\wedge^2\chi$ (see formula (22) in [BZ, §4.6]) so it suffices to show that $\text{Irr}(\wedge^2\chi) \subseteq \text{Irr}(\chi^2)$. Next, $\chi^2 = \wedge^2\chi + \theta$, where θ is the exterior square of χ (see [BZ, Lemma 4.16, formula (17)]) so $\text{Irr}(\wedge^2\chi) \subseteq \text{Irr}(\chi^2)$, as desired. Therefore, if $\text{Irr}(\chi^2) = \{\psi_1, \dots, \psi_n\}$ and

$$(2) \quad \chi^2 = \sum_{j=1}^n a_j \psi_j, \text{ where all } a_j \text{ are positive integers,}$$

then, by (1), we have

$$(3) \quad \chi^{(2)} = \sum_{j=1}^n b_j \psi_j, \text{ where all } b_i \text{ are integers.}$$

Set

$$(4) \quad a = a_1 + \dots + a_n, \quad b = b_1 + \dots + b_n.$$

Let ϵ be a $|G|$ -th primitive root of 1, $\mathcal{G} = \text{Gal}(Q(\epsilon)/Q)$, where Q is the field of rational numbers. The group \mathcal{G} acts in the natural way on the set $\text{Irr}(G)$ as follows: if $\chi \in \text{Irr}(G)$ and $\sigma \in \mathcal{G}$, then $(\sigma\chi)(g) = \sigma(\chi(g))$ for all $g \in G$ (see [BZ, Chapter 3]). Characters $\psi, \psi' \in \text{Irr}(G)$ are said to be *algebraically conjugate* if $\psi' = \sigma\psi$ for some $\sigma \in \mathcal{G}$. In that case, as it is easy to check, $\psi(1) = \psi'(1)$, $\ker(\psi) = \ker(\psi')$ and $Z(\psi) = Z(\psi')$.¹ In what follows we will retain the notation introduced above and in this paragraph.

DEFINITION 1. *A group G with an irreducible character χ possesses a property \mathcal{A} , if it satisfies the following conditions:*

- (A1) $|G|$ is even.
- (A2) χ is faithful.
- (A3) $n \geq 2$ and ψ_j ($j = 1, \dots, n$) are algebraically conjugate with $\psi = \psi_1$ (see decomposition (2)), i.e., $\psi_j = \sigma_j\psi$ for some $\sigma_j \in \mathcal{G}$ ($j = 1, \dots, n$).
- (A4) χ and ψ_1, \dots, ψ_n satisfy (2) and (3).

Our main result is the following

THEOREM 2. *If the group G satisfies condition \mathcal{A} , then the following assertions hold (here, as in part (A3) of the definition, $\psi = \psi_1$):*

- (a) G is nonabelian and $\chi(1) > 1$.
- (b) G has only one involution u .
- (c) $\ker(\psi) = \langle u \rangle$.
- (d) $Z(\psi)$ is abelian.
- (e) Sylow 2-subgroups of G are cyclic.

¹Let us prove the second equality. Take $g \in \ker(\psi)$. We have $\psi'(g) = \sigma(\psi(g)) = \sigma(\psi(1)) = \psi(1)$ so $\ker(\psi) \leq \ker(\psi')$. Since $\sigma^{-1} \in \mathcal{G}$, the reverse inclusion holds as well.

- (f) $G = P \cdot N$, a semidirect product, where $P \in \text{Syl}_2(G)$ and $\{1\} < N \triangleleft G$.
- (g) $\chi(1) = \psi(1)$ and $b = 1$.
- (h) $\chi_N \in \text{Irr}(N)$.
- (i) All $(\psi_j)_N$ are irreducible and nonreal for $j = 1, \dots, n$.
- (j) If $w \in G$ with $o(w) = 8$, then $\psi(w) = m\sqrt{-1}$, where m is an integer dividing $\chi(1)$.

PROOF. (a) It follows from (2) and (4) that $\chi(1)^2 = a\psi(1)$ so $\chi(1) > 1$ since $a \geq n > 1$. Therefore, G is nonabelian.

(b) Let u be an involution in G . Since $\psi(u)$ is a rational integer, it follows that $\psi_j(u) = \psi(u)$ since the ψ_j 's are algebraically conjugate. Therefore, $\chi^{(2)}(u) = b\psi(u)$ (see (3) and (4)). Since $\chi^{(2)}(u) = \chi(u^2) = \chi(1)$, we get

$$(5) \quad \chi(1) = b\psi(u).$$

On the other hand, $\chi(1) = \chi^{(2)}(1) = b\psi(1)$ so, taking into account that $b \neq 0$, we get $\psi(u) = \psi(1)$, by (5), i.e., $u \in \ker(\psi) = \ker(\psi_j)$ for $j = 1, \dots, n$. Therefore, it follows from (2) that $\ker(\chi^2) = \ker(\psi)$ so $u \in \ker(\chi^2)$, i.e., $\chi(u)^2 = \chi(1)^2$. In that case, $\chi(u) = \pm\chi(1)$, and we obtain

$$(6) \quad \chi(u) = -\chi(1)$$

since our character χ is faithful. It follows that $u \in Z(\chi) = Z(G)$ since χ is faithful, i.e., $Z(G)$ contains all involutions of G . Since $Z(G)$ is cyclic (χ is faithful), we conclude that u is the unique involution in G , and (b) is proven.

(c) As we have proved in (b) (see the sentence after formula (5)), $u \in \ker(\psi)$. It suffices to show that u is the unique nonidentity element of $\ker(\psi)$. Take $x \in \ker(\psi)^\#$. Then $\psi_j(x) = \psi(1)$ for all j so, by (2), $x \in \ker(\chi^2)$ so that, again by (2), we have

$$\chi(1)^2 = \chi(x)^2 = a\psi(1), \text{ and hence } \chi(x) = -\chi(1)$$

since χ is faithful. In particular, $x \in Z(\chi)$ so that $\chi(x^2) = \chi(1)$ and $x^2 \in \ker(\chi) = \{1\}$, i.e., x is an involution. It follows from this and (b), that $x = u$. Thus $\ker(\psi) = \{1, u\} = \langle u \rangle$, as required.

(d) It follows from (c), that $Z(\psi)$ is abelian since $Z(\psi)/\ker(\psi)$ is cyclic (in view of irreducibility of ψ) and $|\ker(\psi)| = 2$.

(e) Let $P \in \text{Syl}_2(G)$. By (b), P is either cyclic or generalized quaternion. Assume, by way of contradiction, that P is generalized quaternion. Take in P an element v of order 4. Then, by (b),

$$(7) \quad v^2 = u.$$

Let $j \in \{1, \dots, n\}$. Then, by (A2), $\psi_j = \sigma_j\psi$ so that $\psi_j(v) = (\sigma_j\psi)(v) = \sigma_j(\psi(v))$. We have $\sigma_j\epsilon = \epsilon^{\nu_j}$ for some rational integer ν_j such that $\text{GCD}(\nu_j, |G|) = 1$ (recall that ϵ is the primitive $|G|$ -th root of 1 chosen above); then ν_j is odd in view of (A1). Setting $\nu_j = 2\lambda_j + 1$ and taking into account

that $\psi(v)$ is a sum of powers of ϵ , we get

$$\psi_j(v) = \sigma_j(\psi(v)) = \psi(v^{\nu_j}) = \psi(v^{2\lambda_j+1}) = \psi(u^{\lambda_j}v) = \psi(v)$$

since $u \in \ker(\psi)$, by (c). Thus

$$(8) \quad \psi_j(v) = \psi(v) \quad (j = 1, \dots, n).$$

Then, by (6), (8), (3) and (4), we obtain

$$-\chi(1) = \chi(u) = \chi(v^2) = \chi^{(2)}(v) = \sum_{j=1}^n b_j \psi_j(v) = b\psi(v).$$

On the other hand, $\chi(1) = \chi^{(2)}(1) = b\psi(1)$ so $\psi(v) = -\psi(1)$. It follows from this and (8), that

$$(9) \quad \psi_j(v) = -\psi(1) \quad (j = 1, \dots, n).$$

It follows from (9) that, if T is a representation of G affording the character ψ_j , then $T(v) = -I_{\psi(1)}$, where $I_{\psi(1)}$ is a $\psi(1) \times \psi(1)$ identity matrix. Therefore, by (2), we get

$$\chi(v)^2 = \sum_{j=1}^n a_j \psi_j(v) = -a\psi(1) = -\chi(1)^2,$$

and we conclude that

$$(10) \quad \chi(v) = ci\chi(1),$$

where $c = \pm 1$ and $i = \sqrt{-1}$. It follows from $|\chi(v)| = \chi(1)$ that $v \in Z(\chi) = Z(G)$, a contradiction, since the center of P , which is a generalized quaternion group, has order 2. Thus, $P \in \text{Syl}_2(G)$ is cyclic.

(f) By (e), G is 2-nilpotent so $G = P \cdot N$, and $N > \{1\}$ since G is nonabelian.

(g) Since $G/N \cong P$ is cyclic, then χ is not ramified over N (Burnside; see [BZ, Exercise 7 in Chapter 7]) so we get the following Clifford decomposition:

$$(11) \quad \chi_N = \sum_{k=1}^l \phi_k.$$

It follows from (11) that

$$(12) \quad (\chi^{(2)})_N = \sum_{k=1}^l \phi_k^{(2)}.$$

Since $|N|$ is odd, it follows that $\phi_k^{(2)}$ are distinct irreducible characters of N for all k , and $(\chi^{(2)})_N$ is a character of N .² By (9), we have

$$(\chi^{(2)})_N = \sum_{j=1}^n b_j(\psi_j)_N.$$

Let $\phi_1 = \phi$. Since

$$1 = \langle (\chi^{(2)})_N, \phi^{(2)} \rangle = \sum_{j=1}^n b_j \langle (\psi_j)_N, \phi^{(2)} \rangle,$$

we get $\langle (\psi_s)_N, \phi^{(2)} \rangle \neq 0$ for some $s \in \{1, \dots, n\}$. This means that $\phi^{(2)} \in \text{Irr}((\psi_s)_N)$. By Clifford's theorem, $\text{Irr}((\psi_s)_N)$ is a G -orbit of $\phi^{(2)}$, i.e.,

$$\text{Irr}((\psi_s)_N) = \{\phi_1^{(2)}, \dots, \phi_l^{(2)}\}, \text{ where } \phi_1 = \phi.$$

We conclude that $(\psi_s)_N = \sum_{k=1}^l \phi_k^{(2)}$ so, by (12), $(\chi^{(2)})_N = (\psi_s)_N$. It follows, in particular, that $\chi(1) = \psi_s(1) = \psi(1)$. Since $\chi(1) = \chi^{(2)}(1) = b\psi(1)$, we get $b = 1$, completing the proof.

(h) It follows from (2) and (3) that

$$\chi^2 - \chi^{(2)} = \sum_{j=1}^n (a_j - b_j)\psi_j.$$

Since

$$\frac{1}{2}(\chi^2 - \chi^{(2)}) = \bigwedge^2 \chi \in \text{Char}(G)$$

(see formula (22) in [BZ, §4.6]), we get $a_j \equiv b_j \pmod{2}$ for $j = 1, \dots, n$. Summing up over all j , one obtains $a \equiv b \pmod{2}$ so a is odd since $b = 1$, by (g). As we have noticed in the proof of (a), $\chi(1)^2 = a\psi(1)$. Therefore, since $\chi(1) = \psi(1)$, by (g), we have $\chi(1) = a$, and hence $\chi(1)$ is odd. It follows from (11) that $l = |\text{Irr}(\chi_N)| = |G : I_G(\phi)|$, where $I_G(\phi)$ is the inertia group of ϕ in G , and $\chi(1) = l\phi(1)$ so l is odd. Since $I_G(\phi) \geq N$ and $|G : N| = |P|$ is a power of 2, we get $l = 1$, i.e., $\chi_N \in \text{Irr}(N)$, proving (h).

(i) Since $l = 1$, there exists $s \in \{1, \dots, n\}$ such that $(\psi_s)_N = \phi^{(2)} \in \text{Irr}(N)$. Since all ψ_j are algebraically conjugate, we get $(\psi_j)_N \in \text{Irr}(N)$. If $(\psi_j)_N$ is real for some j , it is the principal character 1_G of G since $|N|$ is odd. It follows that $\psi(1) = \psi_j(1) = 1$. Then, by (g), $\chi(1) = 1$, a contradiction. Thus, all ψ_j are not real.

(j) Let $w \in G$ be of order 8. Setting $v = w^2$, we get $o(v) = 4$. By (b), $v^2 = u$. It follows from (10) that $ci\chi(1) = \chi(v) = \chi(w^2) = \chi^{(2)}(w)$ so, in view

²Indeed, assume that $\phi_i^{(2)} = \phi_j^{(2)}$. Then, for $x \in N$ we have $\phi_i(x^2) = \phi_i^{(2)}(x) = \phi_j^{(2)}(x) = \phi_j(x^2)$, and so $\phi_i = \phi_j$ since $\{x^2 \mid x \in N\} = N$. This proves the first assertion. Now the second assertion is obvious.

of (9), we get

$$(13) \quad \sum_{j=1}^n b_j \psi_j(w) = \chi^{(2)}(w) = ci\chi(1)$$

(here $i = \sqrt{-1}$). Recall, that $\psi_j(w) = \psi(w^{\nu_j})$, where ν_j is odd integer (see the proof of (e)). It follows from $o(w) = 8$ that $\nu_j \in \{1, 3, 5, 7\}$. Since $u \in \ker(\psi)$ and $\psi(v) = -\psi(1)$ (see (9)), we have (consider a representation of G affording the character ψ ; see two line after formula (9))

$$\begin{aligned} \psi(w^3) &= \psi(vw) = -\psi(w), \\ \psi(w^5) &= \psi(uw) = \psi(w), \\ \psi(w^7) &= \psi(uvw) = -\psi(w). \end{aligned}$$

Thus, $\psi_j(w) \in \{\psi(w), -\psi(w)\}$. Setting $\psi_j(w) = c_j\psi(1)$, where $c_j = \pm 1$, one can rewrite (13) in the form

$$(14) \quad ci\chi(1) = d\psi(w),$$

where $d = \sum_{j=1}^n b_j c_j$ is a rational integer. Note that

$$\overline{\psi(w)} = \psi(w^{-1}) = \psi(w^7) = -\psi(w).$$

Therefore, $\psi(w) = im$, where m is a real number. Now, (14) yields $\chi(1) = cdm$. Therefore, $m = c \cdot \frac{\chi(1)}{d}$ is rational. Since $m = -i\psi(w)$ is an algebraic integer, it follows that m is a rational integer so m divides $\chi(1)$. This completes the proof of our theorem. \square

Now we are ready to offer another proof of the following

THEOREM 3 (Isaacs-Zisser [IZ]). *Let $G > \{1\}$ be a group and suppose that there is a faithful $\chi \in \text{Irr}(G)$ such that*

$$(15) \quad \chi^2 = a\psi + b\bar{\psi},$$

where a, b are positive integers and $\psi \in \text{Irr}(G)$. Then G is a direct product of a cyclic 2-group of order not exceeding 4 and a group of odd order.

PROOF. Since $\psi_1 = \psi$ and $\psi_2 = \bar{\psi}$ are algebraically conjugate, one can apply Theorem 2 to our group G . By that theorem, P is cyclic. It remains to prove that $|P| \leq 4$ and P is normal in G . We claim that $|\text{Irr}(\chi^{(2)})| = 1$ (recall that $\chi^{(2)} : x \mapsto \chi(x^2)$). Otherwise, $\chi^{(2)} = b_1\psi_1 + b_2\psi_2$, where b_1, b_2 are nonzero rational integers. It follows that

$$(\chi^{(2)})_N = b_1(\psi_1)_N + b_2(\psi_2)_N$$

(recall that $(\chi^{(2)})_N, (\psi_1)_N, (\psi_2)_N$ are irreducible, by Theorem 2(h,i)). We have $(\chi^{(2)})_N = (\psi_s)_N$ for some $s \in \{1, 2\}$ (see the proof of Theorem 2(g)). Using this and the equality $b_1 + b_2 = 1$, we get $(\psi_1)_N = (\psi_2)_N$. It follows that $(\psi_1)_N = \overline{(\psi_1)_N}$, i.e., character $(\psi_1)_N$ is real, contrary to Theorem 2(i). Thus,

one of numbers b_1, b_2 equals 0 so our claim is proven, i.e., $\chi^{(2)} \in \{\psi_1, \psi_2\}$. Assume, for definiteness, that $\chi^{(2)} = \psi_1$. Thus,

$$\chi^2 = a_1\psi_1 + a_2\psi_2 \text{ and } \chi^{(2)} = \psi_1.$$

As above, set $\psi = \psi_1$. Assume that $|P| > 4$. Take in P an element w of order 8. Set $v = w^2$. Then, by (10),

$$\psi(w) = \chi^{(2)}(w) = \chi(v) = ci\chi(1), \text{ where } c = \pm 1.$$

Since χ and ψ have the same degree, we get $\psi(w) = ci\psi(1)$ so $|\psi(w)| = \psi(1)$ whence $w \in Z(\psi)$. Let $H \in \text{Syl}_2(Z(\psi))$; then H is normal in G so $HN = H \times N$ and w centralizes N . Since w centralizes $P \in \text{Syl}_2(G)$, by Theorem 2(e), we see that $w \in Z(G)$ since $G = \overline{P \cdot N}$, and so $|\chi(w)| = \chi(1)$. Then equalities

$$\psi_2(w) = \overline{\psi_1(w)} = -\psi(w) \text{ and } |\psi(w)| = \chi(1)$$

imply

$$\begin{aligned} \chi(1)^2 &= |\chi(w)|^2 = |a_1\psi(w) + a_2\psi_2(w)| = |(a_1 - a_2)\psi(w)| \\ &= |a_1 - a_2||\psi(w)| = |a_1 - a_2|\chi(1). \end{aligned}$$

Hence $\chi(1) = |a_1 - a_2|$. On the other hand, $\chi(1) = a = a_1 + a_2$. Thus $a_1 + a_2 = |a_1 - a_2|$, a contradiction since $a_1, a_2 > 0$. Thus $|P| \leq 4$.

Let us prove that the subgroup P is normal in G . In view of Theorem 2(b), one may assume that $|P| = 4$ and $P = \langle v \rangle$. Then, by (9), $\psi(v) = -\psi(1)$ so $v \in Z(\psi)$ and hence $P \leq Z(\psi)$. Since P is characteristic in $Z(\psi)$ (indeed, P is a Sylow 2-subgroup of the abelian group $Z(\psi)$; see Theorem 2(d)), it follows that P is normal in G , and the proof is complete. \square

ACKNOWLEDGEMENTS.

We are indebted to the referee for useful remarks.

REFERENCES

- [BZ] Y. Berkovich and E. Zhmud, *Characters of Finite Groups, Parts 1, 2*, AMS, Providence, 1998.
- [BC] H. Blau and D. Chillag, *On powers of characters and powers of conjugacy classes of a finite group*, Proc. Amer. Math. Soc. **98** (1986), 7–10.
- [IZ] I. M. Isaacs and I. Zisser, *Squares of characters with few irreducible constituents in finite groups*, Arch. Math. (Basel) **63** (1994), 197–207.

Received: 2.12.2009.

Revised: 29.12.2009.