

# MEASUREMENT ISSUES AND POOR ADJUSTMENTS FOR PHYSICAL ACTIVITY AND SLEEP UNDERMINE SEDENTARY BEHAVIOUR RESEARCH — THE FOCUS SHOULD SHIFT TO THE BALANCE BETWEEN SLEEP, SEDENTARY BEHAVIOUR, STANDING AND ACTIVITY

Željko Pedišić

*Faculty of Kinesiology, University of Zagreb, Zagreb, Croatia  
Prevention Research Collaboration, Sydney School of Public Health,  
The University of Sydney, Sydney, Australia*

Commentary

UDC: 613.72:796.012.45:159.963.2:796.012.83:531

## Abstract:

This paper critically appraised 54 recent studies linking sedentary behaviour (SB) and health, specifically regarding their assessment of SB and adjustments for physical activity (PA) and sleep. Almost 90% of the studies collected SB data using self-reports and 59% of these measures had not been previously validated. The majority of studies did not adjust for total PA or for both light-intensity PA (LIPA) and moderate-to-vigorous-intensity PA (MVPA), and a large number of studies did not adjust for all domains of PA. It may, therefore, be that these adjustments were not thorough enough to allow for sound conclusions about the independent associations between SB and health outcomes. Sleep time is also likely to act as a confounding variable in associations between SB and health outcomes. Despite that, only three reviewed studies adjusted their analyses for sleep duration. Evidence presented here casts doubt upon the conclusions about independent associations between SB and health outcomes. Given the facts that: 1) the proportions of time allotted to sleep, SB, standing, LIPA and MVPA are perfectly collinear, 2) all these behaviours may be associated with health, and 3) mutual adjustments between all these variables are needed if their independent contributions to health are to be determined, it seems that investigating the balance between times spent in these behaviours is the next logical step in epidemiological research. This paper, therefore, proposes the *Activity Balance Model (AB model)*; a new theoretical framework for investigating associations of sleep duration, SB, standing, LIPA and MVPA with health outcomes.


**Key words:** *sleep duration, sedentary behaviour, standing, LIPA, physical activity, Activity Balance Model*

## Introduction

Sedentary behaviour encompasses all waking activities in a sitting or reclining posture that require energy expenditure of  $\leq 1.5$  METs (e.g. television viewing, passive commuting, occupational sitting) (Sedentary Behaviour Research Network, 2012). The entire 24-hour period of each day is spent in sleep, sedentary behaviour, standing, light-intensity physical activity (LIPA) and moderate to vigorous-intensity physical activity (MVPA). For example, US adults spend on average approximately 36%, 32.9%, 29.5%, and 1.6% of their time at sleep, in sedentary behaviour, LIPA and MVPA, respectively (Bureau of Labor Statistics, 2013; Schuna, Johnson, & Tudor-Locke, 2013). Given the closed nature of a 24-hour block, proportions of time spent in these behaviours are perfectly collinear, that is, every increase in the total time spent in one behaviour necessarily causes a decrease in the total time

spent in one or more remaining behaviours (Figure 1). Accordingly, total sedentary time may show significant associations with MVPA (Booth, et al., 2012; Espinel, Chau, van der Ploeg, & Merom, 2014; Healy, et al., 2008; Spittaels, et al., 2012), LIPA (Espinel, et al., 2014; Healy, et al., 2008; Spittaels, et al., 2012), standing (Katzmarzyk, 2014), and sleep time (Booth, et al., 2012; Buman, et al., 2014; Matthews, et al., 2012).

Studies have suggested that prolonged time spent in sedentary behaviours may be an independent risk factor for: [i] type 2 diabetes; [ii] cardiovascular disease; [iii] colorectal, endometrial, ovarian, and prostate cancer; and [iv] cardiovascular, cancer and all-cause mortality (Bauman, Chau, Ding, & Bennie, 2013; Chau, Grunseit, Chey, et al., 2013; Cong, et al., 2014; Ford & Caspersen, 2012; Grøntved & Hu, 2011; Lynch, 2010; Wilmot, et al., 2012). Sedentary behaviours may also be adversely



Legend: LIPA = light-intensity physical activity; MVPA = moderate to vigorous-intensity physical activity  
 \*No data is available on the time spent standing still. Since the data about sedentary, LIPA and MVPA times are taken from accelerometer-based studies, it is possible that the time spent standing is distributed between sedentary and LIPA times.

Figure 1. Perfectly collinear proportions of time allotted to sleep, sedentary behaviour, LIPA and MVPA in a 24-hour period\*

linked to cardiometabolic biomarkers, insulin resistance, hypertension and mental disorders, but limited and mixed results preclude any definite conclusions (Thorp, Owen, Neuhaus, & Dunstan, 2011).

In order to critically appraise these findings, specifically regarding the assessment of sedentary behaviours and adjustments for physical activity and sleep, 54 recent articles were reviewed (Table 1). These articles come from nine systematic and narrative reviews about health outcomes of sedentary behaviour (Bauman, et al., 2013; Chau, Grunseit, Chey, et al., 2013; Cong, et al., 2014; Ford & Caspersen, 2012; Grøntved & Hu, 2011; Lynch, 2010; Thorp, et al., 2011; van Uffelen, Wong, et al., 2010; Wilmot, et al., 2012). To take into account recent changes in the sedentary behaviour definition and paradigm (Pate, O’Neill, & Lobelo, 2008) and the fact that results of most large-scale studies in this area were reported in the last five years (Bauman, et al., 2013), only the articles published since 2008 were included in this review.

### Assessment of sedentary behaviour

Most of the reviewed studies assessed television viewing/other screen time (48%), total sedentary behaviour (39%), and occupational sedentarism (22%). Half of the studies on occupational sedentarism were based on the classification according to job type. Non-occupational, transport-related, and leisure-time sedentary behaviour were assessed in 4%, 4%, and 7% of the studies, respectively. These percentages clearly show a lack of studies on health outcomes of most domain-specific (excluding occupational sitting) and type-specific (excluding TV viewing) sedentary behaviours. Specific domains and types of sedentary behaviour should, therefore, be of particular interest in future studies.

Almost none of the studies on TV viewing/screen time specifically asked participants if they were sitting/reclining during these activities. It may be that some participants were standing or exercising instead. Hence, according to the current definition of sedentary behaviour (Sedentary Behaviour Research Network, 2012) these studies can only be tentatively considered sedentary behaviour studies.

Furthermore, 89% of the studies collected sedentary behaviour data exclusively using self-reports, whilst 7% used accelerometers and 4% used heart rate monitors. Self-reports of sedentary behaviour used in 59% of the studies had not been previously validated. In some of these studies, authors reported that measurement properties had been established for a similar sedentary behaviour item/questionnaire. However, from the psychometric point of view, this is not considered acceptable, as even slight changes to the questionnaire may significantly change its measurement properties. Therefore, future studies should either use standardized and validated sedentary behaviour questionnaires or conduct their own psychometric assessment.

Two studies included in this review used heart rate monitors and estimated the time spent in sedentary behaviour as all minutes below the flex HR (Ekelund, Brage, Besson, Sharp, & Wareham, 2008; Helmerhorst, Wijndaele, Brage, Wareham, & Ekelund, 2009). Flex HR is calculated as the mean of

Table 1. Methodology of studies on sedentary behaviour and health with regards to measurement issues and adjustments for physical activity and sleep time

| Study                         | Measure of SB |                          |           | Adjustments for PA and sleep time  |            |
|-------------------------------|---------------|--------------------------|-----------|--|------------|
|                               | Instrument    | Domain/type              | Validated | PA variables*  | Sleep time |
| Ekelund, et al., 2008         | HRM           | Total SB (below flex HR) | yes       | Total PA assessed by heart rate monitor  | no         |
| Howard, et al., 2008          | Self-report   | Total SB, TV viewing†    | no‡       | Total PA   | no         |
| Inoue, Iso, et al., 2008      | Self-report   | Total SB                 | no        | Heavy physical work/strenuous exercise, walking/standing, leisure-time sports/exercise | no         |
| Inoue, Yamamoto, et al., 2008 | Self-report   | Total SB                 | no        | Heavy physical work/strenuous exercise, walking/standing, leisure-time sports/exercise | no         |

| Study   | Measure of SB |  |                 | Adjustments for PA and sleep time   |                       |
|---|---------------|--|-----------------|---|-----------------------|
|   | Instrument    | Domain/type  | Validated       | PA variables*   | Sleep time            |
| Landhuis, Poulton, Welch, & Hancox, 2008                            | Self-report   | TV viewing <sup>†</sup>                                | no              | no  | no                    |
| Meyer, et al., 2008   | Self-report   | TV viewing <sup>†</sup>                                | yes             | Sport index from Baecke questionnaire   | no                    |
| Moradi, et al., 2008  | Self-report   | Occupation type  | yes             | no  | no                    |
| Parsons, Manor, & Power, 2008                                       | Self-report   | TV viewing <sup>†</sup>                                | no              | Leisure-time PA   | no                    |
| Patel, et al., 2008   | Self-report   | Non-occupational SB                                    | no <sup>‡</sup> | Recreational MVPA, non-recreational leisure-time PA, exercise                     | no                    |
| Sanchez-Villegas, et al., 2008                                      | Self-report   | TV viewing+computer use <sup>†</sup>                   | yes             | no  | no                    |
| Ekelund, Brage, Griffin, & Wareham, 2009                            | Accelerometer | Total SB (<100 cpm)                                    | yes             | Accelerometer-based LIPA and MVPA   | no                    |
|   | Self-report   | TV viewing <sup>†</sup>                                | no              |   |                       |
| Gierach, et al., 2009   | Self-report   | Total SB, TV viewing <sup>†</sup>                      | no              | no  | no                    |
| Helmerhorst, et al., 2009   | HRM           | Total SB (below flex HR)                               | yes             | MVPA assessed by heart rate monitor   | no                    |
| Katzmarzyk, Church, Craig, & Bouchard, 2009                         | Self-report   | Work+school+home SB                                    | no              | Leisure-time PA   | no                    |
| Krishnan, Rosenberg, & Palmer, 2009                                 | Self-report   | TV viewing <sup>†</sup>                                | no              | Vigorous-intensity exercise, walking for exercise or transport                    | no                    |
| Mathew, et al., 2009  | Self-report   | TV viewing <sup>†</sup>                                | no              | Total PA, household activities  | no                    |
| Mekary, et al., 2009  | Self-report   | Domestic SB+TV viewing                                 | no              | Recreational PA   | no                    |
| Orsini, et al., 2009  | Self-report   | Occupation type  | yes             | Walking+bicycling   | no                    |
| Tonstad, et al., 2009   | Self-report   | TV viewing <sup>†</sup>                                | no              | MVPA  | ≤6, 7, and ≥8 h/night |
| Tudor-Locke, Burton, & Brown, 2009                                  | Self-report   | Occupational SB  | no              | Leisure-time MVPA   | no                    |
| Wijndaele, et al., 2009   | Self-report   | TV viewing <sup>†</sup>                                | no <sup>‡</sup> | Non-occupational MVPA+occupational walking  | no                    |
| Dunstan, et al., 2010   | Self-report   | TV viewing <sup>†</sup>                                | yes             | Exercise  | no                    |
| Ford, et al., 2010  | Self-report   | TV viewing <sup>†</sup>                                | no              | Physical strain at work, non-occupational PA (walking+gardening+bicycling+sports) | no                    |
| Friedenreich, et al., 2010  | Self-report   | Occupational SB  | no              | no  | no                    |
| Gollenberg, et al., 2010  | Self-report   | Occupational SB, TV viewing <sup>†</sup>               | yes             | no  | no                    |
| Lynch, et al., 2010   | Accelerometer | Total SB (<100 cpm)                                    | yes             | Accelerometer-based MVPA  | no                    |
| Patel, et al., 2010   | Self-report   | Non-occupational SB                                    | no <sup>‡</sup> | Recreational MVPA+domestic PA   | no                    |
| van Uffelen, Watson, Dobson, & Brown, 2010                          | Self-report   | Total SB   | no <sup>‡</sup> | Leisure-time MVPA   | no                    |
| Warren, et al., 2010  | Self-report   | TV viewing <sup>†</sup> , riding in a car, sum of both | no              | Total PA self-rated in comparison with peers                                      | no                    |
| Wijndaele, et al., 2010   | Self-report   | TV viewing <sup>†</sup>                                | yes             | Non-occupational MVPA+occupational walking  | no                    |
| Banks, Lim, Seubsman, Bain, & Sleigh, 2011                          | Self-report   | Total SB, TV viewing+computer games <sup>†</sup>       | no              | Exercise-related PA, domestic PA  | no                    |
| Boyle, Fritschi, Heyworth, & Bull, 2011                             | Self-report   | Occupation type  | yes             | Recreational PA, heavy/very heavy occupation type                                 | no                    |
| Hawkes, Lynch, Owen, & Aitken, 2011                                 | Self-report   | TV viewing <sup>†</sup>                                | no <sup>‡</sup> | Non-occupational MVPA+occupational walking  | no                    |
| Lynch, et al., 2011   | Accelerometer | Total SB (<100 cpm)                                    | yes             | Accelerometer-based LIPA and MVPA   | no                    |
| Parent, et al., 2011  | Self-report   | Occupation type  | yes             | Sports and/or outdoor activities, occupational PA                                 | no                    |
| Stamatakis, Hamer, & Dunstan, 2011                                  | Self-report   | Screen time <sup>†</sup>                               | no              | Occupational PA, non-occupational MVPA  | no                    |
| Wijndaele, Brage, Besson, Khaw, Sharp, Luben, Wareham, et al., 2011 | Self-report   | TV viewing <sup>†</sup>                                | yes             | Total PA  | no                    |

| Study  | Measure of SB |  |           | Adjustments for PA and sleep time  |                               |
|--|---------------|--|-----------|--|-------------------------------|
|  | Instrument    | Domain/type  | Validated | PA variables*  | Sleep time                    |
| Wijndaele, Brage, Besson, Khaw, Sharp, Luben, Bhaniani, et al., 2011 | Self-report   | TV viewing†  | yes       | Total PA   | h/day                         |
| Chau, van der Ploeg, Merom, Chey, & Bauman, 2012                     | Self-report   | Occupation type, leisure-time SB   | no‡       | Leisure-time MVPA, occupational activity type (mostly standing, mostly walking, mostly heavy labor)                          | no                            |
| de Heer, Wilkinson, Strong, Bondy, & Koehly, 2012                    | Self-report   | Work+school+home SB  | no        | LIPA, MPA, VPA   | ≤5, 6, 7, 8, 9, and ≥10 h/day |
| Frydenlund, Jorgensen, Toft, Pisinger, & Aadahl, 2012                | Self-report   | Leisure-time SB  | yes       | Total PA   | no                            |
| Gomez-Cabello, et al., 2012  | Self-report   | Total SB   | no        | Walking  | no                            |
| Koster, et al., 2012   | Accelerometer | Total SB (<100 cpm)  | yes       | Accelerometer-based MVPA   | no                            |
| Matthews, et al., 2012   | Self-report   | Total SB, TV viewing†  | no‡       | Leisure time MVPA  | <7, ≥7 h/day§                 |
| Pavey, Geeske Peeters, & Brown, 2012                                 | Self-report   | Total SB   | no‡       | Leisure-time MVPA  | no                            |
| Pinto Pereira, Ki, & Power, 2012                                     | Self-report   | Occupational SB, TV viewing†   | yes       | Leisure-time MVPA  | no                            |
| van der Ploeg, Chey, Korda, Banks, & Bauman, 2012                    | Self-report   | Total SB   | no‡       | Non-occupational MVPA+occupational walking   | no                            |
| Yates, et al., 2012  | Self-report   | Total SB   | yes       | MVPA   | no                            |
| Campbell, Patel, Newton, Jacobs, & Gapstur, 2013                     | Self-report   | Leisure-time SB  | no        | Recreational PA  | no                            |
| Chau, Grunseit, Midthjell, et al., 2013                              | Self-report   | Total SB, TV viewing†, Occupation type   | no‡       | Total PA   | no                            |
| George, Rosenkranz, & Kolt, 2013                                     | Self-report   | Total SB   | no‡       | Non-occupational MVPA+occupational walking   | no                            |
| Pulsford, Stamatakis, Britton, Brunner, & Hillsdon, 2013             | Self-report   | Total SB, occupational SB, TV viewing, non-TV leisure-time SB, leisure-time SB | yes       | MVPA, walking, domestic LIPA§  | no                            |
| Shuval, et al., 2013   | Self-report   | Total SB, transport-related SB, computer use†                                  | yes       | Transport-related MVPA   | no                            |
| Simons, et al., 2013   | Self-report   | Occupational SB  | no        | Occupational PA, walking/cycling for transport, recreational walking/cycling, gardening/doing odd jobs, sports participation | no                            |

Legend: SB = sedentary behaviour; PA = physical activity; LIPA = light-intensity PA; MPA = moderate-intensity PA; VPA = vigorous-intensity PA; MVPA = moderate-to-vigorous intensity PA; HRM = heart rate monitor; Flex HR = mean of the highest resting heart rate and the lowest heart rate during exercise; cpm = accelerometer counts per minute

\*Self-reported if not stated otherwise

†The question about TV viewing/screen time did not specifically ask about the body posture during the activity (or the authors did not report so)

‡Authors reported that measurement properties have been established for a similar measure

§The variable was not included in the final model

the individual's highest resting heart rate and the lowest heart rate during exercise, where the resting heart rate is measured in lying, sitting and standing posture (Spurr, et al., 1988). Although this method may give valid estimates of time spent inactive, it does not differentiate between sitting/reclining and standing, and therefore, cannot be used to estimate

the time spent in sedentary behaviour. Therefore, the studies by Ekelund et al. (2008) and Helmerhorst et al. (2009) cannot be considered sedentary behaviour studies according to the current definition of sedentary behaviour (Sedentary Behaviour Research Network, 2012). New procedures for data processing need to be developed and validated if

heart rate monitors are to be used in sedentary behaviour research.

All the accelerometer-based studies used the common cut-point of <100 counts per minute to detect the time spent in sedentary behaviour. Due to the inability of hip-mounted accelerometers to differentiate between sitting/reclining while awake or taking a nap, sleep time may be misclassified as the time spent in sedentary behaviour. Besides, fidgeting while seated may cause misclassification of sedentary behaviour with physical activity. When evaluated against activPAL inclinometers as the criterion measure, accelerometer-based individual estimates of sedentary behaviour did not show acceptable accuracy (Aguilar-Farías, Brown, & Peeters, 2013; Healy, et al., 2011; Oliver, Schofield, Badland, & Shepherd, 2010). This measurement issue should be taken into account when interpreting results of accelerometer-based sedentary behaviour studies.

Insufficient resources often limit the choice of sedentary behaviour measures in large-scale studies. Despite that, researchers should endeavour to use direct measures of sedentary behaviour that can differentiate between sitting and standing posture (e.g. inclinometers), or, where more appropriate, self-reported measures of known validity and reliability.

### Poor adjustments for physical activity

Total physical activity, LIPA and MVPA may act as confounding variables in associations between sedentary behaviour and health outcomes, because: [i] sedentary behaviour may share as much as 64%, 81% and 25%, respectively, of common variance with these variables (Spittaels, et al., 2012), [ii] well established evidence shows a number of health benefits associated with total physical activity and MVPA (Lee, et al., 2012), and [iii] emerging findings indicate that LIPA may also be associated with different health outcomes (Buman, et al., 2010; Carson, et al., 2013; Friedenreich, Cook, Magliocco, Duggan, & Courneya, 2010; Healy, et al., 2007; Healy, et al., 2008; Pescatello, Murphy, & Costanzo, 2000). Hence, without appropriate adjustments for LIPA and MVPA we cannot draw sound conclusions about independent associations between sedentary behaviour and health.

To demonstrate independent associations between sedentary behaviour and health outcomes, 89% of the reviewed studies adjusted their analyses for one or more physical activity variables. Total physical activity was adjusted in 13%, both non-domain-specific LIPA and MVPA in 7%, and only non-domain-specific MVPA in 11% of the studies. In 39% of the studies analyses were only adjusted for non-occupational physical activity. Out of these studies, 69% assessed only leisure-time physical activities. It is obvious that the majority of studies

did not adjust for total physical activity or for both LIPA and MVPA, and a large number of studies did not adjust for all domains/types of activity that can act as confounders in the relationship between sedentary behaviour and health. It may, therefore, be that these adjustments were not thorough enough to allow for sound conclusions about the independent associations between sedentary behaviour and health outcomes.

In 88%, 8% and 4% of the studies the data on physical activity was collected using self-reports, accelerometers and heart-rate monitors, respectively. More than 30% of the studies used self-reported measures of physical activity with unknown validity and reliability. Furthermore, in most of the studies, prior to entering in the analysis, physical activity responses were collapsed to form dichotomous (insufficiently active / sufficiently active) or ordinal-scale variables. This approach neglects the continuous nature of physical activity variables and does not take into account the between-subject variability of physical activity levels within categories. To improve the quality of adjustments for physical activity, where possible and appropriate, such categorizing should be avoided.

When tested against doubly labelled water in free-living conditions, validity (expressed with Spearman's rank correlation coefficients) of self-reported and accelerometer-based measures of physical activity hardly ever exceeds .60 (Helmerhorst, Brage, Warren, Besson, & Ekelund, 2012; Van Remoortel, et al., 2012). This indicates that in most cases less than 36% of valid physical activity variance is measured, while the rest of the variance (64%) is accounted for by the random measurement error and other factors uncorrelated with the true physical activity level. Therefore, when adjusting analyses of the relationship between sedentary behaviour and health outcomes for self-reported or accelerometer-based physical activity levels, researchers only control for a small amount of true physical activity variance. Accordingly, the great portion of unadjusted physical activity variance may have caused residual confounding and led to increased effect sizes and/or false-positive results in the reviewed studies.

There are still no highly valid measures of physical activity suitable for use in large-scale sedentary behaviour research. Therefore, future studies on health outcomes of sedentary behaviour should aim to reduce the amount of unadjusted physical activity variance by using instruments that assess all domains and intensities of physical activity. Additionally, a recent study indicated that the time spent standing may be an independent health-risk factor (Katzmarzyk, 2014). Thus, this variable should also be considered as a potential confounder in the future studies.

## No adjustments for sleep duration

Sleep time may act as a confounding variable in associations between sedentary behaviour and health outcomes, because: [i] sleep time and the total time spent in sedentary behaviour may show significant associations (Booth, et al., 2012; Buman, et al., 2014; Matthews, et al., 2012), [ii] sleep duration may show significant associations with television viewing time (Basner, et al., 2007; Matthews, et al., 2012; Wijndaele, Brage, Besson, Khaw, Sharp, Luben, Bhaniani, et al., 2011), [iii] short duration of sleep may be associated with an increased risk of obesity (Cappuccio, et al., 2008), weight gain (Patel & Hu, 2008), type 2 diabetes (Cappuccio, D'Elia, Strazzullo, & Miller, 2010a), prevalent hypertension (Knutson, 2010), coronary heart disease and stroke (Cappuccio, Cooper, Delia, Strazzullo, & Miller, 2011), and all-cause mortality (Cappuccio, D'Elia, Strazzullo, & Miller, 2010b), and [iv] long duration of sleep may be associated with increased risk of type 2 diabetes (Cappuccio, et al., 2010a), cardiometabolic syndrome (Knutson, 2010), cardiovascular disease (Cappuccio, et al., 2011), colorectal cancer (Zhao, et al., 2013), cardiovascular and cancer mortality (Gallicchio & Kalesan, 2009), and all-cause mortality (Cappuccio, et al., 2010b). Hence, without adequate adjustments for sleep duration, erroneous conclusions about independent associations between sedentary behaviour and health may be drawn.

Despite that, no reviewed studies on total sedentary behaviour and only two (8%) reviewed studies on health outcomes of TV viewing/screen time (Tonstad, Butler, Yan, & Fraser, 2009; Wijndaele, Brage, Besson, Khaw, Sharp, Luben, Bhaniani, et al., 2011) adjusted their analyses for sleep duration. Matthews et al. (2012) did not adjust their analyses for sleep duration, because adding this covariate to the model did not change the magnitude of associations by more than 10%. In the other reviewed studies, no clear arguments were provided to justify non-adjustment for sleep time in their analyses. The unadjusted sleep time may have caused residual confounding in these studies and led to increased effect sizes and/or false-positive results. Future studies should, therefore, consider sleep duration as a potential confounding variable.


## Activity Balance Model: a new theoretical framework for epidemiological research

Evidence presented here casts doubt upon the conclusions about independent associations between sedentary behaviour and health outcomes. Findings about sedentary behaviour as an independent health risk factor are relatively novel, whilst research about LIPA and standing is in its early

phase. It is, therefore, likely that a large number of prior studies about MVPA and health, did not adjust their analyses for sedentary behaviour, LIPA and the time spent standing. The studies about the association between sleep duration and health could also be put under the same scrutiny. Therefore, further research may be needed to substantiate the hypotheses about sleep duration, sedentary behaviour, LIPA, standing and MVPA as health risk factors. To circumvent the issue of insufficient mutual adjustments, it may be necessary that all these behaviours are analysed together in future studies. Hereafter, a new theoretical framework for investigating health outcomes of these behaviours is presented.

The development of the new theoretical framework was motivated by recent hypotheses, and based on previous research findings. Firstly, as stated above, sound evidence shows that both too short and too long duration of sleep may increase health risks. It is suggested that the optimal amount of sleep for adults is on average 7-8 hours per day, although sleep needs may vary substantially between individuals (National Heart Lung and Blood Institute, 2011). Secondly, global authorities call for a minimum of 150 minutes of moderate or 75 minutes of vigorous-intensity physical activity per week, to reduce risk of several negative health outcomes (World Health Organization, 2010). On the other hand, it is hypothesized that excessive physical activity may also be associated with health risks (O'Keefe, Schnohr, & Lavie, 2013; Patil, et al., 2012). Thirdly, although there are still no official recommendations for the total time spent in sedentary behaviour, there are indications that the upper threshold might be set at around seven hours per day (Chau, Grunseit, Chey, et al., 2013). Finally, recent evidence suggests that LIPA (Buman, et al., 2010; Carson, et al., 2013; Friedenreich, et al., 2010; Healy, et al., 2007; Healy, et al., 2008; Pescatello, et al., 2000) and standing (Katzmarzyk, 2014; Qureshi, et al., 2003) may also be beneficial for health, but due to a lack of evidence, no recommendations for these behaviours can be issued.

Given the facts that: 1) the proportions of time allotted to sleep, sedentary behaviour, standing, LIPA and MVPA are perfectly collinear, 2) all these behaviours may be positively and/or negative associated with health, and 3) mutual adjustments between all these variables are needed if their independent contributions to health are to be determined, it seems that investigating the balance between times spent in these behaviours is the next logical step in epidemiological research. Therefore, this paper proposes the *Activity Balance Model (AB model)*; a theoretical framework for investigating associations of sleep duration, sedentary behaviour, standing, LIPA and MVPA with health outcomes (Figure 2).



Legend: LIPA = light-intensity physical activity; MVPA = moderate to vigorous-intensity physical activity

Figure 2. Activity Balance Model (AB model) – a new theoretical framework for epidemiological research

The AB model includes the times spent asleep, in sedentary behaviour, standing posture, LIPA, and MVPA as independent (exposure) variables and a selected health outcome as the dependent variable. When needed, the model can be adjusted for potential confounders and/or stratified by different groups, to account for hypothesized moderating effects. The data needed for empirical studies based on this model can be collected using motion sensors, self-reports, or a combination of both. In the case when the data is not collected using 24-hour continuous monitoring, estimated times spent in different behaviours for each participant need to be linearly rescaled in the way that their sum gives 24 hours. The accelerometer-based data processed using standard cut-points for different intensity levels does not include separate scores for the time spent standing. Such data can also be analysed based on this model, but its incompleteness should be clearly acknowledged. Furthermore, participants may experience problems in recalling the total time spent in LIPA, because it is comprised of a large number of intermittent daily activities that people may not specifically remember. Valid data on the total time spent in LIPA may, therefore, be difficult to collect using self-reports. However, it is possible to indirectly estimate the total time spent in LIPA by subtracting the sum of times/day spent in all other behaviours from 24 hours (possible negative estimates of LIPA should be replaced with 0). Such estimates of LIPA may, however, be biased, because they are inversely related to the magnitude of participants' under- or overestimation of times spent in other behaviours. Furthermore, an augmented AB model may also include the times

spent in moderate-intensity and vigorous-intensity physical activity separately, instead of MVPA. Likewise, augmented AB models may also include separate domains of physical activity and/or sedentary behaviour. However, this can only be done if the times spent in the included domains constitute the full duration of the corresponding non-domain-specific behaviour.

The AB model can be used to: 1) investigate the independent and joint associations of the times spent asleep, in sedentary behaviour, standing, LIPA and MVPA with health outcomes, 2) identify the type/s of time allocation to these behaviours that are associated with the highest risk of negative health outcomes, and 3) determine the optimal balance between the times spent in these behaviours to prevent negative health outcomes and achieve health benefits.

The independent variables in the AB model account for the whole 24-hour period (or any other period of time), and, therefore, need to be analysed using statistical analyses for compositional data. Statistical procedures designed for unconstrained data (e.g. isothermal substitution) are not suitable for analysing compositional data, and may lead to inappropriate inferences. If the data are from 24-hour continuous monitoring (e.g. using wrist worn motion sensors), the independent variables in the AB model will be perfectly collinear and the regression-based analyses for unconstrained data will not run. These analyses would run, if the data are from self-reports and not previously rescaled to form a 24-hour block. However, this is only due to uncorrelated measurement error and does not negate possible collinearity between the true scores in the independent variables. In such cases, it is likely to expect moderate multicollinearity, which is by itself acceptable. Nevertheless, moderate multicollinearity in conjunction with high measurement error (which often characterizes self-reported duration of sleep, sedentary behaviour and physical activity) may lead to wrong inferences (Grewal, Cote, & Baumgartner, 2004; Zidek, Wong, Le, & Burnett, 1996). Therefore, even in the studies based on self-reports, when analysing mutual associations of the times spent asleep, in sedentary behaviour, standing, LIPA and MVPA, statistical methods for compositional data should be preferred.

In conclusion, a number of findings about the independent associations between sedentary behaviour and health are compromised by poor adjustments for physical activity and sleep duration. To circumvent these issues, future studies on health outcomes of sleep duration, sedentary behaviour, standing, LIPA and MVPA should consider using the newly proposed AB model.

## References

- Aguilar-Farías, N., Brown, W.J., & Peeters, G.M.E.E. (2013). ActiGraph GT3X+ cut-points for identifying sedentary behaviour in older adults in free-living environments. *Journal of Science and Medicine in Sport*, 17(3), 293-299.
- Banks, E., Lim, L., Seubsman, S.A., Bain, C., & Sleigh, A. (2011). Relationship of obesity to physical activity, domestic activities, and sedentary behaviours: Cross-sectional findings from a national cohort of over 70,000 Thai adults. *BMC Public Health*, 11. doi: 10.1186/1471-2458-11-762
- Basner, M., Fomberstein, K.M., Razavi, F.M., Banks, S., William, J.H., & Rosa, R.R. (2007). American time use survey: Sleep time and its relationship to waking activities. *Sleep*, 30(9), 1085-1095.
- Bauman, A.E., Chau, J.Y., Ding, D., & Bennie, J. (2013). Too much sitting and cardio-metabolic risk: An update of epidemiological evidence. *Current Cardiovascular Risk Reports*, 7(4), 293-298.
- Booth, J.N., Bromley, L.E., Darukhanavala, A.P., Whitmore, H.R., Imperial, J.G., & Penev, P.D. (2012). Reduced physical activity in adults at risk for type 2 diabetes who curtail their sleep. *Obesity*, 20(2), 278-284.
- Boyle, T., Fritschi, L., Heyworth, J., & Bull, F. (2011). Long-term sedentary work and the risk of subsite-specific colorectal cancer. *American Journal of Epidemiology*, 173(10), 1183-1191.
- Buman, M.P., Hekler, E.B., Haskell, W.L., Pruitt, L., Conway, T.L., & Cain, K.L. (2010). Objective light-intensity physical activity associations with rated health in older adults. *American Journal of Epidemiology*, 172(10), 1155-1165.
- Buman, M.P., Winkler, E.A.H., Kurka, J.M., Hekler, E.B., Baldwin, C.M., & Owen, N. (2014). Reallocating time to sleep, sedentary behaviors, or active behaviors: Associations with cardiovascular disease risk biomarkers, NHANES 2005-2006. *American Journal of Epidemiology*, 179(3), 323-334.
- Bureau of Labor Statistics. (2013). *American time use survey - 2012 results*. Washington, DC: U.S. Department of Labor.
- Campbell, P.T., Patel, A.V., Newton, C.C., Jacobs, E.J., & Gapstur, S.M. (2013). Associations of recreational physical activity and leisure time spent sitting with colorectal cancer survival. *Journal of Clinical Oncology*, 31(7), 876-885.
- Cappuccio, F. P., Cooper, D., Delia, L., Strazzullo, P., & Miller, M.A. (2011). Sleep duration predicts cardiovascular outcomes: A systematic review and meta-analysis of prospective studies. *European Heart Journal*, 32(12), 1484-1492.
- Cappuccio, F. P., D'Elia, L., Strazzullo, P., & Miller, M. A. (2010a). Quantity and quality of sleep and incidence of type 2 diabetes: A systematic review and meta-analysis. *Diabetes Care*, 33(2), 414-420.
- Cappuccio, F.P., D'Elia, L., Strazzullo, P., & Miller, M.A. (2010b). Sleep duration and all-cause mortality: A systematic review and meta-analysis of prospective studies. *Sleep*, 33(5), 585-592.
- Cappuccio, F.P., Taggart, F.M., Kandala, N.B., Currie, A., Peile, E., & Stranges, S. (2008). Meta-analysis of short sleep duration and obesity in children and adults. *Sleep*, 31(5), 619-626.
- Carson, V., Ridgers, N.D., Howard, B.J., Winkler, E.A.H., Healy, G.N., & Owen, N. (2013). Light-intensity physical activity and cardiometabolic biomarkers in US adolescents. *PLoS ONE*, 8(8).
- Chau, J.Y., Grunseit, A., Midthjell, K., Holmen, J., Holmen, T.L., & Bauman, A.E. (2013). Sedentary behaviour and risk of mortality from all-causes and cardiometabolic diseases in adults: Evidence from the HUNT3 population cohort. *British Journal of Sports Medicine*. doi:10.1136/bjsports-2012-091974
- Chau, J.Y., Grunseit, A.C., Chey, T., Stamatakis, E., Brown, W.J., & Matthews, C.E. (2013). Daily sitting time and all-cause mortality: A meta-analysis. *PLoS ONE*, 8(11). doi: 10.1371/journal.pone.0080000
- Chau, J.Y., van der Ploeg, H.P., Merom, D., Chey, T., & Bauman, A.E. (2012). Cross-sectional associations between occupational and leisure-time sitting, physical activity and obesity in working adults. *Preventive Medicine*, 54(3-4), 195-200.
- Cong, Y.J., Gan, Y., Sun, H.L., Deng, J., Cao, S.Y., & Xu, X. (2014). Association of sedentary behaviour with colon and rectal cancer: A meta-analysis of observational studies. *British Journal of Cancer*, 110(3), 817-826.
- de Heer, H.D., Wilkinson, A.V., Strong, L.L., Bondy, M.L., & Koehly, L.M. (2012). Sitting time and health outcomes among Mexican origin adults: Obesity as a mediator. *BMC Public Health*, 12. doi: 10.1186/1471-2458-12-896
- Dunstan, D.W., Barr, E.L.M., Healy, G.N., Salmon, J., Shaw, J.E., & Balkau, B. (2010). Television viewing time and mortality: The Australian Diabetes, obesity and Lifestyle Study (AusDiab). *Circulation*, 121(3), 384-391.
- Ekelund, U., Brage, S., Besson, H., Sharp, S., & Wareham, N.J. (2008). Time spent being sedentary and weight gain in healthy adults: Reverse or bidirectional causality? *American Journal of Clinical Nutrition*, 88(3), 612-617.
- Ekelund, U., Brage, S., Griffin, S.J., & Wareham, N.J. (2009). Objectively measured moderate- and vigorous-intensity physical activity but not sedentary time predicts insulin resistance in high-risk individuals. *Diabetes Care*, 32(6), 1081-1086.
- Espinell, P.T., Chau, J.Y., van der Ploeg, H.P., & Merom, D. (2014). Older adults' time in sedentary, light and moderate intensity activities and correlates: Application of Australian Time Use Survey. *Journal of Science and Medicine in Sport*. doi: 10.1016/j.jsams.2014.02.012.
- Ford, E.S., & Caspersen, C.J. (2012). Sedentary behaviour and cardiovascular disease: A review of prospective studies. *International Journal of Epidemiology*, 41(5), 1338-1353.
- Ford, E.S., Schulze, M.B., Kröger, J., Pischon, T., Bergmann, M.M., & Boeing, H. (2010). Television watching and incident diabetes: Findings from the European Prospective Investigation into Cancer and Nutrition-Potsdam Study. *Journal of Diabetes*, 2(1), 23-27.



- Friedenreich, C.M., Cook, L.S., Magliocco, A.M., Duggan, M.A., & Courneya, K.S. (2010). Case-control study of lifetime total physical activity and endometrial cancer risk. *Cancer Causes and Control*, 21(7), 1105-1116.
- Frydenlund, G., Jorgensen, T., Toft, U., Pisinger, C., & Aadahl, M. (2012). Sedentary leisure time behavior, snacking habits and cardiovascular biomarkers: The Inter99 Study. *European Journal of Preventive Cardiology*, 19(5), 1111-1119.
- Gallicchio, L., & Kalesan, B. (2009). Sleep duration and mortality: A systematic review and meta-analysis. *Journal of Sleep Research*, 18(2), 148-158.
- George, E.S., Rosenkranz, R.R., & Kolt, G.S. (2013). Chronic disease and sitting time in middle-aged Australian males: Findings from the 45 and Up study. *International Journal of Behavioral Nutrition and Physical Activity*, 10. doi: 10.1186/1479-5868-10-20
- Gierach, G.L., Chang, S., Brinton, L.A., Lacey Jr., J.V., Hollenbeck, A.R., & Schatzkin, A. (2009). Physical activity, sedentary behavior, and endometrial cancer risk in the NIH-AARP diet and health study. *International Journal of Cancer*, 124(9), 2139-2147.
- Gollenberg, A.L., Pekow, P., Bertone-Johnson, E.R., Freedson, P.S., Markenson, G., & Chasan-Taber, L. (2010). Sedentary behaviors and abnormal glucose tolerance among pregnant Latina women. *Medicine and Science in Sports and Exercise*, 42(6), 1079-1085.
- Gomez-Cabello, A., Pedrero-Chamizo, R., Olivares, P.R., Hernández-Perera, R., Rodriguez-Marroyo, J.A., & Mata, E. (2012). Sitting time increases the overweight and obesity risk independently of walking time in elderly people from Spain. *Maturitas*, 73(4), 337-343.
- Grewal, R., Cote, J.A., & Baumgartner, H. (2004). Multicollinearity and measurement error in structural equation models: Implications for theory testing. *Marketing Science*, 23(4), 519-529.
- Grøntved, A., & Hu, F.B. (2011). Television viewing and risk of type 2 diabetes, cardiovascular disease, and all-cause mortality: A meta-analysis. *JAMA - Journal of the American Medical Association*, 305(23), 2448-2455.
- Hawkes, A.L., Lynch, B.M., Owen, N., & Aitken, J.F. (2011). Lifestyle factors associated concurrently and prospectively with co-morbid cardiovascular disease in a population-based cohort of colorectal cancer survivors. *European Journal of Cancer*, 47(2), 267-276.
- Healy, G.N., Clark, B.K., Winkler, E.A.H., Gardiner, P.A., Brown, W.J., & Matthews, C.E. (2011). Measurement of adults' sedentary time in population-based studies. *American Journal of Preventive Medicine*, 41(2), 216-227.
- Healy, G.N., Dunstan, D.W., Salmon, J., Cerin, E., Shaw, J.E., & Zimmet, P.Z. (2007). Objectively measured light-intensity physical activity is independently associated with 2-h plasma glucose. *Diabetes Care*, 30(6), 1384-1389.
- Healy, G.N., Wijndaele, K., Dunstan, D.W., Shaw, J.E., Salmon, J., & Zimmet, P.Z. (2008). Objectively measured sedentary time, physical activity, and metabolic risk the Australian Diabetes, Obesity and Lifestyle Study (AusDiab). *Diabetes Care*, 31(2), 369-371.
- Helmerhorst, H.J.F., Brage, S., Warren, J., Besson, H., & Ekelund, U. (2012). A systematic review of reliability and objective criterion-related validity of physical activity questionnaires. *International Journal of Behavioral Nutrition and Physical Activity*, 9. doi: 10.1186/1479-5868-9-103
- Helmerhorst, H.J.F., Wijndaele, K., Brage, S., Wareham, N.J., & Ekelund, U. (2009). Objectively measured sedentary time may predict insulin resistance independent of moderate- and vigorous-intensity physical activity. *Diabetes*, 58(8), 1776-1779.
- Howard, R.A., Freedman, D.M., Park, Y., Hollenbeck, A., Schatzkin, A., & Leitzmann, M.F. (2008). Physical activity, sedentary behavior, and the risk of colon and rectal cancer in the NIH-AARP Diet and Health Study. *Cancer Causes and Control*, 19(9), 939-953.
- Inoue, M., Iso, H., Yamamoto, S., Kurahashi, N., Iwasaki, M., & Sasazuki, S. (2008). Daily total physical activity level and premature death in men and women: Results from a large-scale population-based cohort study in Japan (JPHC Study). *Annals of Epidemiology*, 18(7), 522-530.
- Inoue, M., Yamamoto, S., Kurahashi, N., Iwasaki, M., Sasazuki, S., & Tsugane, S. (2008). Daily total physical activity level and total cancer risk in men and women: Results from a large-scale population-based cohort study in Japan. *American Journal of Epidemiology*, 168(4), 391-403.
- Katzmarzyk, P.T. (2014). Standing and mortality in a prospective cohort of Canadian adults. *Medicine and Science in Sports and Exercise*, 46(5), 940-946.
- Katzmarzyk, P.T., Church, T.S., Craig, C.L., & Bouchard, C. (2009). Sitting time and mortality from all causes, cardiovascular disease, and cancer. *Medicine and Science in Sports and Exercise*, 41(5), 998-1005.
- Knutson, K.L. (2010). Sleep duration and cardiometabolic risk: A review of the epidemiologic evidence. *Best Practice and Research: Clinical Endocrinology and Metabolism*, 24(5), 731-743.
- Koster, A., Caserotti, P., Patel, K.V., Matthews, C.E., Berrigan, D., & van Domelen, D.R. (2012). Association of sedentary time with mortality independent of moderate to vigorous physical activity. *PLoS ONE*, 7(6). doi: 10.1371/journal.pone.0037696
- Krishnan, S., Rosenberg, L., & Palmer, J.R. (2009). Physical activity and television watching in relation to risk of type 2 diabetes: The black women's health study. *American Journal of Epidemiology*, 169(4), 428-434.
- Landhuis, C.E., Poulton, R., Welch, D., & Hancox, R.J. (2008). Programming obesity and poor fitness: The long-term impact of childhood television. *Obesity*, 16(6), 1457-1459.

- Lee, I. M., Shiroma, E.J., Lobelo, F., Puska, P., Blair, S.N., & Katzmarzyk, P.T. (2012). Effect of physical inactivity on major non-communicable diseases worldwide: An analysis of burden of disease and life expectancy. *The Lancet*, 380(9838), 219-229.
- Lynch, B.M. (2010). Sedentary behavior and cancer: A systematic review of the literature and proposed biological mechanisms. *Cancer Epidemiology Biomarkers and Prevention*, 19(11), 2691-2709.
- Lynch, B.M., Dunstan, D.W., Healy, G.N., Winkler, E., Eakin, E., & Owen, N. (2010). Objectively measured physical activity and sedentary time of breast cancer survivors, and associations with adiposity: Findings from NHANES (2003-2006). *Cancer Causes and Control*, 21(2), 283-288.
- Lynch, B.M., Dunstan, D.W., Winkler, E., Healy, G.N., Eakin, E., & Owen, N. (2011). Objectively assessed physical activity, sedentary time and waist circumference among prostate cancer survivors: Findings from the National Health and Nutrition Examination Survey (2003-2006). *European Journal of Cancer Care*, 20(4), 514-519.
- Mathew, A., Gajalakshmi, V., Rajan, B., Kanimozhi, V.C., Brennan, P., & Binukumar, B.P. (2009). Physical activity levels among urban and rural women in south India and the risk of breast cancer: A case-control study. *European Journal of Cancer Prevention*, 18(5), 368-376.
- Matthews, C.E., George, S.M., Moore, S.C., Bowles, H.R., Blair, A., & Park, Y. (2012). Amount of time spent in sedentary behaviors and cause-specific mortality in US adults. *American Journal of Clinical Nutrition*, 95(2), 437-445.
- Mekary, R.A., Feskanich, D., Malspeis, S., Hu, F.B., Willett, W.C., & Field, A.E. (2009). Physical activity patterns and prevention of weight gain in premenopausal women. *International Journal of Obesity*, 33(9), 1039-1047.
- Meyer, A.M., Evenson, K.R., Couper, D.J., Stevens, J., Pereria, M.A., & Heiss, G. (2008). Television, physical activity, diet, and body weight status: The ARIC cohort. *International Journal of Behavioral Nutrition and Physical Activity*, 5. doi: 10.1186/1479-5868-5-68
- Moradi, T., Gridley, G., Bjork, J., Dosemeci, M., Ji, B.T., & Berkel, H.J. (2008). Occupational physical activity and risk for cancer of the colon and rectum in Sweden among men and women by anatomic subsite. *European Journal of Cancer Prevention*, 17(3), 201-208.
- National Heart Lung and Blood Institute. (2011). *Your guide to healthy sleep*. Washington, DC: National Institutes of Health. U.S. Department of Health & Human Services.
- O'Keefe, J.H., Schnohr, P., & Lavie, C.J. (2013). The dose of running that best confers longevity. *Heart*, 99(8), 588-590.
- Oliver, M., Schofield, G.M., Badland, H.M., & Shepherd, J. (2010). Utility of accelerometer thresholds for classifying sitting in office workers. *Preventive Medicine*, 51(5), 357-360.
- Orsini, N., Bellocco, R., Bottai, M., Pagano, M., Andersson, S.O., & Johansson, J.E. (2009). A prospective study of lifetime physical activity and prostate cancer incidence and mortality. *British Journal of Cancer*, 101(11), 1932-1938.
- Parent, M.T., Rousseau, M.C., El-Zein, M., Latreille, B., Désy, M., & Siemietycki, J. (2011). Occupational and recreational physical activity during adult life and the risk of cancer among men. *Cancer Epidemiology*, 35(2), 151-159.
- Parsons, T.J., Manor, O., & Power, C. (2008). Television viewing and obesity: A prospective study in the 1958 British birth cohort. *European Journal of Clinical Nutrition*, 62(12), 1355-1363.
- Pate, R.R., O'Neill, J.R., & Lobelo, F. (2008). The evolving definition of "sedentary". *Exercise and Sport Sciences Reviews*, 36(4), 173-178.
- Patel, A.V., Bernstein, L., Deka, A., Feigelson, H.S., Campbell, P.T., & Gapstur, S.M. (2010). Leisure time spent sitting in relation to total mortality in a prospective cohort of US adults. *American Journal of Epidemiology*, 172(4), 419-429.
- Patel, A.V., Feigelson, H.S., Talbot, J.T., McCullough, M.L., Rodriguez, C., & Patel, R.C. (2008). The role of body weight in the relationship between physical activity and endometrial cancer: Results from a large cohort of US women. *International Journal of Cancer*, 123(8), 1877-1882.
- Patel, S.R., & Hu, F.B. (2008). Short sleep duration and weight gain: A systematic review. *Obesity*, 16(3), 643-653.
- Patil, H.R., O'Keefe, J.H., Lavie, C.J., Magalski, A., Vogel, R.A., & McCullough, P.A. (2012). Cardiovascular damage resulting from chronic excessive endurance exercise. *Missouri Medicine*, 109(4), 312-321.
- Pavey, T.G., Geeske Peeters, G.M.E.E., & Brown, W.J. (2012). Sitting-time and 9-year all-cause mortality in older women. *British Journal of Sports Medicine*. doi: <http://dx.doi.org/10.1136/bjsports-2012-091676>
- Pescatello, L.S., Murphy, D., & Costanzo, D. (2000). Low-intensity physical activity benefits blood lipids and lipoproteins in older adults living at home. *Age and Ageing*, 29(5), 433-439.
- Pinto Pereira, S.M., Ki, M., & Power, C. (2012). Sedentary behaviour and biomarkers for cardiovascular disease and diabetes in mid-life: The role of television-viewing and sitting at work. *PLoS ONE*, 7(2). doi: 10.1371/journal.pone.0031132
- Pulsford, R.M., Stamatakis, E., Britton, A.R., Brunner, E.J., & Hillsdon, M.M. (2013). Sitting behavior and obesity: Evidence from the Whitehall II study. *American Journal of Preventive Medicine*, 44(2), 132-138.
- Qureshi, H.N., Najib, U., Chaudhary, U.J., Sallahuddin, S., Shah, I.A., & Akram, J. (2003). Effect of long standing time on bone mineral density in middle-aged males. *Asian Oceanian Journal of Radiology*, 8(2), 81-83.
- Sanchez-Villegas, A., Ara, I., Guillen-Grima, F., Bes-Rastrollo, M., Varo-Cenarruzabeitia, J.J., & Martinez-Gonzalez, M.A. (2008). Physical activity, sedentary index, and mental disorders in the sun cohort study. *Medicine and Science in Sports and Exercise*, 40(5), 827-834.

- Schuna, J.M., Johnson, W.D., & Tudor-Locke, C. (2013). Adult self-reported and objectively monitored physical activity and sedentary behavior: NHANES 2005-2006. *International Journal of Behavioral Nutrition and Physical Activity*, 10(126), 1-12.
- Sedentary Behaviour Research Network. (2012). Letter to the editor: Standardized use of the terms “sedentary” and “sedentary behaviours”. *Applied Physiology, Nutrition and Metabolism*, 37(3), 540-542.
- Shuval, K., Leonard, T., Murdoch, J., Caughy, M.O., Kohl, H.W., & Skinner, C.S. (2013). Sedentary behavior and obesity in a low-income, ethnic-minority population. *Journal of Physical Activity & Health*, 10(1), 132-136.
- Simons, C.C.J.M., Hughes, L.A.E., Van Engeland, M., Goldbohm, R.A., Van Den Brandt, P.A., & Weijenberg, M.P. (2013). Physical activity, occupational sitting time, and colorectal cancer risk in the Netherlands cohort study. *American Journal of Epidemiology*, 177(6), 514-530.
- Spittaels, H., Van Cauwenberghe, E., Verbestel, V., De Meester, F., Van Dyck, D., & Verloigne, M. (2012). Objectively measured sedentary time and physical activity time across the lifespan: A cross-sectional study in four age groups. *International Journal of Behavioral Nutrition and Physical Activity*, 9. doi: 10.1186/1479-5868-9-149
- Spurr, G.B., Prentice, A.M., Murgatroyd, P.R., Goldberg, G.R., Reina, J.C., & Christman, N.T. (1988). Energy expenditure from minute-by-minute heart-rate recording: Comparison with indirect calorimetry. *American Journal of Clinical Nutrition*, 48(3), 552-559.
- Stamatakis, E., Hamer, M., & Dunstan, D.W. (2011). Screen-based entertainment time, all-cause mortality, and cardiovascular events: Population-based study with ongoing mortality and hospital events follow-up. *Journal of the American College of Cardiology*, 57(3), 292-299.
- Thorp, A.A., Owen, N., Neuhaus, M., & Dunstan, D.W. (2011). Sedentary behaviors and subsequent health outcomes in adults: A systematic review of longitudinal studies, 1996-2011. *American Journal of Preventive Medicine*, 41(2), 207-215.
- Tonstad, S., Butler, T., Yan, R., & Fraser, G.E. (2009). Type of vegetarian diet, body weight, and prevalence of type 2 diabetes. *Diabetes Care*, 32(5), 791-796.
- Tudor-Locke, C., Burton, N.W., & Brown, W.J. (2009). Leisure-time physical activity and occupational sitting: Associations with steps/day and BMI in 54-59 year old Australian women. *Preventive Medicine*, 48(1), 64-68.
- van der Ploeg, H.P., Chey, T., Korda, R.J., Banks, E., & Bauman, A. (2012). Sitting time and all-cause mortality risk in 222 497 Australian adults. *Archives of Internal Medicine*, 172(6), 494-500.
- Van Remoortel, H., Giavedoni, S., Raste, Y., Burtin, C., Louvaris, Z., & Gimeno-Santos, E. (2012). Validity of activity monitors in health and chronic disease: A systematic review. *International Journal of Behavioral Nutrition and Physical Activity*, 9. doi: 10.1186/1479-5868-9-84
- van Uffelen, J.G.Z., Watson, M.J., Dobson, A.J., & Brown, W.J. (2010). Sitting time is associated with weight, but not with weight gain in mid-aged Australian women. *Obesity*, 18(9), 1788-1794.
- van Uffelen, J.G.Z., Wong, J., Chau, J.Y., van der Ploeg, H.P., Riphagen, I., & Gilson, N.D. (2010). Occupational sitting and health risks: A systematic review. *American Journal of Preventive Medicine*, 39(4), 379-388.
- Warren, T.Y., Barry, V., Hooker, S.P., Sui, X., Church, T.S., & Blair, S.N. (2010). Sedentary behaviors increase risk of cardiovascular disease mortality in men. *Medicine and Science in Sports and Exercise*, 42(5), 879-885.
- Wijndaele, K., Brage, S., Besson, H., Khaw, K.T., Sharp, S.J., & Luben, R. (2011). Television viewing and incident cardiovascular disease: Prospective associations and mediation analysis in the EPIC Norfolk study. *PLoS ONE*, 6(5). doi: 10.1371/journal.pone.0020058
- Wijndaele, K., Brage, S., Besson, H., Khaw, K.T., Sharp, S.J., & Luben, R. (2011). Television viewing time independently predicts all-cause and cardiovascular mortality: The EPIC Norfolk study. *International Journal of Epidemiology*, 40(1), 150-159.
- Wijndaele, K., Healy, G.N., Dunstan, D.W., Barnett, A.G., Salmon, J., & Shaw, J.E. (2010). Increased cardiometabolic risk is associated with increased TV viewing time. *Medicine and Science in Sports and Exercise*, 42(8), 1511-1518.
- Wijndaele, K., Lynch, B.M., Owen, N., Dunstan, D.W., Sharp, S., & Aitken, J.F. (2009). Television viewing time and weight gain in colorectal cancer survivors: A prospective population-based study. *Cancer Causes & Control*, 20(8), 1355-1362.
- Wilmot, E.G., Edwardson, C.L., Achana, F.A., Davies, M.J., Gorely, T., & Gray, L.J. (2012). Sedentary time in adults and the association with diabetes, cardiovascular disease and death: Systematic review and meta-analysis. *Diabetologia*, 55(11), 2895-2905.
- World Health Organization. (2010). *Global recommendations on physical activity for health*. Geneva, CH: World Health Organization.
- Yates, T., Khunti, K., Wilmot, E.G., Brady, E., Webb, D., & Srinivasan, B. (2012). Self-reported sitting time and markers of inflammation, insulin resistance, and adiposity. *American Journal of Preventive Medicine*, 42(1), 1-7.
- Zhao, H., Yin, J.Y., Yang, W.S., Qin, Q., Li, T.T., & Shi, Y. (2013). Sleep duration and cancer risk: A systematic review and meta-analysis of prospective studies. *Asian Pacific Journal of Cancer Prevention*, 14(12), 7509-7515.
- Zidek, J.V., Wong, H., Le, N.D., & Burnett, R. (1996). Causality, measurement error and multicollinearity in epidemiology. *Environmetrics*, 7(4), 441-451.

## PROBLEMI U MJERENJU I NEADEKVATNA PARCIJALIZACIJA UTJECAJA TJELESNE AKTIVNOSTI I SPAVANJA UMANJUJU VJERODOSTOJNOST ISTRAŽIVANJA SEDENTARNOG PONAŠANJA – TREBALO BI SE PREUSMJERITI NA RAVNOTEŽU IZMEĐU SPAVANJA, SEDENTARNOG PONAŠANJA, STAJANJA I AKTIVNOSTI

U ovome su radu kritički ocijenjena 54 istraživanja o povezanosti sedentarnog ponašanja i zdravlja, pri čemu je fokus na metodama prikupljanja podataka o sedentarnom ponašanju i parcijalizaciji utjecaja tjelesne aktivnosti i vremena provedenoga u spavanju. Gotovo 90% pregledanih istraživanja prikupilo je podatke o sedentarnom ponašanju koristeći anketne metode. Za 59% tih instrumenata prethodno nisu utvrđene mjerne karakteristike. Većina istraživanja nije parcijalizirala utjecaj tjelesne aktivnosti svih razina intenziteta ili ukupne tjelesne aktivnosti, a veći broj istraživanja nije parcijalizirao utjecaj tjelesne aktivnosti u svim domenama. Stoga se može zaključiti da te parcijalizacije nisu bile dovoljno temeljite da bi osigurale donošenje vjerodostojnih zaključaka o nezavisnom utjecaju sedentarnog ponašanja na zdravlje. Vrijeme provedeno u spavanju je također jedan od mogućih čimbenika koji posreduju u povezanosti između sedentarnog ponašanja i zdravlja. Unatoč tome, samo su tri istraživanja uključena u ovaj pregled parcijalizirala utjecaj vremena provedenoga u spavanju. Taj metodološki propust dodatno dovodi u pitanje za-

ključke o izravnom utjecaju sedentarnog ponašanja na zdravlje. Ako se u obzir uzme: 1) da su udjeli vremena provedenoga u spavanju, sedentarnom ponašanju, stajanju, tjelesnoj aktivnosti niskog intenziteta i tjelesnoj aktivnosti umjerenog do visokog intenziteta potpuno linearno zavisni, 2) svi ti oblici ponašanja mogu biti povezani sa zdravljem i 3) međusobna parcijalizacija utjecaja je nužna da bi se utvrdili nezavisni utjecaji tih varijabli na zdravlje, nameće se zaključak da je istraživanje ravnoteže između vremena provedenoga u tim oblicima ponašanjima sljedeći logični korak u epidemiološkim istraživanjima. Stoga je u ovome radu predložen „Activity Balance Model (AB model)“ – novi teorijski okvir za istraživanja zdravstvenih utjecaja vremena provedenog u spavanju, sedentarnom ponašanju, stajanju, tjelesnoj aktivnosti niskog intenziteta i tjelesnoj aktivnosti umjerenog do visokog intenziteta.

**Ključne riječi:** spavanje, sedentarno ponašanje, stajanje, tjelesna aktivnost, Activity Balance Model

Submitted: May 30, 2014

Accepted: June 4, 2014

Željko Pedišić, PhD  
University of Zagreb, Faculty of Kinesiology  
Horvacanski zavoj 15, HR-10000 Zagreb, Croatia  
Phone: +385 1 3658 732  
Fax: +385 1 3658 654  
E-mail: zeljko.pedisc@kif.hr