

Human exposure to endocrine disrupting chemicals as a prenatal risk factor for cryptorchidism

Ivana Fratrić¹, Dragana Živković¹, Saša Vukmirović²

This review describes the most recent data on the effects of endocrine disrupting compounds on reproductive tract development, as well as controversies in the field. One of the most frequent conditions affected by endocrine disrupting compounds is cryptorchidism. Recent reports regarding the cause of this disorder continue to increase our understanding of this common and important problem. Endocrine disruptors are defined as exogenous substances with the ability to disrupt normal endocrine homeostasis and reproduction, and they include xenoestrogens, synthetic and natural hormones, phyto- and mycoestrogens, and other substances affecting endocrine signaling. Human exposure to endocrine disrupting chemicals is widespread. Epidemiological studies suggest associations between prenatal exposure to endocrine disrupting chemicals and numerous malformations of androgen dependent tissues. Animal models and epidemiological evidence link exposure to androgen disrupting chemicals with cryptorchidism, reduced sperm cell counts, increasing infertility, and testicular and prostate cancers. Since male sexual differentiation is androgen dependent, it is highly susceptible to endocrine disruptors. Whether the level of exposure contributes to the increasing prevalence of cryptorchidism is an ongoing debate. Further, there appears to be increased sensitivity to these agents during critical developmental periods when male differentiation is at its peak. Differences in the interpretation of the available studies underlie the disparate conclusions of scientific and regulatory body panels on the potential toxicological effects of endocrine disrupting chemicals at the current levels of human exposure. This review will highlight the evidence for endocrine disrupting chemicals that act through interference with the androgen receptor and lead to cryptorchidism.

Keywords: cryptorchidism; endocrine disruptors

Cryptorchidism, defined as the absence of at least one testis in the scrotum, is a frequent condition in the pediatric population. The prevalence of cryptorchidism at birth is 2.5%-9% (1). It affects 1%-1.9% of three-month-old boys and 0.8%-1.5% of 18-month-old boys (2), and may involve the use of considerable medical and economic resources.

Risk factors for cryptorchidism include genetic predisposition, preterm birth, low birth weight and prenatal exposure to endocrine disrupting chemicals (EDCs) or tobacco in either the mother or the father (3, 4). Mutations in the gene for insulin-like factor 3 (*INSL3*) and its receptors have been recognized as the causes of cryptorchidism in some cases, but some chromosomal alterations, above all the Klinefelter syndrome, are also frequently involved.

The role of *INSL3* in testicular descent is mainly related to its effects on gubernaculum differentiation during the transabdominal phase. The transabdominal stage occurs between the 10th and 23rd week of gestation in human fetuses and the inguinoscrotal phase starts at around 26 gestational weeks and ends between 28 weeks of gestation and birth. That is the reason why cryptorchidism is more frequent in

¹ Institut za zdravstvenu zaštitu dece i omladine Vojvodine, Medicinski fakultet, Univerzitet u Novom Sadu, Srbija

² Medicinski fakultet, Univerzitet u Novom Sadu, Srbija

Correspondence to:

Ivana Fratrić, Institut za zdravstvenu zaštitu dece i omladine Vojvodine, Medicinski fakultet, Univerzitet u Novom Sadu, Srbija,
e-mail: ivana.fratric@gmail.com

Primljeno/Received: 17. 8. 2014., Prihvaćeno/Accepted: 2. 3. 2015.

preterm births. The inguinoscrotal phase is androgen dependent, and exposure to endocrine disrupting substances may affect normal androgen production and consequently interrupt normal descent of testes.

Familial cases of cryptorchidism have been described, and a family history of cryptorchidism is a risk factor for undescended testes (5); initial studies suggested that 6.2% of the brothers and 1.5%-4.0% of the fathers of patients with cryptorchidism had undescended testes.

Environmental factors acting as endocrine disruptors of testicular descent have been suggested to contribute to cryptorchidism and its increased incidence in recent years. Endocrine disruptors are defined as exogenous substances with an ability to disrupt normal endocrine homeostasis and reproduction, and they include xenoestrogens (industrial chemicals), synthetic and natural hormones, phyto- and mycoestrogens, and other substances affecting endocrine signaling (6). There are several ways how endocrine disrupting chemicals can alter endocrine functions:

1. by acting as either agonists or antagonists for the receptors for estrogens and androgens;
2. by altering the synthesis, transport and/or catabolism of endogenous hormones; and
3. by modifying the levels of expression and/or functioning of hormone receptors.

Endocrine disrupting chemicals are present in the air (7), water, soil, as well as in food products (soybeans, legumes) and plants (phytoestrogens in fruits, vegetables and grasses). Some derive from household products, while others are agricultural or industrial chemicals. Exposure can occur directly at the workplace or *via* consumer products such as food items, plastic and paints or indirectly *via* air, water and soil. The toxicity of such chemicals for different species of living organisms is determined by their individual chemical properties, duration and frequency of exposure, and their pharmacokinetics (absorption, distribution, transformation and elimination). The adverse effect of exposure at the developmental stage is irreversible, whereas most post-developmental effects are reversible (8, 9).

The Centers for Disease Control and Prevention (CDC) have classified the hormone-modulating effects of EDCs as estrogenic (approximately 48%), anti-androgenic (19%), anti-estrogenic (15%), androgenic (4%), those with effect on thyroid hormone status (42%) and others (10).

Germ, Sertoli and Leydig cells are all highly sensitive to the toxic effects of the variety of xenobiotics. Attenuation of fetal Leydig cell steroidogenesis is associated with development of congenital abnormalities such as cryptorchidism and hypospadias as a result of incomplete masculinization.

Phthalates are one of the proven EDCs. They are widely used as plasticizers (to soften plastics), as well as solvents in inks, and are present, among other places, in food packaging and certain cosmetics (11). The exact biochemical or molecular mechanisms underlying the effects of diethylhexyl phthalate (DEHP) on the testis remain unclear. The mechanisms that have been proposed include alterations in testicular levels of zinc and zinc dependent enzymatic activities; elevated oxidative stress in the testis; modulation of estrogenic activity *via* interactions with estrogen receptors; and changes in the pathways involving peroxisome proliferator-activated receptors (PPARs) or other cellular molecular event pathways. Furthermore, in male rats, inhibition of steroidogenesis in fetal Leydig cells by phthalates gives rise to numerous malformations of androgen dependent tissues, including reduction in the anogenital distance, hypospadias, epididymal agenesis and testicular atrophy (12). These abnormalities are suggestive of suboptimal virilization of the Wolffian duct and urogenital sinus. Finally, recent investigations have revealed that treatment with phthalates attenuates the transcription of several key genes whose protein products are involved both in cholesterol transport and the biosynthesis of testosterone (13, 14).

Studies published during the last few decades have indicated that the birth prevalence of cryptorchidism may have increased in some regions (15, 16). The short time interval in which this high prevalence occurred indicates that environmental factors may contribute greatly to this effect (17). Several organochlorine pesticides can cause adverse effects in the male reproductive system in animals (18, 19). In animal studies, estrogen exposure during pregnancy resulted in the development of cryptorchidism and hypospadias in male offspring (20, 21).

An increased incidence of cryptorchidism in male panthers has been attributed to EDCs in the environment, such as 1,1-dichloro-2,2-bis(4-chlorophenyl)ethane (p,p'-DDE) (22). Dose dependent in utero exposure of male rats and rabbits to dichlorodiphenyldichloroethylene (DDT) resulted in reduced anogenital distance, hypospadias, cryptorchidism and epididymal agenesis (23-25).

Increased rates of orchidopexy in the areas with extensive pesticide use in agriculture have been reported (26). In their study, *Hosie et al.* demonstrated higher pesticide levels in fat tissue samples from boys operated on for cryptorchidism than in children operated on for other reasons (27).

Damgaard et al. showed in their study that simultaneous exposure to more than one chemical at low concentrations represented a risk factor for congenital cryptorchidism. In this study, the levels of persistent organochlorine pesticides in breast milk were compared with the birth prevalence of cryptorchidism. Breast milk was chosen as a surrogate bio-

marker of previous maternal exposure to persistent pesticides because these compounds accumulate in lipid tissue and thereby in breast milk (28). Concentrations of compounds in breast milk are a suitable proxy for fetal exposure during pregnancy. Primiparae, slim women, and smokers tend to have higher pesticide levels (29). *Hosie et al.* compared the levels of pesticides (DDT and metabolites, toxaphene, HCH, HCB, PCA, PeCB, and several chlorinated cyclodienes such as heptachlor) in fat tissue biopsies from 18 cryptorchid boys and 30 controls. Pesticide concentrations were higher in cryptorchid cases than in controls, but reaching significance for only a few (HCB and heptachloroepoxide).

There are numerous studies reporting associations between cryptorchidism and parental pesticide exposure, but most of these were registry-based and retrospective.

Distinction between maternal and paternal exposure is made only in some studies; *Pierik et al.* found that cryptorchidism was significantly associated with paternal but not maternal pesticide exposure (16). *Weidner et al.* (30) describe a significantly increased risk of cryptorchidism in sons of female gardeners. This study included 6 177 males born live in Denmark between 1983 and 1992 who were discharged from the hospital with the diagnosis of cryptorchidism and 1 345 males born with hypospadias. These results were coherent with earlier studies reporting an increased frequency of orchidopexy in the areas with extensive use of pesticides in Spain (26) and an increased occurrence of cryptorchidism in boys born on Norwegian farms where pesticides had been used (31). Unlike these studies that made no discrimination between paternal and maternal exposure, the study performed in Denmark showed no effect of paternal work in gardening on cryptorchidism. The inclusion of paternal exposure serves as a control because factors related to the lifestyle of agricultural families would lead to an apparent effect of both maternal and paternal exposure, while an effect seen only or predominantly in relation to maternal exposure provides a higher degree of evidence of a true effect (30).

Exposure to pesticides can occur *via* numerous exposure pathways, including household use of pesticide products, dietary exposure to pesticide residues and occupational exposure. Biological monitoring studies indicate that pesticide exposure is widespread in pregnant women in New York City, Salinas Valley in California, the Netherlands, and Norway (32-36). Pesticides have also been detected in amniotic fluid, umbilical cord blood, meconium, and infant urine, indicating the exposure of the fetus to pesticides (37, 38). In one of the most recent studies conducted in France, bisphenol A measured in cord blood correlated negatively with *INSL3* but not with testosterone (39). The differences in dimethyl (DM) metabolites in pregnant women across the

world may be due to differences in pesticide use among different countries and differences in diet among populations. Up to 15% of tested food has been reported to have pesticide residues above maximum residue levels. The exact percent depends on the country where studies have been conducted (14% in the Netherlands and up to 6.1% in US) (36). These comparisons have limited value because of the differences in maximum residue levels between Europe and United States and differences in analytical methods between countries. Consumption of more fresh fruit and vegetables with pesticide residues is correlated to higher DM metabolites found in this population. As this habit is usually connected with Mediterranean population, it is expected to find higher total DM metabolite concentrations in this population compared with other populations of pregnant women. Unlike DM metabolites, which are higher in women taking more fresh fruit and vegetables, diethyl (DE) metabolites are likely to be higher in the samples of women using more household pesticides. The primary OP pesticides registered for use in residential setting usually metabolize to DE metabolites. DE metabolites are less stable and more difficult to detect (35). The finding that women with higher education levels have higher DM exposure levels (35) may be due to significant differences in eating habits between lower and higher socioeconomic groups, but it may also be in relation with the place of residence and household used pesticides. Up to 10% of certain OP insecticides may be stored in adipose tissue and gradually released over time into the bloodstream and undergoing metabolism and excretion (40). A study in sons of Danish pregnant women working in greenhouses showed adverse effects of maternal occupational pesticide exposure on reproductive development, despite current greenhouse safeguards and special measures to protect pregnant women (41). Another study conducted more recently showed that increasing concentrations of organotin compounds, found in diet rich in fish products, had positive association with cryptorchidism in Denmark, unlike the opposite result in Finland (42).

Data on the possible mixture effects of the specific organochlorine pesticides *in vitro* are limited. A combination of 10 compounds including endosulfan, dieldrin, methoxychlor, and some DDT metabolites demonstrated a cumulative effect (43, 44).

Prenatal exposure to persistent organochlorine pesticides may adversely affect testicular descent in boys. Exposure to endocrine disruptors during fetal, neonatal life or prepubertal life can influence pubertal timing and reproduction (45).

As the exact etiology of cryptorchidism is not known, influencing the risk factors may be the way to decrease the incidence of undescended testes.

The research in the field of endocrine disruptors as the risk factors for cryptorchidism is very important. It should be followed by providing wide information to the public. Prolonged exposure of the developing male, during both fetal and postnatal life, to endocrine disruptors could reduce Sertoli cell number and sperm output in adult life (46). It has been shown that there is a reduction in the number, as well as delayed maturation of the germ cells in gonads of patients with cryptorchidism born in Vojvodina compared to patients from either Switzerland or USA (47). Vojvodina is mainly an agricultural region, and the use of pesticides is common and also not adequately governed by the regulations. The ecological approach used did not demonstrate a cause-effect relationship between environmental pesticide contamination and the decrease in the number of germ cells in cryptorchid patients from Vojvodina. Nor did it identify the route of exposure in pregnant women, which might have been occupational, environmental (e.g., drinking water) or dietary. To clarify these questions, it would have been necessary to obtain accurate information on the actual exposure of mothers during the vulnerable period in the months prior to conception and during early gestation.

Changes in testicular histology of cryptorchid patients inspired us to design a prospective study that should tell us whether the exposure to pesticides in Vojvodina is equal to the studies in other areas, as well as give us an opportunity to compare the extent of exposure between pregnant women who gave birth to healthy boys and those who gave birth to boys with undescended testis. Future studies in the field of endocrine disruptors, which are globally ubiquitous, could help design appropriate risk assessment as a tool offering an opportunity to systematically assess the potential risk posed by EDCs (48).

NOVČANA POTPORA/FUNDING

This paper was conducted as a part of the Project "Xenobiotic interactions and their effects on systems in biomedicine" supported by the Serbian Ministry of education, science and technological development. Project number: III41012.

ETIČKO ODOBRENJE/ETHICAL APPROVAL

Nije potrebno/None

DOPRINOSI AUTORA/DECLARATION OF AUTHORSHIP

Svi autori jednako su doprinijeli izradi rada/All authors have equally contributed to a manuscript writing

SUKOB INTERESA/CONFLICT OF INTEREST

Autori su popunili the *Unified Competing Interest form* na www.icmje.org/coi_disclosure.pdf (dostupno na zahtjev) obrazac i izjavljuju: nemaju potporu niti jedne organizacije za objavljeni rad; nemaju finansijsku potporu niti jedne organizacije koja bi mogla imati interes za objavu ovog rada u posljednje 3 godine; nemaju drugih veza ili aktivnosti koje bi mogle utjecati na objavljeni rad./All authors have completed the *Unified Competing Interest form* at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare: no support from any organization for the sub-

mitted work; no financial relationships with any organizations that might have an interest in the submitted work in the previous 3 years; no other relationships or activities that could appear to have influenced the submitted work.

REFERENCES

- Boisen KA, Kaleva M, Main KM, Virtanen HE, Haavisto A-M, Schmidt IM, et al. Difference in prevalence of congenital cryptorchidism in infant between two Nordic countries. *Lancet*. 2004;363:1264-9. [http://dx.doi.org/10.1016/S0140-6736\(04\)15998-9](http://dx.doi.org/10.1016/S0140-6736(04)15998-9)
- Preiksa RT, Zilaitiene B, Matulevicius V, et al. Higher than expected prevalence of congenital cryptorchidism in Lithuania. *Hum Repr*. 2005;20:1928-32. <http://dx.doi.org/10.1093/humrep/deh887>
- Thorup J, Cortes D, Petersen BL. The incidence of bilateral cryptorchidism is increased and the fertility potential is reduced in sons born to mothers who have smoked during pregnancy. *J Urol*. 2006;176:734-7. <http://dx.doi.org/10.1016/j.juro.2006.03.042>
- Damgaard IN, Skakkebaek NE, Jorma Toppari J, et al. Persistent pesticides in human breast milk and cryptorchidism. *Environ Health Perspect*. 2006;114:1133-8. <http://dx.doi.org/10.1289/ehp.8741>
- Weidner IS, Moller H, Jensen TK, Skakkebaek NE. Risk factors for cryptorchidism and hypospadias. *J Urol*. 1999;161:1606-9. [http://dx.doi.org/10.1016/S0022-5347\(05\)68992-6](http://dx.doi.org/10.1016/S0022-5347(05)68992-6)
- Vidaeff AC, Sever LE. In utero exposure to environmental estrogens and male reproductive health: a systematic review of biological and epidemiologic evidence. *Reprod Toxicol*. 2005;20:5-20. <http://dx.doi.org/10.1016/j.reprotox.2004.12.015>
- Annamalai J, Namasivayam V. Endocrine disrupting chemicals in the atmosphere: Their effects on humans and wildlife. *Environ Int*. 2015;76:78-97. <http://dx.doi.org/10.1016/j.envint.2014.12.006>
- Colborn T, Smolen MJ, Rolland R. Environmental neurotoxic effects: the search for new protocols in functional teratology. *Toxicol Ind Health*. 1998;14:9-23. <http://dx.doi.org/10.1177/074823379801400104>
- McLachlan JA. Environmental signaling: what embryos and evolution teach us about endocrine disrupting chemicals. *Endocr Rev*. 2001;22:319-41. <http://dx.doi.org/10.1210/edrv.22.3.0432>
- Choi SM, Lee BM. An alternative mode of action of endocrine-disrupting chemicals and chemoprevention. *J Toxicol Environ Health B Crit Rev*. 2004;7:451-63. <http://dx.doi.org/10.1080/10937400490512500>
- Thomas GH. Quantitative determination and confirmation of identity of trace amounts of dialkyl phthalates in environmental samples. *Environ Health Perspect*. 1973;3:23-8. <http://dx.doi.org/10.1289/ehp.730323>
- Gray LE, Jr., Ostby J, Furr J, Price M, Veeramachaneni DN, Parks L. Perinatal exposure to the phthalates DEHP, BBP, and DINP, but not DEP, DMP, or DOTP, alters sexual differentiation of the male rat. *Toxicol Sci*. 2000;58:350-65. <http://dx.doi.org/10.1093/toxsci/58.2.350>
- Barlow NJ, Philips SL, Wallace DG, Sar M, Gaido KW, Foster PM. Quantitative changes in gene expression in fetal rat testes following exposure to di(n-butyl) phthalate. *Toxicol Sci*. 2003;73:431-41. <http://dx.doi.org/10.1093/toxsci/kfg087>
- Thompson CJ, Ross SM, Gaido KW. Di(n-butyl) phthalate impairs cholesterol transport and steroidogenesis in the fetal rat testis through a rapid and reversible mechanism. *Endocrinology*. 2004;145:1227-37. <http://dx.doi.org/10.1210/en.2003-1475>
- Boisen KA, Kaleva M, Main KM, Virtanen HE, Haavisto AM, Schmidt IM, et al. Difference in prevalence of congenital cryptorchidism in infants between two Nordic countries. *Lancet*. 2004;363:1264-9. [http://dx.doi.org/10.1016/S0140-6736\(04\)15998-9](http://dx.doi.org/10.1016/S0140-6736(04)15998-9)
- Pierik FH, Burdorf A, Deddens JA, Juttman RE, Weber RF. Maternal and paternal risk factors for cryptorchidism and hypospadias: a case-control study in newborn boys. *Environ Health Perspect*. 2004;112:1570-6. <http://dx.doi.org/10.1289/ehp.7243>

17. Sharpe RM, Skakkebaek NE. Male reproductive disorders and the role of endocrine disruption: advances in understanding and identification of areas for future research. *Pure Appl Chem*. 2003;75:2023-38. <http://dx.doi.org/10.1351/pac200375112023>
18. Edwards TM, Moore BC, Guillette LJ Jr. Reproductive dysgenesis in wildlife – a comparative view. *Int J Androl*. 2006;29:109-21. <http://dx.doi.org/10.1111/j.1365-2605.2005.00631.x>
19. Vos JG, Dybing E, Greim HA, Ladefoged O, Lambre C, Tarazona JV, et al. Health effects of endocrine-disrupting chemicals on wildlife, with special reference to the European situation. *Crit Rev Toxicol*. 2000;30:71-133. <http://dx.doi.org/10.1080/10408440091159176>
20. Grocock CA, Charlton HM, Pike MC. Role of the fetal pituitary in cryptorchidism induced by exogenous maternal oestrogen during pregnancy in mice. *J Reprod Fertil*. 1988;83:295-300. <http://dx.doi.org/10.1530/jrf.0.0830295>
21. Vorherr H, Messer RH, Vorherr UF, Jordan SW, Kornfeld M. Teratogenesis and carcinogenesis in rat offspring after transplacental and transmammary exposure to diethylstilbestrol. *Biochem Pharmacol*. 1979;28:1865-77. [http://dx.doi.org/10.1016/0006-2952\(79\)90638-5](http://dx.doi.org/10.1016/0006-2952(79)90638-5)
22. Facemir CF, Gross TS, Gulliette LJ Jr. Reproductive impairment in the Florida panther: nature or nurture. *Environ Health Perspect*. 1995;103 (suppl 4):79-86. <http://dx.doi.org/10.2307/3432416>
23. Gray LE Jr, Ostby J, Furr J, Wolf CJ, Lambright C, Parks L, et al. Effects of environmental antiandrogens on reproductive development in experimental animals. *Hum Reprod Update*. 2001;7:248-64. <http://dx.doi.org/10.1093/humupd/7.3.248>
24. Gray LE Jr, Ostby J, Furr J, et al. Toxicant-induced hypospadias in the male rat. *Adv Exp Med Biol*. 2004;545:217-41. http://dx.doi.org/10.1007/978-1-4419-8995-6_14
25. Wolf C, Gray LE Jr, Lambright C, et al. Administration of potentially antiandrogenic pesticides (procymidone, linuron, iprodione, chlozolinate, p,p'-DDE, and ketoconazole) and toxic substances (dibutyl- and diethylhexyl phthalate, PCB 169, and ethane dimethane sulphonate) during sexual differentiation produces diverse profiles of reproductive malformations in the male rat. *Toxicol Ind Health*. 1999;15:94-118. <http://dx.doi.org/10.1177/074823379901500109>
26. Garcia-Rodriguez J, Garcia-Martin M, Noguera-Ocena M, et al. Exposure to pesticides and cryptorchidism: geographical evidence of a possible association. *Environ Health Perspect*. 1996;104:1090-5. <http://dx.doi.org/10.1289/ehp.961041090>
27. Hosie S, Loff S, Witt K, Niessen K, Waag KL. Is there a correlation between organochlorine compounds and undescended testes? *Eur J Pediatr Surg*. 2000;10:304-9. <http://dx.doi.org/10.1055/s-2008-1072381>
28. Damgaard IN, Skakkebaek NE, Jorma Toppari J, et al. Persistent pesticides in human breast milk and cryptorchidism. *Environ Health Perspect*. 2006;114:1133-8. <http://dx.doi.org/10.1289/ehp.8741>
29. Jensen AA, Slorach SA. *Chemical Contaminants in Human Milk*. Boston, MA: CRC Press. 1991.
30. Weidner IS, Moller H, Jensen TK, Skakkebaek NE. Cryptorchidism and hypospadias in sons of gardeners and farmers. *Environ Health Perspect*. 1998;106:793-6. <http://dx.doi.org/10.1289/ehp.98106793>
31. Kristensen P, Irgens LM, Andersen A, Bye AS, Sundheim L. Birth defects among offspring of Norwegian farmers, 1967-1991. *Epidemiology*. 1997;8:537-44. <http://dx.doi.org/10.1097/00001648-199709000-00011>
32. Whyatt RM, Barr DB, Camann DE, et al. Contemporary-use pesticides in personal air samples during pregnancy and blood samples at delivery among urban minority mothers and newborns. *Environ Health Perspect*. 2003;111:749-56. <http://dx.doi.org/10.1289/ehp.5768>
33. Berkowitz GS, Obel J, Deych E, et al. Exposure to indoor pesticides during pregnancy in a multiethnic, urban cohort. *Environ Health Perspect*. 2003;111:79-84. <http://dx.doi.org/10.1289/ehp.5619>
34. Bradman A, Eskenazi B, Barr DB, et al. Organophosphate urinary metabolite levels during pregnancy and after delivery in women living in an agricultural community. *Environ Health Perspect*. 2005;113:1802-7. <http://dx.doi.org/10.1289/ehp.7894>
35. Ye X, Pierik FH, Hauser R, et al. Urinary metabolite concentrations of organophosphorous pesticides, bisphenol A, and phthalates among women in Rotterdam, the Netherlands: the Generation R study. *Environ Res*. 2008;108:260-7. <http://dx.doi.org/10.1016/j.envres.2008.07.014>
36. Ye X, Pierik FH, Angerer J, et al. Levels of metabolites of organophosphate pesticides, phthalates, and bisphenol A in pooled urine specimens from pregnant women participating in the Norwegian Mother and Child Cohort Study (MoBa). *Int J Hyg Environ Health*. 2009;212:481-91. <http://dx.doi.org/10.1016/j.ijheh.2009.03.004>
37. Bradman A, Barr DB, Claus Henn BG, Drumheller T, Curry C, Eskenazi B. Measurement of pesticides and other toxicants in amniotic fluid as a potential biomarker of prenatal exposure: a validation study. *Environ Health Perspect*. 2003;111:1779-82. <http://dx.doi.org/10.1289/ehp.6259>
38. Ostrea Jr EM, Bielawski DM, Posecion Jr NC, et al. Combined analysis of prenatal (maternal hair and blood) and neonatal (infant hair, cord blood and meconium) matrices to detect fetal exposure to environmental pesticides. *Environ Res*. 2009;109:116-22. <http://dx.doi.org/10.1016/j.envres.2008.09.004>
39. Chevalier N, Brucker-Davis F, Lahlou N, et al. A negative correlation between insulinlike peptide 3 and bisphenol A in human cord blood suggests an effect of endocrine disruptors on testicular descent during fetal development. *Human Reproduction*, 2015;30:447-53. <http://dx.doi.org/10.1093/humrep/deu340>
40. Needham LL. Assessing exposure to organophosphorus pesticides by biomonitoring in epidemiologic studies of birth outcomes. *Environ Health Perspect* 2005;113:494-8. <http://dx.doi.org/10.1289/ehp.7490>
41. Andersen HR, Schmidt IM, Grandjean P, et al. Impaired reproductive development in sons of women occupationally exposed to pesticides during pregnancy. *Environ Health Perspect*. 2008;116:566-72. <http://dx.doi.org/10.1289/ehp.10790>
42. Rantakokko P, Main K, Wohlfart-Veje C, et al. Association of placenta organotin concentrations with congenital cryptorchidism and reproductive hormone levels in 280 newborn boys from Denmark and Finland. *Hum Rep*. 2013;28:1647-60. <http://dx.doi.org/10.1093/humrep/det040>
43. Soto AM, Chung KL, Sonnenschein C. The pesticides endosulfan, toxaphene, and dieldrin have estrogenic effects on human estrogen-sensitive cells. *Environ Health Perspect*. 1994;102:380-3. <http://dx.doi.org/10.1289/ehp.94102380>
44. Soto AM, Sonnenschein C, Chung KL, Fernandez MF, Olea N, Olea-Serrano F. The E-SCREEN assay as a tool to identify estrogens: an update on estrogenic environmental pollutants. *Environ Health Perspect*. 1995;103 (suppl 7):173-8. <http://dx.doi.org/10.1289/ehp.95103s7113>
45. Parent AS, Franssen D, Fudvoye J, Gerard ABourguignon JP. Developmental variations in environmental influences including endocrine disruptors on pubertal timing and neuroendocrine control: Revision of human observations and mechanistic insight from rodents. *Front Neuroendocrinol*. 2015; <http://dx.doi.org/10.1016/j.yfrne.2014.12.004>
46. Živković D, Fratrić I. Disturbances of sperm maturation and minipuberty: Is there a connection? *BioMed Res Int*. 2014; <http://dx.doi.org/10.1155/2014/912746>
47. Živković D, Varga J, Konstantinidis G, Vlaski J, Snyder HM, Hadziselimović F. Regional differences in maturation of germ cells of cryptorchid testes: role of environment. *Acta Paediatr*. 2009;98:1339-43. <http://dx.doi.org/10.1111/j.1651-2227.2009.01325.x>
48. Futran Fuhrman V, Tal A, Arnon S. Why endocrine disrupting chemicals (EDCs) challenge traditional risk assessment and how to respond. *J Hazard Mater*. 2015;286:589-611. <http://dx.doi.org/10.1016/j.jhazmat.2014.12.012>

SAŽETAK

Izloženost endokrinim disruptorima kao prenatalni čimbenik rizika za kriptorhizam

I. Fratrić, D. Živković, S. Vukmirović

Ovaj pregledni članak donosi najnovije podatke o učincima endokrinih disruptora na razvoj reproduktivnog sustava, kao i o proturječjima u ovom području. Kriptorhizam je jedno od najčešćih stanja na koja utječu endokrini disruptori. Novija izvješća o uzroku ovoga poremećaja sve više doprinose razumijevanju ovoga čestog i važnog problema. Endokrini disruptori definiraju se kao egzogene tvari sposobne prekinuti normalnu endokrinu homeostazu i reprodukciju, a obuhvaćaju ksenoestrogene, sintetske i prirodne hormone, fito- i mikoestrogene te druge tvari koje utječu na endokrine signale. Ljudi su uvelike izloženi takvim kemikalijama. Epidemiološke studije ukazuju na udruženost prenatalne izloženosti endokrinim disruptorima i brojnih malformacija tkiva ovisnih o androgenima. Životinjski modeli i epidemiološki dokazi povezuju izloženost androgenim disruptorima s kriptorhizmom, smanjenim brojem spermija, povećanjem neplodnosti te s rakom testisa i prostate. Muška seksualna diferencijacija ovisna je o androgenima, a time i veoma osjetljiva na endokrine disruptore. I dalje traju rasprave o tome doprinosi li razina izloženosti rastućoj učestalosti kriptorhizma. Nadalje, izgleda da je osjetljivost na ova sredstva povećana tijekom kritičnih razvojnih razdoblja obilježenih vršnom diferencijacijom kod muškog spola. Razlike u tumačenju dostupnih studija obilježavaju neujednačene zaključke znanstvenih i regulatornih panela o potencijalnim toksikološkim učincima endokrinih disruptora uz današnju razinu izloženosti ljudi. Ovaj pregledni rad pokazuje dokaze za endokrine disruptore koji djeluju kroz interferenciju s androgenim receptorima i dovode do kriptorhizma.

Ključne riječi: kriptorhizam; endokrini disruptori