

Phenolic and Aroma Composition of White Wines Produced by Prolonged Maceration and Maturation in Wooden Barrels

Igor Lukić^{1*}, Nikolina Jedrejčić¹, Karin Kovačević Ganić², Mario Staver³ and Đordano Peršurić¹

¹Institute of Agriculture and Tourism, Karla Huguesa 8, HR-52440 Poreč, Croatia

²University of Zagreb, Faculty of Food Technology and Biotechnology, Pierottijeva 6, HR-10000 Zagreb, Croatia

³Polytechnic of Rijeka, Department of Agriculture, Karla Huguesa 6, HR-52440 Poreč, Croatia

Received: February 24, 2015

Accepted: July 8, 2015

Summary

To investigate the phenolic and aroma composition of Malvazija istarska (*Vitis vinifera* L.) white wines produced by an unconventional technology comprising prolonged maceration followed by maturation in wooden barrels, representative samples were subjected to analysis by UV/Vis spectrometry, high-performance liquid chromatography, and gas chromatography-mass spectrometry. When compared to standard wines, the investigated samples contained higher levels of dry extract, volatile acidity, lactic acid, phenols, colour intensity, antioxidant activity, majority of monoterpenes, C₁₃-norisoprenoids, methanol, higher alcohols, ethyl acetate, branched-chain esters and esters of hydroxy and dicarboxylic acids, ethylphenols, furans, and acetals, as well as lower levels of malic acid, β-damascenone, straight-chain fatty acids, ethyl and acetate esters. It was estimated that maceration had a stronger influence on phenols, and maturation on volatile aromas. Despite different vintages and technological details, the investigated wines showed a relative homogeneity in the composition, representing a clear and distinctive type.

Key words: white wine, prolonged maceration, wine maturation, phenols, wine aroma

Introduction

White wine is most commonly produced with or without short-term, often pre-fermentative maceration, applied mainly to obtain more complex flavour due to the extraction of grape aromas into the must, and to simultaneously keep phenolic compounds at acceptable levels (1–4). To achieve and preserve desirable concentrations of esters with fruity-flowery aroma, fermentation in standard white winemaking is performed by exogenous yeasts in stainless steel tanks, at relatively low temperatures between 12 and 18 °C. Certain alternative practices in white winemaking are also being utilised, such as fermentation by endogenous yeast microflora (5), fermentation and ageing in wooden barrels (6), prolonged maceration during and after fermentation (7), as well as fermentation

and ageing in amphorae (8,9). Their effects have been investigated, but independently and sporadically, covering a limited number of analysed components.

Nowadays, in an effort to expand and differentiate their offer and to increase the competitiveness on the market, wine producers in the Mediterranean countries venture into producing different types of wine by technologies that are far from standard. Some of the largest deviations from the standard white winemaking are expressed through various low-intervention approaches and philosophies, such as the production of so-called biodynamic, natural or orange wines, and similar. Wines produced using these approaches, although considered alternative, have become attractive to consumers, and after receiving a lot of attention from wine experts and enthusi-

*Corresponding author: Phone: +385 (0)52 408 327; Fax: +385 (0)52 431 659; E-mail: igor@iptpo.hr

asts today are a relevant phenomenon. Such wines are also important from an economic point of view, since they usually achieve medium to high prices, mostly because of low yields and extensive manual work (8).

Such technologies often imply many strict rules, including rigorous selection of grapes picked by hand, limited doses of sulphur dioxide, usage of exclusively indigenous yeasts, maturation on lees until bottling, as well as avoiding yeast nutrients, clarification and filtration, treatments with herbicides, pesticides and other chemicals, and any other correction of physicochemical parameters. Unconventional prolonged maceration during and after fermentation followed by maturation in wooden barrels are almost regularly included, and are the logical choice because they enable implementation of low-intervention principles such as spontaneous fermentation, clarification and stabilisation, as well as treatment with lower doses of sulphur dioxide. In return, these rather atypical practices strongly contribute to the creation of more particular and distinguished products.

It is reasonable to assume that the mentioned combination of techniques result in wines with quite a unique composition, incorporating characteristics of both white and red wines. The effects of maceration known up to date, such as increased amounts of phenols and primary aromas, higher dry extract, and reduced acidity, were determined in experiments with short-term, usually prefermentative white grape mash maceration (1–4,10). Published studies that have investigated prolonged maceration in white winemaking are only a few (7,11,12), and the chemical composition of such wines is almost completely unknown.

The aim of this study is to characterise white wines obtained by the described unconventional technology on the basis of phenolic and aroma compound composition. Wines included were produced in the Istria region of Croatia where, after Italian and Slovenian winemakers who started such trends, a group of innovative producers adopted a number of the previously mentioned principles. Such wines were compared to standard white wines from the same variety. It was considered that the results of this study, although set in the local environment, would provide a significant contribution to the knowledge about global diversity and variability of technological procedures applied in white winemaking, as well as that of the chemical composition of white wine.

Materials and Methods

Wine samples

Wine samples selected for this study were produced from the grapes of Malvazija istarska (*Vitis vinifera* L.). Six samples of wines produced by prolonged maceration during and after fermentation and maturation in wooden barrels, at the peak of their technological maturity, were consigned by local producers together with the largest part of the production documentation. Grape clusters were harvested manually from vines with reduced crop load, and were crushed and mashed without the addition of sulphur dioxide and pectolytic enzymes. During maceration, mashes were fermented with the action of endog-

enous yeasts, without the addition of yeast activators, pectolytic enzymes, or any other commercial substrates. All wines underwent spontaneous malolactic fermentation. The majority of samples underwent spontaneous stabilisation and clarification during maturation on lees, which lasted until bottling or at least until the first racking. Clarification of one sample was performed with the use of natural fish swimming bladder. Minimum SO₂ doses (<3 g/L) were added prior to bottling. No correction of any other physicochemical parameter was conducted. Other details on the production of wines obtained by prolonged maceration and maturation in wooden barrels are presented in Table 1.

Six samples of standard young Malvazija istarska wines produced by standard white winemaking from grapes harvested in 2010 were collected from the regional market. Basic data on the production collected from the producers through interviews confirmed that standard practices were applied: grape mashes were treated with usual doses of SO₂ (10–15 g per 100 kg) and pectolytic enzymes (2–5 g per 100 kg). Mashes were subjected or not to short-term maceration with a maximum duration of 24 h at temperatures below 20 °C. Fermentation was conducted with commercial *Saccharomyces cerevisiae* wine yeasts in stainless steel tanks of various volumes at temperatures between 17 and 19 °C. Commercial yeast activators were utilised. Protein stabilisation (in all wines) and tartrate stabilisation (cold stabilisation of two wines and metatartaric acid stabilisation of three wines) were performed. Six samples of standard Malvazija istarska wines from the market produced in 2009 by standard white winemaking, matured in bottle for a period of one year at variable temperatures between 17 and 20 °C in the dark, in the minivinification cellar of the Institute of Agriculture and Tourism, Poreč, Croatia, were also included in the investigation. All samples were analysed in June 2011.

Chemical standards and standard solutions

Pure standards of individual phenolic and volatile aroma compounds were purchased from Merck (Darmstadt, Germany) and Fluka (Buchs, Switzerland). Stock standard solutions were prepared in ethanol. Working standard solutions were prepared by diluting stock standard solutions in synthetic wine.

Analysis of standard physicochemical parameters and organic acids

Standard physicochemical wine parameters were determined according to the EEC regulation No. 2676/90 (13). Organic acids were identified and quantified using a Varian HPLC system (Varian Inc., Harbour City, CA, USA) with a ProStar 320 solvent delivery module and a ProStar 310 UV-Vis detector. A MetaCarb 87H 300 column (300 mm×7.8 mm i.d., 5 µm particle size) was used. Samples were diluted ten times with an 11 % ethanol solution, and filtered through 0.45-µm nylon filters. Isocratic elution with a 0.0125 M sulphuric acid in deionized water was applied. The flow was 0.7 mL/min and the temperature was set at 65 °C. Chromatograms were recorded at 214 nm. Acids were identified by comparing their retention times with those of pure standards. Quantification was

Table 1. Production parameters of Malvazija istarska wines produced by prolonged maceration during and after fermentation followed by maturation in wooden barrels

Production parameter	NMI-1	NMI-2	NMI-3	NMI-4	NMI-5	NMI-6
Harvest	2003	2007	2007	2008	2008	2009
Simultaneous maceration and fermentation						
Tank/barrel material	oak wood	clay amphora	oak wood	stainless steel	stainless steel	oak wood
<i>t</i> /day	21	14	16	15	17	20
Cap management	PD	PD	PD, S	PO	PD, S	PD, S
Temperature/°C	ambient	18–20	18–20	18–20	27	<30
Post-fermentative maceration						
Tank/barrel material	oak wood	clay amphora	oak wood	–	stainless steel	oak wood
<i>t</i> /day	150	2	100	–	3	40
Cap management	–	–	–	–	PD	PD, S
Temperature/°C	ambient	20–22	18–20	–	27	<30
Maturation						
Tank/barrel material	oak wood	oak wood	oak wood	oak wood	oak wood	oak wood
<i>t</i> (in wooden barrel)/month	30	12	30	24	17	12
<i>t</i> (in glass bottle)/month	54	30	12	6	13	6
<i>t</i> (total)/month	84	42	42	30	30	18

NMI-1 to NMI-6=samples of wines produced by prolonged maceration and maturation in six replicates, PD=punching down, S=stirring, PO=pumping over

performed using standard calibration curves. The accuracy of the method was checked by the addition of the solution of chemical standards (each acid at 1 g/L) to a wine sample, and it ranged from 97 to 101 % with standard deviation not surpassing 3 % ($N=5$). Linearity was checked by analysing five different concentrations of standard solutions of each acid ranging from 0.1 to 10 g/L and $R^2>0.9999$ was obtained in all cases.

Analysis of phenols, colour parameters and antioxidant activity

The concentrations of total phenols, total flavonoids, non-anthocyanin flavonoids and total proanthocyanidins, and vanillin index were determined according to Di Stefano *et al.* (14) using a Cary 50 UV/Vis spectrophotometer (Varian Inc.). Total phenols were determined using the Folin-Ciocalteu reagent and absorbance measurement at 760 nm with gallic acid as a calibration standard. The concentration of total and non-anthocyanin flavonoids and proanthocyanidins (high-molecular-mass proanthocyanidins) were measured at 280, 550 and 525 nm, respectively, while vanillin index (flavan-3-ol monomers and oligomers) was determined at 500 nm.

Colour intensity was measured at 420 nm, which corresponds to the brownish hues resulting from chemical and enzymatic browning reactions (7).

Antioxidant activity was determined according to Brand-Williams *et al.* (15). A reaction between wine antioxidants and 0.094 mM 2-diphenyl-1-picrylhydrazyl (DPPH) in methanol was induced and their antioxidant activity was evaluated by measuring the absorbance at 515 nm after 60 min at 20 °C in a water bath. Calibration curves were prepared using a Trolox (6-hydroxy-2,5,7,8-tetramethylchro-

man-2-carboxylic acid) reagent solutions.

Phenolic acids and flavan-3-ols were analysed using the already described HPLC system. Samples were diluted and filtered. A ChromSep C₁₈ column (250 mm×4.6 mm i.d., 5 µm particle size) with an OmniSpher 5 C₁₈ guard column was used. A gradient of solvents A (water/phosphoric acid at 99.5:0.5, by volume) and B (acetonitrile/water/phosphoric acid at 50:49.5:0.5, by volume) was applied as follows: 0–2 min, 100 % A isocratic; 2–7 min, 80 % A linear; 7–25 min, 60 % A linear; 25–31 min, 60 % A isocratic; 31–40, 0 % A linear; 40–42 min, 0 % A isocratic; 42–47 min, 100 % A linear, and 47–57 min, 100 % A isocratic. The flow was 1.0 mL/min and the temperature was 30 °C. Chromatograms were recorded at 280 nm. Identification was performed by comparing retention times with those of pure standards. Quantification was performed using standard calibration curves. The accuracy of the method was checked by the addition of the solution of chemical standards (each phenol at 10 mg/L) to a wine sample, and it ranged from 95 to 99 % with standard deviation not surpassing 5 % ($N=5$). Linearity was checked by the analysis of standard solutions of each phenol at five different concentrations ranging from 1 to 20 mg/L, and $R^2>0.998$ was obtained in all cases.

Analysis of volatile aroma compounds

Minor volatile aroma compounds were isolated using headspace solid-phase microextraction (HS-SPME) according to the modified method proposed by Noguerol-Pato *et al.* (16), and analysed by gas chromatography-mass spectrometry (GC-MS).

SPME fibre holder and 50/30 nm divinylbenzene-carboxen-polydimethylsiloxane (DVB-CAR-PDMS) fibres were

purchased from Supelco (Bellefonte, PA, USA). All wine samples were adjusted to 11 % of ethanol (by volume). Compounds with high analytical responses were analysed from a 15-fold diluted wine sample. A sample (4 mL) was placed in a 10-mL glass vial. Internal standard (3-octanol, 30 mg/L, 50 μ L) and sodium chloride (1.4 g) were added. The vial was sealed with a Teflon-faced septum cap, and the sample was preconditioned at 35 °C for 15 min. Microextraction lasted for 40 min at 35 °C with stirring (1100 rpm). For desorption, the fibre was inserted into a GC-MS injector port for 10 min (3-minute splitless mode).

Identification and quantification of minor volatiles was performed using a Varian 3900 GC coupled with a Varian Saturn 2100T ion trap mass spectrometer (Varian Inc.). The column used was a 60 m \times 0.25 mm i.d., 0.25 μ m film thickness Rtx-WAX (Restek, Bellefonte, PA, USA). Initial oven temperature was 40 °C, then increased at 2 °C/min to 240 °C, and then kept at 240 °C for 10 min. Injector, transfer line and ion trap temperatures were 245, 180 and 120 °C, respectively. Mass spectra were acquired in electron impact mode (70 eV) at 1 scan/s, full scan with a range of 30–450 *m/z*. The carrier gas was helium (1 mL/min). Identification was performed by comparing retention times and mass spectra with those of pure standards when available, and with mass spectra from NIST05 library (National Institute of Standards and Technology, Gaithersburg, MD, USA). Linear retention indices (relative to *n*-alkanes) were calculated and compared to those from literature. When standards were available, standard calibration curves (based on quantification ions) were constructed. For other compounds semi-quantitative analysis was carried out, and their concentrations were expressed in μ g/L of equivalents of compounds with similar chemical structure for which standards were available, assuming a response factor equal to one.

Major volatiles: acetaldehyde, ethyl acetate, methanol, 1-propanol, 1-butanol, isobutanol and isoamyl alcohol were analysed by a direct injection after distillation on a Varian 3350 GC (Varian Inc.) with a flame-ionisation detector and the same Rtx-WAX column. Initial oven temperature was 40 °C, increased after 4 min at 5 °C/min to 90 °C, then at 15 °C/min to 235 °C, and then kept for 10 min. Injector and detector temperatures were 160 and 240 °C, respectively. The carrier gas was helium (1.1 mL/min). Internal standard was 1-pentanol, with the concentration in the analytical sample of 162.18 mg/L. A volume of 2 μ L was injected in the split mode (1:20). Compounds were identified by comparing their retention times to those of pure standards. Calibration curves were constructed. The accuracy of the method was checked by the addition of the solution of chemical standards (each compound at 30 mg/L) to a wine sample, and ranged from 95 to 99 % with standard deviation not surpassing 5 % (*N*=5). Linearity was checked by the analysis of standard solutions of each compound at five different concentrations ranging from 10 to 500 mg/L, and $R^2 > 0.998$ was obtained in all cases.

Odour activity values (OAV) of volatile aroma compounds were calculated as the quotients of their concentration and the corresponding odour perception threshold from the literature.

Statistical analysis

All analyses were performed in duplicates, and mean values were used in further data elaboration. Mean values of concentrations and their standard deviations were calculated from six replicates, *i.e.* six samples of each investigated group of wines. One-way analysis of variance (ANOVA) was carried out using Microsoft Excel (Microsoft, Seattle, WA, USA), and least significant difference (LSD) test was used to compare the mean values at the level of significance of $p < 0.05$.

Results and Discussion

Physicochemical parameters of wines and concentration of organic acids

The results of the standard physicochemical analyses are presented in Table 2. As expected, the highest total dry extract was found in the wines produced by prolonged maceration followed by maturation in wooden barrels. Higher volatile acidity in these wines probably resulted from the oxidative conditions during maceration and maturation.

The technology of vinification of these wines included spontaneous malolactic fermentation (Table 1), which explains the lowest malic and the highest lactic acid concentration in these wines (Table 2). Similar can be concluded for the low concentration of citric acid in the wines, since it may also serve as a substrate for the growth of malolactic bacteria.

Phenolic compounds, colour parameters and antioxidant activity of wines

Concentration of total phenols, flavonoids and proanthocyanidins, and vanillin index

The average concentration of total phenols in white wines produced by standard technology with or without one-year maturation (Table 3) was consistent with the concentrations previously found in standard Malvazija istarska (4,17) and other white wines (1,18).

Wines produced by prolonged maceration and maturation contained significantly higher concentrations of total phenols, which is probably primarily a result of their extraction during maceration. The concentrations of total phenols in these wines were lower than those previously found in white wines produced by prolonged maceration but without maturation (7,19). This is possibly a result of the reduction of phenols during maturation of wines produced by prolonged maceration due to hydrolysis, oxidation, complexation, as well as adsorption on yeast cells and other macromolecules followed by precipitation, which was described by Baiano *et al.* (8) and Recamales *et al.* (18). Strong negative correlation found between the concentration of total phenols and the total duration of the maturation of wines produced by prolonged maceration and maturation (*N*=6), with the correlation coefficient of -0.74 , supports this thesis. Markedly higher concentrations of flavonoids, and especially proanthocyanidins in these wines (Table 3) are probably the result of their good solubility in ethanol and extraction from grapes into wine during maceration after finished alcoholic fermentation, as implied in a recent study (20). The extraction of certain

Table 2. Ranges of standard physicochemical parameters and the concentration of major organic acids in Malvazija istarska wines produced by prolonged maceration during and after fermentation followed by maturation in wooden barrels (NMI), standard white wine production technology followed by maturation in bottle for one year (SMI-1), and standard white wine production technology without maturation (SMI)

Parameter	NMI		SMI-1		SMI	
	Mean value±S.D.	Range	Mean value±S.D.	Range	Mean value±S.D.	Range
Relative density	0.991±0.001	0.989–0.992	0.992±0.001	0.991–0.994	0.9907±0.0006	0.9896–0.9911
φ (alcohol)/%	(14.3±0.3) ^a	13.9–14.8	(12.9±0.8) ^b	11.4–13.7	(13.0±0.5) ^b	12.3–13.6
γ (total dry extract)/(g/L)	(24.7±2.7) ^a	19.6–27.6	(22.3±1.1) ^b	21.7–24.5	(19.7±1.0) ^c	18.9–21.4
γ (reducing sugars)/(g/L)	2.4±0.7	1.3–3.4	2.1±0.6	1.7–3.2	1.6±0.4	1.1–2.2
γ (ash)/(g/L)	2.8±0.6	2.2–3.8	2.6±0.4	2.12–3.08	2.5±0.4	2.0–3.2
pH	3.6±0.2	3.4–3.8	3.5±0.1	3.4–3.7	3.6±0.2	3.4–3.9
Total acidity/(g/L)	(5.2±0.3) ^a	4.8–5.6	(5.7±0.7) ^a	4.8–6.6	(4.5±0.4) ^b	4.0–5.0
Volatile acidity/(g/L)	(0.7±0.1) ^a	0.6–0.9	(0.24±0.07) ^b	0.2–0.4	(0.4±0.1) ^b	0.2–0.6
γ (free SO ₂)/(mg/L)	(9±2) ^c	7–14	(15±5) ^b	10–24	(23±4) ^a	19–30
γ (total SO ₂)/(mg/L)	(67±10) ^c	56–86	(94±14) ^b	79–109	(126±30) ^a	89–180
γ (organic acids)/(mg/L)						
Citric acid	(164±100) ^b	14–281	(278±94) ^{ab}	149–397	(392±134) ^a	246–566
Tartaric acid	(1718±210) ^b	1511–2053	(1615±181) ^b	1325–1819	(2210±223) ^a	1825–2496
Malic acid	(103±93) ^c	0–217	(2588±373) ^a	2114–3095	(1791±215) ^b	1498–2137
Lactic acid	(1931±677) ^a	1444–3272	(692±593) ^b	161–1506	(333±33) ^b	285–383

Different superscript lowercase letters in a row present statistically significant differences between mean values at $p < 0.05$ obtained by one-way ANOVA and least significant difference (LSD) test, S.D.=standard deviation ($N=6$)

Table 3. Concentration and ranges of phenols, colour parameters and antioxidant activity of Malvazija istarska wines produced by prolonged maceration during and after fermentation followed by maturation in wooden barrels (NMI), standard white wine production technology followed by maturation in bottle for one year (SMI-1), and standard white wine production technology without maturation (SMI)

Parameter	γ /(mg/L)					
	NMI		SMI-1		SMI	
	Mean value±S.D.	Range	Mean value±S.D.	Range	Mean value±S.D.	Range
Phenols						
Total phenols	(660±124) ^a	476–800	(362±55) ^b	285–436	(290±70) ^b	220–390
Total flavonoids	(386±107) ^a	296–585	(90±51) ^b	25–147	(111±30) ^b	78–156
Proanthocyanidins	(146±82) ^a	34–281	(32±19) ^b	13–68	(11±5) ^b	4–19
Vanillin index	(91±48) ^a	19–144	(80±50) ^a	49–180	(15±5) ^b	8–21
Hydroxycinnamic acids						
Caffeic acid	11.8±5.0	8.1–19.0	6.2±5.5	1.6–16.9	5.3±4.2	1.4–12.4
<i>p</i> -Coumaric acid	2.3±1.0	1.2–3.9	1.3±0.8	0.7–2.9	1.5±0.7	0.7–2.7
Ferulic acid	(2.2±1.2) ^a	0.6–3.5	(0.7±0.4) ^b	0.2–1.0	(1.0±0.2) ^b	0.8–1.2
Hydroxybenzoic acids						
Protocatechuic acid	(14.9±6.9) ^a	3.6–22.2	(3.4±1.5) ^b	1.1–5.7	(6.4±1.1) ^b	5.5–8.6
Vanillic acid	(12.3±11.8) ^a	0.0–29.9	(0.8±0.4) ^b	0.4–1.3	(4.7±2.1) ^{ab}	2.1–7.8
Syringic acid	(2.3±1.3) ^a	0.6–4.3	(0.7±0.4) ^b	0.3–1.4	(0.7±0.7) ^b	0.2–1.9
Flavan-3-ols						
(-)-Epicatechin	(17.3±14.2) ^a	3.4–42.3	(2.6±2.6) ^b	0.9–7.7	(2.9±1.7) ^b	1.7–6.2
(+)-Catechin	(31.6±20.7) ^a	1.6–64.8	(7.8±5.2) ^b	3.2–17.6	(7.9±4.4) ^b	3.0–15.8
Colour intensity ($A_{420\text{nm}}$)	(0.40±0.07) ^a	0.30–0.52	(0.12±0.09) ^b	0.09–0.14	(0.12±0.01) ^b	0.10–0.13
Antioxidant activity (DPPH [•] as TE/(mmol/L))	(4.7±0.7) ^a	4.0–5.8	(1.5±0.5) ^b	0.9–2.2	(0.8±0.2) ^c	0.6–1.2

Different superscript lowercase letters in a row present statistically significant differences between mean values at $p < 0.05$ obtained by one-way ANOVA and least significant difference (LSD) test, S.D.=standard deviation ($N=6$). TE=Trolox equivalent

phenols from the wood of the barrel probably contributed as well.

Concentration of phenolic acids and flavan-3-ols in wine samples

Except maceration, other possible causes of the increase of the concentration of phenolic acids in the wines produced by prolonged maceration and maturation in relation to standard wines produced with or without maturation (Table 3) comprise hydrolysis of corresponding esters and release of their free forms (18,21), as well as their extraction from wood (22,23). Competitive reactions that could have reduced the concentration of phenolic acids include their esterification with tartaric acid to form quinones, hydrolysis, oxidation, and complexation (7,23).

Higher average concentrations of flavan-3-ols in wines produced by prolonged maceration and maturation (Table 3) are in accordance with the findings of a number of authors who noted an increase of flavan-3-ol concentration after a certain maceration period (1,24). However, in this work strong negative correlation between (+)-catechin and (–)-epicatechin concentration and total maceration duration was found in macerated wines ($N=6$), with the correlation coefficients of -0.85 and -0.71 , respectively. Apart from that, (+)-catechin content was found to be inversely proportional to total maturation duration ($R^2=-0.54$). On the basis of the results of Ortega *et al.* (23), it was presumed that the decrease in flavan-3-ol concentration during prolonged maceration and maturation was partially caused by its condensation with acetaldehyde as well as oxidation and condensation with glyoxylic acid.

Colour parameters of wines

Wines produced by prolonged maceration and matured in wooden barrels had significantly higher colour intensities of brown hues measured at 420 nm in relation to standard wines produced with or without maturation (Table 3). This was most likely the result of enzymatic as well as oxygen-assisted chemical browning reactions during fermentation, maceration and maturation, as well as the extraction of different compounds from the wood material into the wine, as noted by others (7,22,23).

Antioxidant activity of wines

Almost six times higher average value of the antioxidant activity found in the wines produced by prolonged maceration and maturation in relation to standard wines produced with or without maturation (Table 3) is comparable even with those found in certain red wines (25). High positive correlation coefficients between antioxidant activity and the concentration of total phenols, flavonoids and proanthocyanidins were determined in the wines produced by prolonged maceration and maturation ($N=6$): 0.9018, 0.9490 and 0.8741, respectively, and in standard wines produced without maturation ($N=6$): 0.9672, 0.6723, and 0.9756, respectively.

Concentration of volatile aroma compounds in the produced wines

Volatile monoterpenes

The composition of monoterpenes in standardly produced wines was in a fair agreement with that previously reported for Malvazija istarska young wine (4,26), while

in the wines produced by prolonged maceration and maturation it was significantly different (Table 4).

Maceration generally favours the extraction of monoterpenes into must, while during maturation of wine their glycosidic precursors hydrolyse and release free volatile forms (4), which probably contributed to the higher content of linalool in the wines produced by prolonged maceration and maturation (Table 4). It is possible that a part of monoterpene glycosides decreased during fermentation and prolonged maceration due to precipitation, absorption to yeast cells and solids, as well as hydrolysis, as observed by Zoecklein *et al.* (27), and because of oxidation, acid-catalysed conversions, and evaporation during maturation, as noted by other authors (28,29). In this context, it is interesting to point out high negative correlation coefficients ranging from -0.68 to -0.78 ($N=6$) found between the concentration of the major monoterpenols linalool, nerol, citronellol and geraniol and the duration of maturation of macerated wines.

The highest amount of α -terpineol in these wines (Table 4) is in line with previous studies which reported that this compound is formed by oxidation of other monoterpenols and increases during wine ageing (28–30).

C_{13} -norisoprenoids

The highest concentration of β -damascenone was found in standard wines, while the highest levels of other C_{13} -norisoprenoids were found in the wines produced by prolonged maceration and maturation (Table 4). This result is in line with the findings by Oliveira *et al.* (29), who explained a decrease in β -damascenone concentration during maturation as a consequence of its rapid release from its precursor 3,6,9-trihydroxymegastigma-6,7-diene and reaction with sulphur dioxide. Other authors found a positive correlation of vitispirane, 1,1,6-trimethyl-1,2-dihydronaphthalene (TDN) and actinidol concentration with wine ageing (29,31,32). Except for β -damascenone and actinidols, the correlation between C_{13} -norisoprenoid concentration and total maturation duration in the wines produced by prolonged maceration and maturation was very strong with the regression coefficients ranging from 0.83 to 0.93 ($N=6$). A positive correlation was found between all analysed C_{13} -norisoprenoids except β -damascenone and the duration of maceration of the wines, with the coefficients ranging from 0.74 to 0.95 ($N=6$).

Methanol and higher alcohols

The highest methanol content was found in the wines produced by prolonged maceration and maturation (Table 4), most probably as a result of prolonged maceration. Although high, the levels of methanol in these wines were under the maximum acceptable limit set by the International Organisation of Vine and Wine (OIV) (33).

Elevated concentrations of higher alcohols found in macerated wines are probably the result of the influence of various parameters which were demonstrated to favour their formation: lengthy maceration with aeration, lower partial pressure of CO_2 in wooden barrels in relation to stainless steel tanks, higher fermentation temperature, malolactic fermentation, and slow spontaneous clarification (6,34). Câmara *et al.* (35) partly explained the increase in higher alcohols during ageing in wooden bar-

Table 4. Concentration and ranges of volatile aroma compounds in Malvazija istarska wines produced by prolonged maceration during and after fermentation followed by maturation in wooden barrels (NMI), standard white wine production technology followed by maturation in bottle for one year (SMI-1), and standard white wine production technology without maturation (SMI)

Compound	LRI	ID	γ /($\mu\text{g/L}$)					
			NMI		SMI-1		SMI	
			Mean value \pm S.D.	Range	Mean value \pm S.D.	Range	Mean value \pm S.D.	Range
<i>Monoterpenes</i>								
2-Ethenyltetrahydro-2,6,6-trimethyl-2H-pyran	1111	RI, MS	(47.1 \pm 32.7) ^a	18.1–90.8	(9.7 \pm 4.8) ^b	4.8–18.1	(0.5 \pm 1.2) ^b	0.0–3.0
Limonene	1196	S, RI, MS	(26.7 \pm 7.4) ^a	17.2–36.8	(13.1 \pm 3.6) ^b	10.1–18.9	(8.1 \pm 1.9) ^b	6.1–11.3
α -Terpinolene	1281	RI, MS	(8.4 \pm 2.8) ^a	4.6–12.4	(1.2 \pm 3.0) ^b	0.0–7.3	(3.8 \pm 4.5) ^b	0.0–9.2
Terpineol (stereoisomer)	1446	MS	(114.7 \pm 38.2) ^a	47.6–150.8	(24.0 \pm 19.3) ^b	5.3–60.2	(13.4 \pm 11.0) ^b	0.0–26.1
Nerol oxide	1467	RI, MS	(40.0 \pm 26.9) ^a	7.8–74.6	(9.7 \pm 6.4) ^b	5.5–22.5	(3.9 \pm 4.1) ^b	0.0–11.5
α -Terpinene	1506	RI, MS	(20.7 \pm 7.9) ^a	14.1–35.1	(3.6 \pm 0.6) ^b	2.9–4.3	n.d.	–
Linalool	1542	S, RI, MS	(37.4 \pm 23.5) ^a	13.1–75.2	(20.3 \pm 4.4) ^b	16.1–28.5	(15.3 \pm 4.9) ^b	8.2–22.9
HO-trienol	1601	RI, MS	22.8 \pm 17.8	11.4–58.1	15.4 \pm 6.2	8.9–25.0	13.2 \pm 5.4	7.4–20.6
α -Terpineol	1684	S, RI, MS	(44.0 \pm 18.0) ^a	18.3–61.2	(19.2 \pm 9.0) ^b	8.6–30.9	(5.3 \pm 2.3) ^c	2.0–8.9
Citronellol	1758	S, RI, MS	2.2 \pm 2.9	0.0–6.4	1.6 \pm 0.4	1.2–2.4	2.0 \pm 1.0	0.5–3.2
Nerol	1791	S, RI, MS	3.1 \pm 2.0	1.0–5.2	2.2 \pm 1.0	1.1–4.1	4.1 \pm 2.1	1.7–7.9
Geraniol	1838	S, RI, MS	8.1 \pm 8.6	0.8–23.8	4.9 \pm 2.2	2.7–8.1	6.8 \pm 3.1	3.2–11.8
<i>C₁₃-norisoprenoids</i>								
Vitispirane (isomer I)	1521	RI, MS	(37.1 \pm 36.3) ^a	5.0–107.0	(5.6 \pm 4.2) ^b	1.9–13.7	(1.3 \pm 1.1) ^b	0.0–2.6
Actinidol ethyl ether (isomer I)	1690	RI, MS	(21.4 \pm 17.4) ^a	3.0–49.2	(6.3 \pm 5.6) ^b	0.8–16.3	(0.3 \pm 0.4) ^b	0.0–0.9
Actinidol ethyl ether (isomer II)	1723	RI, MS	12.9 \pm 11.5	0.0–31.2	5.2 \pm 4.3	0.8–12.5	n.d.	–
TDN	1731	RI, MS	12.1 \pm 14.7	1.3–41.3	3.2 \pm 2.9	0.5–8.2	1.8 \pm 0.9	1.0–2.8
β -Damascenone	1809	RI, MS	(1.4 \pm 0.9) ^b	0.5–2.9	(1.5 \pm 0.2) ^b	1.2–1.8	(2.4 \pm 0.6) ^a	1.4–3.2
TPB	1816	RI, MS	(3.3 \pm 3.3) ^a	0.5–9.5	(0.7 \pm 1.0) ^b	0.0–2.6	n.d.	–
Actinidol (isomer I)	1914	RI, MS	(5.6 \pm 4.1) ^a	0.0–10.5	(3.3 \pm 2.4) ^{ab}	0.7–6.7	(0.4 \pm 0.9) ^b	0.0–2.2
Actinidol (isomer II)	1927	RI, MS	(8.8 \pm 5.3) ^a	1.0–14.5	(5.7 \pm 3.3) ^a	1.4–10.4	(0.8 \pm 1.4) ^b	0.0–3.4
<i>C₆-alcohols</i>								
1-Hexanol	1356	S, RI, MS	1046.7 \pm 622.5	353.2–1796.9	1200.1 \pm 504.1	581.8–2077.3	1021.2 \pm 692.9	385.2–2109.4
<i>trans</i> -3-Hexen-1-ol	1361	S, RI, MS	(31.6 \pm 17.6) ^b	11.9–52.8	(104.2 \pm 62.7) ^a	43.2–222.2	(55.4 \pm 20.9) ^b	36.1–93.1
<i>cis</i> -3-Hexen-1-ol	1379	S, RI, MS	49.4 \pm 36.1	12.5–105.9	65.6 \pm 29.1	29.8–117.8	71.7 \pm 21.6	45.4–105.1
Methanol*	<1000	S	(218.8 \pm 23.0) ^a	180.3–239.2	(87.1 \pm 29.0) ^b	62.3–133.2	(79.0 \pm 15.8) ^b	60.3–102.5
<i>Higher alcohols</i>								
1-Propanol*	1025	S, MS	18.4 \pm 13.8	7.5–44.0	21.0 \pm 6.3	15.9–33.3	15.5 \pm 3.2	11.1–19.3
Isobutanol*	1100	S, RI, MS	(48.3 \pm 15.8) ^a	30.2–75.3	(18.5 \pm 3.3) ^b	13.5–23.2	(19.4 \pm 5.0) ^b	12.9–26.8
Isoamyl alcohol*	1206	S, RI, MS	(243.5 \pm 37.6) ^a	210.7–301.6	(159.6 \pm 27.1) ^b	124.9–192.6	(149.8 \pm 17.0) ^b	121.8–169.1
1-Octen-4-ol	1535	RI, MS	(1.0 \pm 0.5) ^a	0.6–1.8	(0.2 \pm 0.2) ^b	0.0–0.6	n.d.	–
2-Phenylethanol*	1893	S, RI, MS	(40.3 \pm 11.0) ^a	26.3–57.8	(26.3 \pm 7.6) ^b	16.8–34.9	(16.7 \pm 4.0) ^b	12.1–22.4
<i>Fatty acids</i>								
Isobutyric acid*	1554	S, RI, MS	1.5 \pm 0.9	1.0–3.2	1.2 \pm 0.3	0.8–1.4	1.0 \pm 0.5	0.5–1.9
Butyric acid*	1612	S, RI, MS	(0.9 \pm 0.2) ^b	0.6–1.2	(2.1 \pm 0.5) ^a	1.4–3.0	(1.8 \pm 0.3) ^a	1.5–2.2
Hexanoic acid*	1830	S, RI, MS	(1.1 \pm 0.5) ^b	0.4–1.7	(5.8 \pm 1.6) ^a	3.3–8.1	(5.1 \pm 1.5) ^a	3.4–7.5
Octanoic acid*	2043	S, RI, MS	(0.7 \pm 0.6) ^b	0.0–1.6	(6.2 \pm 1.2) ^a	4.1–7.6	(5.3 \pm 1.1) ^a	3.7–6.9
Decanoic acid*	2257	S, RI, MS	(0.1 \pm 0.0) ^b	0.1–0.2	(1.0 \pm 0.5) ^a	0.4–1.6	(1.7 \pm 0.7) ^a	0.8–2.6
Ethyl acetate*	<1000	S, RI	(75.1 \pm 8.6) ^a	61.8–85.4	(34.5 \pm 5.1) ^c	26.9–41.0	(44.6 \pm 6.4) ^b	36.2–53.2
<i>Ethyl esters of fatty acids</i>								
Ethyl butyrate	1030	S, RI, MS	(141.7 \pm 62.6) ^c	53.5–247.0	(436.7 \pm 73.8) ^b	390.7–580.5	(535.5 \pm 96.9) ^a	372.3–636.6
Ethyl 2-methylbutyrate	1049	S, RI, MS	(12.9 \pm 5.7) ^a	8.0–23.4	(4.5 \pm 2.1) ^b	1.7–6.9	(2.4 \pm 0.3) ^b	2.1–2.7
Ethyl 3-methylbutyrate	1065	S, RI, MS	(29.2 \pm 17.2) ^a	13.1–61.7	(8.9 \pm 2.8) ^{ab}	4.1–12.4	(4.1 \pm 0.9) ^b	2.8–5.0

Table 4. – continued

Compound	LRI	ID	$\gamma/(\mu\text{g/L})$					
			NMI		SMI-1		SMI	
			Mean value \pm S.D.	Range	Mean value \pm S.D.	Range	Mean value \pm S.D.	Range
Ethyl hexanoate	1236	S, RI, MS	(145.7 \pm 84.3) ^b	41.0–266.3	(546.6 \pm 80.4) ^a	443.0–679.7	(668.7 \pm 134.8) ^a	514.5–875.8
Ethyl 2-hydroxyisovalerate	1418	MS	1.5 \pm 0.8	0.7–2.9	n.d.	–	n.d.	–
Ethyl octanoate	1435	S, RI, MS	(128.1 \pm 74.2) ^b	42.1–227.0	(804.8 \pm 222.8) ^a	528.2–1124.2	(1131.4 \pm 269.7) ^a	829.5–1586.9
Ethyl decanoate	1637	S, RI, MS	(76.9 \pm 56.3) ^c	31.8–182.9	(493.1 \pm 173.3) ^b	289.0–764.2	(709.2 \pm 163.5) ^a	554.4–981.4
<i>Acetate esters</i>								
Isobutyl acetate	1009	S, RI, MS	(35.7 \pm 6.1) ^b	27.9–44.9	(40.0 \pm 9.0) ^b	26.2–53.1	(79.3 \pm 37.4) ^a	32.5–132.9
Isoamyl acetate*	1120	S, RI, MS	(0.1 \pm 0.1) ^c	0.0–0.3	(1.8 \pm 0.3) ^b	1.3–2.3	(3.9 \pm 1.4) ^a	2.1–5.3
Hexyl acetate	1272	S, RI, MS	(2.2 \pm 1.8) ^c	1.0–5.9	(49.0 \pm 11.9) ^b	30.2–62.0	(126.2 \pm 41.2) ^a	76.8–164.1
2-Phenethyl acetate*	1803	S, RI, MS	(0.2 \pm 0.1) ^c	0.0–0.3	(1.3 \pm 0.1) ^b	1.3–1.4	(2.0 \pm 0.4) ^a	1.4–2.4
<i>Esters of hydroxy and dicarboxylic acids</i>								
Ethyl lactate*	1341	S, RI, MS	(143.9 \pm 44.3) ^a	90.1–192.9	(20.7 \pm 7.9) ^b	8.1–32.0	(12.4 \pm 6.7) ^b	6.6–23.1
Isoamyl lactate	1561	RI, MS	(461.4 \pm 89.9) ^a	362.3–616.8	(134.3 \pm 25.3) ^b	108.7–175.3	(157.4 \pm 25.8) ^b	122.8–191.6
Diethyl malonate	1570	RI, MS	(38.5 \pm 32.9) ^a	0.0–79.9	(34.9 \pm 27.2) ^a	14.0–85.7	(1.6 \pm 3.9) ^b	0.0–9.5
Ethyl methyl succinate	1622	RI, MS	(123.3 \pm 31.7) ^a	89.8–158.7	(16.8 \pm 14.3) ^b	0.0–35.1	(1.9 \pm 4.7) ^b	0.0–11.4
Diethyl succinate*	1667	S, RI, MS	(13.9 \pm 4.1) ^a	8.3–18.2	(3.1 \pm 0.7) ^b	2.4–4.2	(0.8 \pm 0.8) ^b	0.2–2.4
Acetaldehyde*	<1000	S	45.5 \pm 32.0	15.1–98.6	92.4 \pm 61.4	27.6–169.6	40.9 \pm 4.6	34.9–46.9
<i>Volatile phenols</i>								
4-Ethylguaiaicol	2009	RI, MS	(445.7 \pm 392.2) ^a	25.5–916.5	(2.2 \pm 3.0) ^b	0.0–7.7	(6.3 \pm 15.5) ^b	0.0–38.0
4-Propylguaiaicol	2088	RI, MS	43.8 \pm 45.0	0.0–124.9	n.d.	–	n.d.	–
Eugenol	2148	S, RI, MS	13.0 \pm 12.2	3.9–37.0	n.d.	–	n.d.	–
4-Ethylphenol	2156	S, RI, MS	(119.0 \pm 127.8) ^a	3.3–332.0	(0.7 \pm 0.3) ^b	0.3–1.0	(0.8 \pm 0.6) ^b	0.2–1.9
4-Vinylguaiaicol	2175	RI, MS	117.1 \pm 73.4	44.3–208.1	284.9 \pm 209.9	130.5–698.7	449.2 \pm 494.9	124.5–1426.9
4-Vinylphenol	2366	RI, MS	83.4 \pm 102.3	0.0–250.1	84.3 \pm 98.4	0.0–270.3	190.8 \pm 211.4	0.0–572.6
<i>Benzenoids</i>								
Styrene	1253	RI, MS	(5.9 \pm 2.0) ^a	3.6–8.9	(1.7 \pm 0.6) ^b	0.8–2.5	(7.2 \pm 2.7) ^a	3.3–10.5
Benzaldehyde	1508	S, RI, MS	(4.2 \pm 2.7) ^{ab}	0.7–8.6	(0.5 \pm 0.4) ^b	0.0–1.1	(14.0 \pm 14.0) ^a	2.3–39.8
Methyl salicylate	1759	RI, MS	45.5 \pm 60.3	9.0–164.7	9.2 \pm 4.4	5.6–17.7	19.0 \pm 7.5	10.9–29.2
Ethyl benzeneacetate	1773	RI, MS	(9.1 \pm 2.3) ^a	6.1–12.6	(4.6 \pm 2.0) ^b	2.0–7.2	(1.8 \pm 0.7) ^c	1.0–3.0
2-Ethoxybenzyl alcohol	1922	MS	1.8 \pm 4.4	0.00–10.77	n.d.	–	n.d.	–
Ethyl cinnamate	2111	S, RI, MS	0.5 \pm 0.4	0.3–1.2	0.4 \pm 0.1	0.3–0.5	0.6 \pm 0.3	0.3–1.1
<i>Furans</i>								
Furfuryl ether	1284	MS	(248.0 \pm 260.7) ^a	19.9–677.1	1.1 \pm 1.2) ^b	0.0–3.0	n.d.	–
Furfural	1451	RI, MS	(2.1 \pm 0.6) ^a	1.3–2.6	(0.6 \pm 0.3) ^b	0.3–1.1	(0.1 \pm 0.1) ^b	0.0–0.3
2-Acetylfuran	1491	RI, MS	18.8 \pm 6.3	9.1–27.2	n.d.	–	n.d.	–
5-Methylfurfural	1556	RI, MS	(95.5 \pm 89.7) ^a	11.2–262.8	(1.0 \pm 1.2) ^b	0.0–2.5	(0.4 \pm 1.0) ^b	0.0–2.5
Ethyl 2-furoate	1609	RI, MS	29.0 \pm 17.8	6.6–53.4	18.7 \pm 12.2	5.8–40.3	10.2 \pm 8.3	0.0–23.5
Furfuryl alcohol	1644	RI, MS	59.8 \pm 42.8	4.3–109.1	n.d.	–	n.d.	–
<i>Lactones</i>								
γ -Butyrolactone	1606	RI, MS	(163.1 \pm 29.2) ^a	130.8–213.5	(57.9 \pm 11.3) ^b	41.6–71.3	(58.9 \pm 18.4) ^b	39.1–92.0
<i>trans</i> -Oak lactone	1868	RI, MS	164.7 \pm 110.1	77.8–376.9	n.d.	–	n.d.	–
<i>cis</i> -Oak lactone	1938	RI, MS	604.6 \pm 617.9	152.1–1793.2	n.d.	–	n.d.	–
γ -Nonalactone	2008	RI, MS	50.7 \pm 35.4	10.2–115.0	21.5 \pm 21.2	0.0–59.9	37.8 \pm 23.0	13.8–77.5
γ -Decalactone	2149	RI, MS	(55.8 \pm 51.7) ^a	11.1–143.1	(17.2 \pm 15.3) ^b	0.0–38.2	n.d.	–
γ -Undecalactone	2206	RI, MS	49.8 \pm 29.4	0.0–79.2	25.3 \pm 9.9	14.0–41.9	19.2 \pm 22.0	0.0–47.7
<i>Acetals</i>								
2,4,5-Trimethyl-1,3-dioxolane	<1000	RI, MS	8.0 \pm 19.5	0.0–47.7	n.d.	–	n.d.	–
1,1-Diethoxy-3-methylbutane	1065	RI, MS	21.8 \pm 24.1	0.0–53.8	n.d.	–	n.d.	–

Table 4. – continued

Compound	LRI	ID	$\gamma/(\mu\text{g/L})$					
			NMI		SMI-1		SMI	
			Mean value \pm S.D.	Range	Mean value \pm S.D.	Range	Mean value \pm S.D.	Range
2-Cyclohexyl-4,5-dimethyl-1,3-dioxolane I	1091	MS	3.5 \pm 8.6	0.0–21.0	n.d.	–	n.d.	–
2-Cyclohexyl-4,5-dimethyl-1,3-dioxolane II	1151	MS	4.8 \pm 11.7	0.0–28.7	n.d.	–	n.d.	–
<i>Sulphur compounds</i>								
Dihydro-2-methyl-3(2H)-thiophenone	1510	MS	n.d.	–	25.2 \pm 18.3	0.0–49.9	38.2 \pm 17.9	19.2–67.5
Methionol	1700	RI, MS	(114.3 \pm 53.2) ^a	76.7–217.9	(69.8 \pm 14.5) ^b	46.9–86.5	(67.5 \pm 14.4) ^b	46.7–86.7

LRI=linear retention index; identification of compounds (ID): S=retention time and mass spectrum consistent with those of the pure standard and with NIST05 mass spectra electronic library, RI=retention index consistent with that found in literature, MS=mass spectra consistent with those from NIST05 mass spectra electronic library or literature. The compounds for which pure standards were not available (without symbol S in the ID column) were quantified semi-quantitatively, and their concentrations expressed as equivalents of compounds with similar chemical structure assuming a response factor of 1.

Different superscript lowercase letters in a row present statistically significant differences between mean values at $p < 0.05$ obtained by one-way ANOVA and least significant difference (LSD) test. TDN=1,1,6-trimethyl-1,2-dihydronaphthalene, TPB=*trans*-1-(2,3,6-trimethylphenyl)buta-1,3-diene, S.D.=standard deviation ($N=6$), ^aconcentrations expressed in mg/L

rels as a result of hydrolysis of their esters. The same authors hypothesized the possibility of the oxidative deamination of residual free amino acid precursors during maturation, so it is possible that additional quantities of higher alcohols in the wines produced by prolonged maceration and maturation were formed in the barrel and the bottle by the Ehrlich mechanism.

Fatty acids

Evidence exists that maceration and fermentation of unfiltered must may result in the inhibition of fatty acid biosynthesis in yeast cells, which consequently have to assimilate them directly from must, resulting in their lower concentrations in wine (36). Higher oxygen availability in wooden barrels slows down the release of middle-chain fatty acids from yeast cell walls (6,37). A decrease in the concentration of straight-chain fatty acids during maturation was observed in several studies (35,38). It was tentatively explained as a result of the action of residual microorganisms (39), while the possibility of acid-catalysed autoreduction to yield aldehydes has also been considered (28). All of the mentioned probably contributed to the significantly reduced straight-chain fatty acid content in the wines produced by prolonged maceration and maturation in relation to standard wines (Table 4).

Volatile esters

Significantly lower ester concentrations were found in the wines produced by prolonged maceration and maturation in comparison with the standard wines produced with or without maturation, except for ethyl acetate (Table 4). Evaporation due to higher temperatures, cap manipulation, and barrel porosity during fermentation and maceration could certainly have had a negative effect. Furthermore, the wines produced by prolonged maceration were in contact with lees, which have the ability to absorb esters (6), as well as to release esterases that hydrolyse them (40). During maturation, straight-chain esters hydrolyse (35,41), which is most probably the main cause of

their decrease in the wines produced by prolonged maceration and maturation. Higher concentration of ethyl acetate in these wines is in line with higher levels common for aged wines (35).

Within the group of the wines produced by prolonged maceration and maturation ($N=6$), high positive correlation coefficients were found between the concentration of branched short-chain esters and the duration of maceration as well as maturation, ranging from 0.66 to 0.79, and from 0.79 to 0.98, respectively. The highest concentrations of ethyl lactate and diethyl succinate, the main malolactic fermentation esters, were found in these wines (Table 4). Esterification during maturation in order to reach equilibrium concentrations was probably the main cause.

Volatile phenols, benzenoids, furans, lactones and acetals

Higher level of ethylphenols and 4-propylphenol in the wines produced by prolonged maceration and maturation (Table 4) was probably a result of the activity of *Brettanomyces* and *Dekkera* yeast species which favour aerobic conditions during maturation in wood (41,42). Besides that, it was previously found that the formation of ethylphenols positively correlates with maceration duration and temperature (41,42). Among benzenoids, 2-ethoxybenzyl alcohol was found only in the wines produced by prolonged maceration and maturation (Table 4). Significantly higher concentration of furans in these wines (Table 4) is in agreement with previous findings, where furans were found to increase during maturation (35), which is favoured by the presence of lees and oxygen (6). The exclusive occurrence of the so-called oak lactones, together with the increased amounts of other identified lactones in the wines produced by prolonged maceration and maturation (Table 4), most probably originated from wood, as reported previously (42). Acetals are formed during fermentation, but their content increases most significantly during oxidative conditions of maturation

(35,42). In this work, four acetals were tentatively identified solely in these wines (Table 4). Among them, 2,4,5-trimethyl-1,3-dioxolane and 1,1-diethoxy-3-methylbutane have already been reported as important odourants in wines subjected to oxidation and in matured wines (42).

The impact of the key volatile aroma compounds

Aroma compounds with odour activity values (OAV) higher than 1, calculated on the basis of odour perception thresholds found in literature (43–47), and therefore with the direct impact on the aroma of the investigated wines, are listed in Table 5. Other compounds with OAV generally <1 are not presented.

Although fruity esters, such as ethyl octanoate and hexanoate, were the compounds with the strongest impact in all the investigated groups of wines, their OAVs were much higher in standard wine samples. In fact, these wines were characterised by a typical aroma profile

of young white wines dominated by fruity esters and fatty acids, complemented by a significant contribution of β -damascenone, which corresponds to previous findings on Malvazija istarska wines (26). Besides a notable impact of straight-chain esters and β -damascenone, the most influential odourants in the wines produced by prolonged maceration and maturation were the compounds that originate from wood (lactones and eugenol), compounds often associated with fermentative maceration (higher alcohols, linalool), compounds mostly or exclusively generated by species other than yeasts (ethyl acetate, 4-ethylguaiacol, ethyl lactate) or in (semi)aerobic conditions and at higher temperatures (ethyl acetate, ethyl 3-methylbutyrate, isoamyl alcohol), as well as compounds whose concentration typically increases with ageing (ethyl acetate, ethyl 3-methylbutyrate, isoamyl alcohol, TDN, ethyl lactate). The aroma of the wines produced by prolonged maceration and maturation could be less intense due to

Table 5. Odour perception thresholds, odour descriptions, and odour activity values (OAV) of the key volatile aroma compounds in Malvazija istarska wines produced by prolonged maceration during and after fermentation followed by maturation in wooden barrels (NMI), standard white wine production technology followed by maturation in bottle for one year (SMI-1), and standard white wine production technology without maturation (SMI)

Compound	Odour perception threshold	Odour description	OAV		
	$\mu\text{g/L}$		NMI	SMI-1	SMI
More characteristic of NMI wines*					
<i>cis</i> -Oak lactone	35 ^c	coconut, burnt wood	17.3 \pm 17.7	–	–
4-Ethylguaiacol	33 ^d	toasted bread, smoky, clove	13.5 \pm 11.9	0.1 \pm 0.1	0.2 \pm 0.5
Ethyl acetate	7500 ^c	varnish, fruity	10.0 \pm 1.1	4.6 \pm 0.7	6.0 \pm 0.9
Ethyl 3-methylbutyrate	3 ^d	berry, blackberry	9.7 \pm 5.7	3.0 \pm 0.9	1.4 \pm 0.1
Isoamyl alcohol	30000 ^d	solvent, alcoholic	8.1 \pm 1.3	5.3 \pm 0.9	5.0 \pm 0.6
TDN	2 ^b	petrol, kerosene	6.0 \pm 7.3	1.6 \pm 1.5	0.9 \pm 0.4
2-Phenylethanol	10000 ^c	rose	4.0 \pm 1.1	2.6 \pm 0.8	1.7 \pm 0.4
Linalool	15 ^a	floral, rose, sweet	2.5 \pm 1.6	1.4 \pm 0.3	1.0 \pm 0.3
Eugenol	6 ^d	cloves, cinnamon	2.2 \pm 2.0	–	–
γ -Nonalactone	30 ^d	coconut	1.7 \pm 1.2	0.7 \pm 0.7	1.3 \pm 0.8
Ethyl lactate	100000 ^c	buttery	1.4 \pm 0.4	0.2 \pm 0.1	0.1 \pm 0.1
<i>trans</i> -Oak lactone	122 ^c	coconut, burnt wood	1.4 \pm 0.9	–	–
More characteristic of SMI wines					
Ethyl octanoate	2 ^c	fruity, soapy	64.0 \pm 47.1	402.4 \pm 111.4	595.7 \pm 134.9
Ethyl hexanoate	5 ^c	fruity, green apple	29.1 \pm 16.9	109.3 \pm 16.1	133.7 \pm 27.0
Isoamyl acetate	30 ^d	banana	4.0 \pm 3.7	60.3 \pm 11.3	129.3 \pm 47.0
β -Damascenone	0.05 ^a	stewed apple, dried plum, honey, lilac	27.0 \pm 17.4	29.0 \pm 4.6	47.0 \pm 12.0
Ethyl butyrate	20 ^d	banana, pineapple	7.1 \pm 3.1	21.8 \pm 3.7	26.8 \pm 4.9
Hexanoic acid	420 ^d	cheese, fatty, rancid	2.6 \pm 1.1	13.7 \pm 3.8	12.2 \pm 3.6
Octanoic acid	500 ^d	fatty, rancid	1.4 \pm 1.1	12.4 \pm 2.4	10.6 \pm 2.2
Butyric acid	173 ^d	cheese, fatty, rancid	5.1 \pm 1.2	12.3 \pm 3.1	10.3 \pm 1.9
2-Phenethyl acetate	250 ^d	flowery, rose	0.6 \pm 0.3	5.3 \pm 0.2	7.9 \pm 1.5
Ethyl decanoate	200 ^c	fruity, soapy	0.4 \pm 0.3	2.5 \pm 0.9	3.6 \pm 0.8
Decanoic acid	1000 ^d	fatty, rancid	0.1 \pm 0.0	1.0 \pm 0.5	1.7 \pm 0.7

OAV values are expressed as mean \pm standard deviation ($N=6$), TDN=1,1,6-trimethyl-1,2-dihydronaphthalene

*Compounds more characteristic of NMI wines are listed in a decreasing order of the mean OAV values in NMI wines, while compounds more characteristic of SMI wines are listed in a decreasing order of the mean OAV values in SMI wines

Odour detection thresholds from references: ^a(43), ^b(44), ^c(45), ^d(46), ^e(47)

lower OAVs in general (Table 5). However, judging by the odour descriptors of the key components which introduced certain atypical nuances into the aroma profile of Malvazija istarska (Table 5), it is probably more complex than that of standard wines.

Conclusions

The present investigation showed that the application of an unconventional combination of oenological practices in the production of white wines, including prolonged maceration during and after fermentation followed by maturation in wooden barrels, results in unusual composition, which in all probability strongly reflects on their sensory properties. Such wines exhibited elevated levels of dry extract, volatile acidity, lactic acid and phenols, as well as higher colour intensities when compared to standard wines. In contrast to a typical aroma profile of young white wines dominated by fruity esters, unconventional wines contained increased concentrations of the majority of monoterpenes, C₁₃-norisoprenoids, methanol, higher alcohols, ethyl acetate, branched ethyl esters and esters from hydroxy and dicarboxylic acids, ethylphenols, furans, and acetals, while the content of straight-chain ethyl and acetate esters was drastically reduced. In a number of aspects, the composition of such wines resembled that of aged red wine. Besides the measured drift in colour intensity at 420 nm, possible repercussions on their organoleptic characteristics might include milder acidity due to the replacement of malic with lactic acid, fuller body with more intense bitterness and astringency originating from the increased levels of phenols, and a less pronounced but more complex aroma, which should all be confirmed by detailed and systematic sensory evaluation in further research. Judging by analogy with the existing knowledge, it can be roughly stated that maceration had a major impact on phenols, while volatile aromas were more significantly altered by maturation. The unconventional wine samples showed a solid level of homogeneity in the composition representing a distinctive wine type, despite different vintages and technological details. Accentuated antioxidant activity found offers the possibility to raise awareness about their value in a nutritional sense, and develop new marketing strategies until now mainly reserved for red wines.

References

- Darias-Martín JJ, Rodríguez O, Díaz E, Lamuela-Raventós RM. Effect of skin contact on the antioxidant phenolics in white wine. *Food Chem.* 2000;71:483–7. [http://dx.doi.org/10.1016/S0308-8146\(00\)00177-1](http://dx.doi.org/10.1016/S0308-8146(00)00177-1)
- Selli S, Cabaroglu T, Canbas A, Erten H, Nurgel C. Effect of skin contact on the aroma composition of the musts of *Vitis vinifera* L. cv. Muscat of Bornova and Narince grown in Turkey. *Food Chem.* 2003;81:341–7. [http://dx.doi.org/10.1016/S0308-8146\(02\)00428-4](http://dx.doi.org/10.1016/S0308-8146(02)00428-4)
- Gómez-Míguez MJ, González-Miret ML, Hernanz D, Fernández MA, Vicario IM, Heredia FJ. Effects of prefermentative skin contact conditions on colour and phenolic content of white wines. *J Food Eng.* 2007;78:238–45. <http://dx.doi.org/10.1016/j.jfoodeng.2005.09.021>
- Radeka S, Herjavec S, Peršurić Đ, Lukić I, Sladonja B. Effect of different maceration treatments on free and bound varietal aroma compounds in wine of *Vitis vinifera* L. cv. Malvazija istarska bijela. *Food Technol Biotechnol.* 2008;46:86–92.
- Di Maro E, Ercolini D, Coppola S. Yeast dynamics during spontaneous wine fermentation of the Catalanesca grape. *Int J Food Microbiol.* 2007;117:201–10. <http://dx.doi.org/10.1016/j.ijfoodmicro.2007.04.007>
- Liberatore MT, Pati S, Del Nobile MA, La Notte E. Aroma quality improvement of Chardonnay white wine by fermentation and ageing in barrique on lees. *Food Res Int.* 2010;43:996–1002. <http://dx.doi.org/10.1016/j.foodres.2010.01.007>
- Lomolino G, Zocca F, Spettoli P, Zanin G, Lante A. A preliminary study on changes in phenolic content during Bianchetta Trevigiana winemaking. *J Food Comp Anal.* 2010;23:575–9. <http://dx.doi.org/10.1016/j.jfca.2010.04.001>
- Baiano A, Varva G, De Gianni A, Viggiani I, Terracone C, Del Nobile MA. Influence of type of amphora on physico-chemical properties and antioxidant capacity of 'Falanghina' white wines. *Food Chem.* 2014;146:226–33. <http://dx.doi.org/10.1016/j.foodchem.2013.09.069>
- Díaz C, Laurie VF, Molina AM, Bücking M, Fischer R. Characterization of selected organic and mineral components of qvevri wines. *Am J Enol Vitic.* 2013;64:532–7. <http://dx.doi.org/10.5344/ajev.2013.13027>
- Ribéreau-Gayon P, Glories Y, Maujean A, Dubourdieu D, editors. *Handbook of enology: The chemistry of wine – stabilization and treatments*, vol. 2. Chichester, UK: John Wiley & Sons Ltd.; 2000. <http://dx.doi.org/10.1002/0470010398>
- Bavčar D, Baša Češnik H, Čuš F, Vanzo A, Gašperlin L, Košmerl T. Impact of alternative skin contact procedures on the aroma composition of white wine. *S Afr J Enol Vitic.* 2011;32:190–203.
- Di Lecce G, Boselli E, D'Ignazi G, Frega NG. Evolution of phenolics and glutathione in Verdicchio wine obtained with maceration under reductive conditions. *LWT – Food Sci Technol.* 2013;53:54–60. <http://dx.doi.org/10.1016/j.lwt.2013.03.006>
- EEC Regulation No. 2676/90. Community methods for the analysis of wine. *Off J.* 1990;L272:1–192.
- Di Stefano R, Cravero MC, Gentilizi N. Methods of determination of polyphenols in wine. *L'Enotecnico.* 1989;5:83–9 (in Italian).
- Brand-Williams W, Cuvelier ME, Berset C. Use of a free radical method to evaluate antioxidant activity. *LWT – Food Sci Technol.* 1995;28:25–30. [http://dx.doi.org/10.1016/S0023-6438\(95\)80008-5](http://dx.doi.org/10.1016/S0023-6438(95)80008-5)
- Noguerol-Pato R, González-Barreiro C, Cancho-Grande B, Simal-Gándara J. Quantitative determination and characterisation of the main odourants of Mencía monovarietal red wines. *Food Chem.* 2009;117:473–84. <http://dx.doi.org/10.1016/j.foodchem.2009.04.014>
- Kovačević Ganić K, Peršurić Đ, Komes D, Dragović-Uzelac V, Banović M, Piljac J. Antioxidant activity of Malvasia istriana grape juice and wine. *Ital J Food Sci.* 2006;18:187–97.
- Recamales AF, Sayago A, González-Miret ML, Hernanz D. The effect of time and storage conditions on the phenolic composition and colour of white wine. *Food Res Int.* 2006;39:220–9. <http://dx.doi.org/10.1016/j.foodres.2005.07.009>
- Ružić I, Škerget M, Knez Ž, Runje M. Phenolic content and antioxidant potential of macerated white wines. *Eur Food Res Technol.* 2011;233:465–72. <http://dx.doi.org/10.1007/s00217-011-1535-4>
- Hernández-Jiménez A, Kennedy JA, Bautista-Ortín AB, Gómez-Plaza E. Effect of ethanol on grape seed proanthocyan-

- din extraction. *Am J Enol Vitic.* 2012;63:57–61.
<http://dx.doi.org/10.5344/ajev.2011.11053>
21. Hernanz D, Gallo V, Recamales ÁF, Meléndez-Martémez AJ, González-Miret ML, Heredia, FJ. Effect of storage on the phenolic content, volatile composition and colour of white wines from the varieties Zalema and Colombard, *Food Chem.* 2009;113:530–7.
<http://dx.doi.org/10.1016/j.foodchem.2008.07.096>
 22. Ortega AF, Lopez-Toledano A, Mayen M, Merida J, Medina M. Changes in color and phenolic compounds during oxidative aging of sherry white wine. *J Food Sci.* 2003;68:2461–8.
<http://dx.doi.org/10.1111/j.1365-2621.2003.tb07046.x>
 23. Ortega AF, Mayen M, Medina M. Study of colour and phenolic compounds in two models of oxidative ageing for sherry type white wines. *Food Control.* 2008;19:949–56.
<http://dx.doi.org/10.1016/j.foodcont.2007.09.008>
 24. Hernanz D, Recamales ÁF, González-Miret ML, Gomez-Miguez MJ, Vicario IM, Heredia FJ. Phenolic composition of white wines with a prefermentative maceration at experimental and industrial scale. *J Food Eng.* 2007;80:327–35.
<http://dx.doi.org/10.1016/j.jfoodeng.2006.06.006>
 25. Stratil P, Kubáň V, Fojtová J. Comparison of the phenolic content and total antioxidant activity in wines as determined by spectrophotometric methods. *Czech J Food Sci.* 2008;26: 242–53.
 26. Lukić I, Plavša T, Sladonja B, Radeka S, Peršurić Đ. Aroma compounds as markers of wine quality in the case of Malvazija istarska young wine. *J Food Quality.* 2008;31:717–35.
<http://dx.doi.org/10.1111/j.1745-4557.2008.00230.x>
 27. Zoecklein BW, Marcy JE, Jasinski Y. Effect of fermentation, storage sur lie or post-fermentation thermal processing on White Riesling (*Vitis vinifera* L.) glycoconjugates. *Am J Enol Vitic.* 1997;48:397–402.
 28. Ferreira V, Escudero A, Fernández P, Cacho JF. Changes in the profile of volatile compounds in wines stored under oxygen and their relationship with the browning process. *Z Lebensm Unters Forsch A.* 1997;205:392–6.
<http://dx.doi.org/10.1007/s002170050187>
 29. Oliveira JM, Oliveira P, Baumes RL, Maia O. Changes in aromatic characteristics of Loureiro and Alvarinho wines during maturation. *J Food Compos Anal.* 2008;21:695–707.
<http://dx.doi.org/10.1016/j.jfca.2008.08.002>
 30. Marais J. Terpene concentrations and wine quality of *Vitis vinifera* L. cv. Gewurtztraminer as affected by grape maturity and cellar practices. *Vitis.* 1987;26:231–45.
 31. Loscos N, Hernández-Orte P, Cacho J, Ferreira V. Evolution of the aroma composition of wines supplemented with grape flavor precursors from different varieties during accelerated wine ageing. *Food Chem.* 2010;120:205–16.
<http://dx.doi.org/10.1016/j.foodchem.2009.10.008>
 32. Silva Ferreira AC, de Pinho PG. Nor-isoprenoids profile during port wine ageing – influence of some technological parameters. *Anal Chim Acta.* 2004;513:169–76.
<http://dx.doi.org/10.1016/j.aca.2003.12.027>
 33. OIV-MA-C1-01: R2011. Maximum acceptable limits of various substances contained in wine. Compendium of International Methods of Analysis of Wines and Must. International Organization of Vine and Wine (OIV); 2011.
 34. Vidrih R, Hribar J. Synthesis of higher alcohols during cider processing. *Food Chem.* 1999;67:287–94.
[http://dx.doi.org/10.1016/S0308-8146\(99\)00136-3](http://dx.doi.org/10.1016/S0308-8146(99)00136-3)
 35. Câmara JS, Alves MA, Marques JC. Changes in volatile composition of Madeira wines during their oxidative ageing. *Anal Chim Acta.* 2006;563:188–97.
<http://dx.doi.org/10.1016/j.aca.2005.10.031>
 36. Valero E, Millán C, Ortega JM. Influence of pre-fermentative treatment on the fatty acid content of *Saccharomyces cerevisiae* (M330-9) during alcoholic fermentation of grape must. *J Biosci Bioeng.* 2001;91:117–22.
[http://dx.doi.org/10.1016/S1389-1723\(01\)80052-9](http://dx.doi.org/10.1016/S1389-1723(01)80052-9)
 37. Torija MJ, Beltran G, Novo M, Poblet M, Guillamón JM, Mas A, Rozès N. Effects of fermentation temperature and *Saccharomyces* species on the cell fatty acid composition and presence of volatile compounds in wine. *Int J Food Microbiol.* 2003;85:127–36.
[http://dx.doi.org/10.1016/S0168-1605\(02\)00506-8](http://dx.doi.org/10.1016/S0168-1605(02)00506-8)
 38. Hernández-Orte P, Lapeña AC, Escudero A, Astrain J, Baron C, Pardo I, et al. Effect of micro-oxygenation on the evolution of aromatic compounds in wines: malolactic fermentation and ageing in wood. *LWT – Food Sci Technol.* 2009;42:391–401.
<http://dx.doi.org/10.1016/j.lwt.2008.05.020>
 39. Ancín C, Ayestarán B, García A, Garrido J. Evolution of fatty acid contents in Garnacha and Viura musts during fermentation and the aging of wine. *Z Lebensm Unters Forsch A.* 1998;206:144–7.
<http://dx.doi.org/10.1007/s002170050230>
 40. Ancín-Azpilicueta C, González-Marco A, Jiménez-Moreno N. Evolution of esters in aged Chardonnay wines obtained with different vinification methods. *J Sci Food Agric.* 2009;89: 2446–51.
<http://dx.doi.org/10.1002/jsfa.3744>
 41. Garde Cerdán T, Rodríguez Mozaz S, Ancín Azpilicueta C. Volatile composition of aged wine in used barrels of French oak and of American oak. *Food Res Int.* 2002;35:603–10.
[http://dx.doi.org/10.1016/S0963-9969\(01\)00151-X](http://dx.doi.org/10.1016/S0963-9969(01)00151-X)
 42. Perestrelo R, Barros AS, Câmara JS, Rocha SM. In-depth search focused on furans, lactones, volatile phenols, and acetals as potential age markers of Madeira wines by comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometry combined with solid phase microextraction. *J Agric Food Chem.* 2011;59:3186–204.
<http://dx.doi.org/10.1021/jf104219t>
 43. Guth H. Quantitation and sensory studies of character impact odorants of different white wine varieties. *J Agric Food Chem.* 1997;45:3027–32.
<http://dx.doi.org/10.1021/jf970280a>
 44. Sacks GL, Gates MJ, Ferry FX, Lavin EH, Kurtz AJ, Acree TE. Sensory threshold of 1,1,6-trimethyl-1,2-dihydronaphthalene (TDN) and concentrations in young Riesling and non-Riesling wines. *J Agric Food Chem.* 2012;60:2998–3004.
<http://dx.doi.org/10.1021/jf205203b>
 45. Moreno JA, Zea L, Moyano L, Medina M. Aroma compounds as markers of the changes in sherry wines subjected to biological ageing. *Food Control* 2005;16:333–8.
<http://dx.doi.org/10.1016/j.foodcont.2004.03.013>
 46. Ferreira V, López R, Cacho JF. Quantitative determination of the odorants of young red wines from different grape varieties. *J Sci Food Agric.* 2000;80:1659–67.
[http://dx.doi.org/10.1002/1097-0010\(20000901\)80:11<1659::AID-JSFA693>3.0.CO;2-6](http://dx.doi.org/10.1002/1097-0010(20000901)80:11<1659::AID-JSFA693>3.0.CO;2-6)
 47. Zea L, Moyano L, Moreno JA, Medina M. Aroma series as fingerprints for biological ageing in fino sherry-type wines. *J Sci Food Agric.* 2007;87:2319–26.
<http://dx.doi.org/10.1002/jsfa.2992>