

# STATIONARY SOLUTIONS FOR THE CAHN-HILLIARD EQUATION

JUNCHENG WEI AND MATTHIAS WINTER

ABSTRACT. We study the Cahn-Hilliard equation in a bounded domain without any symmetry assumptions. We assume that the mean curvature of the boundary has a nongenerate critical point. Then we show that there exists a spike-like stationary solution whose global maximum lies on the boundary. Our method is based on Lyapunov-Schmidt reduction and the Brouwer fixed-point theorem.

RÉSUMÉ. Nous étudions l'équation de Cahn et Hilliard dans une domaine ouverte sans supposer aucunes conditions de symétrie pour la domaine. Nous supposons que la courbature moyenne sur la frontière a un point critique non dégénéré. Nous montrons qu'il existe une solution stationnaire avec un pic qui atteint son maximum sur la frontière de la domaine. Notre méthode utilise la réduction de Lyapunov et Schmidt et le théorème du point fixe de Brouwer. (Titre: Solutions stationnaires pour l'équation de Cahn et Hilliard).

## 1. INTRODUCTION

The Cahn-Hilliard equation [5] is an accepted macroscopic field-theoretical model of processes such as phase separation in a binary alloy. In its original form it is derived from a Helmholtz free energy

$$E(u) = \int_{\Omega} [F(u(x)) + \frac{1}{2}\epsilon^2|\nabla u(x)|^2]dx$$

where  $\Omega$  is the region occupied by the body,  $u(x)$  is a conserved order parameter representing for example the concentration of one of the components, and  $F(u)$  is the free energy density which has a double well structure at low temperatures (see Figure 1). The most commonly used model is for  $F(u) = (1 - u^2)^2$ .

The constant  $\epsilon$  is proportional to the range of intermolecular forces and the gradient term is a contribution to the free energy coming from spatial

---

1991 *Mathematics Subject Classification.* Primary 35B40, 35B45; Secondary 35J40.

*Key words and phrases.* Phase Transition, Nonlinear Elliptic Equations.

fluctuations of the order parameter. Moreover the mass  $\bar{m} = \frac{1}{|\Omega|} \int_{\Omega} u dx$  is constant. Thus a stationary solution of  $E(u)$  under  $\bar{m} = \frac{1}{|\Omega|} \int_{\Omega} u dx$  takes the following form

$$\begin{cases} \epsilon^2 \Delta u - f(u) = \sigma_{\epsilon} & \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega, \\ \int_{\Omega} u = \bar{m} |\Omega| \end{cases} \quad (1.1)$$

where  $f(u) = F'(u)$  (see Figure 2) and  $\sigma_{\epsilon}$  is a constant.

There have been numerous studies of the Cahn-Hilliard equation. The global minimizer of  $E(u)$  has a transition layer. More precisely there exists an open set  $\Gamma \subset \Omega$  such that  $u_{\epsilon}$  is a global minimizer then  $u_{\epsilon} \rightarrow 1$  on  $\Omega \setminus \bar{\Gamma}$ ,  $u_{\epsilon} \rightarrow -1$  on  $\Gamma$  and  $\partial\Gamma \cap \bar{\Omega}$  is a minimal surface and has constant mean curvature, see [16]. The dynamics of the interface have been studied extensively, see for example [2], [3], [23]. Also local minimizers of  $E(u)$  have been studied and their transition layer structure has been established in [6] and [13]. In particular, Chen and Kowalczyk in [6] used boundary mean curvature to construct local minimizers (therefore transition layer solutions) for equation (1.1).

In this paper we are concerned with solutions of (1.1) with spike layers. In the one dimensional case, Bates and Fife [4] studied nucleation phenomena for the Cahn-Hilliard equation and proved the existence of three monotone nondecreasing stationary solutions when  $\bar{m}$  is in the metastable region ( $\sqrt{1/3} < \bar{m} < 1$ ), (a) the constant solution  $u \equiv \bar{m}$ , (b) a boundary spike layer solution where the layer is located at the left-hand endpoint, (c) a transition layer solution with a layer in the interior of the material.

Motivated by the results of [4], we shall construct a boundary spike layer solution to (1.1) for  $\epsilon \ll 1$  in the higher dimensional case when  $\bar{m}$  is in the metastable region.

The existence of spike layer solutions as well as the location and the profile of the peaks for other problems arising in various models such as chemotaxis, pattern formation, chemical reactor theory, etc. have been studied by Lin, Ni, Pan, and Takagi [14, 17, 18, 19] for the Neumann problem and by Ni and Wei [20] for the Dirichlet problem. However, they do not have the volume constraint and the nonlinearity is simpler than here. To our knowledge the

present paper is the first to establish this kind of results for the Cahn-Hilliard equation in higher dimensions without any symmetry assumptions on  $\Omega$ .

Naturally these stationary solutions are essential for the understanding of the dynamics of the corresponding evolution process. While Bates and Fife [4] prove some results in this direction for the one dimensional case these questions are open for higher dimensions.

In [11] in the one dimensional case the number of all stationary solutions is counted by arguments using transversality.

First we make the following transformation.

$$\begin{aligned} v &= \bar{m} - u, \\ g(v) &= -f(\bar{m}) + f(\bar{m} - v). \end{aligned}$$

Rewrite

$$g'(0) = -m, \quad g(v) = -mv + h(v).$$

Then equation (1.1) becomes

$$\begin{cases} \epsilon^2 \Delta v - mv + h(v) - \frac{1}{|\Omega|} \int_{\Omega} h(v) = 0 & \text{in } \Omega, \\ \frac{\partial v}{\partial \nu} = 0 & \text{on } \partial\Omega. \end{cases} \quad (1.2)$$

(Figure 3 shows qualitatively how the graph of  $g$  looks like.)

To accommodate more general  $g$  we assume that

- (1)  $g'(0) < 0$ ,  $g(0) = 0$ ,  $g \in C^3(R, R)$ .
- (2)  $g(v)$  has only two zeroes for  $v > 0$ ,  $0 < a_1 < a_2$  and

$$\int_0^{a_2} g(s) ds > 0, \quad g'(a_2) < 0.$$

- (3) The function  $v \rightarrow \frac{g(v)}{v-v_0}$  is nonincreasing in the interval  $(v_0, a_2)$  where  $v_0$  is defined as the unique number in  $(a_1, a_2)$  such that  $\int_0^{v_0} g(s) ds = 0$ .

- (4)  $|h'(v)|, |h''(v)| \leq C$  for any  $v$ .

**Remarks:**

(1) Condition (3) can be weakened further. For example, the conditions in [7] will be enough since we just need the uniqueness and weak nondegeneracy of the ground state solutions of (1.3).

(2) Condition (4) is not a restriction physically since in the physical world  $v$  is always bounded. Hence we can modify  $h$  near infinity so that  $h$  satisfies (4).

It is easy to see that for  $f(u) = -2u(1 - u^2)$  conditions (1), (2), (3), and (4) are satisfied. Our main result can be stated as follows.

**Theorem 1.1.** *Let  $\Omega$  be a bounded smooth domain in  $R^N (N \geq 2)$  and  $P_0 \in \partial\Omega$  be such that  $\nabla_{P_0} H(P_0) = 0$  and  $(\nabla_{P_0}^2 H(P_0)) \neq 0$  where  $H(P_0)$  is the mean curvature of  $P_0 \in \partial\Omega$  and  $\nabla_{P_0}$  is the tangential derivative at  $P_0$ . Then for  $\epsilon \ll 1$  there exists a solution  $v_\epsilon$  of (1.2) such that  $v_\epsilon \rightarrow 0$  in  $C_{loc}^1(\bar{\Omega} \setminus P_0)$ ,  $v_\epsilon$  has only one local (hence global) maximum point  $P_\epsilon$  and  $P_\epsilon \in \partial\Omega, P_\epsilon \rightarrow P_0, v_\epsilon(P_\epsilon) \rightarrow V(0) > 0$ . Moreover*

$$\epsilon^{-N} \left\{ \int_{\Omega} \epsilon^2 \left| \nabla v_\epsilon - \nabla V \left( \frac{x - P_\epsilon}{\epsilon} \right) \right|^2 + \int_{\Omega} \left| v_\epsilon - V \left( \frac{x - P_\epsilon}{\epsilon} \right) \right|^2 \right\} \rightarrow 0$$

as  $\epsilon \rightarrow 0$  where  $V(y)$  is the unique solution of

$$\begin{cases} \Delta V - mV + h(V) = 0, \\ V(0) = \max_{y \in R^N} V(y), V > 0, \\ V(y) \rightarrow 0 \text{ at } \infty. \end{cases} \quad (1.3)$$

(By the results of [9] and [24], (1.3) has a unique radial solution).

The method of our construction evolves from that of [8], [21] and [22] on the semi-classical (i.e. for small parameter  $h$ ) solution of the nonlinear Schrödinger equation

$$\frac{h^2}{2} \Delta U - (V - E)U + U^p = 0 \quad (1.4)$$

in  $R^N$  where  $V$  is a potential function and  $E$  is a real constant. The method of Lyapunov-Schmidt reduction was used in [8], [21] and [22] to construct solutions of (1.4) close to nondegenerate critical points of  $V$  for  $h$  sufficiently small.

Following the strategy of [8], [21] and [22] we shall construct a solution  $v_\epsilon$  of (1.2) with maximum near a given nondegenerate critical point of the mean curvature  $P_0$  on  $\partial\Omega$ . Heuristically we rescale (1.2) to obtain

$$\begin{cases} \Delta u_\epsilon - m u_\epsilon + h(u_\epsilon) - \frac{1}{|\Omega_{\epsilon,P}|} \int_{\Omega_{\epsilon,P}} h(u_\epsilon) = 0 & \text{in } \Omega_{\epsilon,P}, \\ \frac{\partial u_\epsilon}{\partial \nu_\epsilon} = 0 & \text{on } \partial\Omega_{\epsilon,P} \end{cases} \quad (1.5)$$

where  $u_\epsilon(z) = v_\epsilon(x)$  for  $z = (x - P)/\epsilon, z \in \Omega_{\epsilon,P}$  and  $\Omega_{\epsilon,P} = \{z \in R^N \mid \epsilon z + P \in \Omega\}$  and  $\nu_\epsilon$  is the unit outer normal to  $\partial\Omega_{\epsilon,P}$ .

Taking the limit  $\epsilon \rightarrow 0$ ,  $u_\epsilon \rightarrow V$  where  $V$  is the unique solution of

$$\begin{cases} \Delta w - mw + h(w) = 0 & \text{in } R_+^N, \\ w > 0 & \text{in } R_+^N, \\ \frac{\partial w}{\partial y_N} = 0 & \text{on } R^{N-1} \times \{0\} \end{cases} \quad (1.6)$$

with  $V(0) = \max_{R_+^N} V$ . Therefore the ground state solution  $V$  restricted to  $R_+^N$  can be an approximate solution for  $u_\epsilon$ . Since the linearized problem arising from (1.6) has the  $(N-1)$ -dimensional kernel  $\text{span}\{\frac{\partial V}{\partial y_1}, \dots, \frac{\partial V}{\partial y_{N-1}}\}$  we first “solve” (1.6) up to this kernel and then use the nondegeneracy of  $H(P_0)$  to take care of the kernel separately.

The paper is organized as follows. Notation, preliminaries and some useful estimates are explained in Section 2. Section 3 contains the setup of our problem and we solve (1.2) up to approximate kernel and cokernel, respectively. Finally in Section 4 we solve the reduced problem.

**Acknowledgement.** The first author would like to thank Professor Wei-Ming Ni for his enlightening discussions. Part of the work is inspired by some related work by Professor Wei-Ming Ni and Professor Y.-G. Oh. This research was done while the second author visited the Department of Mathematics, The Chinese University of Hong Kong. It is supported by a Direct Grant from The Chinese University of Hong Kong and by a grant of the European Union (contract ERBCHBICT930744).

## 2. TECHNICAL ANALYSIS

In this section we introduce a projection and derive some useful estimates.

Throughout the paper we shall use the letter  $C$  to denote a generic positive constant which may vary from term to term. We denote  $R_+^N = \{(x', x_N) | x_N > 0\}$ . Let  $V$  be the unique solution of (1.3).

Let  $P \in \partial\Omega$ . We can define a diffeomorphism straightening the boundary in a neighborhood of  $P$ . After rotation of the coordinate system we may assume that the inward normal to  $\partial\Omega$  at  $P$  is pointing in the direction of the positive  $x_N$ -axis. Denote  $x' = (x_1, \dots, x_{N-1})$ ,  $B'(R_0) = \{x' \in R^{N-1} | |x'| < R_0\}$  and  $\Omega_1 = \Omega \cap B(P, R_0) = \{(x', x_N) \in B(P, R_0) | x_N - P_N > \rho(x' - P')\}$  where  $B(P, R_0) = \{x \in R^N | |x - P| < R_0\}$ . Then, since  $\partial\Omega$  is smooth, we can find a constant  $R_0 > 0$  such that  $\partial\Omega \cap \overline{\Omega_1}$  can be represented by the

graph of a smooth function  $\rho_P : B'(R_0) \rightarrow R$  where  $\rho_P(0) = 0, \nabla \rho_P(0) = 0$ . From now on we omit the use of  $P$  in  $\rho_P$  and write  $\rho$  instead if this can be done without causing confusion. The sum of the principal curvatures of  $\partial\Omega$  at  $P$  is  $H(P) = \sum_{i=1}^{N-1} \rho_{ii}(0)$  where

$$\rho_i = \frac{\partial \rho}{\partial x_i}, \quad i = 1, \dots, N-1$$

and higher derivatives will be defined in the same way. By Taylor expansion we have

$$\begin{aligned} \rho(x' - P') &= \frac{1}{2} \sum_{i,j=1}^{N-1} \rho_{ij}(0)(x_i - P_i)(x_j - P_j) \\ &+ \frac{1}{6} \sum_{i,j,k=1}^{N-1} \rho_{ijk}(0)(x_i - P_i)(x_j - P_j)(x_k - P_k) + \mathcal{O}(|x' - P'|^4) \end{aligned}$$

In the following we use  $\rho_\alpha$  to denote the multiple differentiation  $\frac{\partial^{|\alpha|} \rho}{\partial x^\alpha}$ , where  $\alpha$  is a multiple index.

For  $x \in \partial\Omega$ , let  $\nu(x)$  denote the unit outward normal at  $x$  and  $\partial/\partial\nu$  the normal derivative. Let  $(\tau_1(x), \dots, \tau_{N-1}(x))$  denote  $(N-1)$  linearly independent tangential vectors and  $(\frac{\partial}{\partial\tau_1}, \dots, \frac{\partial}{\partial\tau_{N-1}})$  the tangential derivatives.

In our coordinate system, for  $x \in \omega_1 := \partial\Omega \cap B(P, R_0)$ , we have

$$\begin{aligned} \nu(x) &= \frac{1}{\sqrt{1 + |\nabla_{x'} \rho|^2}} (\nabla_{x'} \rho, -1), \\ \frac{\partial}{\partial\nu} &= \frac{1}{\sqrt{1 + |\nabla_{x'} \rho|^2}} \left\{ \sum_{j=1}^{N-1} \rho_j \frac{\partial}{\partial x_j} - \frac{\partial}{\partial x_N} \right\} \Big|_{x_N - P_N = \rho(x' - P')}, \\ \tau_i(x) &= (0, \dots, 1, \dots, 0, \rho_i(x')), \\ \frac{\partial}{\partial\tau_i} &= \frac{1}{\sqrt{1 + |\nabla_{x'} \rho|^2}} \left\{ \frac{\partial}{\partial x_i} + \rho_i \frac{\partial}{\partial x_N} \right\} \Big|_{x_N - P_N = \rho(x' - P')}. \end{aligned}$$

For a smooth bounded domain  $U$  we now introduce a projection  $P_U$  of  $H^2(U)$  onto  $\{v \in H^2(U) | \partial v / \partial \nu = 0 \text{ at } \partial U\}$  as follows: For  $v \in H^2(U)$  let  $w = P_U v$  be the unique solution of the boundary value problem

$$\begin{cases} \Delta w - mw + h(v) = 0 & \text{in } U, \\ \frac{\partial w}{\partial \nu} = 0 & \text{on } \partial U. \end{cases}$$

Let  $h_{\epsilon, P}(x) = V\left(\frac{x-P}{\epsilon}\right) - P_{\Omega_{\epsilon, P}} V\left(\frac{x-P}{\epsilon}\right)$  where

$$\Omega_{\epsilon, P} = \{z \in R^n | P + \epsilon z \in \Omega\}.$$

Then  $h_{\epsilon, P}$  satisfies

$$\begin{cases} \epsilon^2 \Delta v - mv = 0 & \text{in } \Omega, \\ \frac{\partial v}{\partial \nu} = \frac{\partial V}{\partial \nu} & \text{on } \partial \Omega. \end{cases} \quad (2.1)$$

We denote

$$\|v\|_\epsilon^2 = \epsilon^{-N} \int_\Omega [\epsilon^2 |\nabla v|^2 + mv^2].$$

For  $x \in \Omega_1$  set now

$$\begin{cases} \epsilon y' = x' - P', \\ \epsilon y_N = x_N - P_N - \rho(x' - P'). \end{cases} \quad (2.2)$$

Furthermore, for  $x \in \Omega_1$  we introduce the transformation

$$\begin{cases} T_i(x') = x_i, & i = 1, \dots, N-1 \\ T_N(x') = x_N - P_N - \rho(x' - P'). \end{cases} \quad (2.3)$$

Note that then

$$y = \frac{1}{\epsilon} T(x).$$

The Laplace operator and the boundary derivative operator become

$$\epsilon^2 \Delta_x = \Delta_y + |\nabla_{x'} \rho|^2 \frac{\partial^2}{\partial y_N^2} - 2 \sum_{i=1}^{N-1} \rho_i \frac{\partial^2}{\partial y_i \partial y_N} - \epsilon \Delta_{x'} \rho \frac{\partial}{\partial y_N} \quad \text{for } x \in \Omega_1, \quad (2.4)$$

$$\sqrt{1 + |\nabla_{x'} \rho|^2} \frac{\partial}{\partial \nu_x} = \frac{1}{\epsilon} \left\{ \sum_{j=1}^{N-1} \rho_j \frac{\partial}{\partial y_j} - (1 + |\nabla_{x'} \rho|^2) \frac{\partial}{\partial y_N} \right\} \quad \text{for } x \in \omega_1. \quad (2.5)$$

Let  $v_1$  be the unique solution of

$$\begin{cases} \Delta v - mv = 0 & \text{in } R_+^N, \\ \frac{\partial v}{\partial y_N} = -\frac{V'}{|y|} \frac{1}{2} \sum_{i,j=1}^{N-1} \rho_{ij}(0) y_i y_j & \text{on } \partial R_+^N \end{cases} \quad (2.6)$$

where  $V'$  is the radial derivative of  $V$ , i.e.  $V' = V_r(r)$ , and  $r = \left| \frac{x-P}{\epsilon} \right|$ . Let  $v_2$  be the unique solution of

$$\begin{cases} \Delta v - mv - 2 \sum_{i,j=1}^{N-1} \rho_{ij}(0) y_i \frac{\partial^2 v_1}{\partial y_j \partial y_N} = 0 & \text{in } R_+^N, \\ \frac{\partial v}{\partial y_N} = \sum_{i,j=1}^{N-1} \rho_{ij}(0) y_i \frac{\partial v_1}{\partial y_j} & \text{on } \partial R_+^N. \end{cases} \quad (2.7)$$

Let  $v_3$  be the unique solution of

$$\begin{cases} \Delta v - mv = 0 & \text{in } R_+^N, \\ \frac{\partial v}{\partial y_N} = -\frac{V'}{|y|} \frac{1}{3} \sum_{i,j,k=1}^{N-1} \rho_{ijk}(0) y_i y_j y_k & \text{on } \partial R_+^N. \end{cases} \quad (2.8)$$

Note that  $v_1, v_2$  are even functions in  $y' = (y_1, \dots, y_{N-1})$  and  $v_3$  is an odd function in  $y' = (y_1, \dots, y_{N-1})$  (i.e.  $v_1(y', y_N) = v_1(-y', y_N), v_3(y', y_N) = -v_3(-y', y_N)$ ). Moreover, it is easy to see that  $|v_1|, |v_2|, |v_3| \leq C e^{-\mu|y|}$  for

some  $0 < \mu < \sqrt{m}$ . Let  $\chi(x)$  be a smooth cutoff function such that  $\chi(x) = 1$ ,  $x \in B(0, R_0 - \delta)$  and  $\chi(x) = 0$  for  $x \in B(0, R_0)^C$  (for a positive number  $\delta$ .) Set

$$h_{\epsilon, P}(x) = \epsilon v_1(y)\chi(x - P) + \epsilon^2(v_2(y)\chi(x - P) + v_3(y)\chi(x - P)) + \epsilon^3\Psi_{\epsilon, P}(x).$$

Then we have

**Proposition 2.1.**

$$\|\Psi_{\epsilon, P}\|_{\epsilon} \leq C.$$

To prove Proposition 2.1, we begin with

**Lemma 2.2.** *Let  $u$  be a solution of*

$$\begin{cases} \epsilon^2 \Delta u - mu + f = 0 & \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} = g & \text{on } \partial\Omega, \end{cases}$$

*Assume that  $\int_{\Omega} |f|^2 \leq C\epsilon^N$ ,  $\int_{\partial\Omega} |g|^2 \leq C\epsilon^{N-1}$ . Then*

$$\|u\|_{\epsilon} \leq C.$$

*Proof:* Multiplying the equation by  $u$ , we have

$$\epsilon^2 \int_{\Omega} |\nabla u|^2 + m \int_{\Omega} u^2 = \int_{\Omega} fu + \epsilon^2 \int_{\partial\Omega} gu.$$

Lemma 2.2 follows easily by the following interpolation inequality (the proof of it is delayed to Appendix A),

$$\|u\|_{L^2(\partial\Omega_{\epsilon, P})} \leq C\|u\|_{\epsilon}$$

where  $\Omega_{\epsilon, P} = \{z \mid x = P + \epsilon z \in \Omega\}$  for a fixed  $P \in \partial\Omega$ .  $\square$

*Proof of Proposition 2.1:* We first compute the equation for  $\Psi_{\epsilon, P}(x)$ :

$$\begin{aligned} & -\epsilon^2 \Delta \Psi_{\epsilon, P}(x) + m\Psi_{\epsilon, P}(x) \\ &= \frac{1}{\epsilon^3} \left[ \epsilon^2 \left\{ \Delta_x (\epsilon v_1 \chi + \epsilon^2 (v_2 \chi + v_3 \chi)) \right\} - m\epsilon v_1 \chi - m\epsilon^2 v_2 \chi - m\epsilon^2 v_3 \chi \right] \\ &= \frac{1}{\epsilon^2} \left[ \left\{ \Delta_y v_1 + |\nabla_{x'} \rho|^2 \frac{\partial^2 v_1}{\partial y_N^2} - 2 \sum_{i=1}^{N-1} \rho_i \frac{\partial^2 v_1}{\partial y_i \partial y_N} - \epsilon \Delta_{x'} \rho \frac{\partial v_1}{\partial y_N} - mv_1 \right\} \chi \right. \\ & \quad + \epsilon \left\{ \Delta_y v_2 + |\nabla_{x'} \rho|^2 \frac{\partial^2 v_2}{\partial y_N^2} - 2 \sum_{i=1}^{N-1} \rho_i \frac{\partial^2 v_2}{\partial y_i \partial y_N} - \epsilon \Delta_{x'} \rho \frac{\partial v_2}{\partial y_N} - mv_2 \right\} \chi \\ & \quad \left. + \epsilon \left\{ \Delta_y v_3 + |\nabla_{x'} \rho|^2 \frac{\partial^2 v_3}{\partial y_N^2} - 2 \sum_{i=1}^{N-1} \rho_i \frac{\partial^2 v_3}{\partial y_i \partial y_N} - \epsilon \Delta_{x'} \rho \frac{\partial v_3}{\partial y_N} - mv_3 \right\} \chi \right] \end{aligned}$$



$$\begin{aligned}
& \left. + E_\epsilon(\chi) \right] \\
& = \frac{1}{\epsilon^2} \left[ \chi \left\{ |\nabla \rho|^2 \frac{\partial^2 v_1}{\partial y_N^2} - \epsilon \Delta \rho \frac{\partial v_1}{\partial y_N} - 2 \sum_{i,j=1}^{N-1} (\rho_i - \epsilon \rho_{ij}(0) y_j) \frac{\partial^2 v_1}{\partial y_i \partial y_N} \right\} \right. \\
& \quad + \chi \left\{ \epsilon |\nabla \rho|^2 \frac{\partial^2 v_2}{\partial y_N^2} - 2\epsilon \sum_{i=1}^{N-1} \rho_i \frac{\partial^2 v_2}{\partial y_i \partial y_N} - \epsilon^2 \Delta \rho \frac{\partial v_2}{\partial y_N} \right\} \\
& \quad \left. + \chi \left\{ \epsilon |\nabla \rho|^2 \frac{\partial^2 v_3}{\partial y_N^2} - 2\epsilon \sum_{i=1}^{N-1} \rho_i \frac{\partial^2 v_3}{\partial y_i \partial y_N} - \epsilon^2 \Delta \rho \frac{\partial v_3}{\partial y_N} \right\} \right] \\
& \quad + \frac{1}{\epsilon^2} E_\epsilon(\chi) \\
& \quad = f_\epsilon
\end{aligned}$$

where  $E_\epsilon(\chi)$  denotes all the terms involving derivatives of  $\chi$ . Since  $|v_1|, |v_2|, |v_3| \leq \exp(-\mu|y|)$  for some  $\mu < \sqrt{m}$  we have  $f_\epsilon \in L^2(\Omega_{\epsilon,P})$  and  $\int_{\Omega_{\epsilon,P}} f_\epsilon^2 \leq C$ . On the other hand, for  $x \in \partial\Omega$  it holds that

$$\frac{\partial \Psi_{\epsilon,P}}{\partial \nu}(x) = \frac{1}{\epsilon^3} \left\{ \frac{\partial V}{\partial \nu} - \epsilon \frac{\partial(v_1 \chi)}{\partial \nu} - \epsilon^2 \left( \frac{\partial(v_2 \chi)}{\partial \nu} + \frac{\partial(v_3 \chi)}{\partial \nu} \right) \right\}.$$

Note that

$$\begin{aligned}
& \frac{\partial V}{\partial \nu} \sqrt{1 + |\nabla \rho|^2} = V' \frac{\langle x - P, \nu \rangle}{\epsilon |x - P|} \sqrt{1 + |\nabla \rho|^2} \\
& = V' \left( \frac{x - P}{\epsilon} \right) \frac{1}{\epsilon |x - P|} \left\{ \frac{1}{2} \sum_{i,j=1}^{N-1} \rho_{ij}(0) (x_i - P_i)(x_j - P_j) \right. \\
& \quad \left. + \frac{1}{3} \sum_{i,j,k=1}^{N-1} \rho_{ijk}(0) (x_i - P_i)(x_j - P_j)(x_k - P_k) + \mathcal{O}(|x' - P'|^4) \right\} \\
& = \frac{V'(y)}{|y|} \left\{ \frac{1}{2} \sum_{i,j=1}^{N-1} \rho_{ij}(0) y_i y_j + \frac{\epsilon}{3} \sum_{i,j,k=1}^{N-1} \rho_{ijk}(0) y_i y_j y_k \right\} + \mathcal{O}(\epsilon^2 \exp(-\mu|z|)).
\end{aligned}$$

Furthermore,

$$\begin{aligned}
& \sqrt{1 + |\nabla \rho|^2} \frac{\partial v_1}{\partial \nu} = \frac{1}{\epsilon} \left\{ \sum_{k=1}^{N-1} \rho_k \frac{\partial v_1}{\partial y_k} - (1 + |\nabla \rho|^2) \frac{\partial v_1}{\partial y_N} \right\}, \\
& \epsilon \frac{\partial \Psi_{\epsilon,P}}{\partial \nu}(x) = \frac{1}{\sqrt{1 + |\nabla \rho|^2} \epsilon^2} \left[ \frac{V'}{|y|} \left\{ \frac{1}{2} \sum_{i,j=1}^{N-1} \rho_{ij} y_i y_j + \frac{\epsilon}{3} \sum_{i,j,k=1}^{N-1} \rho_{ijk}(0) y_i y_j y_k \right\} \right. \\
& \quad \left. + \mathcal{O}(\epsilon^2 (\exp(-\mu|y|))) \right]
\end{aligned}$$

$$\begin{aligned}
& +\chi \left\{ -\sum_{k=1}^{N-1} \rho_k \frac{\partial v_1}{\partial y_k} + \frac{\partial v_1}{\partial y_N} + |\nabla \rho|^2 \frac{\partial v_1}{\partial y_N} \right. \\
& \quad \left. -\epsilon \sum_{k=1}^{N-1} \rho_k \frac{\partial v_2}{\partial y_k} + \epsilon \frac{\partial v_2}{\partial y_N} + \epsilon |\nabla \rho|^2 \frac{\partial v_2}{\partial y_N} \right. \\
& \quad \left. -\epsilon \sum_{k=1}^{N-1} \rho_k \frac{\partial v_3}{\partial y_k} + \epsilon \frac{\partial v_3}{\partial y_N} + |\nabla \rho|^2 \frac{\partial v_3}{\partial y_N} \right\} + E_\epsilon(\chi) \Big] \\
& = g_\epsilon(y)
\end{aligned}$$

where again  $E_\epsilon(\chi)$  denotes all the terms involving derivatives of  $\chi$ . This implies

$$g_\epsilon \leq C \exp(-\mu|z|) \quad \text{for } |z'| \leq \frac{R_0 - \delta}{\epsilon}.$$

Therefore

$$\left| \epsilon \frac{\partial \Psi_{\epsilon, P}}{\partial \nu}(x) \right| \leq C \exp(-\mu|z|) \quad \text{for } z = \frac{x - P}{\epsilon}.$$

Let  $\tilde{\Psi}_{\epsilon, P}(z) = \Psi_{\epsilon, P}(x)$ ,  $x = P + \epsilon z$ . Then  $\tilde{\Psi}_{\epsilon, P}$  satisfies

$$\begin{aligned}
\Delta \tilde{\Psi}_{\epsilon, P} - \tilde{\Psi}_{\epsilon, P} + f_\epsilon &= 0 \quad \text{in } \Omega_{\epsilon, P}, \\
\frac{\partial \tilde{\Psi}_{\epsilon, P}}{\partial \nu_\epsilon} &= g_\epsilon \quad \text{on } \partial \Omega_{\epsilon, P}
\end{aligned}$$

where  $f_\epsilon \in L^2(\Omega_{\epsilon, P})$ ,  $g_\epsilon \in L^2(\partial \Omega_{\epsilon, P})$  and both the corresponding norms are bounded independent of  $\epsilon$ . Hence by Lemma 2.2

$$\|\Psi_{\epsilon, P}\|_\epsilon \leq C.$$

Therefore Proposition 2.1 is proved.  $\square$

We next analyze  $\partial/\partial \tau_{P_j} P_{\Omega_{\epsilon, P}} V\left(\frac{x-P}{\epsilon}\right)$ . After choosing a suitable coordinate system we can assume that  $\partial/\partial \tau_{P_j} = \partial/\partial P_j$ . Then  $\partial/\partial P_j h_{\epsilon, P}(x)$  satisfies

$$\begin{aligned}
\epsilon^2 \Delta v - mv &= 0 \quad \text{in } \Omega, \\
\frac{\partial v}{\partial \nu} &= \frac{\partial}{\partial \nu} \frac{\partial}{\partial P_j} V\left(\frac{x-P}{\epsilon}\right) \quad \text{on } \partial \Omega.
\end{aligned}$$

We compute

$$(1 + |\nabla_{x'} \rho|^2) \frac{\partial}{\partial \nu} \frac{\partial}{\partial P_j} V\left(\frac{x-P}{\epsilon}\right) = \sum_{i=1}^{N-1} \frac{\partial}{\partial x_i} \frac{\partial}{\partial P_j} V\left(\frac{x-P}{\epsilon}\right) \rho_i$$

$$\begin{aligned}
& -\frac{\partial}{\partial x_N} \frac{\partial}{\partial P_j} V \left( \frac{x-P}{\epsilon} \right) \\
& = - \left[ \sum_{i=1}^{N-1} \frac{\partial^2 V}{\partial x_i \partial x_j} \left( \frac{x-P}{\epsilon} \right) \rho_i - \frac{\partial^2 V}{\partial x_N \partial x_j} \left( \frac{x-P}{\epsilon} \right) \right].
\end{aligned}$$

Now we have (let  $x = P + \epsilon z$ )

$$\begin{aligned}
& \frac{\partial V}{\partial z_j}(z) = V' \frac{z_j}{|z|}, \\
& \frac{\partial^2 V}{\partial z_i \partial z_j} = V'' \frac{z_i z_j}{|z|^2} + V' \left\{ \frac{\delta_{ij}}{|z|} - \frac{z_i z_j}{|z|^3} \right\}, \\
& \frac{\partial^2 V((x-P)/\epsilon)}{\partial x_N \partial x_j} = \frac{1}{\epsilon^2} \left\{ V'' \frac{z_j \rho/\epsilon}{|z|^2} - V' \frac{z_j \rho/\epsilon}{|z|^3} \right\} \\
& = \left\{ V'' \frac{y_j}{|y|^2} - V' \frac{y_j}{|y|^3} \right\} \frac{1}{\epsilon^3} \rho + \text{h.o.t.}, \\
& \frac{\partial^2 V((x-P)/\epsilon)}{\partial x_i \partial x_j} = \frac{1}{\epsilon^2} \left\{ V'' \frac{z_i z_j}{|z|^2} + V' \left\{ \frac{\delta_{ij}}{|z|} - \frac{z_i z_j}{|z|^3} \right\} \right\} \rho_i, \\
& (1 + |\nabla \rho|^2) \frac{\partial}{\partial \nu} \frac{\partial}{\partial P_j} V \left( \frac{x-P}{\epsilon} \right) = \\
& - \left[ \frac{1}{\epsilon^2} \left\{ V'' \frac{y_i y_j}{|y|^2} + V' \left\{ \frac{\delta_{ij}}{|y|} - \frac{y_i y_j}{|y|^3} \right\} \right\} \epsilon \sum_{k=1}^{N-1} \rho_{ik} y_k \right. \\
& \left. - \frac{1}{\epsilon^3} \left\{ V'' \frac{y_j}{|y|^2} - V' \frac{y_j}{|y|^3} \right\} \frac{\epsilon^2}{2} \sum_{k,l=1}^{N-1} \rho_{kl} y_k y_l \right] + \text{h.o.t.} \\
& = \frac{1}{\epsilon} \left[ \frac{1}{2} \sum_{k,l=1}^{N-1} \rho_{kl} \left( \frac{V''}{|y|^2} - \frac{V'}{|y|^3} \right) y_k y_j y_l + \frac{V'}{|y|} \sum_{k=1}^{N-1} \rho_{jk} y_k \right] + \text{h.o.t.}
\end{aligned}$$

Let

$$\left[ \frac{\partial V}{\partial \tau_{P_j}} - \frac{\partial P_{\Omega_{\epsilon,P}} V}{\partial \tau_{P_j}} \right] \left( \frac{x-P}{\epsilon} \right) = w_1(y) \chi(x-P) + \epsilon w_2^\epsilon(x).$$

Here  $w_1$  is the unique solution of

$$\begin{cases} \Delta v - mv = 0 & \text{in } R_+^N, \\ \frac{\partial v}{\partial \nu} = -\frac{1}{2} \left( \frac{V''}{|y|^2} - \frac{V'}{|y|^3} \right) \sum_{k,l=1}^{N-1} \rho_{kl}(0) y_k y_l y_j - \frac{V'}{|y|} \sum_{k=1}^{N-1} \rho_{jk}(0) y_k & \text{on } \partial R_+^N. \end{cases} \tag{2.9}$$

Note that  $|w_1| \leq C \exp(-\mu|y|)$  for some  $\mu < \sqrt{m}$  and  $w_1$  is an odd function in  $y'$ . Then  $w_2$  satisfies

$$\begin{cases} \epsilon^2 \Delta w_2 - m w_2 + \frac{1}{\epsilon} [\epsilon^2 \Delta w_1 \chi_1 - w_1 \chi_1] = 0, \\ \frac{\partial w_2}{\partial \nu} = \frac{1}{\epsilon} \left( \frac{\partial}{\partial \nu} \frac{\partial V}{\partial \tau_{P_j}} - \frac{\partial}{\partial \nu} [w_1(y) \chi(x - P)] \right) \end{cases} \quad (2.10)$$

Note that  $|w_2| \leq C \exp(-\mu|y|)$  for some  $\mu < \sqrt{m}$ . Similar to the proof of Proposition 2.1, we have

**Proposition 2.3.**

$$\left[ \frac{\partial V}{\partial \tau_{P_j}} - \frac{\partial P_{\Omega_{\epsilon, P}} V}{\partial \tau_{P_j}} \right] \left( \frac{x - P}{\epsilon} \right) = w_1(y) \chi(x - P) + \epsilon w_2^\epsilon(x)$$

where  $w_1$  is defined above and

$$\|w_2^\epsilon\|_\epsilon \leq C.$$

Finally, let

$$L_0 = \Delta - m + h'(V).$$

We have

**Lemma 2.4.**

$$\text{Ker}(L_0) \cap H_N^2(R_+^N) = \text{span} \left\{ \frac{\partial V}{\partial y_1}, \dots, \frac{\partial V}{\partial y_{N-1}} \right\}.$$

where  $H_N^2(R_+^N) = \{u \in H^2(R_+^N), \frac{\partial u}{\partial y_N} = 0 \text{ on } \partial R_+^N\}$ .

*Proof.* See Lemma 4.2 in [19].  $\square$

### 3. REDUCTION TO FINITE DIMENSIONS

Let  $P \in \Omega$  and

$$\Omega_{\epsilon, P} = \{z \in R^N | \epsilon z + P \in \Omega\}.$$

Let  $H_N^2(\Omega_{\epsilon, P})$  be a Hilbert space defined by

$$H_N^2(\Omega_{\epsilon, P}) = \left\{ u \in H^2(\Omega_{\epsilon, P}) \mid \frac{\partial u}{\partial \nu_\epsilon} = 0 \text{ on } \partial \Omega_{\epsilon, P} \right\}.$$

For  $u \in H_N^2(\Omega_{\epsilon, P})$ , set

$$S_\epsilon(u) = \Delta u - m u + h(u) - \frac{1}{|\Omega|} \int_\Omega h(u).$$

Then solving equation (1.2) is equivalent to

$$S_\epsilon(u) = 0, u \in H_N^2(\Omega_{\epsilon, P}).$$

To this end, we first study the linearized operator

$$\tilde{L}_\epsilon : u(z) \mapsto \Delta u(z) - mu(z) + h'(P_{\Omega_{\epsilon,P}} V(z))u(z),$$

$$H_N^2(\Omega_{\epsilon,P}) \rightarrow L^2(\Omega_{\epsilon,P}).$$

$\tilde{L}_\epsilon$  is not invertible due to the approximate kernel

$$\mathcal{K}_{\epsilon,P} = \text{span} \left\{ \frac{\partial P_{\Omega_{\epsilon,P}} V(z)}{\partial \tau_{P_j}} \Big|_{j=1, \dots, N-1} \right\}$$

in  $H_N^2(\Omega_{\epsilon,P})$ . It is easy to see (integration by parts) that the cokernel of  $\tilde{L}_\epsilon$  coincides with its kernel. We choose approximate cokernel and kernel as follows:

$$\mathcal{C}_{\epsilon,P} = \mathcal{K}_{\epsilon,P} = \text{span} \left\{ \frac{\partial P_{\Omega_{\epsilon,P}} V(z)}{\partial \tau_{P_j}} \Big|_{j=1, \dots, N-1} \right\}.$$

Let  $\pi_{\epsilon,P}$  denote the projection in  $L^2(\Omega_{\epsilon,P})$  onto  $\mathcal{C}_{\epsilon,P}^\perp$ . Our goal in this section is to show that the equation

$$\pi_{\epsilon,P} \circ S_\epsilon(P_{\Omega_{\epsilon,P}} V + \Phi_{\epsilon,P}) = 0$$

has a unique solution  $\Phi_{\epsilon,P} \in \mathcal{K}_{\epsilon,P}^\perp$  if  $\epsilon$  is small enough.

As a preparation in the following two propositions we show invertibility of the corresponding linearized operator.

**Proposition 3.1.** *Let  $L_{\epsilon,P} = \pi_{\epsilon,P} \circ \tilde{L}_\epsilon$ . There exist positive constants  $\bar{\epsilon}, \lambda$  such that for all  $\epsilon \in (0, \bar{\epsilon})$*

$$\|L_{\epsilon,P} \Phi\|_{L^2(\Omega_{\epsilon,P})} \geq \lambda \|\Phi\|_{H^2(\Omega_{\epsilon,P})} \quad (3.1)$$

for all  $\Phi \in \mathcal{K}_{\epsilon,P}^\perp$ .

**Proposition 3.2.** *There exists a positive constant  $\bar{\epsilon}$  such that for all  $\epsilon \in (0, \bar{\epsilon})$  and  $P \in \partial\Omega$  the map*

$$L_{\epsilon,P} = \pi_{\epsilon,P} \circ \tilde{L}_\epsilon : \mathcal{K}_{\epsilon,P}^\perp \rightarrow \mathcal{C}_{\epsilon,P}^\perp$$

is surjective.

*Proof of Proposition 3.1:* We will follow the method used in [8], [21] and [22]. Suppose that (3.1) is false. Then there exist sequences  $\{\epsilon_k\}$ ,  $\{P_k\}$ , and

$\{\Phi_k\}$  with  $P_k \in \partial\Omega$ ,  $\Phi_k \in \mathcal{K}_{\epsilon_k, P_k}^\perp$  such that

$$\|L_{\epsilon_k, P_k} \Phi_k\|_{L^2} \rightarrow 0, \quad (3.2)$$

$$\|\Phi_k\|_{H^2} = 1, \quad k = 1, 2, \dots \quad (3.3)$$

We omit the argument  $\Omega_{\epsilon_k, P_k}$  where this can be done without confusion.

Denote

$$e_{k,j} = \frac{\partial}{\partial \tau_{P_j}} P_{\Omega_{\epsilon, P}} V / \left\| \frac{\partial}{\partial \tau_{P_j}} P_{\Omega_{\epsilon, P}} V \right\|_{L^2}.$$

Note that

$$\langle e_{k,i}, e_{k,j} \rangle = \delta_{ij} + \mathcal{O}(\epsilon_k) \quad \text{as } k \rightarrow \infty$$

by Proposition 2.3 and because of the symmetry of the function  $w_1$ , which was defined in (2.9), where  $\delta_{ij}$  is the Kronecker symbol. Furthermore because of (3.2),

$$\|\tilde{L}_{\epsilon_k} \Phi_k\|_{L^2}^2 - \sum_{j=1}^{N-1} \left( \int_{\Omega_{\epsilon_k, P_k}} \tilde{L}_{\epsilon_k} \Phi_k e_{k,j} \right)^2 \rightarrow 0 \quad (3.4)$$

as  $k \rightarrow \infty$ . Let  $\Omega_1$ ,  $\chi$ ,  $\rho$  and  $T$  be as defined in Section 2. Then  $T$  has an inverse  $T^{-1}$  such that

$$T^{-1} : T(B(P, R_0) \cap \bar{\Omega}) \rightarrow B(P, R_0) \cap \bar{\Omega}.$$

Recall that  $\epsilon y = T(x)$ . We introduce a new sequence  $\{\varphi_k\}$  by

$$\varphi_k(y) = \chi(T^{-1}(\epsilon_k y)) \Phi_k \left( T^{-1}(\epsilon_k y) \right) \quad (3.5)$$

for  $y \in R_+^N$ . Since  $T$  and  $T^{-1}$  have bounded derivatives it follows from (3.3) and the smoothness of  $\chi$  that

$$\|\varphi_k\|_{H^2(R_+^N)} \leq C$$

for all  $k$  sufficiently large. Therefore there exists a subsequence, again denoted by  $\{\varphi_k\}$  which converges weakly in  $H^2(R_+^N)$  to a limit  $\varphi_\infty$  as  $k \rightarrow \infty$ .

We are now going to show that  $\varphi_\infty \equiv 0$ . As a first step we deduce

$$\int_{R_+^N} \varphi_\infty \frac{\partial V}{\partial P_j} = 0, \quad j = 1, \dots, N-1. \quad (3.6)$$

This statement is shown as follows (note that  $\det DT = \det DT^{-1} = 1$ )

$$\int_{R_+^N} \varphi_k(y) \left[ \frac{\partial P_{\Omega_{\epsilon, P}} V}{\partial \tau_{P_{k,j}}} \left( \frac{T^{-1}(\epsilon_k y) - P_k}{\epsilon_k} \right) \right] dy$$

$$\begin{aligned}
&= \epsilon_k^{-N} \int_{\Omega_1} \chi(x) \Phi_k(x) \frac{\partial P_{\Omega_{\epsilon,P}} V \left( \frac{x-P_k}{\epsilon_k} \right)}{\partial \tau_{P_{k,j}}} dx \\
&= \epsilon_k^{-N} \int_{\Omega} \Phi_k(x) \frac{\partial P_{\Omega_{\epsilon,P}} V \left( \frac{x-P_k}{\epsilon_k} \right)}{\partial \tau_{P_{k,j}}} \\
&\quad - \epsilon_k^{-N} \int_{\Omega \setminus \Omega_1} \Phi_k(x) \frac{\partial P_{\Omega_{\epsilon,P}} V \left( \frac{x-P_k}{\epsilon_k} \right)}{\partial \tau_{P_{k,j}}} \\
&\quad - \epsilon_k^{-N} \int_{\Omega_1} [1 - \chi(x)] \Phi_k(x) \frac{\partial P_{\Omega_{\epsilon,P}} V \left( \frac{x-P_k}{\epsilon_k} \right)}{\partial \tau_{P_{k,j}}} \\
&= 0 - \epsilon_k^{-N} \int_{\Omega \setminus \Omega_1} \Phi_k(x) \left[ \frac{\partial V}{\partial P_{k,j}} - \frac{\partial P_{\Omega_{\epsilon,P}} V}{\partial \tau_{P_{k,j}}} \right] \left( \frac{x - P_k}{\epsilon_k} \right) \\
&\quad - \epsilon_k^{-N} \int_{\Omega_1} [1 - \chi(x)] \Phi_k(x) \left[ \frac{\partial V \left( \frac{x-P_k}{\epsilon_k} \right)}{\partial P_{k,j}} - \frac{\partial P_{\Omega_{\epsilon,P}} V}{\partial \tau_{P_{k,j}}} \right] \left( \frac{x - P_k}{\epsilon_k} \right) \\
&\quad \quad - \epsilon_k^{-N} \int_{\Omega \setminus \Omega_1} \Phi_k(x) \frac{\partial V \left( \frac{x-P_k}{\epsilon_k} \right)}{\partial P_{k,j}} \\
&\quad \quad - \epsilon_k^{-N} \int_{\Omega_1} [1 - \chi(x)] \Phi_k(x) \frac{\partial V \left( \frac{x-P_k}{\epsilon_k} \right)}{\partial P_{k,j}}
\end{aligned}$$

where  $\Omega_1$  is as defined in section 2. In the last expression the first two terms tend to zero as  $k \rightarrow \infty$  since  $\epsilon_k^{-N} \Phi_k$  is bounded in  $L^2(\Omega)$  and  $[\dots] \rightarrow 0$  strongly in  $L^2(\Omega)$ . The last two terms tend to zero as  $k \rightarrow \infty$  because of the exponential decay of  $\partial V / \partial P_{k,j}$  at infinity.

We conclude

$$\limsup_{k \rightarrow \infty} \left| \int_{R_+^N} \varphi_k(y) \frac{\partial P_{\Omega_{\epsilon,P}} V \left( \frac{T^{-1}(\epsilon_k y) - P_k}{\epsilon_k} \right)}{\partial \tau_{P_{k,j}}} \right| = 0, \quad j = 1, \dots, N-1. \quad (3.7)$$

This implies (3.6).

Let  $\mathcal{K}_0$  and  $\mathcal{C}_0$  be the kernel and cokernel, respectively, of the linear operator  $S'_0(V)$  which is the Fréchet derivative at  $V$  of

$$\begin{aligned}
S_0(v) &= \Delta v - mv + h(v), \\
S_0 &: H_N^2(R_+^N) \rightarrow L^2(R_+^N), \\
H_N^2(R_+^N) &= \left\{ u \in H_N^2(R_+^N) \left| \frac{\partial u}{\partial y_N} = 0 \right. \right\}.
\end{aligned}$$

Note that

$$\mathcal{K}_0 = \mathcal{C}_0 = \text{span} \left\{ \frac{\partial V}{\partial y_j} \mid j = 1, \dots, N-1 \right\}.$$

Equation (3.6) implies that  $\varphi_\infty \in \mathcal{K}_0^\perp$ . By the exponential decay of  $V$  and by (3.2) we have after possibly taking a further subsequence that

$$\Delta\varphi_\infty - m\varphi_\infty + h'(V)\varphi_\infty = 0,$$

i.e.  $\varphi_\infty \in \mathcal{K}_0$ . Therefore  $\varphi_\infty = 0$ .

Hence

$$\varphi_k \rightharpoonup 0 \quad \text{weakly in } H^2(R_+^N) \quad (3.8)$$

as  $k \rightarrow \infty$ . By the definition of  $\varphi_k$  we get  $\Phi_k \rightharpoonup 0$  in  $H^2$  and

$$\|h'(P_{\Omega_{\epsilon_k, P_k}} V)\Phi_k\|_{L^2} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Furthermore,

$$\|(\Delta - m)\Phi_k\|_{L^2} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Since

$$\begin{aligned} \int_{\Omega_{\epsilon_k, P_k}} |\nabla\Phi_k|^2 + m\Phi_k^2 &= \int_{\Omega_{\epsilon_k, P_k}} [(m - \Delta)\Phi_k]\Phi_k \\ &\leq C\|(\Delta - m)\Phi_k\|_{L^2} \end{aligned}$$

we have that

$$\|\Phi_k\|_{H^1} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

In summary:

$$\|\Delta\Phi_k\|_{L^2} \rightarrow 0 \text{ and } \|\Phi_k\|_{H^1} \rightarrow 0. \quad (3.9)$$

From (3.9) and the following elliptic regularity estimate (for a proof see Appendix B)

$$\|\Phi_k\|_{H^2} \leq C(\|\Delta\Phi_k\|_{L^2} + \|\Phi_k\|_{H^1}) \quad (3.10)$$

for  $\Phi_k \in H_N^2$  we imply that

$$\|\Phi_k\|_{H^2} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

This contradicts the assumption

$$\|\Phi_k\|_{H^2} = 1$$

and the proof of Proposition 3.1 is completed.  $\square$



*Proof of Proposition 3.2:* Assume that the statement is not true. Then there exist sequences  $\{\epsilon_k\}, \{P_k\}$  such that  $\epsilon_k \rightarrow 0$  as  $k \rightarrow \infty$  and  $P_k \in \partial\Omega$  and such that for all  $k$ ,  $L_{\epsilon_k, P_k} : \mathcal{K}_{\epsilon_k, P_k}^\perp \rightarrow \mathcal{C}_{\epsilon_k, P_k}^\perp$  is not surjective. Let  $\mathcal{K}_{\epsilon, P}$  and  $\mathcal{C}_{\epsilon, P}$  be the kernel and cokernel of  $\tilde{L}_\epsilon$ , respectively. Then  $\pi_{\epsilon_k, P_k} : \mathcal{C}_{\epsilon_k, P_k}^\perp \rightarrow \mathcal{C}_{\epsilon_k, P_k}^\perp$  is not surjective, i.e. for all  $k$  there exists a  $\Phi_k \in \mathcal{C}_{\epsilon_k, P_k}^\perp$  with  $\Phi_k \neq 0$  such that  $\Psi + \Phi_k \notin \mathcal{C}_{\epsilon_k, P_k}^\perp$  for all  $\Psi \in \mathcal{C}_{\epsilon_k, P_k}^\perp$ . This is equivalent to  $\Phi_k \in \mathcal{C}_{\epsilon_k, P_k}$  and  $\Phi_k \neq 0$ . Because we can assume that w.l.o.g.  $\Phi_k = 1$  this can be rewritten as follows. For all  $k$  there exists a  $\Phi_k \in \mathcal{C}_{\epsilon_k, P_k}$  such that

$$\|\Phi_k\|_{L^2} = 1, \quad (3.11)$$

$$\int_{\Omega_{\epsilon_k, P_k}} \Phi_k \frac{\partial P_{\Omega_{\epsilon, P}} V}{\partial \tau_{P_k, j}} = 0 \quad j = 1, \dots, N-1.$$

Now since

$$\Delta \Phi_k - m \Phi_k + h'(P_{\Omega_{\epsilon, P}} V) \Phi_k = 0$$

and because of the elliptic estimate (3.10) it follows that

$$\|\Phi_k\|_{H^2} \leq C$$

for some constant  $C$  independent of  $k$ . Extract a subsequence (again denoted by  $\{\Phi_k\}$ ) such that  $\varphi_k$  as defined in (3.5) converges weakly in  $H^2(R_+^N)$  to  $\varphi_\infty$  as  $k \rightarrow \infty$  and  $\varphi_\infty$  satisfies

$$\begin{aligned} \Delta \varphi_\infty - m \varphi_\infty + h'(V) \varphi_\infty &= 0 \quad \text{in } R_+^N, \\ \frac{\partial \varphi_\infty}{\partial y_n} &= 0 \quad \text{in } R^{N-1} \times \{0\} \end{aligned} \quad (3.12)$$

with

$$\int_{R_+^N} \varphi_\infty \frac{\partial V}{\partial y_j} = 0 \quad j = 1, \dots, N-1. \quad (3.13)$$

From (3.12) we deduce that  $\varphi_\infty$  belongs to the kernel of  $S'_0(V)$  and (3.13) implies that  $\varphi_\infty$  lies in the orthogonal complement of the kernel of  $S'_0(V)$ .

Therefore  $\varphi_\infty = 0$ . As in the proof of Proposition 3.1 we show by the elliptic regularity estimate (3.10) that  $\|\Phi_k\|_{H^2} \rightarrow 0$  as  $k \rightarrow \infty$ . This contradicts (3.11) and the proof of Proposition 3.2 is finished.  $\square$

We are now in a position to solve the equation

$$\pi_{\epsilon, P} \circ S_\epsilon(P_{\Omega_{\epsilon, P}} V + \Phi_{\epsilon, P}) = 0. \quad (3.14)$$

Since  $L_{\epsilon,P}|_{\mathcal{K}_{\epsilon,P}^\perp}$  is invertible (call the inverse  $L_{\epsilon,P}^{-1}$ ) we can rewrite

$$\Phi = -(L_{\epsilon,P}^{-1} \circ \pi_{\epsilon,P})(S_\epsilon(P_{\Omega_{\epsilon,P}}V)) - (L_{\epsilon,P}^{-1} \circ \pi_{\epsilon,P})N_{\epsilon,P}(\Phi) \equiv M_{\epsilon,P}(\Phi) \quad (3.15)$$

where

$$N_{\epsilon,P}(\Phi) = S_\epsilon(P_{\Omega_{\epsilon,P}}V + \Phi) - [S_\epsilon(P_{\Omega_{\epsilon,P}}V) + S'_\epsilon(P_{\Omega_{\epsilon,P}}V)\Phi]$$

and the operator  $M_{\epsilon,P}$  is defined by the last equation for  $\Phi \in H_N^2(\Omega_{\epsilon,P})$ . We are going to show that the operator  $M_{\epsilon,P}$  is a contraction on

$$B_{\epsilon,\delta} \equiv \{\Phi \in H^2(\Omega_{\epsilon,P}) \mid \|\Phi\|_{H^2(\Omega_{\epsilon,P})} < \delta\}$$

if  $\delta$  is small enough. We have

$$\begin{aligned} \|M_{\epsilon,P}(\Phi)\|_{H^2(\Omega_{\epsilon,P})} &\leq \lambda^{-1}(\|\pi_{\epsilon,P}N_{\epsilon,P}(\Phi)\|_{L^2(\Omega_{\epsilon,P})} + \|\pi_{\epsilon,P}(P_{\Omega_{\epsilon,P}}V - V)\|_{L^2(\Omega_{\epsilon,P})}) \\ &\leq \lambda^{-1}C(c(\delta)\delta + \epsilon) \end{aligned}$$

where  $\lambda > 0$  is independent of  $\delta > 0$  and  $c(\delta) \rightarrow 0$  as  $\delta \rightarrow 0$ . Similarly we show

$$\|M_{\epsilon,P}(\Phi) - M_{\epsilon,P}(\Phi')\|_{H^2(\Omega_{\epsilon,P})} \leq \lambda^{-1}C(\epsilon + c(\delta)\delta)\|\Phi - \Phi'\|_{H^2(\Omega_{\epsilon,P})}$$

where  $c(\delta) \rightarrow 0$  as  $\delta \rightarrow 0$ . Therefore  $M_{\epsilon,P}$  is a contraction on  $B_\delta$ . The existence of a fixed point  $\Phi_{\epsilon,P}$  now follows from the Contraction Mapping Principle and  $\Phi_{\epsilon,P}$  is a solution of (3.15).

Because of

$$\begin{aligned} \|\Phi_{\epsilon,P}\|_{H^2(\Omega_{\epsilon,P})} &\leq \lambda^{-1}(\|N_{\epsilon,P}(\Phi_{\epsilon,P})\|_{L^2(\Omega_{\epsilon,P})} + \|P_{\Omega_{\epsilon,P}}V - V\|_{L^2}) \\ &\leq \lambda^{-1}(c\epsilon + c(\delta)\|\Phi_{\epsilon,P}\|_{H^2(\Omega_{\epsilon,P})}) \end{aligned}$$

we have

$$(1 - \lambda^{-1}c(\delta))\|\Phi_{\epsilon,P}\|_{H^2} \leq C\epsilon.$$

We have proved

**Lemma 3.3.** *There exists  $\bar{\epsilon} > 0$  such that for every pair of  $\epsilon, P$  with  $0 < \epsilon < \bar{\epsilon}$  and  $P \in \partial\Omega$  there exists a unique  $\Phi_{\epsilon,P} \in \mathcal{K}_{\epsilon,P}^\perp$  satisfying  $S_\epsilon(P_{\Omega_{\epsilon,P}}V + \Phi_{\epsilon,P}) \in \mathcal{C}_{\epsilon,P}$  and*

$$\|\Phi_{\epsilon,P}\|_{H^2(\Omega_{\epsilon,P})} \leq C\epsilon. \quad (3.16)$$

We need another statement about the asymptotic behavior of the function  $\Phi_{\epsilon,P}$  as  $\epsilon \rightarrow 0$ , which gives an expansion in  $\epsilon$  and is stated as follows.

**Proposition 3.4.**

$$\Phi_{\epsilon,P}(x) = \epsilon(\Phi_0(y)\chi(x-P)) + \epsilon^2\Psi_{\epsilon,P}(x) \quad (3.17)$$

where

$$\|\Psi_{\epsilon,P}\|_{\epsilon} \leq C$$

and  $\Phi_0$  is the unique solution of

$$\begin{aligned} \Delta\Phi_0 - m\Phi_0 + h'(V)\Phi_0 - h'(V)v_1 &= 0, & \text{in } R_+^N, \\ \frac{\partial\Phi_0}{\partial y_N} &= 0 & \text{on } \partial R_+^N, \end{aligned} \quad (3.18)$$

$$\Phi_0 \text{ is orthogonal to the kernel of } L_0 \quad (3.18)$$

where  $L_0 = \Delta - m + h'(V)$ ,  $L_0 : H_N^2(R_+^N) \rightarrow L^2(R_+^N)$ .

*Proof.* Note that the kernel of  $L_0$  is

$$\left\{ \frac{\partial V}{\partial y_j} \Big| j = 1, \dots, N-1 \right\}.$$

Furthermore we have

$$|\Phi_0| \leq C \exp(-\mu|y|) \quad \text{for } \mu < \sqrt{m}.$$

The notations for  $\Omega_1$ ,  $\chi$ ,  $\rho$  and  $T$  are as in section 2. Our strategy is to decompose  $\Psi_{\epsilon,P}$  into three parts and show that each of them is bounded in  $\|\cdot\|_{H^1(\Omega_{\epsilon,P})}$  as  $\epsilon \rightarrow 0$ . That means we make the ansatz

$$\Psi_{\epsilon,P}(x) = \Psi_{\epsilon}^1(x) + \Psi_{\epsilon}^{2,1}(x) + \Psi_{\epsilon}^{2,2}(x)$$

where the functions  $\Psi_{\epsilon}^1$ ,  $\Psi_{\epsilon}^{2,1}$ ,  $\Psi_{\epsilon}^{2,2}$  will be defined as follows. Let  $\Psi_{\epsilon}^1$  be the unique solution of

$$\begin{aligned} \epsilon^2\Delta\Psi_{\epsilon}^1 - m\Psi_{\epsilon}^1 &= 0 & \text{in } \Omega, \\ \frac{\partial\Psi_{\epsilon}^1}{\partial\nu} &= g^{\epsilon} & \text{on } \partial\Omega \end{aligned} \quad (3.19)$$

where

$$g_{\epsilon}(x) = -\frac{\partial}{\partial\nu_x}[\Phi_0(y)\chi(x)].$$

Since  $\|g_\epsilon\|_{L^2} \leq C$  there exists a constant  $C > 0$  such that

$$\|\Psi_\epsilon^1\|_{H^1} \leq C. \quad (3.20)$$

Define  $\Psi_\epsilon^{2,1}$  by

$$\Psi_\epsilon^{2,1} = -\frac{1}{\epsilon}\tilde{\pi}\Phi_0(x)\chi - \tilde{\pi}\Psi_\epsilon^1 \quad (3.21)$$

where  $\tilde{\pi}$  is the projection in  $L^2(\Omega_{\epsilon,P})$  onto  $\mathcal{K}_{\epsilon,P}$ . Because of the exponential decay of  $\Phi_0$ , the smoothness of  $\chi$  and and by (3.20) it follows that

$$\|\Psi_\epsilon^{2,1}\|_\epsilon \leq C. \quad (3.22)$$

Finally, define  $\Psi_\epsilon^{2,2}(x)$  to be the unique solution in  $H_N^2(\Omega)$  of the following equation

$$\epsilon^2\Delta\Psi_\epsilon^{2,2} - m\Psi_\epsilon^{2,2} + h'(P_{\Omega_{\epsilon,P}}V)\Psi_\epsilon^{2,2} = -\frac{1}{\epsilon^2}f_\epsilon \quad \text{in } \Omega, \quad (3.23)$$

$$\frac{\partial\Psi_\epsilon^{2,2}}{\partial\nu} = 0 \quad \text{on } \partial\Omega \quad (3.24)$$

where

$$f_\epsilon = \tilde{L}_\epsilon(\Phi_{\epsilon,P} - \epsilon\Phi_0\chi - \epsilon^2(\Psi_\epsilon^1 + \Psi_\epsilon^{2,1})).$$

Note that the right-hand side of the last equation lies in  $\mathcal{C}_{\epsilon,P}^\perp$  since

$$\Phi_{\epsilon,P} - \epsilon\Phi_0\chi - \epsilon^2(\Psi_\epsilon^1 + \Psi_\epsilon^{2,1}) \in H_N^2.$$

This is clear for  $\Phi_{\epsilon,P}$  by definition. By construction we have that  $-\epsilon\Phi_0\chi - \epsilon^2(\Psi_\epsilon^1 + \Psi_\epsilon^{2,1})$  satisfies the Neumann boundary condition. By (3.18) and the smoothness of  $\chi$  we conclude that  $\Phi_0\chi \in H^2$ . By (3.19),  $\Psi_\epsilon^1 \in H^2$ . Finally, since  $e_j \in H^2$  where

$$e_j = \frac{\partial V}{\partial\tau_{P_j}} / \left\| \frac{\partial V}{\partial\tau_{P_j}} \right\|_{L^2(\Omega_{\epsilon,P})} \quad j = 1, \dots, N-1$$

we have  $\Psi_\epsilon^{2,1} \in H^2$ . Therefore  $f_\epsilon \in \mathcal{C}_{\epsilon,P}^\perp$ . Furthermore, the following lemma is true.

**Lemma 3.5.**

$$\|f_\epsilon\|_{L^2(\Omega_{\epsilon,P})} \leq C\epsilon^2.$$

*Proof.* We have

$$\begin{aligned} f_\epsilon &= S'_\epsilon(P_{\Omega_\epsilon, P} V)(\Phi_\epsilon - \epsilon\Phi_0\chi - \epsilon^2(\Psi_\epsilon^1 + \Psi_\epsilon^{2,1})) \\ &= -h(P_{\Omega_\epsilon, P} V) + h(V) + \epsilon h'(V)v_1\chi + N'_{\epsilon, P}(\Phi_\epsilon) \end{aligned}$$

where

$$\begin{aligned} N'_{\epsilon, P}(\Phi) &= \frac{1}{|\Omega|} \int_\Omega h'(P_{\Omega_\epsilon, P} V)\Phi_\epsilon + \frac{1}{|\Omega|} \int_\Omega [h(P_{\Omega_\epsilon, P} V) - h(V)] \\ &\quad - [h(P_{\Omega_\epsilon, P} V + \Phi_\epsilon) - h(P_{\Omega_\epsilon, P} V) - h'(P_{\Omega_\epsilon, P} V)\Phi_\epsilon] \\ &\quad + \frac{1}{|\Omega|} \int_\Omega [h(P_{\Omega_\epsilon, P} V + \Phi_\epsilon) - h(P_{\Omega_\epsilon, P} V) - h'(P_{\Omega_\epsilon, P} V)\Phi_\epsilon] \\ &\quad + \epsilon\Phi_0(y)[\Delta - m + h'(P_{\Omega_\epsilon, P} V)]\chi(x) + \epsilon \langle \nabla_x \Phi_0(y), \nabla_x \Phi(x) \rangle \\ &\quad + \epsilon^2 h'(P_{\Omega_\epsilon, P} V)\Psi_\epsilon^1 + \epsilon^2 [\Delta - m + h'(P_{\Omega_\epsilon, P} V)]\Psi_\epsilon^{2,1}. \end{aligned}$$

Note that

$$\begin{aligned} &\| -h(P_{\Omega_\epsilon, P} V) + h(V) + \epsilon h'(V)v_1\chi(x) \|_{L^2} \\ &\leq \| -h(P_{\Omega_\epsilon, P} V) + h(V) + \epsilon h'(V)v_1 \|_{L^2} \\ &\quad + \| \epsilon(-h'(V)v_1 + h'(V)v_1\chi) \|_{L^2} \\ &\leq C(\epsilon^2 + \exp(-\mu R_0)) \end{aligned}$$

by the definition of  $\chi$  and the exponential decay of  $V$ . Furthermore

$$\|N'_{\epsilon, P}(\Phi)\|_{L^2} \leq C\epsilon^2.$$

This proves Lemma 3.5.  $\square$

By Lemma 3.5 and the invertibility of

$$\tilde{L}_\epsilon : H_N^2 \cap \mathcal{K}_{\epsilon, P}^\perp \rightarrow \mathcal{C}_{\epsilon, P}^\perp$$

Proposition 3.4 follows.  $\square$

## 4. THE REDUCED PROBLEM

In this section we solve the reduced problem and prove our main theorem.

By Lemma 3.3 there exists a unique solution  $\Phi_{\epsilon,P} \in \mathcal{K}_{\epsilon,P}^\perp$  such that

$$\begin{aligned} S_\epsilon(u_\epsilon) &= S_\epsilon \left( P_{\Omega_{\epsilon,P}} V \left( \frac{x-P}{\epsilon} \right) + \Phi_{\epsilon,P} \right) \\ &= \epsilon^2 \Delta u_\epsilon - m u_\epsilon + h(u_\epsilon) - \frac{1}{|\Omega|} \int_\Omega h(u_\epsilon) \in \mathcal{C}_{\epsilon,P}. \end{aligned}$$

Our idea is to find  $P$  such that

$$S_\epsilon(u_\epsilon) \perp \mathcal{C}_{\epsilon,P}.$$

Let

$$\begin{aligned} W_{\epsilon,j}(P) &= \frac{1}{\epsilon^{N+1}} \int_\Omega \left( S_\epsilon(u_\epsilon) \frac{\partial P_{\Omega_{\epsilon,P}} V}{\partial \tau_{P_j}} \right), \\ W_\epsilon(P) &= (W_{\epsilon,1}(P), \dots, W_{\epsilon,N-1}(P)). \end{aligned}$$

Then  $W_\epsilon(P)$  is a continuous map of  $P$ .

Let us now calculate  $W_\epsilon(P)$ . First of all, from condition (4) on  $h$ , we have

$$|h(t)| \leq C t^2.$$

Therefore

$$\int_\Omega h(u_\epsilon) \leq C \epsilon^N.$$

Hence by Proposition 2.3

$$\begin{aligned} \frac{1}{\epsilon^{N+1}} \int_\Omega \left( \int_\Omega h(u_\epsilon) \right) \frac{\partial P_{\Omega_{\epsilon,P}} V}{\partial \tau_{P_j}} &= \int_\Omega h(u_\epsilon) \frac{1}{\epsilon^{N+1}} \int_\Omega \frac{\partial P_{\Omega_{\epsilon,P}} V}{\partial \tau_{P_j}} \\ &= \mathcal{O}(\epsilon^N) \left( \frac{1}{\epsilon^{N+1}} \int_\Omega \left( \frac{\partial V}{\partial P_j} + w_1(y) \chi(x-P) + \epsilon w_2^\epsilon(x) \right) \right) \\ &= \mathcal{O}(\epsilon^N) \frac{1}{\epsilon^{N+1}} \left[ \mathcal{O}(\exp(-\sigma/\epsilon)) + \epsilon \int_\Omega w_2^\epsilon \right] \\ &= \mathcal{O}(\epsilon^{N/2}) \end{aligned}$$

because

$$\frac{1}{\epsilon^N} \int_\Omega w_2^\epsilon \leq \frac{1}{\epsilon^{N/2}} \|w_2^\epsilon\|_{L^2(\Omega)}$$

and Proposition 2.3. On the other hand, since

$$\epsilon^2 \Delta \frac{\partial P_{\Omega_{\epsilon,P}} V}{\partial \tau_{P_j}} - m \frac{\partial P_{\Omega_{\epsilon,P}} V}{\partial \tau_{P_j}} + h'(V) \frac{\partial V}{\partial P_j} = 0,$$

we conclude

$$\begin{aligned}
& \int_{\Omega} [\epsilon^2 \Delta u_{\epsilon} - m u_{\epsilon} + h(u_{\epsilon})] \frac{\partial P_{\Omega_{\epsilon}, P} V}{\partial \tau_{P_j}} \\
&= \int_{\Omega} \left\{ h(u_{\epsilon}) \frac{\partial P_{\Omega_{\epsilon}, P} V \left( \frac{x-P}{\epsilon} \right)}{\partial \tau_{P_j}} + \left[ \epsilon^2 \Delta \frac{\partial P_{\Omega_{\epsilon}, P} V}{\partial \tau_{P_j}} - m \frac{\partial P_{\Omega_{\epsilon}, P} V}{\partial \tau_{P_j}} \right] u_{\epsilon} \right\} \\
&= \int_{\Omega} \left[ h(u_{\epsilon}) \frac{\partial P_{\Omega_{\epsilon}, P} V \left( \frac{x-P}{\epsilon} \right)}{\partial \tau_{P_j}} - h'(V) \frac{\partial V \left( \frac{x-P}{\epsilon} \right)}{\partial P_j} u_{\epsilon} \right] \\
&= \int_{\Omega} \left[ h(P_{\Omega_{\epsilon}, P} V + \Phi_{\epsilon, P}) - h(P_{\Omega_{\epsilon}, P} V) - h'(P_{\Omega_{\epsilon}, P} V) \Phi_{\epsilon, P} \right] \frac{\partial P_{\Omega_{\epsilon}, P} V}{\partial \tau_{P_j}} \\
&\quad + \int_{\Omega} \left[ h'(P_{\Omega_{\epsilon}, P} V) \frac{\partial P_{\Omega_{\epsilon}, P} V}{\partial \tau_{P_j}} - h'(V) \frac{\partial V}{\partial P_j} \right] \Phi_{\epsilon, P} \\
&\quad + \int_{\Omega} \left[ h(P_{\Omega_{\epsilon}, P} V) - h(V) \right] \frac{\partial P_{\Omega_{\epsilon}, P} V}{\partial \tau_{P_j}} \\
&= I_{\epsilon}^1 + I_{\epsilon}^2 + J_{\epsilon}
\end{aligned}$$

where  $I_{\epsilon}^1$ ,  $I_{\epsilon}^2$ , and  $J_{\epsilon}$  are defined by the last equality. We first calculate  $I_{\epsilon}^2$ .

$$\begin{aligned}
I_{\epsilon}^2 &= \int_{\Omega} \left[ h'(P_{\Omega_{\epsilon}, P} V) \frac{\partial P_{\Omega_{\epsilon}, P} V}{\partial \tau_{P_j}} - h'(V) \frac{\partial V}{\partial \tau_{P_j}} \right] (\epsilon \Phi_0(y) \chi(x-P) + \epsilon^2 \Psi_{\epsilon}(x)) dx \\
&= \int_{\Omega} \left[ h'(P_{\Omega_{\epsilon}, P} V) \frac{\partial P_{\Omega_{\epsilon}, P} V}{\partial \tau_{P_j}} - h'(V) \frac{\partial V}{\partial \tau_{P_j}} \right] \epsilon \Phi_0 \chi \\
&\quad + \epsilon^2 \int_{\Omega} \left[ h'(P_{\Omega_{\epsilon}, P} V) \frac{\partial P_{\Omega_{\epsilon}, P} V}{\partial \tau_{P_j}} - h'(V) \frac{\partial V}{\partial \tau_{P_j}} \right] \Psi_{\epsilon} \\
&= \epsilon I_{\epsilon}^{2,1} + \epsilon^2 I_{\epsilon}^{2,2}.
\end{aligned}$$

Note that

$$\begin{aligned}
& h'(P_{\Omega_{\epsilon}, P} V) \frac{\partial P_{\Omega_{\epsilon}, P} V}{\partial \tau_{P_j}} - h'(V) \frac{\partial V}{\partial \tau_{P_j}} \\
&= \left[ h'(P_{\Omega_{\epsilon}, P} V) - h'(V) \right] \frac{\partial P_{\Omega_{\epsilon}, P} V}{\partial \tau_{P_j}} + h'(V) \left[ \frac{\partial P_{\Omega_{\epsilon}, P} V}{\partial \tau_{P_j}} - \frac{\partial V}{\partial \tau_{P_j}} \right]
\end{aligned}$$

and

$$\begin{aligned}
& \int_{\Omega} \left[ h'(P_{\Omega_{\epsilon}, P} V) - h'(V) \right] \frac{\partial P_{\Omega_{\epsilon}, P} V}{\partial \tau_{P_j}} \Phi_0 \chi \\
&= \int_{\Omega} h''(V) (P_{\Omega_{\epsilon}, P} V - V) \frac{\partial P_{\Omega_{\epsilon}, P} V}{\partial \tau_{P_j}} \Phi_0 + \int_{\Omega} h'''(v_1) (P_{\Omega_{\epsilon}, P} V - V)^2 \frac{\partial P_{\Omega_{\epsilon}, P} V}{\partial \tau_{P_j}} \Phi_0 \\
&\quad + \mathcal{O}(\exp(-\delta/\epsilon)) \\
&= \mathcal{O}(\epsilon^{N+1})
\end{aligned}$$

since  $\Phi_0$  is even and  $V - P_{\Omega_{\epsilon,P}}V = \epsilon V_1$  where  $V_1$  is even. By Proposition 2.1

$$\int |\Psi_\epsilon|^2 \leq C\epsilon^N.$$

Hence

$$|I_\epsilon^{2,2}| \leq \mathcal{O}(\epsilon^N).$$

So

$$|I_\epsilon^2| \leq \mathcal{O}(\epsilon^{N+2}).$$

We next compute  $I_\epsilon^1$ .

$$\begin{aligned} I_\epsilon^1 &= \int_\Omega h''(P_{\Omega_{\epsilon,P}}V) \Phi_{\epsilon,P}^2 \frac{\partial P_{\Omega_{\epsilon,P}}V}{\partial \tau_{P_j}} + \int_\Omega h'''(v_1 + P_{\Omega_{\epsilon,P}}V) \Phi_{\epsilon,P}^3 \frac{\partial P_{\Omega_{\epsilon,P}}V}{\partial \tau_{P_j}} \\ &= \int_\Omega h''(P_{\Omega_{\epsilon,P}}V) \epsilon^2 [\Phi_0^2 \chi^2 + 2\epsilon \Phi_0 \chi \Psi_{\epsilon,P} + \epsilon^2 \Psi_{\epsilon,P}^2] \frac{\partial P_{\Omega_{\epsilon,P}}V}{\partial \tau_{P_j}} + \mathcal{O}(\epsilon^{N+2}) \\ &= \mathcal{O}(\epsilon^{N+2}) \end{aligned}$$

since  $\Phi_0$  is even. Finally, we compute the term  $J_\epsilon$ .

$$\begin{aligned} J_\epsilon &= \int_\Omega [h(P_{\Omega_{\epsilon,P}}V) - h(V)] \frac{\partial P_{\Omega_{\epsilon,P}}V}{\partial \tau_{P_j}} \\ &= \int_\Omega h'(V)(P_{\Omega_{\epsilon,P}}V - V) \frac{\partial P_{\Omega_{\epsilon,P}}V}{\partial \tau_{P_j}} + h''(V)(P_{\Omega_{\epsilon,P}}V - V)^2 \frac{\partial P_{\Omega_{\epsilon,P}}V}{\partial \tau_{P_j}} + \mathcal{O}(\epsilon^{N+2}) \\ &= \epsilon \int_\Omega h'(V)(v_1 \chi + \epsilon(v_2 \chi + v_3 \chi) + \epsilon^2 \Psi_\epsilon) \left( \frac{\partial V}{\partial P_j} + w_1 + \epsilon w_2^\epsilon(x) \right) \\ &\quad + \epsilon^2 \int_\Omega h''(V)(v_1^2 \chi^2 + \epsilon(\Psi_\epsilon^1)^2) \frac{\partial P_{\Omega_{\epsilon,P}}V}{\partial \tau_{P_j}} + \mathcal{O}(\epsilon^{N+2}) \\ &= \epsilon^2 \int_\Omega h'(V)v_3 \frac{\partial V}{\partial P_j} + \mathcal{O}(\epsilon^{N+2}) \\ &= -\epsilon^{N+1} \left( \int_{\Omega_{\epsilon,P}} h'(V)v_3 \frac{\partial V}{\partial y_j} \right) + \mathcal{O}(\epsilon^{N+2}) \\ &= -\epsilon^{N+1} \int_{R_+^N} h'(V)v_3 \frac{\partial V}{\partial y_j} + \mathcal{O}(\epsilon^{N+2}). \end{aligned}$$

But

$$\begin{aligned} \int_{R_+^N} h'(V)v_3 \frac{\partial V}{\partial y_j} &= - \int_{R_+^N} \left( \Delta \frac{\partial V}{\partial y_j} - m \frac{\partial V}{\partial y_j} \right) v_3 \\ &= \int_{\partial R_+^N} \frac{\partial v_3}{\partial y_N} \frac{\partial V}{\partial y_j} - v_3 \frac{\partial}{\partial y_N} \frac{\partial V}{\partial y_j} \end{aligned}$$



$$\begin{aligned}
&= -\frac{1}{3} \int_{R^{N-1}} \left( \frac{V'}{|y|} \right)^2 \sum_{k,l,m=1}^{N-1} \rho_{klm}(0) y_k y_l y_m y_j dy \\
&= -\frac{1}{3} \int_{R^{N-1}} \left( \frac{V'}{|y|} \right)^2 \sum_{k,l,m=1}^{N-1} y_k y_l y_m y_j \rho_{klm}(0) dy \\
&= -\frac{1}{3} \int_{R^{N-1}} \left( \frac{V'}{|y|} \right)^2 y_j^2 \sum_{l,m=1}^{N-1} y_l y_m \rho_{jlm}(0) dy \\
&\quad = \nu \rho_{jkk}(0) \\
&\quad = \nu \nabla_j H(P)
\end{aligned}$$

where

$$\nu = - \sum_{k=1}^{N-1} \frac{1}{3} \int_{R^{N-1}} \left( \frac{V'}{|y|} \right)^2 y_j^2 y_k^2 dy \neq 0.$$

Combining  $I_\epsilon^1$ ,  $I_\epsilon^2$ ,  $J_\epsilon$ , we obtain

$$W_\epsilon(P) = \nu \nabla_{P_0} H(P) + W'_\epsilon(P)$$

where  $W'_\epsilon(P)$  is continuous in  $P$  and  $W'_\epsilon(P) = \mathcal{O}(\epsilon)$  uniformly in  $P$ . Suppose at  $P_0$ , we have  $\det(\nabla_j \nabla_k H(P_0)) \neq 0$  then standard Brouwer's fixed point theorem shows that for  $\epsilon \ll 1$  there exists a  $P_\epsilon$  such that  $W_\epsilon(P_\epsilon) = 0$ ,  $P_\epsilon \rightarrow P_0$ .

Thus we have proved the following proposition.

**Proposition 4.1.** *For  $\epsilon$  sufficiently small there exist points  $P_\epsilon$  with  $P_\epsilon \rightarrow P_0$  such that  $W_\epsilon(P_\epsilon) = 0$ .*

By Lemma 3.3 and Proposition 4.1 we have

$$S_\epsilon(v_\epsilon) = 0,$$

i.e.

$$\begin{aligned}
\epsilon^2 \Delta v_\epsilon - m v_\epsilon + h(u_\epsilon) - \frac{1}{|\Omega|} \int_\Omega h(v_\epsilon) &= 0 \quad \text{in } \Omega, \\
\frac{\partial v_\epsilon}{\partial \nu} &= 0 \quad \text{on } \partial\Omega.
\end{aligned}$$

Hence  $\int_\Omega v_\epsilon = 0$ . Let  $u_\epsilon = \bar{m} - v_\epsilon$ . We have

$$\begin{aligned}
\epsilon^2 \Delta u_\epsilon - f(u_\epsilon) &= \sigma_\epsilon, \\
\partial u_\epsilon / \partial \nu &= 0 \quad \text{on } \partial\Omega
\end{aligned}$$

$$\int_{\Omega} u_{\epsilon} = \bar{m}|\Omega|,$$

i.e.  $u_{\epsilon}$  is a solution of the Cahn-Hilliard equation. Moreover

$$\left\| v_{\epsilon} - V\left(\frac{x - P_{\epsilon}}{\epsilon}\right) \right\|_{\epsilon} \rightarrow 0$$

and  $P_{\epsilon} \rightarrow P_0 \in \partial\Omega$ .

Finally, we study the shape of the solutions  $v_{\epsilon}$ . Let  $P_{\epsilon}$  be any local maximum point of  $v_{\epsilon}$ . Then by (1.1),

$$mv_{\epsilon} - h(v_{\epsilon}) + \frac{1}{|\Omega|} \int_{\Omega} h(v_{\epsilon}) \leq 0.$$

But  $\epsilon^{-N} \int_{\Omega} h(v_{\epsilon}) \rightarrow \int_{R_+^N} h(V) > 0$ , hence

$$mv_{\epsilon} - h(v_{\epsilon}) < 0.$$

So  $v_{\epsilon}(P_{\epsilon}) \geq a_1 > 0$ . On the other hand, from our construction,

$$\|v_{\epsilon}\|_{\epsilon}^2 \rightarrow \frac{1}{2} \left( \int_{R^N} |\nabla V|^2 + mV^2 \right).$$

Similar proof as in Theorem 1.2 of [18], we conclude  $P_{\epsilon} \in \partial\Omega$  and there is only one such  $P_{\epsilon}$ .

## Appendix A: Trace Inequality

**Lemma A.1** *Let  $0 < \epsilon \leq 1$ . Then*

$$(A.1) \quad \|\Phi\|_{L^2(\partial\Omega_{\epsilon,P})} \leq C \|\Phi\|_{H^1(\Omega_{\epsilon,P})}$$

for all  $\Phi \in H^1(\Omega)$  where the constant  $C$  is independent of  $\epsilon$ .

Note that the constant  $C$  in (A.1) is required to be independent of  $\epsilon$ . Therefore Lemma A.1 is special although trace inequalities are quite standard.

*Proof of Lemma A.1.* For  $\Phi \in H^1(\Omega_{\epsilon,P})$  define  $\Psi \in H^1(\Omega)$  by a linear transformation:

$$\Psi(x) = \Phi(z) \quad \text{where } z = \frac{x - P}{\epsilon}.$$

Observe that  $\|\Phi\|_{L^2(\partial\Omega_{\epsilon,P})}^2 = \epsilon^{1-N} \|\Psi\|_{L^2(\partial\Omega)}^2$ ,  $\|\Phi\|_{L^2(\Omega_{\epsilon,P})}^2 = \epsilon^{-N} \|\Psi\|_{L^2(\Omega)}^2$ , and  $\|\nabla\Phi\|_{L^2(\Omega_{\epsilon,P})}^2 = \epsilon^{2-N} \|\nabla\Psi\|_{L^2(\Omega)}^2$ . Therefore (and after translation) (A.1) is equivalent to

$$(A.2) \quad \|\Psi\|_{L^2(\partial\Omega)}^2 \leq C(\epsilon \|\nabla\Psi\|_{L^2(\Omega)}^2 + \frac{1}{\epsilon} \|\Psi\|_{L^2(\Omega)}^2)$$

for all  $\Psi \in H^1(\Omega)$  and  $0 < \epsilon \leq 1$  where  $C$  is independent of  $\epsilon$ . The proof of (A.2) is standard and is omitted here (see for example the proof of Theorem 3.1 in [1]).  $\square$

### Appendix B: An elliptic regularity estimate

In this section we prove the following inequality

$$(B.1) \quad \|\Phi\|_{H^2(\Omega_{\epsilon,P})} \leq C(\|\Delta\Phi\|_{L^2(\Omega_{\epsilon,P})} + \|\Phi\|_{H^1(\Omega_{\epsilon,P})})$$

for all  $\Phi \in H_N^2(\Omega_{\epsilon,P})$ ,  $0 < \epsilon \leq \epsilon_0$  where  $\Omega_{\epsilon,P}$  is as defined in Section 2 and  $C$  is a constant independent of  $\epsilon$ . For a point  $P$  on  $\partial\Omega$  we can find a constant  $R_0 > 0$  and a smooth function  $\rho : B'(R_0) \rightarrow \mathbb{R}$  such that in  $B(P, R_0)$  the boundary  $\partial\Omega$  is described by the graph of  $\rho$  where  $\rho$  satisfies  $\rho(0) = 0$ ,  $\nabla\rho(0) = 0$  (compare Section 2). Furthermore there exists a map  $\eta = T(\xi)$  with  $DT(0) = I$  (the identity map) from a neighborhood  $U_P$  of  $P$  onto a ball  $B(0, R_1)$  (compare Section 3). By a linear transformation we naturally get a map  $T^\epsilon$  from  $U_P^\epsilon = \{(x - P)/\epsilon | x \in U_P\}$  onto a ball  $B(R_1/\epsilon)$  with center at 0. We set  $y = \eta/\epsilon$ . Then the Laplace operator becomes  $\epsilon^2\Delta_x = \Delta_y + A^\epsilon$  where

$$A^\epsilon = |\nabla_{x'}\rho|^2 \frac{\partial^2}{\partial y_N^2} - 2 \sum_{i=1}^{N-1} \rho_i \frac{\partial^2}{\partial y_i \partial y_N} - \epsilon \Delta_{x'} \rho \frac{\partial}{\partial y_N}.$$

Observe that for given  $\delta > 0$  we can find  $R_1 > 0$  and  $\epsilon_0$  such that for  $0 < \epsilon \leq \epsilon_0$

$$(B.2) \quad \|\nabla_{x'}\rho\|_{L^\infty(B(R_0/\epsilon))} \leq \delta, \|\rho\|_{L^\infty(B(R_0/\epsilon))} \leq \delta, \|\epsilon\Delta_{x'}\rho\|_{L^\infty(B(R_0/\epsilon))} \leq \delta.$$

In the same way we transform

$$\begin{aligned} \epsilon \frac{\partial}{\partial y_N} &= \{1 + |\nabla_{x'}\rho|^2\}^{-1/2} \left\{ \sum_{k=1}^{N-1} \rho_k \frac{\partial}{\partial y_k} - (1 + |\nabla_{x'}\rho|^2) \frac{\partial}{\partial y_N} \right\} \\ &= -\frac{\partial}{\partial y_N} + B^\epsilon \end{aligned}$$

where  $B^\epsilon$  is a differential operator on  $B(R_1/\epsilon) \cup \{y_N = 0\}$  with coefficients which are bounded in  $L^\infty$  for  $0 < \epsilon \leq \epsilon_0$  (compare section 2). From  $\{U_P | P \in \partial\Omega\}$  we select a finite subcovering of  $\partial\Omega$  and denote it by  $\{U_1, \dots, U_n\}$ . Choosing  $U_0 = \Omega$  the set  $\{U_0, \dots, U_n\}$  is a finite covering of  $\bar{\Omega}$  consisting of open sets. We keep this covering fixed from now on. Let  $\{\theta_0, \dots, \theta_n\}$  be a partition of unity subordinate to this open covering. Denote  $\theta_i^\epsilon(y) =$

$\theta_i \circ T^{-1}(\epsilon y)$ . Since

$$u = \sum_{i=0}^n \theta_i^\epsilon u$$

we have

$$(B.3) \quad \|u\|_{H^2(\Omega_{\epsilon,P})}^2 \leq \|\theta_0^\epsilon u\|_{H^2(\Omega_{\epsilon,P})}^2 + \sum_{i=1}^n \|\theta_i^\epsilon u\|_{H^2(\Omega_{\epsilon,P})}^2.$$

Since  $\theta_0^\epsilon$  has compact support in  $R^N$

$$\|\theta_0^\epsilon u\|_{H^2(R^N)}^2 = \|\Delta(\theta_0^\epsilon u)\|_{L^2(R^N)}^2 + \|\theta_0^\epsilon u\|_{H^1(R^N)}^2$$

(see for example [10], Corollary 9.10). Because of

$$\Delta(\theta_0^\epsilon) = \theta_0^\epsilon \Delta u + 2\nabla u \cdot \theta_0^\epsilon + u \Delta \theta_0^\epsilon$$

and

$$\|\nabla \theta_0^\epsilon\|_{L^\infty(R^N)} \leq C\epsilon, \quad \|\Delta \theta_0^\epsilon\|_{L^\infty(R^N)} \leq C\epsilon^2,$$

we obtain

$$(B.4) \quad \|\theta_0^\epsilon u\|_{H^2(\Omega_{\epsilon,P})}^2 \leq C(\|\theta_0^\epsilon \Delta u\|_{L^2(\Omega_{\epsilon,P})}^2 + \|u\|_{H^1(\Omega_{\epsilon,P})}^2).$$

We are now going to estimate  $\theta_i^\epsilon u$ ,  $i = 1, \dots, n$ . Note that

$$(B.5) \quad \frac{1}{C} \|(\theta_i^\epsilon u)^*\|_{H^k(R_+^N)} \leq \|\theta_i^\epsilon u\|_{H^k(\Omega_{\epsilon,P})} \leq C \|(\theta_i^\epsilon u)^*\|_{H^k(R_+^N)}$$

where  $k = 0, 1$ , or  $2$  and

$$v^*(y) \equiv v\left(\frac{1}{\epsilon} T^{-1}(\epsilon y)\right)$$

for  $v \in H^2(U_i^\epsilon)$ . Then

$$(B.6) \quad \|(\theta_i^\epsilon u)^*\|_{H^2(R_+^N)}^2 \leq \overline{C} \left( \|\Delta(\theta_i^\epsilon u)^*\|_{L^2(R_+^N)}^2 + \left\| \frac{\partial}{\partial y_N} (\theta_i^\epsilon u)^* \right\|_{H^{1/2}(R^{N-1} \times \{0\})}^2 + \|(\theta_i^\epsilon u)^*\|_{H^1(R_+^N)}^2 \right)$$

(see for example [15], Theorem 4.1). Now (B.2) implies that

$$\|A^\epsilon(\theta_i^\epsilon u)^*\|_{L^2(R_+^N)}^2 \leq \delta^2 \|(\theta_i^\epsilon u)^*\|_{H^1(R_+^N)}^2.$$

Therefore from (B.6)

$$\begin{aligned} & (1 - \overline{C}\delta^2) \|(\theta_i^\epsilon u)^*\|_{H^2(R_+^N)}^2 \\ & \leq \overline{C} \left( \|(\Delta + A^\epsilon)(\theta_i^\epsilon u)^*\|_{L^2(R_+^N)}^2 + \left\| \frac{\partial}{\partial y_N} (\theta_i^\epsilon u)^* \right\|_{H^{1/2}(R^{N-1} \times \{0\})}^2 + \|(\theta_i^\epsilon u)^*\|_{H^1(R_+^N)}^2 \right). \end{aligned}$$

For the operator  $B^\epsilon$  we can calculate in an analogous way. The trace theorem implies

$$\begin{aligned} & (1 - \tilde{C}\delta^2) \|(\theta_i^\epsilon u)^*\|_{H^2(\mathbb{R}_+^N)}^2 \\ & \leq C \left( \|(\Delta + A^\epsilon)(\theta_i^\epsilon u)^*\|_{L^2(\mathbb{R}_+^N)}^2 + \left\| \left( \frac{\partial}{\partial y_N} + B^\epsilon \right) (\theta_i^\epsilon u)^* \right\|_{H^{1/2}(\mathbb{R}^{N-1} \times \{0\})}^2 \right. \\ & \quad \left. + \|(\theta_i^\epsilon u)^*\|_{H^1(\mathbb{R}_+^N)}^2 \right). \end{aligned}$$

Since  $\tilde{C}$  is by construction independent of  $\epsilon$  we can choose  $\delta$  so small that  $1 - \tilde{C}\delta^2 \geq 1/2$ . This implies

$$\begin{aligned} \text{(B.7)} \quad \|\theta_i^\epsilon u\|_{H^2(\Omega_{\epsilon,P})}^2 & \leq C \left( \|\Delta(\theta_i^\epsilon u)\|_{L^2(\Omega_{\epsilon,P})}^2 + \left\| \frac{\partial}{\partial \nu_\epsilon}(\theta_i^\epsilon u) \right\|_{H^{1/2}(\partial\Omega_{\epsilon,P})}^2 \right. \\ & \quad \left. + \|\theta_i^\epsilon u\|_{H^1(\Omega_{\epsilon,P})}^2 \right). \end{aligned}$$

Similarly as before

$$\text{(B.8)} \quad \|\Delta(\theta_i^\epsilon u)\|_{L^2(\Omega_{\epsilon,P})}^2 \leq C(\|\theta_i^\epsilon \Delta u\|_{L^2(\Omega_{\epsilon,P})}^2 + \|u\|_{H^1(\Omega_{\epsilon,P})}^2)$$

and

$$\text{(B.9)} \quad \left\| \frac{\partial}{\partial \nu_\epsilon}(\theta_i^\epsilon u) \right\|_{H^{1/2}(\partial\Omega_{\epsilon,P})}^2 \leq C\|u\|_{H^1(\Omega_{\epsilon,P})}^2$$

because of  $\partial u / \partial \nu_\epsilon = 0$ . Combining (B.7) - (B.9) we get

$$\text{(B.10)} \quad \|\theta_i^\epsilon u\|_{H^2(\Omega_{\epsilon,P})}^2 \leq C(\|\theta_i^\epsilon \Delta u\|_{L^2(\Omega_{\epsilon,P})}^2 + \|u\|_{H^1(\Omega_{\epsilon,P})}^2).$$

We conclude, using (B.3), (B.4) and (B.10), that

$$\begin{aligned} \|u\|_{H^2(\Omega_{\epsilon,P})}^2 & \leq C \left( \sum_{i=0}^n \|\theta_i^\epsilon \Delta u\|_{L^2(\Omega_{\epsilon,P})}^2 + (n+1)\|u\|_{H^1(\Omega_{\epsilon,P})}^2 \right) \\ & \leq C_n(\|\Delta u\|_{L^2(\Omega_{\epsilon,P})}^2 + \|u\|_{H^1(\Omega_{\epsilon,P})}^2) \end{aligned}$$

where  $C_n$  depends on  $n$ . Since  $n$  is independent of  $\epsilon$  the proof of (B.1) is finished.  $\square$

## REFERENCES

- [1] S. Agmon, *Lectures on Elliptic Boundary Value Problems*, Von Nostrand, Princeton, 1965.
- [2] Alikakos, N., Bates, P.W., and Chen, X., Convergence of the Cahn-Hilliard equation to the Hele-Shaw model, *Arch. Rat. Mech. Anal.*, 1994, Vol. **128**, pp. 165-205.
- [3] Alikakos, N., Bates, P.W. and Fusco, G., Slow motion for the Cahn-Hilliard equation in one space dimension, *J. Diff. Eqns.*, 1991, Vol. **90**, pp. 81-134.
- [4] Bates, P.W. and Fife, P.C., The dynamics of nucleation for the Cahn-Hilliard equation, *SIAM J. Appl. Math.*, 1993, Vol. **53**, pp. 990-1008.

- [5] Cahn, J.W. and Hilliard, J.E., Free energy of a nonuniform system, I. Interfacial free energy, *J. Chem. Phys.*, 1958, Vol **28**, pp. 258-267.
- [6] Chen, X. and Kowalczyk, M., Existence of equilibria for the Cahn-Hilliard equation via local minimizers of the perimeter, preprint.
- [7] Dancer, E.N., A note on asymptotic uniqueness for some nonlinearities which change sign, preprint.
- [8] Floer, A. and Weinstein, A., Nonspreading wave packets for the cubic Schrödinger equation with a bounded potential, 1986, *J. Funct. Anal.*, Vol. **69**, pp. 397-408.
- [9] Gidas, B., Ni, W.-M., and Nirenberg, L., Symmetry of positive solutions of nonlinear elliptic equations in  $R^n$ , *Mathematical Analysis and Applications*, Part A, Adv. Math. Suppl. Studies Vol. **7A**, pp. 369-402, Academic Press, New York, 1981.
- [10] Gilbarg, D. and Trudinger, N.S., *Elliptic Partial Differential Equations of Second Order*, 2nd edition, Springer, Berlin, 1983.
- [11] Grinfeld, M. and Novick-Cohen, A., Counting stationary solutions of the Cahn-Hilliard equation by transversality arguments, *Proc. Roy. Soc. Edinburgh Sect. A*, 1995, Vol. **125**, pp. 351-370.
- [12] Helffer, B. and Sjöstrand, J., Multiple wells in the semi-classical limit I, *Comm. PDE*, 1984, Vol. **9**, pp. 337-408.
- [13] Kohn, R.V. and Sternberg, P., Local minimizers and singular perturbations, *Proc. Roy. Soc. Edinburgh Sect. A*, 1989, Vol. **111**, pp. 69-84.
- [14] Lin, C.-S., Ni, W.-M., and Takagi, I., Large amplitude stationary solutions to a chemotaxis systems, *J. Diff. Eqns.*, 1988, Vol. **72**, pp. 1-27.
- [15] Lions, J.L. and Magenes, E., *Non-Homogeneous Boundary Value Problems and Applications*, Vol **I**, Springer-Verlag, New York/Heidelberg/Berlin, 1972.
- [16] Modica, L., The gradient theory of phase transitions and the minimal interface criterion, *Arch. Rational Mech. Anal.*, 1989, Vol. **107**, 71-83.
- [17] Ni, W.-M., Pan, X., and Takagi, I., Singular behavior of least-energy solutions of a semilinear Neumann problem involving critical Sobolev exponents, *Duke Math. J.*, 1992, Vol. **67**, pp. 1-20.
- [18] Ni, W.-M. and Takagi, I., On the shape of least energy solution to a semilinear Neumann problem, *Comm. Pure Appl. Math.*, 1991, Vol. **41**, pp. 819-851.
- [19] Ni, W.-M. and Takagi, I., Locating the peaks of least energy solutions to a semilinear Neumann problem, *Duke Math. J.*, 1993, Vol. **70**, pp. 247-281.
- [20] Ni, W.-M. and Wei, J., On the location and profile of spike-layer solutions to singularly perturbed semilinear Dirichlet problems, *Comm. Pure Appl. Math.*, 1995, Vol. **48**, pp. 731-768.
- [21] Oh, Y.G., Existence of semi-classical bound states of nonlinear Schrödinger equations with potentials of the class  $(V)_a$ , 1988, *Comm. PDE*, Vol. **13(12)**, pp. 1499-1519.
- [22] Oh, Y.G., On positive multi-lump bound states of nonlinear Schrödinger equations under multiple-well potentials, 1990, *Comm. Math. Phys.*, Vol. **131**, pp. 223-253.
- [23] Pego, R.L., Front migration in the nonlinear Cahn-Hilliard equation, *Proc. Roy. Soc. London A*, 1989, Vol. **422**, pp. 261-278.
- [24] Peletier, L.A. and Serrin, J., Uniqueness of positive solutions of semilinear equations in  $R^n$ , *Arch. Rational Mech. Anal.*, 1983, Vol. **81**, pp. 181-197.

DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN,  
HONG KONG

MATHEMATISCHES INSTITUT A, UNIVERSITÄT STUTTGART, PFAFFENWALDRING 57,  
D-70569 STUTTGART, GERMANY