

Revised Version

MULTIPLE BOUNDARY PEAK SOLUTIONS FOR SOME SINGULARLY PERTURBED NEUMANN PROBLEMS

CHANGFENG GUI, JUNCHENG WEI, AND MATTHIAS WINTER

ABSTRACT. We consider the problem

$$\begin{cases} \varepsilon^2 \Delta u - u + f(u) = 0 & \text{in } \Omega \\ u > 0 \text{ in } \Omega, \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega, \end{cases}$$

where Ω is a bounded smooth domain in R^N , $\varepsilon > 0$ is a small parameter and f is a superlinear, subcritical nonlinearity. It is known that this equation possesses boundary spike solutions such that the spike concentrates, as ε approaches zero, at a critical point of the mean curvature function $H(P)$, $P \in \partial\Omega$. It is also known that this equation has multiple boundary spike solutions at multiple nondegenerate critical points of $H(P)$ or multiple local maximum points of $H(P)$.

In this paper, we prove that for any fixed positive integer K there exist boundary K -peak solutions at a local minimum point of $H(P)$. This implies that for any smooth and bounded domain there always exist boundary K -peak solutions.

We first use the Liapunov-Schmidt method to reduce the problem to finite dimensions. Then we use a maximizing procedure to obtain multiple boundary spikes.

1. INTRODUCTION

The aim of this paper is to construct a family of multiple boundary peak solutions to the following singularly perturbed elliptic problem

$$\begin{cases} \varepsilon^2 \Delta u - u + u^p = 0 & \text{in } \Omega, \\ u > 0 \text{ in } \Omega \text{ and } \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where $\Delta = \sum_{i=1}^N \frac{\partial^2}{\partial x_i^2}$ is the Laplace operator, Ω is a bounded smooth domain in R^N , $\varepsilon > 0$ is a constant, the exponent p satisfies $1 < p < \frac{N+2}{N-2}$ for $N \geq 3$ and $1 < p < \infty$ for $N = 2$ and $\nu(x)$ denotes the normal derivative at $x \in \partial\Omega$.

Equation (1.1) is known as the stationary equation of the Keller-Segal system in chemotaxis. It can also be seen as the limiting stationary equation

1991 *Mathematics Subject Classification.* Primary 35B40, 35B45; Secondary 35J40.

Key words and phrases. Multiple Boundary Spikes, Nonlinear Elliptic Equations.

of the so-called Gierer-Meinhardt system in biological pattern formation, see [33] for more details.

In the pioneering papers of [17], [20] and [21], Lin, Ni and Takagi established the existence of least-energy solutions and showed that for ε sufficiently small the least-energy solution has only one local maximum point P_ε and $P_\varepsilon \in \partial\Omega$. Moreover, $H(P_\varepsilon) \rightarrow \max_{P \in \partial\Omega} H(P)$ as $\varepsilon \rightarrow 0$, where $H(P)$ is the mean curvature of P at $\partial\Omega$. In [22], Ni and Takagi constructed boundary spike solutions for axially symmetric domains. The second author in [33] studied the general domain case and showed that for single boundary spike solutions, the boundary spike must approach a critical point of the mean curvature; on the other hand, for any nondegenerate critical point of $H(P)$, one can construct boundary spike solutions whose spike approaches that point. The first author in [11] constructed multiple boundary spike layer solutions at multiple local maximum points of $H(P)$ while the second and third authors in [36] constructed multiple boundary spike layer solutions at multiple nondegenerate critical points of $H(P)$. Later these results were improved by Y. Y. Li in [16] in a unified approach. When $p = \frac{N+2}{N-2}$, similar results for the boundary spike layer solutions have been obtained in [1], [2], [3], [12], [19], [26], [27], [28], [30] etc.

In this paper, we study the existence of multiple boundary peak solutions at a local minimum point of $H(P)$.

More precisely, we consider the problem

$$\begin{cases} \varepsilon^2 \Delta u - u + f(u) = 0 & \text{in } \Omega, \\ u > 0 & \text{in } \Omega \text{ and } \frac{\partial u}{\partial \nu} = 0 & \text{in } \partial\Omega. \end{cases} \quad (1.2)$$

We will assume that $f : R^+ \rightarrow R$ is of class $C^{1+\sigma}$ and satisfies the following conditions

- (f1) $f(t) \equiv 0$ for $t \leq 0$ and $f(t) \rightarrow +\infty$ as $t \rightarrow \infty$
- (f2) There exist some constants $1 < p_1, p_2, p_3 < \left(\frac{N+4}{N-4}\right)_+$ ($= \infty$ if $N \leq 4$; $= \frac{N+4}{N-4}$ if $N > 4$) such that $f(0) = 0, f'(0) = 0$ and

$$f(u) = O(|u|^{p_1}), f'(u) = O(|u|^{p_2-1}) \text{ as } |u| \rightarrow \infty,$$

$$|f_u(u + \phi) - f_u(u)| \leq \begin{cases} C|\phi|^{p_3-1} & \text{if } p_3 > 2 \\ C(|\phi| + |\phi|^{p_3-1}) & \text{if } p_3 \leq 2. \end{cases}$$

(f3) The equation

$$\begin{cases} \Delta w - w + f(w) = 0 & \text{in } R^N, \\ w > 0, w(0) = \max_{z \in R^N} w(z), \\ w \rightarrow 0 & \text{at } \infty \end{cases} \quad (1.3)$$

has a unique solution $w(y)$ (by the results of [9], w is radial, i.e., $w = w(r)$ and $w' < 0$ for $r = |y| \neq 0$) and w is nondegenerate. Namely the operator

$$L := \Delta - 1 + f'(w) \quad (1.4)$$

is invertible in the space $H_r^2(R^N) := \{u = u(|y|) \in H^2(R^N)\}$.

Two important examples of f are the following.

Example 1 (chemotaxis and pattern formation): $f(u) = u^p$ where $1 < p < (\frac{N+2}{N-2})_+ (= \infty \text{ if } N = 2; = \frac{N+2}{N-2} \text{ if } N > 2)$. It is easy to see that f satisfies (f1), (f2) and (f3). This problem arises from the Keller-Segal model in chemotaxis and the Gierer-Meinhardt system in pattern formation (see [20], [21] and the references therein).

Example 2 (population dynamics and chemical reaction theory): $f(u) = u(u - a)(1 - u)$ where $0 < a < \frac{1}{2}$. This is a famous model from population dynamics and chemical reaction theory (see [5], [14], [29]). If $N \leq 8$ then by the result of [8], f satisfies (f1)-(f3).

Other nonlinearities satisfying (f1), (f2) and (f3) can be found in [6].

Let $\Lambda \subset \partial\Omega$ be an open set such that

$$\min_{P \in \partial\Gamma} H(P) > \min_{P \in \Gamma} H(P). \quad (1.5)$$

We now state the main result in this paper.

Theorem 1.1. *Assume that condition (1.5) holds. Let f satisfy assumptions (f1)-(f3). Then for ε sufficiently small problem (1.2) has a solution u_ε which possesses exactly K local maximum points $Q_1^\varepsilon, \dots, Q_K^\varepsilon$ with $\mathbf{Q}^\varepsilon =$*

$(Q_1^\varepsilon, \dots, Q_K^\varepsilon) \in \Gamma \times \dots \times \Gamma$. Moreover $H(Q_i^\varepsilon) \rightarrow \min_{P \in \Gamma} H(P)$, $w(\frac{|Q_k^\varepsilon - Q_l^\varepsilon|}{\varepsilon}) \rightarrow 0$, $i, k, l = 1, \dots, K, k \neq l$ as $\varepsilon \rightarrow 0$. Furthermore, we have

$$u_\varepsilon(x) \leq a \exp\left(-\frac{b \min_{i=1, \dots, K} (|x - Q_i^\varepsilon|)}{\varepsilon}\right) \quad (1.6)$$

for certain positive constants a, b .

Theorem 1.1 can be derived from a more general theorem as follows.

Theorem 1.2. *Let $\Gamma_i, i = 1, \dots, K$ be open sets in $\partial\Omega$ such that*

$$\min_{P \in \partial\Gamma_i} H(P) > \min_{P \in \Gamma_i} H(P), i = 1, \dots, K.$$

Let f satisfy assumptions (f1)-(f3). Then for ε sufficiently small problem (1.2) has a solution u_ε which possesses exactly K local maximum points $Q_1^\varepsilon, \dots, Q_K^\varepsilon$ with $\mathbf{Q}^\varepsilon = (Q_1^\varepsilon, \dots, Q_K^\varepsilon) \in \Gamma_1 \times \dots \times \Gamma_K$. Moreover $H(Q_i^\varepsilon) \rightarrow \min_{P \in \Gamma_i} H(P)$, $w(\frac{|Q_k^\varepsilon - Q_l^\varepsilon|}{\varepsilon}) \rightarrow 0$, $i, k, l = 1, \dots, K, k \neq l$ as $\varepsilon \rightarrow 0$. Furthermore, we have

$$u_\varepsilon(x) \leq a \exp\left(-\frac{b \min_{i=1, \dots, K} (|x - Q_i^\varepsilon|)}{\varepsilon}\right) \quad (1.7)$$

for certain positive constants a, b .

More details about the asymptotic behavior of u_ε can be found in the proof of Theorem 1.2.

We have the following interesting corollary.

Corollary 1.3. *For any smooth and bounded domain and any fixed positive integer $K \in \mathbb{Z}$, there always exists a boundary K -peaked solution of (1.2) if ε is small enough.*

Theorem 1.1 is the first result in proving the existence of multiple boundary spike solutions for problem (1.2) in any smooth bounded domain. Note that the boundary spikes can approach the same point on the boundary when Λ has a strictly local minimum point of $H(P)$. This is new and interesting in its own right.

We shall only prove Theorem 1.2. To introduce the main idea of the proof of Theorem 1.2, we need to give some necessary notations and definitions first.

Let w be the unique solution of (1.3). It is known (see [9]) that w is radially symmetric, decreasing and

$$\lim_{|y| \rightarrow \infty} w(y) e^{|y|} |y|^{\frac{N-1}{2}} = c_0 > 0.$$

Associated with problem (1.2) is the following energy functional

$$J_\varepsilon(u) = \frac{1}{2} \int_\Omega (\varepsilon^2 |\nabla u|^2 + u^2) - \int_\Omega F(u)$$

where $F(u) = \int_0^u f(s) ds$ and $u \in H^1(\Omega)$.

For any smooth bounded domain U we set $P_U w$ to be the unique solution of

$$\begin{cases} \Delta u - u + f(w) = 0 & \text{in } U, \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } U. \end{cases} \quad (1.8)$$

Let $\eta > 0$ be a small number. Let Γ_i be as in Theorem 1.2. Set

$$\Lambda = \{\mathbf{P} = (P_1, \dots, P_K) \in \Gamma_1 \times \dots \times \Gamma_K, w\left(\frac{|P_k - P_l|}{\varepsilon}\right) < \eta\varepsilon, k, l = 1, \dots, K, k \neq l\}.$$

For $P \in \partial\Omega$, we set

$$\Omega_\varepsilon = \{y : \varepsilon y \in \Omega\}, \quad \Omega_{\varepsilon, P} = \{y : \varepsilon y + P \in \Omega\}.$$

Fix $\mathbf{P} = (P_1, P_2, \dots, P_K) \in \bar{\Lambda}$. We set

$$Pw_i(y) = P_{\Omega_{\varepsilon, P_i}} w\left(y - \frac{P_i}{\varepsilon}\right), \quad w_i(y) = w\left(y - \frac{P_i}{\varepsilon}\right), \quad y \in \Omega_\varepsilon,$$

$$u = \sum_{i=1}^K Pw_i + \Phi_{\varepsilon, \mathbf{P}} \in H^2(\Omega_\varepsilon),$$

$$\mathcal{K}_{\varepsilon, \mathbf{P}} = \text{span} \left\{ \frac{\partial Pw_i}{\partial \tau_{P_i, i_j}}, i = 1, \dots, K, j = 1, \dots, N-1 \right\} \subset H^2(\Omega_\varepsilon),$$

$$\mathcal{C}_{\varepsilon, \mathbf{P}} = \text{span} \left\{ \frac{\partial Pw_i}{\partial \tau_{P_i, i_j}}, i = 1, \dots, K, j = 1, \dots, N-1 \right\} \subset L^2(\Omega_\varepsilon)$$

where τ_{P_i, i_j} are the $(N-1)$ tangential derivatives at P_i (without loss of generality we assume that the inward normal derivative at P_i is e_N and denote τ_{P_i, i_j} as $\tau_{P_i, j}$ in the rest of the paper.)

We first solve for $\Phi_{\varepsilon, \mathbf{P}}$ in $\mathcal{K}_{\varepsilon, \mathbf{P}}^\perp$ up to $\mathcal{C}_{\varepsilon, \mathbf{P}}^\perp$ by using the Liapunov-Schmidt reduction method. This method evolves from that of [7], [24] and [25] on

the semi-classical (i.e. for small parameter h) solution of the nonlinear Schrödinger equation

$$\frac{h^2}{2}\Delta U - (V - E)U + U^p = 0 \quad (1.9)$$

in R^N where V is a potential function and E is a real constant. The method of Liapunov-Schmidt reduction was used in [7], [24] and [25] to construct solutions of (1.9) close to nondegenerate critical points of V for h sufficiently small.

Then we show that $\Phi_{\varepsilon, \mathbf{P}}$ is C^1 in \mathbf{P} . After that, we define a new function

$$M_\varepsilon(\mathbf{P}) = J_\varepsilon\left(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon, \mathbf{P}}\right). \quad (1.10)$$

We maximize $M_\varepsilon(\mathbf{P})$ over $\bar{\Lambda}$. Condition (1.5) ensures that $M_\varepsilon(\mathbf{P})$ attains its maximum inside Λ . We show that the resulting solution has the properties of Theorem 1.2.

The paper is organized as follows. Notation, preliminaries and some useful estimates are explained in Section 2. Section 3 contains the setup of our problem and we solve (1.2) up to approximate kernel and cokernel, respectively. We set up and solve a maximizing problem in Section 4. Finally, in Section 5, we show that the solution to the maximizing problem is indeed a solution of (1.2) and satisfies all the properties of Theorem 1.2.

Throughout this paper, unless otherwise stated, the letter C will always denote various generic constants which are independent of ε , for ε sufficiently small. $\delta > 0$ is a very small number. $o(1)$ means $|o(1)| \rightarrow 0$ as $\varepsilon \rightarrow 0$.

Acknowledgement. The research of the second author is supported by an Earmarked Grant from RGC of Hong Kong.

2. TECHNICAL ANALYSIS

In this section we introduce a projection and derive some useful estimates. Throughout the paper we shall use the letter C to denote a generic positive constant which may vary from term to term. We denote $R_+^N = \{(x', x_N) | x_N > 0\}$. Let w be the unique solution of (1.3).

Set

$$I(w) = \frac{1}{2} \int_{R^N} (|\nabla w|^2 + w^2) - \int_{R^N} F(w),$$

Let $P \in \partial\Omega$. We can define a diffeomorphism straightening the boundary in a neighborhood of P . After rotation of the coordinate system we may assume that the inward normal to $\partial\Omega$ at P is pointing in the direction of the positive x_N -axis. Denote $x' = (x_1, \dots, x_{N-1})$, $B'(R_0) = \{x' \in R^{N-1} \mid |x'| < R_0\}$, $B(P, R_0) = \{x \in R^N \mid |x - P| < R_0\}$, and $\Omega_0 = \Omega \cap B(P, R_0) = \{(x', x_N) \in B(P, R_0) \mid x_N - P_N > \rho(x' - P')\}$. Then, since $\partial\Omega$ is smooth, we can find a constant $R_0 > 0$ such that $\partial\Omega \cap \overline{\Omega_0}$ can be represented by the graph of a smooth function $\rho_P : B'(R_0) \rightarrow R$ where $\rho_P(0) = 0$, $\nabla \rho_P(0) = 0$.

From now on we omit the use of P in ρ_P and write ρ instead if this can be done without causing confusion. The mean curvatures of $\partial\Omega$ at P is $H(P) = \frac{1}{n-1} \sum_{i=1}^{N-1} \rho_{ii}(0)$ where

$$\rho_i = \frac{\partial \rho}{\partial x_i}, \quad i = 1, \dots, N-1$$

and higher derivatives are defined in the same way. By Taylor expansion we have

$$\begin{aligned} \rho(x' - P') &= \frac{1}{2} \sum_{i,j=1}^{N-1} \rho_{ij}(0)(x_i - P_i)(x_j - P_j) \\ &+ \frac{1}{6} \sum_{i,j,k=1}^{N-1} \rho_{ijk}(0)(x_i - P_i)(x_j - P_j)(x_k - P_k) + O(|x' - P'|^4) \end{aligned}$$

Recall that for a smooth bounded domain U the *projection* P_U of $H^2(U)$ onto $\{v \in H^2(U) \mid \partial v / \partial \nu = 0 \text{ at } \partial U\}$ is defined as follows: For $v \in H^2(U)$ let $\omega = P_U v$ be the unique solution of the boundary value problem

$$\begin{cases} \Delta \omega - \omega + f(v) = 0 & \text{in } U, \\ \frac{\partial \omega}{\partial \nu} = 0 & \text{on } \partial U. \end{cases} \quad (2.1)$$

Let $h_{\varepsilon, P}(x) = w\left(\frac{x-P}{\varepsilon}\right) - P_{\Omega_{\varepsilon, P}} w\left(\frac{x-P}{\varepsilon}\right)$, $x \in \Omega$ where

$$\Omega_{\varepsilon, P} = \{z \in R^n \mid \varepsilon z + P \in \Omega\}.$$

Then $h_{\varepsilon, P}$ satisfies

$$\begin{cases} \varepsilon^2 \Delta v - v = 0 & \text{in } \Omega, \\ \frac{\partial v}{\partial \nu} = \frac{\partial}{\partial \nu} w\left(\frac{x-P}{\varepsilon}\right) & \text{on } \partial\Omega. \end{cases} \quad (2.2)$$

We denote

$$\|v\|_\varepsilon^2 = \varepsilon^{-N} \int_\Omega [\varepsilon^2 |\nabla v|^2 + v^2].$$

For $x \in \Omega_0$ set now

$$\begin{cases} \varepsilon y' = x' - P', \\ \varepsilon y_N = x_N - P_N - \rho(x' - P'). \end{cases} \quad (2.3)$$

Furthermore, for $x \in \Omega_0$ we introduce the transformation T by

$$\begin{cases} T_i(x') = x_i, & i = 1, \dots, N-1, \\ T_N(x') = x_N - P_N - \rho(x' - P'). \end{cases} \quad (2.4)$$

Note that then

$$y = \frac{1}{\varepsilon} T(x).$$

Let v_1 be the unique solution of

$$\begin{cases} \Delta v - v = 0 & \text{in } R_+^N, \\ \frac{\partial v}{\partial y_N} = -\frac{w'}{|y|} \frac{1}{2} \sum_{i,j=1}^{N-1} \rho_{ij}(0) y_i y_j & \text{on } \partial R_+^N \end{cases} \quad (2.5)$$

where w' is the radial derivative of w , i.e. $w' = w_r(r)$, and $r = \left| \frac{x-P}{\varepsilon} \right|$.

Note that v_1 is an even functions in $y' = (y_1, \dots, y_{N-1})$. Moreover, it is easy to see that $|v_1| \leq C e^{-\mu|y|}$ for some $0 < \mu < 1$.

Let $\chi(x)$ be a smooth cutoff function such that $\chi(x) = 1$, $x \in B(0, 0.8R_0)$ and $\chi(x) = 0$ for $x \in B(0, R_0)^C$.

In fact we set R_0 be such that $w(\frac{R_0}{\varepsilon}) = 0.9\eta\varepsilon$.

Note that this χ is as good as the cut-off function in [33].

Set

$$h_{\varepsilon,P}(x) = \varepsilon v_1(y) \chi(x - P) + \varepsilon^2 \Psi_{\varepsilon,P}(x), x \in \Omega.$$

Then we have

Proposition 2.1.

$$\|\Psi_{\varepsilon,P}\|_\varepsilon \leq C.$$

Proof. Proposition 2.1 was proved in [35] by Taylor expansion and a rigorous estimate for the remainder using estimates for elliptic partial differential equations. \square

Similarly, we know from [35] that

Proposition 2.2.

$$\left[\frac{\partial w}{\partial \tau_{P_j}} - \frac{\partial P_{\Omega_\varepsilon, P} w}{\partial \tau_{P_j}} \right] \left(\frac{x - P}{\varepsilon} \right) = w_1(y) \chi(x - P) + \varepsilon w_2^\varepsilon(x), x \in \Omega$$

where $\varepsilon y = T(x)$ and w_1 satisfies

$$\begin{cases} \Delta v - v = 0 & \text{in } R_+^N, \\ \frac{\partial v}{\partial y_N} = -\frac{1}{2} \left(\frac{w''}{|y|^2} - \frac{w'}{|y|^3} \right) \sum_{k,l=1}^{N-1} \rho_{kl}(0) y_k y_l y_j - \frac{w'}{|y|} \sum_{k=1}^{N-1} \rho_{jk}(0) y_k & \text{on } \partial R_+^N \end{cases} \quad (2.6)$$

and

$$\|w_2^\varepsilon\|_\varepsilon \leq C.$$

Note that $|w_1| \leq C \exp(-\mu|y|)$ for some $\mu < 1$ and w_1 is an odd function in y' . Finally, let

$$L_0 = \Delta - 1 + f'(w).$$

We have

Lemma 2.3.

$$\text{Ker}(L_0) \cap H_N^2(R_+^N) = \text{span} \left\{ \frac{\partial w}{\partial y_1}, \dots, \frac{\partial w}{\partial y_{N-1}} \right\}.$$

where $H_N^2(R_+^N) = \{u \in H^2(R_+^N), \frac{\partial u}{\partial y_N} = 0 \text{ on } \partial R_+^N\}$.

Proof. See Lemma 4.2 in [21]. \square

Next we state some useful lemmas about the interactions of two w 's.

Lemma 2.4. *Let $\mathbf{P} = (P_1, \dots, P_K) \in \bar{\Lambda}$. Then we have*

$$\int_{\Omega_\varepsilon} f(w_k) w_l = (\gamma_{kl} + o(1)) w \left(\frac{|P_k - P_l|}{\varepsilon} \right), k, l = 1, \dots, K, k \neq l \quad (2.7)$$

where $\gamma_{kl} \in \Sigma$ and Σ is defined as follows

$$\Sigma = \left\{ \int_{R_+^N} f(w(y)) e^{\langle b, y \rangle} dy \mid b \in R^N, |b| = 1 \right\}. \quad (2.8)$$

Furthermore, if $w \left(\frac{|P_k - P_l|}{\varepsilon} \right) = \eta \varepsilon$, we have $\gamma_{kl} \in \Sigma_1$ where

$$\Sigma_1 = \left\{ \int_{R_+^N} f(w(y)) e^{\langle b, y \rangle} dy \mid b = (b_1, \dots, b_N) \in R^N, b_N = 0, |b| = 1 \right\}. \quad (2.9)$$

Proof: Note that as $\frac{|P_k - P_l|}{\varepsilon} \rightarrow \infty$ we have

$$w\left(\frac{|P_k - P_l|}{\varepsilon}\right) = \left(\frac{|P_k - P_l|}{\varepsilon}\right)^{-\frac{N-1}{2}} e^{-\frac{|P_k - P_l|}{\varepsilon}} (1 + o(1)). \quad (2.10)$$

Hence if we straighten the boundary at P_k we have

$$\begin{aligned} \int_{\Omega_\varepsilon} f(w_k) w_l &= \int_{R_+^N} f(w(y)) w\left(y - \frac{P_l - P_k}{\varepsilon}\right) (1 + o(1)) \\ &= w\left(\frac{|P_k - P_l|}{\varepsilon}\right) (1 + o(1)) \int_{R_+^N} f(w(y)) w\left(y - \frac{P_l - P_k}{\varepsilon}\right) w^{-1}\left(\frac{|P_k - P_l|}{\varepsilon}\right) \\ &= w\left(\frac{|P_k - P_l|}{\varepsilon}\right) (1 + o(1)) \int_{R_+^N} f(w(y)) e^{\langle b, y \rangle} dy \end{aligned}$$

for some $b = \lim_{\varepsilon \rightarrow \frac{|P_k - P_l|}{\varepsilon}} \frac{P_k - P_l}{|P_k - P_l|} \in R^N$, $|b| = 1$.

Note that if $w\left(\frac{|P_k - P_l|}{\varepsilon}\right) = \eta\varepsilon$, we have $P_k \rightarrow P_l$ and $b_N = 0$ where b_N is the N -th component of b . □

Note: $\gamma_{kl} = \gamma_{lk}$.

Next we are going to show three technical lemmas.

The first lemma is about some relations of several integrals associated with w in R_{N-1} .

Let

$$\gamma_1 = \frac{1}{N+1} \int_{R^{N-1}} |\nabla w|^2 |y'|^2 dy', \quad (2.11)$$

We have

Lemma 2.5.

$$\frac{N-3}{2} \gamma_1 = \int_{R^{N-1}} F(w(|y'|)) |y'|^2 dy' - \frac{1}{2} \int_{R^{N-1}} |w|^2 |y'|^2 dy' \quad (2.12)$$

$$\begin{aligned} (N+1) \gamma_1 &= \frac{N-1}{2} \int_{R^{N-1}} |w|^2 dy' - \int_{R^{N-1}} |w|^2 |y'|^2 dy' \\ &\quad + \int_{R^{N-1}} f(w) w |y'|^2 dy'. \end{aligned} \quad (2.13)$$

Proof:

Let $y = (y', y_N)$. The operators Δ and ∇ below are with respect to $y \in R^N$, and the integrations are with respect to $y' = (y', 0) \in R^{N-1}$. We will also use r for $|y'|$.

By straightforward computations we have

$$\begin{aligned}
& \int_{R^{N-1}} |y'|^2 \Delta w (\nabla w \cdot y) dy' \\
&= \omega_{N-2} \int_0^\infty (w''(r) + \frac{N-1}{r} w'(r)) w'(r) r^{N+1} dr \\
&= \frac{N-3}{2} \omega_{N-2} \int_0^\infty r^N (w'(r))^2 dr \\
&= \frac{(N+1)(N-3)}{2} \gamma_1,
\end{aligned} \tag{2.14}$$

and

$$\begin{aligned}
& \int_{R^{N-1}} |y'|^2 w (\nabla w \cdot y) dy' \\
&= \omega_{N-2} \int_0^\infty w'(r) w(r) r^{N+1} dr \\
&= -\frac{N+1}{2} \omega_{N-2} \int_0^\infty r^N w^2 dr \\
&= -\frac{N+1}{2} \int_{R^{N-1}} w^2 |y'|^2 dy'
\end{aligned} \tag{2.15}$$

and

$$\begin{aligned}
& \int_{R^{N-1}} |y'|^2 f(w) (\nabla w \cdot y) dy' \\
&= \omega_{N-2} \int_0^\infty f(w) w'(r) r^{N+1} dr \\
&= -(N+1) \omega_{N-2} \int_0^\infty r^N F(w) dr \\
&= -(N+1) \int_{R^{N-1}} F(w) |y'|^2 dy'.
\end{aligned} \tag{2.16}$$

Since w satisfies

$$\Delta w - w + f(w) = 0, y \in R^N \tag{2.17}$$

by multiplying (2.17) by $|y'|^2 (\nabla w \cdot y)$ and integrating it with respect to y' in R^{N-1} we obtain (2.12).

$$\begin{aligned}
\int_{R^{N-1}} |y'|^2 w \Delta w dy' &= \omega_{N-2} \int_0^\infty w(r) (w''(r) + \frac{N-1}{r} w'(r)) r^N dr \\
&= -\omega_{N-2} \int_0^\infty r^N (w'(r))^2 + \frac{N-1}{2} \omega_{N-2} \int_0^\infty r^{N-2} w^2 dr \\
&= -(N+1) \gamma_1 + \frac{N-1}{2} \int_{R^{N-1}} w^2 dy'.
\end{aligned}$$

Multiply (2.17) by $|y'|^2 w$ and integrate it in R^{N-1} . Then (2.13) is derived.

This proves Lemma 2.5. □

Lemma 2.6. *For any function $G(t)$ in $C^{1+\sigma}([0, \infty))$ with $G(0) = G'(0) = 0$, we have*

$$\int_{\Omega_{\varepsilon, P}} G(w(y)) dy = \int_{R_+^N} G(w(y)) dy - \varepsilon H(P) \frac{1}{2} \int_{R^{N-1}} G(w(y', 0)) |y'|^2 dy' + o(\varepsilon).$$

Proof:

Since w decays exponentially in y at infinity, we have

$$\begin{aligned} & \int_{\Omega_{\varepsilon, P}} G(w(y)) dy = \int_{(\Omega_0)_{\varepsilon, P}} G(w(y)) dy + o(\varepsilon) \\ &= \int_{B^+(\frac{R_0}{\varepsilon})} G(w(y)) dy - \int_{B^+(\frac{R_0}{\varepsilon}) \setminus (\Omega_0)_{\varepsilon, P}} G(w(y)) dy + o(\varepsilon) \\ &= \int_{R_+^N} G(w(y)) dy - \int_{|y'| \leq \frac{R_0}{\varepsilon}} \int_0^{\frac{1}{\varepsilon} \rho(\varepsilon y')} G(w(y', y_N)) dy_N dy' + o(\varepsilon) \\ &= \int_{R_+^N} G(w(y)) dy - \int_{|y'| \leq \frac{R_0}{\varepsilon}} \int_0^{\frac{1}{\varepsilon} \rho(\varepsilon y')} G(w(y', 0)) dy_N dy' \\ &\quad + \int_{|y'| \leq \frac{R_0}{\varepsilon}} \int_0^{\frac{1}{\varepsilon} \rho(\varepsilon y')} (G(w(y', y_N)) - G(w(y', 0))) dy_N dy' + o(\varepsilon) \\ &= \int_{R_+^N} G(w(y)) dy - \int_{|y'| \leq \frac{R_0}{\varepsilon}} G(w(y', 0)) \frac{\rho(\varepsilon y')}{\varepsilon} dy' \\ &\quad + \int_{|y'| \leq \frac{R_0}{\varepsilon}} O\left(|w(y', 0)|^\sigma \left(\frac{\rho(\varepsilon y')}{\varepsilon}\right)^2\right) dy' + o(\varepsilon) \\ &= \int_{R_+^N} G(w(y)) dy - \frac{1}{2} \varepsilon \int_{|y'| \leq \frac{R_0}{\varepsilon}} G(w(y', 0)) \sum_{i, j=1}^{N-1} \rho_{ij}(0) y_i y_j dy' \\ &\quad + \int_{|y'| \leq \frac{R_0}{\varepsilon}} O(\varepsilon^2 |y'|^3) dy' + o(\varepsilon) \\ &= \int_{R_+^N} G(w(y)) dy - \varepsilon H(P) \frac{1}{2} \int_{|y'| \leq \frac{R_0}{\varepsilon}} G(w(y', 0)) |y'|^2 dy' + o(\varepsilon) \\ &= \int_{R_+^N} G(w(y)) dy - \varepsilon H(P) \frac{1}{2} \int_{R^{N-1}} G(w(y', 0)) |y'|^2 dy' + o(\varepsilon) \quad (2.18) \end{aligned}$$

where

$$B^+\left(\frac{R_0}{\varepsilon}\right) = B\left(\frac{R_0}{\varepsilon}\right) \cap R_+^N$$

and

$$(\Omega_0)_{\varepsilon, P} = \{y | \varepsilon y + P \in \Omega_0\}.$$

Hence Lemma 2.6 is proven. □

Lemma 2.7.

$$\int_{R_+^N} f(w)(P_{\Omega_{\varepsilon,P}} w - w) = \varepsilon H(P) \frac{N-1}{4} \int_{R^{N-1}} |w|^2 dy' + o(\varepsilon).$$

Proof:

Using (2.5), (2.17) and the exponential decay of w and v_1 , we have

$$\begin{aligned} \int_{R_+^N} f(w(y))v_1(y)dy &= \int_{R_+^N} (w - \Delta w)v_1(y)dy \\ &= \int_{R_+^N} (w(v_1 - \Delta v_1) + \int_{R^{N-1}} (v_1 \frac{\partial w}{\partial y_N} - w \frac{\partial v_1}{\partial y_N})dy') \\ &= \frac{1}{2} \int_{R^{N-1}} w(r)w'(r)r^{-1} \sum_{i,j=1}^{N-1} \rho_{ij}(0)y_i y_j dy' \\ &= \frac{1}{2} \int_{R^{N-1}} w(r)w'(r)r^{-1} \sum_{i=1}^{N-1} \rho_{ii}(0)|y_i|^2 dy' \\ &= \frac{1}{2} H(P) \omega_{N-2} \int_0^\infty w(r)w'(r)r^{N-1} dr \\ &= -\frac{N-1}{4} H(P) \int_{R^{N-1}} w^2 dy'. \end{aligned} \tag{2.19}$$

In view of Proposition 2.1, Lemma 2.7 follows immediately. □

The next lemma is the key result in this section.

Lemma 2.8. *For any $\mathbf{P} = (P_1, \dots, P_K) \in \bar{\Lambda}$ and ε sufficiently small*

$$\begin{aligned} J_\varepsilon(\sum_{i=1}^K P w_i) &= \varepsilon^N \left[\frac{K}{2} I(w) - \varepsilon(\gamma_1 + o(1)) \sum_{i=1}^K H(P_i) \right. \\ &\quad \left. - \frac{1}{2} \sum_{k,l=1, k \neq l}^K (\gamma_{kl} + o(1)) w\left(\frac{|P_k - P_l|}{\varepsilon}\right) + o(\varepsilon) \right], \end{aligned} \tag{2.20}$$

where γ_1 is defined in (2.11) $\gamma_{kl} = \gamma_{lk} \in \Sigma$ and Σ is defined by (2.8) and γ_1 is defined in (2.11).

Furthermore, if $w(\frac{|P_k - P_l|}{\varepsilon}) = \eta\varepsilon$, we have $\gamma_{kl} \in \Sigma_1$ where Σ_1 is defined by (2.9).

Proof:

We shall prove the case when $K = 2$. The other cases are similar.

Since $\mathbf{P} = (P_1, P_2) \in \Lambda$, we have that $w(\frac{|P_1 - P_2|}{\varepsilon}) < \eta\varepsilon$.

First we look at the case $K = 1$. Note that by Proposition 2.1, Lemma 2.6 and Lemma 2.7 we have

$$\begin{aligned}
& \varepsilon^2 \int_{\Omega} |\nabla(P_{\Omega_{\varepsilon, P}} w(\frac{x - P}{\varepsilon}))|^2 + \int_{\Omega} |P_{\Omega_{\varepsilon, P}} w(\frac{x - P}{\varepsilon})|^2 \\
&= \varepsilon^N \int_{\Omega_{\varepsilon, P}} f(w) P_{\Omega_{\varepsilon, P}} w \\
&= \varepsilon^N \left(\int_{\Omega_{\varepsilon, P}} f(w) w + \int_{\Omega_{\varepsilon, P}} f(w) (P_{\Omega_{\varepsilon, P}} w - w) \right) \\
&= \varepsilon^N \left(\int_{R_+^N} f(w) w - \varepsilon H(P) \frac{1}{2} \int_{R^{N-1}} f(w) w |y'|^2 dy' \right. \\
&\quad \left. + \varepsilon H(P) \frac{N-1}{4} \int_{R^{N-1}} |w|^2 dy' + o(\varepsilon) \right).
\end{aligned} \tag{2.21}$$

Similarly we have

$$\begin{aligned}
& \int_{\Omega} F(P_{\Omega_{\varepsilon, P}} w(\frac{x - P}{\varepsilon})) dx \\
&= \varepsilon^N \left(\int_{\Omega_{\varepsilon, P}} F(w) dy + \int_{\Omega_{\varepsilon, P}} (F(P_{\Omega_{\varepsilon, P}} w) - F(w)) dy \right) \\
&= \varepsilon^N \left(\int_{R_+^N} F(w) dy - \varepsilon H(P) \frac{1}{2} \int_{R^{N-1}} f(w) w |y'|^2 dy' \right. \\
&\quad \left. + \varepsilon H(P) \frac{N-1}{4} \int_{R^{N-1}} |w|^2 dy' + o(\varepsilon) \right).
\end{aligned} \tag{2.22}$$

Then

$$J_{\varepsilon}(P_{\Omega_{\varepsilon, P}} w) = \varepsilon^N \left(\frac{1}{2} I(w) - \gamma_1 H(P) + o(\varepsilon) \right).$$

For the case $K = 2$, we can write

$$\begin{aligned}
\int_{\Omega} F(Pw_1 + Pw_2) &= \int_{\Omega_1} F(Pw_1 + Pw_2) + \int_{\Omega_2} F(Pw_1 + Pw_2) + \int_{\Omega_3} F(Pw_1 + Pw_2) \\
&= I_1 + I_2 + I_3
\end{aligned}$$

where $I_i, i = 1, 2, 3$ are defined at the last equality and

$$\begin{aligned}
\Omega_1 &= \{|x - P_1| \leq \frac{1}{2}|P_1 - P_2|\}, \quad \Omega_2 = \{|x - P_2| \leq \frac{1}{2}|P_1 - P_2|\}, \\
\Omega_3 &= \Omega \setminus (\Omega_1 \cup \Omega_2).
\end{aligned}$$

For I_3 , we have

$$|\varepsilon^{-N} I_3| \leq C \int_{(\Omega_3)_\varepsilon} (w_1 + w_2)^{2+\sigma} = O(w(\frac{|P_1 - P_2|}{\varepsilon})^{1+0.5\sigma}) = O(\varepsilon^{1+0.5\sigma}).$$

For I_1 , using $w(\frac{|P_1 - P_2|}{\varepsilon}) = O(\varepsilon)$ we have

$$\begin{aligned} \varepsilon^{-N} I_1 &= \int_{(\Omega_1)_\varepsilon} (F(Pw_1) + f(Pw_1)Pw_2) + O(\varepsilon^{1+0.5\sigma}) \\ &= \int_{\Omega_\varepsilon} F(Pw_1) + \int_{(\Omega_1)_\varepsilon} f(w_1)w_2 + O(\varepsilon^{1+0.5\sigma}) \\ &= \int_{R_+^N} F(w) - \varepsilon H(P_1) \left(\frac{1}{2} \int_{R^{N-1}} F(w)|y'|^2 dy' - \frac{N-1}{4} \int_{R^{N-1}} |w|^2 dy' \right) \\ &\quad + \int_{(\Omega_1)_\varepsilon} f(w_1)w_2 + O(\varepsilon^{1+0.5\sigma}). \end{aligned}$$

Similarly,

$$\begin{aligned} \varepsilon^{-N} I_2 &= \int_{R_+^N} F(w) - \varepsilon H(P_2) \left(\frac{1}{2} \int_{R^{N-1}} F(w)|y'|^2 dy' - \frac{N-1}{4} \int_{R^{N-1}} |w|^2 dy' \right) \\ &\quad + \int_{(\Omega_2)_\varepsilon} f(w_2)w_1 + O(\varepsilon^{1+0.5\sigma}). \end{aligned}$$

Hence

$$\begin{aligned} &\varepsilon^{-N} J_\varepsilon \left(\sum_{i=1}^K Pw_i \right) \\ &= \int_{\Omega_\varepsilon} \left[\frac{1}{2} \left(\sum_{i=1}^2 (|\nabla Pw_i|^2 + (Pw_i)^2) \right) + \nabla Pw_1 \nabla Pw_2 + Pw_1 Pw_2 \right] \\ &\quad - \int_{\Omega_\varepsilon} F(Pw_1 + Pw_2) \\ &= \int_{\Omega_\varepsilon} \left[\frac{1}{2} \left(\sum_{i=1}^2 (|\nabla Pw_i|^2 + (Pw_i)^2) \right) \right] + \int_{\Omega_\varepsilon} f(w_1)Pw_2 \\ &\quad - \int_{\Omega_\varepsilon} F(Pw_1 + Pw_2) \\ &= 2I(w) - \gamma_1 \varepsilon \sum_{i=1}^2 H(P_i) + \int_{\Omega_\varepsilon} f(w_1)Pw_2 \\ &\quad - \int_{(\Omega_1)_\varepsilon} f(w_1)w_2 - \int_{(\Omega_2)_\varepsilon} f(w_2)w_1 + o(w(\frac{|P_1 - P_2|}{\varepsilon})) + O(\varepsilon^{1+0.5\sigma}) \\ &= 2I(w) - \gamma_1 \varepsilon \sum_{i=1}^2 H(P_i) - (\gamma_{12} + o(1))w(\frac{|P_1 - P_2|}{\varepsilon}) + O(\varepsilon^{1+0.5\sigma}). \end{aligned} \tag{2.23}$$

Here we have used Lemma 2.4

$$\int_{\Omega_\varepsilon} f(w_1)w_2 = (\gamma_{12} + o(1))w\left(\frac{|P_1 - P_2|}{\varepsilon}\right)$$

and similarly

$$\int_{(\Omega_1)_\varepsilon} f(w_1)w_2 = (\gamma_{12} + o(1))w\left(\frac{|P_1 - P_2|}{\varepsilon}\right),$$

$$\int_{(\Omega_2)_\varepsilon} f(w_2)w_1 = (\gamma_{21} + o(1))w\left(\frac{|P_1 - P_2|}{\varepsilon}\right) = (\gamma_{12} + o(1))w\left(\frac{|P_1 - P_2|}{\varepsilon}\right).$$

□

3. LIAPUNOV-SCHMIDT REDUCTION

In this section, we reduce problem (1.2) to finite dimensions by the Liapunov-Schmidt method. We first introduce some notations.

Let $H_N^2(\Omega_\varepsilon)$ be the Hilbert space defined by

$$H_N^2(\Omega_\varepsilon) = \left\{ u \in H^2(\Omega_\varepsilon) \left| \frac{\partial u}{\partial \nu_\varepsilon} = 0 \text{ on } \partial\Omega_\varepsilon \right. \right\}.$$

Define

$$S_\varepsilon(u) = \Delta u - u + f(u)$$

for $u \in H_N^2(\Omega_\varepsilon)$. Then solving equation (1.2) is equivalent to

$$S_\varepsilon(u) = 0, u \in H_N^2(\Omega_\varepsilon).$$

Fix $\mathbf{P} = (P_1, \dots, P_K) \in \bar{\Lambda}$. To study (1.2) we first consider the linearized operator

$$\tilde{L}_\varepsilon : u \mapsto \Delta u - u + f' \left(\sum_{i=1}^K Pw_i \right) u,$$

$$H_N^2(\Omega_\varepsilon) \rightarrow L^2(\Omega_\varepsilon).$$

It is easy to see (integration by parts) that the cokernel of \tilde{L}_ε coincides with its kernel. Choose approximate cokernel and kernel as

$$\begin{aligned} \mathcal{C}_{\varepsilon, \mathbf{P}} &= \mathcal{K}_{\varepsilon, \mathbf{P}} \\ &= \text{span} \left\{ \frac{\partial Pw_i}{\partial \tau_{P_{i,j}}} \left| i = 1, \dots, K, j = 1, \dots, N-1 \right. \right\}. \end{aligned}$$

Let $\pi_{\varepsilon, \mathbf{P}}$ denote the projection from $L^2(\Omega_\varepsilon)$ onto $\mathcal{C}_{\varepsilon, \mathbf{P}}^\perp$. Our goal in this section is to show that the equation

$$\pi_{\varepsilon, \mathbf{P}} \circ S_\varepsilon \left(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon, \mathbf{P}} \right) = 0$$

has a unique solution $\Phi_{\varepsilon, \mathbf{P}} \in \mathcal{K}_{\varepsilon, \mathbf{P}}^\perp$ if ε is small enough and $\mathbf{P} = (P_1, \dots, P_K) \in \bar{\Lambda}$.

As a preparation in the following two propositions we show the invertibility of the corresponding linearized operator.

Proposition 3.1. *Let $L_{\varepsilon, \mathbf{P}} = \pi_{\varepsilon, \mathbf{P}} \circ \tilde{L}_\varepsilon$. There exist positive constants $\bar{\varepsilon}, \bar{\Lambda}$ such that for all $\varepsilon \in (0, \bar{\varepsilon})$ and $\mathbf{P} = (P_1, \dots, P_K) \in \bar{\Lambda}$*

$$\|L_{\varepsilon, \mathbf{P}} \Phi\|_{L^2(\Omega_\varepsilon)} \geq \lambda \|\Phi\|_{H^2(\Omega_\varepsilon)} \quad (3.1)$$

for all $\Phi \in \mathcal{K}_{\varepsilon, \mathbf{P}}^\perp$.

Proposition 3.2. *For any $\varepsilon \in (0, \tilde{\varepsilon})$ and $\mathbf{P} = (P_1, \dots, P_K) \in \bar{\Lambda}$ the map*

$$L_{\varepsilon, \mathbf{P}} = \pi_{\varepsilon, \mathbf{P}} \circ \tilde{L}_\varepsilon : \mathcal{K}_{\varepsilon, \mathbf{P}}^\perp \rightarrow \mathcal{C}_{\varepsilon, \mathbf{P}}^\perp$$

is surjective.

Proof of Proposition 3.1: We will follow the method used in [7], [24], [25], and [35]. Suppose that (3.1) is false. Then there exist sequences $\{\varepsilon_k\}$, $\{\mathbf{P}_k\} = \{(P_{1,k}, \dots, P_{K,k})\}$, and $\{\Phi_k\}$ ($i = 1, 2, \dots, K$, $k = 1, 2, \dots$) with $\varepsilon_k > 0$, $\mathbf{P}_k \in \bar{\Lambda}$, $\Phi_k \in \mathcal{K}_{\varepsilon_k, \mathbf{P}_k}^\perp$ such that

$$\varepsilon_k \rightarrow 0, \quad (3.2)$$

$$\mathbf{P}_k \rightarrow \mathbf{P} \in \bar{\Lambda}, \quad (3.3)$$

$$\|L_{\varepsilon_k, \mathbf{P}_k} \Phi_k\|_{L^2(\Omega_{\varepsilon_k})} \rightarrow 0, \quad (3.4)$$

$$\|\Phi_k\|_{H^2(\Omega_{\varepsilon_k})} = 1, \quad k = 1, 2, \dots \quad (3.5)$$

For $j = 1, 2, \dots, N-1$ denote

$$e_{ij,k} = \frac{\partial}{\partial \tau_{(P_i,k)_j}} Pw_{i,k} / \left\| \frac{\partial}{\partial \tau_{(P_i,k)_j}} Pw_{i,k} \right\|_{L^2(\Omega_{\varepsilon_k})}$$

where

$$Pw_{i,k}(y) = P_{\Omega_{\varepsilon_k, P_i,k}} w\left(y - \frac{P_{i,k}}{\varepsilon_k}\right), \quad y \in \Omega_{\varepsilon_k}.$$

Note that

$$\langle e_{i_1 j_1, k}, e_{i_2 j_2, k} \rangle = \delta_{i_1 i_2} \delta_{j_1 j_2} + \mathcal{O}(\varepsilon_k) \quad \text{as } k \rightarrow \infty$$

by Proposition 2.2, the symmetry of the function w and the fact that $\mathbf{P} \in \bar{\Lambda}$ (recall that $w(\frac{|P_k - P_l|}{\varepsilon}) \leq \eta\varepsilon$). Here $\delta_{i_1 i_2}$ is the Kronecker symbol. Furthermore, because of (3.4),

$$\|\tilde{L}_{\varepsilon_k} \Phi_k\|_{L^2}^2 - \sum_{i=1}^K \sum_{j=1}^{N-1} \left(\int_{\Omega_{\varepsilon_k}} \tilde{L}_{\varepsilon_k} \Phi_k e_{ij, k} \right)^2 \rightarrow 0 \quad (3.6)$$

as $k \rightarrow \infty$. Let Ω_0 , χ , ρ and T be as defined in Section 2. (Note that we allow $R_0 \rightarrow 0$ but $\frac{R_0}{\varepsilon} \rightarrow \infty$). Then T has an inverse T^{-1} such that

$$T^{-1} : T(B(P, R_0) \cap \bar{\Omega}) \rightarrow B(P, R_0) \cap \bar{\Omega}.$$

Recall that $\varepsilon y = T(x)$. We use the notation $T^{(i)}$ if P is replaced by P_i . We introduce new sequences $\{\varphi_{i, k}\}$ by

$$\varphi_{i, k}(y) = \chi\left(\frac{1}{\varepsilon_k} (T^{(i)})^{-1}(\varepsilon_k y)\right) \Phi_k\left((T^{(i)})^{-1}(\varepsilon_k y)\right) \quad (3.7)$$

for $y \in R_+^N$. Since $T^{(i)}$ and $(T^{(i)})^{-1}$ have bounded derivatives it follows from (3.5) and the smoothness of χ that

$$\|\varphi_{i, k}\|_{H^2(R_+^N)} \leq C$$

for all k sufficiently large. Since also

$$\|\varphi_{i, k}\|_{H^2(R_+^N \setminus B(0, R))} \rightarrow 0 \quad \text{as } R \rightarrow \infty$$

uniformly in k for all k large enough there exists a subsequence, again denoted by $\{\varphi_{i, k}\}$ which converges weakly in $H^2(R_+^N)$ to a limit $\varphi_{i, \infty}$ as $k \rightarrow \infty$. We are now going to show that $\varphi_{i, \infty} \equiv 0$. As a first step we deduce

$$\int_{R_+^N} \varphi_{i, \infty} \frac{\partial w}{\partial y_j} = 0, \quad j = 1, \dots, N-1. \quad (3.8)$$

This statement is shown as follows (note that $\det DT = \det DT^{-1} = 1$)

$$\begin{aligned} & \int_{R_+^N} \varphi_{i, k}(y) \left[\frac{\partial P w_{i, k}}{\partial \tau_{(P_i, k)_j}} \left(\frac{(T^{(i)})^{-1}(\varepsilon_k y)}{\varepsilon_k} \right) \right] dy \\ &= \varepsilon_k^{-N} \int_{\Omega_0} \chi(x - P_{i, k}) \Phi_k\left(\frac{x}{\varepsilon_k}\right) \frac{\partial P w_{i, k}}{\partial \tau_{(P_i, k)_j}}\left(\frac{x - P_{i, k}}{\varepsilon_k}\right) dx \end{aligned}$$

$$\begin{aligned}
&= \varepsilon_k^{-N} \int_{\Omega} \Phi_k\left(\frac{x}{\varepsilon_k}\right) \frac{\partial Pw_{i,k}}{\partial \tau_{(P_{i,k})_j}}\left(\frac{x - P_{i,k}}{\varepsilon_k}\right) \\
&\quad - \varepsilon_k^{-N} \int_{\Omega \setminus \Omega_0} \Phi_k\left(\frac{x}{\varepsilon_k}\right) \frac{\partial Pw_{i,k}}{\partial \tau_{(P_{i,k})_j}}\left(\frac{x - P_{i,k}}{\varepsilon_k}\right) \\
&\quad - \varepsilon_k^{-N} \int_{\Omega_0} [1 - \chi(x - P_{i,k})] \Phi_k\left(\frac{x}{\varepsilon_k}\right) \frac{\partial Pw_{i,k}}{\partial \tau_{(P_{i,k})_j}}\left(\frac{x - P_{i,k}}{\varepsilon_k}\right) \\
&= 0 - \varepsilon_k^{-N} \int_{\Omega \setminus \Omega_0} \Phi_k\left(\frac{x}{\varepsilon_k}\right) \left[\frac{\partial w}{\partial (P_{i,k})_j} \left(\frac{x - P_{i,k}}{\varepsilon_k}\right) - \frac{\partial Pw_{i,k}}{\partial \tau_{(P_{i,k})_j}} \left(\frac{x - P_{i,k}}{\varepsilon_k}\right) \right] \\
&\quad - \varepsilon_k^{-N} \int_{\Omega_0} [1 - \chi(x - P_{i,k})] \Phi_k\left(\frac{x}{\varepsilon_k}\right) \left[\frac{\partial w}{\partial (P_{i,k})_j} \left(\frac{x - P_{i,k}}{\varepsilon_k}\right) - \frac{\partial Pw_{i,k}}{\partial \tau_{(P_{i,k})_j}} \left(\frac{x - P_{i,k}}{\varepsilon_k}\right) \right] \\
&\quad - \varepsilon_k^{-N} \int_{\Omega \setminus \Omega_0} \Phi_k\left(\frac{x}{\varepsilon_k}\right) \frac{\partial w}{\partial (P_{i,k})_j} \left(\frac{x - P_{i,k}}{\varepsilon_k}\right) \\
&\quad - \varepsilon_k^{-N} \int_{\Omega_0} [1 - \chi(x - P_{i,k})] \Phi_k\left(\frac{x}{\varepsilon_k}\right) \frac{\partial w}{\partial (P_{i,k})_j} \left(\frac{x - P_{i,k}}{\varepsilon_k}\right)
\end{aligned}$$

where Ω_0 is as defined in section 2. In the last expression the first two terms tend to zero as $k \rightarrow \infty$ since $\varepsilon_k^{-N} \Phi_k$ is bounded in $L^2(\Omega)$ and the term in the square bracket converges to 0 strongly in $L^2(\Omega)$. The last two terms tend to zero as $k \rightarrow \infty$ because of the exponential decay of $\partial w / \partial (P_{i,k})_j$ at infinity.

We conclude

$$\begin{aligned}
\limsup_{k \rightarrow \infty} \left| \int_{R_+^N} \varphi_{i,k}(y) \left[\frac{\partial Pw_{i,k}}{\partial \tau_{(P_{i,k})_j}} \left(\frac{(T^{(i)})^{-1}(\varepsilon_k y)}{\varepsilon_k} \right) \right] \right| &= 0 \\
i = 1, \dots, K, \quad j = 1, \dots, N - 1. & \tag{3.9}
\end{aligned}$$

This implies (3.8).

Let \mathcal{K}_0 and \mathcal{C}_0 be the kernel and cokernel, respectively, of the linear operator $S'_0(w)$ which is the Fréchet derivative at w of

$$\begin{aligned}
S_0(v) &= \Delta v - v + f(v), \\
S_0 &: H_N^2(R_+^N) \rightarrow L^2(R_+^N)
\end{aligned}$$

where

$$H_N^2(R_+^N) = \left\{ u \in H_N^2(R_+^N) \left| \frac{\partial u}{\partial y_N} = 0 \right. \right\}.$$

Note that

$$S'_0(w)v = \Delta v - v + f'(w)v,$$

$$\mathcal{K}_0 = \mathcal{C}_0 = \text{span} \left\{ \frac{\partial w}{\partial y_j} \mid j = 1, \dots, N-1 \right\}.$$

Equation (3.8) implies that $\varphi_{i,\infty} \in \mathcal{K}_0^\perp$. By the exponential decay of w and by (3.4) we have after possibly taking a further subsequence that

$$\Delta \varphi_{i,\infty} - \varphi_{i,\infty} + f'(w)\varphi_{i,\infty} = 0,$$

i.e. $\varphi_{i,\infty} \in \mathcal{K}_0$. Therefore $\varphi_{i,\infty} = 0$.

Hence

$$\varphi_{i,k} \rightharpoonup 0 \quad \text{weakly in } H^2(\mathbb{R}_+^N) \quad \text{as } k \rightarrow \infty. \quad (3.10)$$

By the definition of $\varphi_{i,k}$ we get $\Phi_k \rightharpoonup 0$ in H^2 and

$$\|\Phi_k\|_{L^2(\Omega_{\varepsilon_k})} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Furthermore,

$$\|f'(\sum_{i=1}^K Pw_i)\Phi_k\|_{L^2(\Omega_{\varepsilon_k})} \rightarrow 0$$

and therefore

$$\|(\Delta - 1)\Phi_k\|_{L^2(\Omega_{\varepsilon_k})} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Since

$$\begin{aligned} \int_{\Omega_{\varepsilon_k}} |\nabla \Phi_k|^2 + \Phi_k^2 &= \int_{\Omega_{\varepsilon_k}} [(1 - \Delta)\Phi_k]\Phi_k \\ &\leq C\|(\Delta - 1)\Phi_k\|_{L^2(\Omega_{\varepsilon_k})} \end{aligned}$$

we have that

$$\|\Phi_k\|_{H^1(\Omega_{\varepsilon_k})} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

In summary:

$$\|\Delta \Phi_k\|_{L^2(\Omega_{\varepsilon_k})} \rightarrow 0 \quad \text{and} \quad \|\Phi_k\|_{H^1(\Omega_{\varepsilon_k})} \rightarrow 0. \quad (3.11)$$

From (3.11) and the following elliptic regularity estimate (for a proof see Appendix B in [35])

$$\|\Phi_k\|_{H^2(\Omega_{\varepsilon_k})} \leq C(\|\Delta \Phi_k\|_{L^2(\Omega_{\varepsilon_k})} + \|\Phi_k\|_{H^1(\Omega_{\varepsilon_k})}) \quad (3.12)$$

for $\Phi_k \in H_N^2(\Omega_{\varepsilon_k})$ we deduce that

$$\|\Phi_k\|_{H^2(\Omega_{\varepsilon_k})} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

This contradicts the assumption

$$\|\Phi_k\|_{H^2(\Omega_{\varepsilon_k})} = 1$$

and the proof of Proposition 3.1 is completed. \square

Proof of Proposition 3.2:

We define a linear operator T from $L^2(\Omega_\varepsilon)$ to itself by

$$T = \pi_{\varepsilon, \mathbf{P}} \circ \tilde{L} \circ \pi_{\varepsilon, \mathbf{P}}$$

Its domain of definition is $H_N^2(\Omega_\varepsilon)$. By the theory of elliptic equations and by integration by parts it is easy to see that T is a (unbounded) self-adjoint operator on $L^2(\Omega_\varepsilon)$ and a closed operator. The L^2 estimates of elliptic equations imply that the range of T is closed in $L^2(\Omega_\varepsilon)$. Then by the Closed Range Theorem ([37], page 205), we know that the range of T is the orthogonal complement of its kernel which is, by Proposition 3.1, $\mathcal{K}_{\varepsilon, \mathbf{P}}$. This leads to Proposition 3.2. \square

We are now in a position to solve the equation

$$\pi_{\varepsilon, P} \circ S_\varepsilon \left(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon, \mathbf{P}} \right) = 0. \quad (3.13)$$

Since $L_{\varepsilon, \mathbf{P}}|_{\mathcal{K}_{\varepsilon, \mathbf{P}}^\perp}$ is invertible (call the inverse $L_{\varepsilon, \mathbf{P}}^{-1}$) we can rewrite

$$\begin{aligned} \Phi &= -(L_{\varepsilon, \mathbf{P}}^{-1} \circ \pi_{\varepsilon, \mathbf{P}}) \left(S_\varepsilon \left(\sum_{i=1}^K Pw_i \right) \right) \\ &\quad - (L_{\varepsilon, \mathbf{P}}^{-1} \circ \pi_{\varepsilon, \mathbf{P}}) N_{\varepsilon, \mathbf{P}}(\Phi) \\ &\equiv G_{\varepsilon, \mathbf{P}}(\Phi) \end{aligned} \quad (3.14)$$

where

$$\begin{aligned} N_{\varepsilon, \mathbf{P}}(\Phi) &= S_\varepsilon \left(\sum_{i=1}^K Pw_i + \Phi \right) \\ &\quad - \left[S_\varepsilon \left(\sum_{i=1}^K Pw_i \right) + S'_\varepsilon \left(\sum_{i=1}^K Pw_i \right) \Phi \right] \end{aligned}$$

and the operator $G_{\varepsilon, \mathbf{P}}$ is defined by the last equation for $\Phi \in H_N^2(\Omega_\varepsilon)$. We are going to show that the operator $G_{\varepsilon, \mathbf{P}}$ is a contraction on

$$B_{\varepsilon, \delta} \equiv \{ \Phi \in H^2(\Omega_\varepsilon) \mid \|\Phi\|_{H^2(\Omega_\varepsilon)} < \delta \}$$

if δ is small enough.

In fact we have the following lemma

Lemma 3.3. *For ε sufficiently small, we have*

$$|N_{\varepsilon, \mathbf{P}}| \leq C(|\Phi_{\varepsilon, \mathbf{P}}|^{1+\sigma} + |\Phi_{\varepsilon, \mathbf{P}}|^{p_1}), \quad (3.15)$$

$$\|S_\varepsilon(\sum_{i=1}^K Pw_i)\|_{L^2(\Omega_\varepsilon)} \leq C\varepsilon^{\frac{1+\sigma}{2}}. \quad (3.16)$$

Proof: (3.16) follows from the mean value theorem.

To prove (3.17), we divide the domain into $(K+1)$ parts: let $\Omega = \cup_{i=1}^{K+1} \Omega_i$ where

$$\Omega_i = \{|x - P_i| \leq \frac{1-\delta}{2} \min_{k \neq i} |P_k - P_l|\}, i = 1, \dots, K, \Omega_{K+1} = \Omega \setminus \cup_{i=1}^K \Omega_i.$$

Note that

$$S_\varepsilon(\sum_{i=1}^K Pw_i) = f(\sum_{i=1}^K Pw_i) - \sum_{i=1}^K f(w_i).$$

We now estimate $S_\varepsilon(\sum_{i=1}^K Pw_i)$ in each domain.

In Ω_{K+1} , we have

$$|S_\varepsilon(\sum_{i=1}^K Pw_i)| \leq (w_1 + \dots + w_K)^{1+\sigma} \leq O(\varepsilon^{\frac{1+\sigma}{2}}).$$

Hence, using also the fact that $w(y)$ decays exponentially in $|y|$ we obtain

$$\|S_\varepsilon(\sum_{i=1}^K Pw_i)\|_{L^2((\Omega_{K+1})_\varepsilon)} \leq O(\varepsilon^{\frac{1+\sigma}{2}}).$$

In $\Omega_i, i = 1, \dots, K$, we have

$$\begin{aligned} |S_\varepsilon(\sum_{i=1}^K Pw_i)| &\leq \sum_{j \neq i} \left(|f'(w_i)w_j| + |f'(w_i)(Pw_j - w_j)| \right) \\ &+ O(\sum_{j \neq i} (|Pw_j|^{1+\sigma} + |w_j|^{1+\sigma})) + O(|Pw_i - w_i|^{1+\sigma}). \end{aligned}$$

Using Proposition 2.1 and the facts that Pw, w and v_1 decay exponentially, we obtain

$$\|S_\varepsilon(\sum_{i=1}^K Pw_i)\|_{L^2((\Omega_i)_\varepsilon)} \leq O(\varepsilon^{\frac{1+\sigma}{2}}).$$

□

Thus

$$\|G_{\varepsilon, \mathbf{P}}(\Phi)\|_{H^2(\Omega_\varepsilon)} \leq \lambda^{-1}(\|\pi_{\varepsilon, \mathbf{P}} \circ N_{\varepsilon, \mathbf{P}}(\Phi)\|_{L^2(\Omega_\varepsilon)})$$

$$\begin{aligned}
& + \|\pi_{\varepsilon, \mathbf{P}} \circ (S_\varepsilon(\sum_{i=1}^K Pw_i))\|_{L^2(\Omega_\varepsilon)} \\
& \leq \lambda^{-1} C(c(\delta)\delta + \varepsilon^{\frac{1+\sigma}{2}})
\end{aligned}$$

where $\lambda > 0$ is independent of $\delta > 0$ and $c(\delta) \rightarrow 0$ as $\delta \rightarrow 0$. Similarly we show

$$\|G_{\varepsilon, \mathbf{P}}(\Phi) - G_{\varepsilon, \mathbf{P}}(\Phi')\|_{H^2(\Omega_\varepsilon)} \leq \lambda^{-1} Cc(\delta)\|\Phi - \Phi'\|_{H^2(\Omega_\varepsilon)}$$

where $c(\delta) \rightarrow 0$ as $\delta \rightarrow 0$. Therefore $M_{\varepsilon, \mathbf{P}}$ is a contraction on B_δ . The existence of a fixed point $\Phi_{\varepsilon, \mathbf{P}}$ now follows from the Contraction Mapping Principle and $\Phi_{\varepsilon, \mathbf{P}}$ is a solution of (3.14).

Because of

$$\begin{aligned}
\|\Phi_{\varepsilon, \mathbf{P}}\|_{H^2(\Omega_\varepsilon)} & \leq \lambda^{-1} (\|N_{\varepsilon, \mathbf{P}}(\Phi_{\varepsilon, \mathbf{P}})\|_{L^2(\Omega_\varepsilon)} \\
& \quad + \|S_\varepsilon(\sum_{i=1}^K Pw_i)\|_{L^2(\Omega_\varepsilon)}) \\
& \leq \lambda^{-1} C(\varepsilon^{\frac{1+\sigma}{2}} + c(\delta))\|\Phi_{\varepsilon, \mathbf{P}}\|_{H^2(\Omega_\varepsilon)}
\end{aligned}$$

we have

$$\|\Phi_{\varepsilon, \mathbf{P}}\|_{H^2(\Omega_\varepsilon)} \leq C\varepsilon^{\frac{1+\sigma}{2}}.$$

We have proved

Lemma 3.4. *There exists $\bar{\varepsilon} > 0$ such that for every $(N+1)$ -tuple $\varepsilon, P_1, \dots, P_K$ with $0 < \varepsilon < \bar{\varepsilon}$ and $\mathbf{P} = (P_1, \dots, P_K) \in \bar{\Lambda}$ there is a unique $\Phi_{\varepsilon, \mathbf{P}} \in \mathcal{K}_{\varepsilon, \mathbf{P}}^\perp$ satisfying $S_\varepsilon(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon, \mathbf{P}}) \in \mathcal{C}_{\varepsilon, \mathbf{P}}$ and*

$$\|\Phi_{\varepsilon, \mathbf{P}}\|_{H^2(\Omega_\varepsilon)} \leq C\varepsilon^{\frac{1+\sigma}{2}}. \quad (3.17)$$

The next lemma is our main estimate.

Lemma 3.5. *Let $\Phi_{\varepsilon, \mathbf{P}}$ be defined by Lemma 3.4. Then we have*

$$\begin{aligned}
& J_\varepsilon(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon, \mathbf{P}}) \\
& = \varepsilon^N \left[\frac{K}{2} I(w) - \gamma_1 \varepsilon \sum_{i=1}^K H(P_i) \right. \\
& \quad \left. - \frac{1}{2} \sum_{k, l=1, \dots, K, k \neq l} (\gamma_{kl} + o(1)) w \left(\frac{|P_k - P_l|}{\varepsilon} \right) + o(\varepsilon) \right]
\end{aligned} \quad (3.18)$$

where γ_1 and γ_{kl} are defined in Lemma 2.5.

Proof:

In fact for any $\mathbf{P} \in \bar{\Lambda}$, we have

$$\varepsilon^{-N} J_\varepsilon \left(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon, \mathbf{P}} \right) = \varepsilon^{-N} J_\varepsilon \left(\sum_{i=1}^K Pw_i \right) + g_{\varepsilon, \mathbf{P}}(\Phi_{\varepsilon, \mathbf{P}}) + O(\|\Phi_{\varepsilon, \mathbf{P}}\|_{H^2(\Omega_\varepsilon)}^2)$$

where

$$\begin{aligned} & g_{\varepsilon, \mathbf{P}}(\Phi_{\varepsilon, \mathbf{P}}) \\ &= \int_{\Omega_\varepsilon} \sum_{i=1}^K (\nabla Pw_i \nabla \Phi_{\varepsilon, \mathbf{P}} + Pw_i \Phi_{\varepsilon, \mathbf{P}}) - \int_{\Omega_\varepsilon} f \left(\sum_{i=1}^K Pw_i \right) \Phi_{\varepsilon, \mathbf{P}} \\ &= \int_{\Omega_\varepsilon} \left[\sum_{i=1}^K f(w_i) - f \left(\sum_{i=1}^K Pw_i \right) \right] \Phi_{\varepsilon, \mathbf{P}} \\ &\leq \left\| \sum_{i=1}^K f(w_i) - f \left(\sum_{i=1}^K Pw_i \right) \right\|_{L^2} \|\Phi_{\varepsilon, \mathbf{P}}\|_{L^2(\Omega_\varepsilon)} \\ &\leq O(\varepsilon^{1+\sigma}) \end{aligned}$$

by Lemma 3.3 and Lemma 3.4.

Estimate (3.19) now follows from Lemma 2.6 and Lemma 3.4. \square

Finally, we show that $\Phi_{\varepsilon, \mathbf{P}}$ is actually smooth in \mathbf{P} .

Lemma 3.6. *Let $\Phi_{\varepsilon, \mathbf{P}}$ be defined by Lemma 3.4. Then $\Phi_{\varepsilon, \mathbf{P}} \in C^1$ in \mathbf{P} .*

Proof. Recall that $\Phi_{\varepsilon, \mathbf{P}}$ is a solution of the equation

$$\pi_{\varepsilon, \mathbf{P}} \circ S_\varepsilon \left(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon, \mathbf{P}} \right) = 0 \quad (3.19)$$

such that

$$\Phi_{\varepsilon, \mathbf{P}} \in \mathcal{K}_{\varepsilon, P}^\perp. \quad (3.20)$$

By differentiating equation (3.19) twice we easily conclude that the functions Pw_i and $\partial^2 Pw_i / (\partial \tau_{P_{i,j}} \partial \tau_{P_{i,k}})$ are C^1 in \mathbf{P} . This implies that the projection $\pi_{\varepsilon, \mathbf{P}}$ is C^1 in \mathbf{P} . Applying $\partial / \partial \tau_{P_{i,j}}$ gives

$$\pi_{\varepsilon, \mathbf{P}} \circ DS_\varepsilon \left(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon, \mathbf{P}} \right) \left(\sum_{i=1}^K \frac{\partial Pw_i}{\partial \tau_{P_{i,j}}} + \frac{\partial \Phi_{\varepsilon, \mathbf{P}}}{\partial \tau_{P_{i,j}}} \right)$$

$$+\frac{\partial\pi_{\varepsilon,\mathbf{P}}}{\partial\tau_{P_{i,j}}}\circ S_{\varepsilon}\left(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon,\mathbf{P}}\right) = 0. \quad (3.21)$$

where

$$DS_{\varepsilon}\left(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon,\mathbf{P}}\right) = \Delta - 1 + f'\left(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon,\mathbf{P}}\right).$$

We decompose $\frac{\partial\Phi_{\varepsilon,\mathbf{P}}}{\partial\tau_{P_{i,j}}}$ into two parts:

$$\frac{\partial\Phi_{\varepsilon,\mathbf{P}}}{\partial\tau_{P_{i,j}}} = \left(\frac{\partial\Phi_{\varepsilon,\mathbf{P}}}{\partial\tau_{P_{i,j}}}\right)_1 + \left(\frac{\partial\Phi_{\varepsilon,\mathbf{P}}}{\partial\tau_{P_{i,j}}}\right)_2$$

where $\left(\frac{\partial\Phi_{\varepsilon,\mathbf{P}}}{\partial\tau_{P_{i,j}}}\right)_1 \in \mathcal{K}_{\varepsilon,\mathbf{P}}$ and $\left(\frac{\partial\Phi_{\varepsilon,\mathbf{P}}}{\partial\tau_{P_{i,j}}}\right)_2 \in \mathcal{K}_{\varepsilon,\mathbf{P}}^{\perp}$.

We can easily show that $\left(\frac{\partial\Phi_{\varepsilon,\mathbf{P}}}{\partial\tau_{P_{i,j}}}\right)_1$ is continuous in \mathbf{P} since

$$\int_{\Omega_{\varepsilon}} \Phi_{\varepsilon,\mathbf{P}} \frac{\partial Pw_k}{\partial\tau_{P_{k,l}}} = 0, \quad k = 1, \dots, K, \quad l = 1, \dots, N-1$$

and

$$\int_{\Omega_{\varepsilon}} \frac{\partial\Phi_{\varepsilon,\mathbf{P}}}{\partial\tau_{P_{i,j}}} \frac{\partial Pw_k}{\partial\tau_{P_{k,l}}} + \int_{\Omega_{\varepsilon}} \Phi_{\varepsilon,\mathbf{P}} \frac{\partial^2 Pw_k}{\partial\tau_{P_{i,j}} \partial\tau_{P_{k,l}}} = 0$$

$$k, i = 1, \dots, K, \quad l, j = 1, \dots, N-1.$$

We can write equation (3.23) as

$$\begin{aligned} & \pi_{\varepsilon,\mathbf{P}} \circ DS_{\varepsilon}\left(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon,\mathbf{P}}\right) \left(\left(\frac{\partial\Phi_{\varepsilon,\mathbf{P}}}{\partial\tau_{P_{i,j}}}\right)_2\right) \\ & + \pi_{\varepsilon,\mathbf{P}} \circ DS_{\varepsilon}\left(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon,\mathbf{P}}\right) \left(\sum_{i=1}^K \frac{\partial Pw_i}{\partial\tau_{P_{i,j}}} + \left(\frac{\partial\Phi_{\varepsilon,\mathbf{P}}}{\partial\tau_{P_{i,j}}}\right)_1\right) \\ & + \frac{\partial\pi_{\varepsilon,\mathbf{P}}}{\partial\tau_{P_{i,j}}}\circ S_{\varepsilon}\left(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon,\mathbf{P}}\right) = 0. \end{aligned} \quad (3.22)$$

As in the proof of Propositions 3.1 and 3.2, we can show that the operator

$$\pi_{\varepsilon,\mathbf{P}} \circ DS_{\varepsilon}\left(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon,\mathbf{P}}\right)$$

is invertible from $\mathcal{K}_{\varepsilon,\mathbf{P}}^{\perp}$ to $\mathcal{C}_{\varepsilon,\mathbf{P}}^{\perp}$. Then we can take inverse of $\pi_{\varepsilon,\mathbf{P}} \circ DS_{\varepsilon}\left(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon,\mathbf{P}}\right)$ in the above equation and the inverse is continuous in \mathbf{P} .

Since $\frac{\partial Pw_i}{\partial\tau_{P_{i,j}}}, \left(\frac{\partial\Phi_{\varepsilon,\mathbf{P}}}{\partial\tau_{P_{i,j}}}\right)_1 \in \mathcal{K}_{\varepsilon,\mathbf{P}}$ are continuous in \mathbf{P} and so is $\frac{\partial\pi_{\varepsilon,\mathbf{P}}}{\partial\tau_{P_{i,j}}}$, we conclude that $\left(\frac{\partial\Phi_{\varepsilon,\mathbf{P}}}{\partial\tau_{P_{i,j}}}\right)_2$ is also continuous in \mathbf{P} . This is the same as the C^1 dependence of $\Phi_{\varepsilon,\mathbf{P}}$ in \mathbf{P} . The proof is finished. \square

4. THE REDUCED PROBLEM: A MAXIMIZING PROCEDURE

In this section, we study a maximizing problem.

Fix $\mathbf{P} \in \bar{\Lambda}$. Let $\Phi_{\varepsilon, \mathbf{P}}$ be the solution given by Lemma 3.4. We define a new functional

$$M_\varepsilon(\mathbf{P}) = J_\varepsilon\left(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon, \mathbf{P}}\right) : \bar{\Lambda} \rightarrow R. \quad (4.1)$$

We shall prove

Proposition 4.1. *For ε small, the following maximizing problem*

$$\max\{M_\varepsilon(\mathbf{P}) : \mathbf{P} \in \bar{\Lambda}\} \quad (4.2)$$

has a solution $\mathbf{P}^\varepsilon \in \Lambda$.

Proof: Since $J_\varepsilon(\sum_{i=1}^K P_{\Omega_\varepsilon, P_i} w + \Phi_{\varepsilon, \mathbf{P}})$ is continuous in \mathbf{P} , the maximizing problem has a solution. Let $M_\varepsilon(\mathbf{P}^\varepsilon)$ be the maximum where $\mathbf{P}^\varepsilon \in \bar{\Lambda}$.

We claim that $\mathbf{P}^\varepsilon \in \Lambda$.

In fact for any $\mathbf{P} \in \bar{\Lambda}$, by Lemma 3.5, we have

$$M_\varepsilon(\mathbf{P}) = \varepsilon^N \left[\frac{K}{2} I(w) - \varepsilon \gamma \left(\sum_{i=1}^K H(P_i) \right) - \frac{1}{2} \sum_{k, l=1, \dots, K, k \neq l} (\gamma_{kl} + o(1)) w \left(\frac{|P_k - P_l|}{\varepsilon} \right) + o(\varepsilon) \right].$$

Since $M_\varepsilon(\mathbf{P}^\varepsilon)$ is the maximum, we have

$$\begin{aligned} & \gamma_1 \sum_{i=1}^K H(P_i^\varepsilon) + \frac{1}{\varepsilon} \sum_{k \neq l} \left(\frac{1}{2} \gamma_{kl} + o(1) \right) w \left(\frac{|P_k^\varepsilon - P_l^\varepsilon|}{\varepsilon} \right) \\ & \leq \gamma_1 \sum_{i=1}^K H(P_i) + \frac{1}{\varepsilon} \sum_{k \neq l} \left(\frac{1}{2} \gamma_{kl} + o(1) \right) w \left(\frac{|P_k - P_l|}{\varepsilon} \right) + o(1) \end{aligned}$$

for any $\mathbf{P} = (P_1, \dots, P_K) \in \bar{\Lambda}$.

Choose P_i such that $H(P_i) \rightarrow \min_{P \in \Gamma_i} H(P)$ and $w\left(\frac{|P_k - P_l|}{\varepsilon}\right) \rightarrow 0$. This implies that

$$\gamma_1 \sum_{i=1}^K H(P_i^\varepsilon) + \frac{1}{\varepsilon} \sum_{k \neq l} \left(\frac{1}{2} \gamma_{kl} + o(1) \right) w \left(\frac{|P_k^\varepsilon - P_l^\varepsilon|}{\varepsilon} \right) \leq \gamma_1 \sum_{i=1}^K \min_{P \in \Gamma_i} H(P) + \delta$$

for any $\delta > 0$.

Note that $\partial\Lambda \subset \{P_i \in \partial\Gamma_i \text{ or } w(\frac{|P_k - P_l|}{\varepsilon}) = \varepsilon\eta\}$. Hence if $\mathbf{P} \in \partial\Lambda$, we have that either

$$H(P_i) \geq \min_{P \in \partial\Gamma_i} H(P) \geq \min_{P \in \Gamma_i} H(P) + 2\eta_0$$

for some $i = 1, \dots, K$ and $\eta_0 > 0$ (by condition (1.5)) or

$$\frac{1}{\varepsilon} w\left(\frac{|P_k - P_l|}{\varepsilon}\right) = \eta$$

for some $k \neq l$.

Hence if $\mathbf{P} \in \partial\Lambda$ we have

$$\begin{aligned} & \gamma_1 \sum_{i=1}^K H(P_i^\varepsilon) + \frac{1}{\varepsilon} \sum_{k \neq l} \left(\frac{1}{2} \gamma_{kl} + o(1)\right) w\left(\frac{|P_k^\varepsilon - P_l^\varepsilon|}{\varepsilon}\right) \\ & \geq \gamma_1 \sum_{i=1}^K \min_{P \in \Gamma_i} H(P) + \min(\gamma_1 \eta_0, \min_{k \neq l, w(\frac{|P_k - P_l|}{\varepsilon}) = \eta\varepsilon} \gamma_{kl} \eta). \end{aligned}$$

Note that $\min_{k \neq l, w(\frac{|P_k - P_l|}{\varepsilon}) = \eta\varepsilon} \gamma_{kl} \geq \inf_{\tau \in \Sigma_1} \tau \geq \delta_0 > 0$ since for any $\tau \in \Sigma_1$, we have

$$\tau = \int_{R_+^N} f(w) e^{(b,y)} = \frac{1}{2} \int_{R^N} f(w) e^{(b,y)} > 0.$$

A contradiction to (4.3) if we choose δ small enough.

It follows that $\mathbf{P}^\varepsilon \in \Lambda$.

This completes the proof of Proposition 4.1. \square

5. PROOF OF THEOREM 1.2

In this section section, we apply results in Section 3 and Section 4 to prove Theorem 1.1, Theorem 1.2 and Corollary 1.3.

Proofs of Theorem 1.1, Theorem 1.2 and Corollary 1.3

By Lemma 3.4 and Lemma 3.6, there exists ε_0 such that for $\varepsilon < \varepsilon_0$ we have a C^1 map which, to any $\mathbf{P} \in \bar{\Lambda}$, associates $\Phi_{\varepsilon, \mathbf{P}} \in \mathcal{K}_{\varepsilon, \mathbf{P}}^\perp$ such tha

t

$$S_\varepsilon\left(\sum_{i=1}^K P w_i + \Phi_{\varepsilon, \mathbf{P}}\right) = \sum_{k=1, \dots, K; l=1, \dots, N-1} \alpha_{kl} \frac{\partial P w_k}{\partial \tau_{P_{k,l}}} \quad (5.1)$$

for some constants $\alpha_{kl} \in R^{K(N-1)}$.

By Proposition 4.1, we have $\mathbf{P}^\varepsilon \in \Lambda$, achieving the maximum of the maximization problem in Proposition 4.1. Let $\Phi_\varepsilon = \Phi_{\varepsilon, \mathbf{P}^\varepsilon}$ and $u_\varepsilon = \sum_{i=1}^K P_{\Omega_{\varepsilon, P_i^\varepsilon}} w +$

$\Phi_{\varepsilon, \mathbf{P}^\varepsilon}$. Then we have

$$\frac{\partial}{\partial \tau_{P_{i,j}}} \Big|_{\mathbf{P}=\mathbf{P}^\varepsilon} M_\varepsilon(\mathbf{P}^\varepsilon) = 0, i = 1, \dots, K, j = 1, \dots, N-1.$$

Hence we have

$$\begin{aligned} \int_{\Omega_\varepsilon} [\nabla u_\varepsilon \nabla \frac{\partial(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon, \mathbf{P}})}{\partial \tau_{P_{i,j}}} \Big|_{\mathbf{P}=\mathbf{P}^\varepsilon} + u_\varepsilon \frac{\partial(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon, \mathbf{P}})}{\partial \tau_{P_{i,j}}} \Big|_{\mathbf{P}=\mathbf{P}^\varepsilon} \\ - f(u_\varepsilon) \frac{\partial(\sum_{i=1}^K Pw_i + \Phi_{\varepsilon, \mathbf{P}})}{\partial \tau_{P_{i,j}}} \Big|_{\mathbf{P}=\mathbf{P}^\varepsilon}] = 0. \end{aligned}$$

Thus

$$\begin{aligned} \int_{\Omega_\varepsilon} \nabla u_\varepsilon \nabla \frac{\partial(Pw_i + \Phi_{\varepsilon, \mathbf{P}})}{\partial \tau_{P_{i,j}}} \Big|_{\mathbf{P}=\mathbf{P}^\varepsilon} \\ + u_\varepsilon \frac{\partial(Pw_i + \Phi_{\varepsilon, \mathbf{P}})}{\partial \tau_{P_{i,j}}} \Big|_{\mathbf{P}=\mathbf{P}^\varepsilon} - f(u_\varepsilon) \frac{\partial(Pw_i + \Phi_{\varepsilon, \mathbf{P}})}{\partial \tau_{P_{i,j}}} \Big|_{\mathbf{P}=\mathbf{P}^\varepsilon} = 0 \end{aligned}$$

for $i = 1, \dots, K$ and $j = 1, \dots, N-1$.

Therefore we have

$$\sum_{k=1, \dots, K; l=1, \dots, N-1} \alpha_{kl} \int_{\Omega_\varepsilon} \frac{\partial Pw_k}{\partial \tau_{P_{k,l}}} \frac{\partial(Pw_i + \Phi_{\varepsilon, \mathbf{P}})}{\partial \tau_{P_{i,j}}} = 0. \quad (5.2)$$

Since $\Phi_{\varepsilon, \mathbf{P}} \in \mathcal{K}_{\varepsilon, \mathbf{P}}^\perp$, we have that

$$\begin{aligned} \left| \int_{\Omega_\varepsilon} \frac{\partial Pw_k}{\partial \tau_{P_{k,l}}} \frac{\partial \Phi_{\varepsilon, \mathbf{P}}}{\partial \tau_{P_{i,j}}} \right| &= \left| - \int_{\Omega_\varepsilon} \frac{\partial^2 Pw_i}{\partial \tau_{P_{k,l}} \partial \tau_{P_{i,j}}} \Phi_{\varepsilon, \mathbf{P}} \right| \\ &\leq \left\| \frac{\partial^2 Pw_i}{\partial \tau_{P_{k,l}} \partial \tau_{P_{i,j}}} \right\|_{L^2} \|\Phi_{\varepsilon, \mathbf{P}}\|_{L^2} \\ &= O(\varepsilon^{-2 + \frac{1+\sigma}{2}}). \end{aligned}$$

Note that

$$\int_{\Omega_\varepsilon} \frac{\partial Pw_k}{\partial \tau_{P_{k,l}}} \frac{\partial Pw_i}{\partial \tau_{P_{i,j}}} = \frac{1}{\varepsilon^2} \delta_{ik} \delta_{lj} (A + o(1))$$

where

$$A = \int_{R_+^N} \left(\frac{\partial w}{\partial y_1} \right)^2 > 0.$$

Thus equation (5.2) becomes a system of homogeneous equations for α_{kl} and the matrix of the system is nonsingular since it is diagonally dominant. So $\alpha_{kl} \equiv 0, k = 1, \dots, K, l = 1, \dots, N-1$.

Hence $u_\varepsilon = \sum_{i=1}^K P_{\Omega_\varepsilon, P_i^\varepsilon} w + \Phi_{\varepsilon, P_1^\varepsilon, \dots, P_K^\varepsilon}$ is a solution of (1.2).

By our construction, it is easy to see that by the maximum principle $u_\varepsilon > 0$ in Ω . Moreover $\varepsilon^N J_\varepsilon(u_\varepsilon) \rightarrow \frac{K}{2} I(w)$ and u_ε has only K local maximum

points $Q_1^\varepsilon, \dots, Q_K^\varepsilon$ and $Q_i^\varepsilon \in \partial\Omega$. By the structure of u_ε we see that (up to a permutation) $Q_i^\varepsilon - P_i^\varepsilon = o(1)$. This proves Theorem 1.2.

Theorem 1.1 follows from Theorem 1.2 by taking $\Gamma_i = \Gamma, i = 1, \dots, K$.

Finally, we prove Corollary 1.3.

If Ω is not a ball, then $H(P)$ has a local minimum on some open set Γ , Theorem 1.1 can be applied.

If Ω is a ball, Corollary 1.3 follows by minimizing energy in symmetric spaces. See [20] and [22].

□

REFERENCES

- [1] Adimurthi, Mancinni, G., and Yadava, S. L., *The role of mean curvature in a semilinear Neumann problem involving the critical Sobolev exponent*, Comm. P.D.E. 20, 1995, pp. 591-631.
- [2] Adimurthi, Pacella, F., and Yadava, S. L., *Interaction between the geometry of the boundary and positive solutions of a semilinear Neumann problem with critical nonlinearity*, J. Funct. Anal. 113, 1993, pp. 318-350.
- [3] Adimurthi, Pacella, F., and Yadava, S. L., *Characterization of concentration points and L^∞ -estimates for solutions involving the critical Sobolev exponent*, Diff. Int. Eqn. 8(1), 1995, pp. 41-68.
- [4] Agmon, S., *Lectures on Elliptic Boundary Value Problems*, Van Nostrand, Princeton, 1965.
- [5] Aronson, D. G. and Weinberger, H. F., *Multidimensional nonlinear diffusion arising in population genetics*, Adv. in Math. 30, 1978, pp. 33-76.
- [6] Dancer, E.N., *A note on asymptotic uniqueness for some nonlinearities which change sign*, Rocky Mountain Math. J., to appear.
- [7] Floer, A. and Weinstein, A., *Nonspreading wave packets for the cubic Schrödinger equation with a bounded potential*, J. Funct. Anal. 69, 1986, pp. 397-408.
- [8] Gardner, R. and Peletier, L. A., *The set of positive solutions of semilinear equations in large balls*, Proc. Roy. Soc. Edinburgh 104 A, 1986, pp. 53-72.
- [9] Gidas, B., Ni, W.-M., and Nirenberg, L., *Symmetry of positive solutions of nonlinear elliptic equations in R^n* , Mathematical Analysis and Applications, Part A, Adv. Math. Suppl. Studies Vol. 7A, pp. 369-402, Academic Press, New York, 1981.
- [10] Gilbarg, D. and Trudinger, N.S., *Elliptic Partial Differential Equations of Second Order*, 2nd edition, Springer, Berlin, 1983.
- [11] Gui, C., *Multi-peak solutions for a semilinear Neumann problem*, Duke Math. J. 84, 1996, pp. 739-769.
- [12] Gui, C. and Ghoussoub N., *Multi-peak solutions for a semilinear Neumann problem involving the critical Sobolev exponent*, preprint.
- [13] Helffer, B. and Sjöstrand, J., *Multiple wells in the semi-classical limit I*, Comm. PDE 9 1984, pp. 337-408.
- [14] Jang, J., *On spike solutions of singularly perturbed semilinear Dirichlet problems*, J. Diff. Eqns. 114, 1994, pp. 370-395.

- [15] Kwong, M. K., *Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in R^n* , Arch. Rational Mech. Anal. 105, 1989, pp. 243-266.
- [16] Li, Y. Y., *On a singularly perturbed equation with Neumann boundary condition*, Comm. in PDE., to appear.
- [17] Lin, C., Ni, W.-M., and Takagi, I., *Large amplitude stationary solutions to a chemotaxis systems*, J. Diff. Eqns. 72, 1988, pp. 1-27.
- [18] Lions, J.L. and Magenes, E., *Non-Homogeneous Boundary Value Problems and Applications*, Vol I, Springer-Verlag, New York, Berlin, Heidelberg, Tokyo, 1972.
- [19] Ni, W.-M., Pan, X., and Takagi, I., *Singular behavior of least-energy solutions of a semilinear Neumann problem involving critical Sobolev exponents*, Duke Math. J. 67, 1992, pp. 1-20.
- [20] Ni, W.-M. and Takagi, I., *On the shape of least energy solutions to a semilinear Neumann problem*, Comm. Pure Appl. Math. 41, 1991, pp. 819-851.
- [21] Ni, W.-M. and Takagi, I., *Locating the peaks of least energy solutions to a semilinear Neumann problem*, Duke Math. J. 70, 1993, pp. 247-281.
- [22] Ni, W.-M. and Takagi, I., *Point-condensation generated by a reaction-diffusion system in axially symmetric domains*, Japan J. Industrial Appl. Math. 12, 1995, 327-365.
- [23] Ni, W.-M. and Wei, J., *On the location and profile of spike-layer solutions to singularly perturbed semilinear Dirichlet problems*, Comm. Pure Appl. Math. 48, 1995, pp. 731-768.
- [24] Oh, Y. G., *Existence of semi-classical bound states of nonlinear Schrödinger equations with potentials of the class $(V)_a$* , Comm. PDE 13 (12), 1988, pp. 1499-1519.
- [25] Oh, Y. G., *On positive multi-lump bound states of nonlinear Schrödinger equations under multiple-well potentials*, Comm. Math. Phys. 131, 1990, pp. 223-253.
- [26] Pan, X. B., *Condensation of least-energy solutions of a semilinear Neumann problem*, J. Partial Diff. Eqns. 8, 1995, pp. 1-36.
- [27] Pan, X. B., *Condensation of least-energy solutions: the effect of boundary conditions*, Nonlinear Analysis, TMA 24, 1995, pp. 195-222.
- [28] Pan, X. B., *Further study on the effect of boundary conditions*, J. Diff. Eqns. 117, 1995, pp. 446-468.
- [29] Smoller, J. and Wasserman, A., *Global bifurcation of steady-state solutions*, J. Diff. Eqns. 39, 1981, pp. 269-290.
- [30] Wang, Z.-Q., *On the existence of multiple single-peaked solutions for a semilinear Neumann problem*, Arch. Rational Mech. Anal. 120, 1992, pp. 375-399.
- [31] Wei, J., *On the construction of single-peaked solutions of a singularly perturbed semilinear Dirichlet problem*, J. Diff. Eqns. 129, 1996, pp. 315-333.
- [32] Wei, J., *On the effect of the geometry of the domain in a singularly perturbed Dirichlet problem*, Diff. Int. Eqns., to appear.
- [33] Wei, J., *On the boundary spike layer solutions of singularly perturbed semilinear Neumann problem*, J. Diff. Eqns. 134, 1997.
- [34] Wei, J., *On the construction of single interior peak solutions for a singularly perturbed Neumann problem*, submitted.
- [35] Wei, J. and Winter, M., *Stationary solutions for the Cahn-Hilliard equation*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 1997, to appear.
- [36] Wei, J. and Winter, M., *Multiple Boundary Spike Solutions for a Wide Class of Singular Perturbation Problems*, J. London Math. Soc., to appear.

- [37] K. Yosida, *Functional Analysis*, fifth edition, Springer, Berlin, Heidelberg, New York, 1978.
- [38] Zeidler, E., *Nonlinear Functional Analysis and its Applications I, Fixed-Point Theorems*, Springer, New York, Berlin, Heidelberg, Tokyo, 1986.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER,
BC V6T 1Z2, CANADA

E-mail address: `cgui@math.ubc.ca`

DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN,
HONG KONG

E-mail address: `wei@math.cuhk.edu.hk`

MATHEMATISCHES INSTITUT A, UNIVERSITÄT STUTTGART, D-70511 STUTTGART,
GERMANY

E-mail address: `winter@mathematik.uni-stuttgart.de`