

CONSERVATIVE MANAGEMENT OF LOW BACK PAIN

For International Journal of Clinical Research

Andrew Frank and Lorraine De Souza

Front page
INTRODUCTION
EPIDEMIOLOGY
THE NATURE OF LOW BACK PAIN
DIFFERENTIAL DIAGNOSIS
PREVENTION
ACUTE LOW BACK PAIN
 Drugs
 Physical management
CHRONIC LOW BACK PAIN
 Medication
INTRACTABLE LOW BACK PAIN
 Physical management
 Psychological and social management
SERVICE IMPLICATIONS
SUMMARY



Tables

1. Patterns of referred pain from the spine
2. Potential factors predisposing to spinal pain
3. Estimated consequences for the back pain epidemic in the UK in 1993
4. Features of non-specific (mechanical) low back pain
5. Some causes of low back pain
6. Features suggestive of underlying illness
7. Guidelines for radiological investigation
8. Strategies for preventing back pain
9. Ergonomic approaches to minimising industrial back injuries
10. Yellow flags for the psychological risks of chronicity
11. The initial consultation for acute low back pain
12. Medication for acute low back pain
13. Physical barriers to successful outcome
14. Psychological barriers to successful outcome
15. Factors influencing return to work



INTRODUCTION

Back pain is a world-wide experience. Disabling back pain appears to be a problem for western and industrialised societies, possibly related to the development of “welfare” states. Disabling back pain is a major problem in the United Kingdom (UK) - see Fig. 1

¹, the United States of America (USA) ² and elsewhere ³. In this review the term "low back pain" will imply the symptom complex where pain is localised to the lumbar spine or refers into the leg or foot and where other specific conditions causing such pain have been excluded. Other clinically important patterns of referred pain are given in Table 1.

Explanations for back pain may be found in factors shown in Table 2. Whilst chronic pain may stem from disc, facet joints and the supporting soft tissues (particularly the ligaments), psychosocial factors are important. Indeed the concept of pain includes both sensory and emotional components - see Glossary.

Medical training probably inhibits a satisfactory approach by doctors by concentrating on the exclusion of pathology, when patients need understanding of their problems ⁴, alleviation of their symptoms and encouragement that activity is not harmful, but therapeutic ⁵.

The individual's strengths and weaknesses are important e.g. in their positive or negative responses, reactions of family and colleagues, ability to "fight" to prevent recurrences etc. Thus a positive attitude is required from "professional advisors" from the initial acute attack onwards. "Person management" is a prerequisite to back pain management.

The spine is the most complex part of the skeleton, constituting 139 joints, 24 discs, numerous bursae, ligaments and muscles comprising support structures. It is designed to take the weight of the upper part of the body, to allow movement and to protect the spinal cord - and thus it goes wrong more often than any other part of the musculoskeletal system. Patients need to understand this. Pain may arise from any spinal structure except bone, as they all contain nerve endings which could contain pain receptors ⁶.

The difficulties of assessing studies on therapy in low back pain have been extensively reviewed elsewhere ⁷⁻⁹, but benefit has been shown for increasing numbers of patients through randomised controlled trials. The Cochrane Library has been used to assist evaluation of such studies in this review.



EPIDEMIOLOGY

The “back pain epidemic”¹⁰ is of enormous economic proportions (Table 3), costing the National Health Service (NHS) £481-1061 million with indirect costs possibly exceeding £5-10 billion^{1,11,12}. Back pain is one of the commonest causes of inability to work through illness in the UK and makes enormous demands on the NHS and the private sector - see Fig. 2.

In the USA medical costs exceed \$24 billion¹³ and indirect costs may exceed \$30 billion¹⁴. Approximately 200,000 surgical procedures are performed annually. As many as 30,000 operations may be unsuccessful, joining the “high-cost, high-demand and highly emotional subset of the low back disabled”¹⁵. The chances of any individual coming to back surgery are 6 times greater in North America than in Europe.

The point prevalence of back pain is 7-14%,^{16,17} one year prevalence is 36-37% and life time prevalence is 58%¹⁸. Back pain frequently recurs¹⁹⁻²⁶. It is equally prevalent in both sexes²⁷, although its clinical course may be different; disc disease being more prevalent in men²⁸ who have more surgery²⁹ whilst in women it may be more likely to linger into chronicity³⁰. Pregnancy, abortions and parity are associated³¹. A causal relationship with epidural analgesia seems unlikely.

Jobs requiring physically heavy work, static work postures, frequent bending and twisting, lifting & forceful movements and repetitive work and vibration³²⁻³⁵, which includes that from driving vehicles^{28,36,37}, predispose to low back pain³⁸. Nurses doing the most physical work have a greater incidence than construction workers, garbage collectors and truck drivers³⁹. Psychological factors are implicated e.g. monotony and work dissatisfaction^{34,38,40,41}. Other risk factors include smoking^{20,28,36,42,43} (which may be linked with atherosclerosis of the abdominal aorta⁴⁴), road traffic accidents and falls⁴⁵.

Related social factors include a poorer economic situation; drug and alcohol abuse; divorces and family problems; self-reported sexual and physical abuse⁴⁶ and a lower level of education^{38,47}. Health associations include previous hospital admissions/operations²⁰ and a propensity to use health services; cardiovascular and gastrointestinal symptoms⁴⁸.



THE NATURE OF BACK PAIN

Low back pain is classified into that caused by mechanical back pain (that arising from the lumbar segment without direct nerve root involvement); direct nerve root (or rarely the corda equina) involvement from structures within the lumbar segment; and spinal pathology.

Many treatments for low back pain are based on the assumption that pain relates either to excessive mechanical stresses (at work or leisure) on normal structures, normal stresses on degenerate structures or the poorly resolved aftermath of an acute episode⁴⁹. Psychological management for both acute³⁰ and chronic back pain^{1,50} is important. In 1959 it was noted that 'the patient's ability to "live with his back" seems to be more important than the form of treatment'⁵¹.

Patients should be treated optimistically avoiding bed rest and hospitalisation if possible⁵². Waddell views the "back pain epidemic" as partly being contributed to by doctors prescribing rest and partly to social concepts of disability and contrasts the traditional medical model of illness with a biopsychosocial model^{1,50}.

Many episodes of pain do not resolve^{53,54}. Croft has shown that patients with back pain may have persisting symptoms even though they may not return to the general practitioner (GP). The risk of an exacerbation or a recurrent attack is always present. Management must emphasise the importance for patients of adopting strategies that minimise the risks of recurrences.

Chronic pain is multifactorial with obstruction of the epidural veins and fibrosis around the nerve roots giving rise to perineural atrophy⁵⁵. The dorsal horn may be involved accounting for hyperpathia and allodynia, explaining widespread sensitivity of the back and lower limbs to pressure. Sympathetically maintained pain may arise from altered central neuro-modulation within the spinal cord giving rise to syndromes similar to regional pain syndromes⁵⁶.



DIFFERENTIAL DIAGNOSIS

There is a delicate balance between not missing treatable pathology, and avoiding unnecessary investigation which may increase patient's fears about their condition⁵⁷. It is achieved by pattern recognition of common presentations of low back pain (Table 4). Non-specific low back and/or leg pain (NSLBP) should be diagnosed on positive grounds (Table 4). There are 3 important patient groups to consider - those with NSLBP; compression of the nerve root or cauda equina; and back pain caused by other conditions^{1,58}. A marked relationship to body posture is common, often aggravated by sitting or standing and eased by walking. Pain aggravated by walking raises the possibility of vascular claudication, spinal or lateral canal stenosis.

Spinal stenosis is a symptom complex of root pain and sensory or motor symptoms noted when walking and fading after a few minutes sitting or flexing the spine⁵⁹. More complex algorithms can be consulted^{59,60} and fuller reviews of the differential diagnosis are available^{57,61,62} (Table 5).

Thoracic pain is less common than lumbar or cervical pain and requires investigation¹. Vertebral collapse may be due to osteoporosis but myeloma or secondary malignancy need to be excluded. Loss of height is almost universal with age and usually reflects loss of disc height. Most kyphotic spines reflect degenerative processes.

Serious causes of back pain are present only in a small minority of patients referred to hospital and are suspected from "red flags"¹ (Table 6). Some vascular and neurological disorders, osteoarthritis and osteoporosis may not have these stigmata!

Pelvic pathology is hard to exclude on history as many women notice their pain to be worse in the last few days of their menstrual cycle, easing off early in their period. This may also be noted in women with other pains or headaches. Where there is doubt, abdominal and pelvic examinations are required.

Having excluded sinister diseases, exclusion of root or cord compression may require imaging. A scheme for radiological investigation is given in Table 7. Discussion with a musculoskeletal radiologist is invaluable.



PREVENTION

Back pain and subsequent disability (including work loss) can be prevented (Table 8).

Primary prevention eliminates/minimises the risks to health or well being ⁶³. Corporate stress management programmes at three high risk work sites have reduced the number of accidents and related costs ⁶⁴. Low back injuries are the commonest and most costly injuries in terms of compensation and productivity in the US railroad industry and are correctable (Table 9). Low back injuries and lost work days fell to zero and absenteeism fell from 4-1% ⁶⁵. Introducing a back injury prevention programme in California showed a net benefit to be \$161,108 with a return on investment of 179 percent ⁶⁶.

A large study showed no reduction in among postal workers who had been instructed in back pain prevention ⁶⁷. Teaching and social support may not alter behaviour ⁶⁸ but in a work environment may be more effective ⁶⁹.

Secondary prevention is the alleviation of symptoms, minimising residual disability and factors that may cause a recurrence ⁶³.

This topic is addressed in the sections on back schools and intensive rehabilitation. It is the combination of intensive physical training, behavioural principles and attention to the working environment that is effective. Back schools may be more effective in the work environment ⁶⁴.

Tertiary prevention implies the rehabilitation of those with disabilities to optimal function and modification of the workplace to accommodate residual disability ⁶³.

Studies reporting follow-up data of 6 months duration may be considered preventive⁶⁴. Cognitive behavioural therapy is effective for many patients ⁷⁰ and rehabilitative approaches are discussed below.

Back education from specially trained physiotherapists during pregnancy reduced the amount of back pain during pregnancy, sick leave during pregnancy and postpartum back pain ⁷¹.



ACUTE LOW BACK PAIN

Correct management of acute back pain in primary care is crucial to the prevention of long-term back pain and disability¹. Consequently, guidelines for the initial management have been issued by many bodies^{1,2,72-75}. All concentrate on the critical initial triage of patients into three groups - those with non-specific low back pain (table 4), those with nerve root compression or those with other diseases presenting as low back pain (Tables 5,6). They use “red flags” (Table 6) to alert clinicians to serious pathology^{1,73}.

Advice on the history, examination and key management is given in Table 10. Provision of literature prevents unnecessary treatment including surgery⁷⁶; lessons harmful beliefs, and reduces disability⁷⁷. The patient must be taken seriously⁷⁸. In the absence of root signs or suggestions of other pathology, advice that activity is helpful, clicks are not harmful and that pain does not signify serious illness are all important.

Without neurological involvement, bed rest beyond 2 days is counterproductive^{1,75,79}. Analgesia assists this policy, as will the reassurance given by the physical assessment performed by the GP. Bed rest is usually best in the position of maximum comfort^{57,80,81}. Provision of a corset, assessment of furniture heights to assist transfers, domiciliary physiotherapy and a commode by the bed may all assist mobilisation.

Drugs

Medication is important (Table 11) as severe pain inhibits compliance and a speedy return to work. It minimises unnecessary rest to control pain which enhances the ability to regain normal activities quickly. A non-steroidal anti-inflammatory drug (NSAID) is often helpful⁸, particularly to obtain sufficient duration of analgesia for a good sleep⁸². The risks of short courses may be no more than the risks of prescribing stronger opiate derivatives. Care must be taken in smokers, those with a history of dyspepsia or elderly people. Inability to control pain at home without using strong opiates is an indication for admission to hospital in my practice. Further information is available in standard reviews/texts^{2,57,83-86}.

Physical management

Considerable evidence supports the role of physical therapy in acute/subacute back pain. Increased compliance with better results from physiotherapy can be obtained by simultaneous use of complementary literature and by planned review of patients after treatment⁸⁷. Physical therapy⁸⁸; chiropractic^{89,90}; probably osteopathic manipulation⁹¹; manipulation by physical therapists^{9,58,92-95} psychological management (with more supervision and planned withdrawal of treatment (behavioural group)); McKenzie techniques of passive extension and postural correction^{21,23}; graded activity programmes with a behavioural therapy approach⁹⁶ and "behavioural support" in a “sports injury” approach^{97,98} can reduce pain and disability and in some cases work



absence, compensation and disability award costs. Corsets should be reserved for those with failed manual therapy^{49,99}.

As cervical symptoms often co-exist with low back symptoms (Frank et al - unpublished data), advice about physical and ergonomic measures must be appropriate for all parts of the spine. Thus lying prone may help some with low back pain, but aggravate mild or asymptomatic neck problems.

An international group has now produced a series of psychological risk factors for chronicity (Table 12). These “yellow flags” may prevent development of excess pain behaviour / sick role, recurrence/re-injury and the long-term complications of disability - loss of function, independence and work⁶³.

Back pain in elderly people is an important topic which has been reviewed elsewhere⁸¹.



Chronic low back pain

Exercise regimes aim to increase range of movement, strengthen muscles, stretch tightened structures or toughen up physically and mentally^{49,57,100}. Evidence supports the following exercise regimes:- back extension^{87,101-105}; calisthenics¹⁰⁵; mixed exercise regimes¹⁰⁶; isometric flexion exercises^{107,108}; mobilising exercises¹⁰⁸; and McKenzie^{23,109}.

Exercises combined with behavioural methods are more effective¹¹⁰, reduce sickness behaviour¹¹¹ and get people back to work quicker⁹⁶. Increasing the range of spinal movement does not necessarily reduce pain or disability¹¹². Aerobic exercise is beneficial^{113,114} and may improve job satisfaction³⁹.

Intensive outpatient physical retraining consisting of pain relief & mobilization, increasing movement & muscle strengthening and work conditioning reduces work absence. Health care costs were offset by savings in "wages loss cost"^{97,98}.

Many studies^{89,90,92,93} and reviews^{94,115} support the view that manual therapy including manipulation is effective in reducing chronic pain although this view is not universally accepted¹¹⁶. Studies comparing manipulation with exercise (or manipulation + exercise) are currently taking place.

Corsets for unselected patients are less effective than active therapy¹¹⁷. Patients with failed manual therapy, pain on physical activities (particularly if it facilitates remaining at work) and those with scoliosis or unstable spondylolisthesis may benefit.

Traction helps a few patients though demonstration of its benefits in controlled studies is lacking¹¹⁸. Bed rest in traction is not recommended¹.

The background of chiropractic and osteopathy has been reviewed¹¹⁹. Chiropractic has been studied^{89,90,120} and appears effective. The public need to be safeguarded against those using unproved techniques which may be harmful^{121,122}. Doctors are advised not to recommend heterodox treatment unless the practitioner is a member of a recognised professional association¹²³.

Medication

Many people strive to avoid medication "on principle". Whilst understandable, this can be counter-productive and patients must be aware of the side-effects of not taking drugs e.g. constant pain, loss of sleep, irritability and its effects on family life, loss of function at home and work.

Initially, regular analgesia is advised using the simplest effective medication (Table 11). Long-acting preparations may be available. NSAIDs may be useful but long-term they are best avoided in view of their side-effects¹²⁴.

Stronger preparations e.g. Tramadol have been found particularly useful recently¹²⁴ but longer term analgesia encourages sedation, nausea and constipation etc and many patients rightly discontinue their use. Having effective tablets "on hand" is prudent



and helps counter the fear of aggravation with activity and may inhibit the risks of depression⁴⁹, in addition to giving the patient some control of their pain.

Opiate use is controversial^{84,125}. In my opinion, if patients are deemed to need opiates, they should be assessed (probably as an inpatient) to exclude treatable causes of pain, sort out psychosocial issues, treat depression if present and to commence aggressive rehabilitation.

A major advance in chronic pain management is the use of tricyclic antidepressant compounds (TCA)^{84,124,126-128} for pain modulation. They appear more effective in the presence of root compression. Benefits are shown for those with or without depression and for those with a physical basis for their pain. The sedative side effects are helpful in those having disturbed sleep and they may facilitate relaxation. Appropriate warnings about sedation and a dry mouth must be given. Patients will need reassurance that TCAs are not addictive or being prescribed for psychiatric purposes. Amitriptyline, Dothiepin (Dosulepin) and Lofepramine have all been used. The role of serotonin antagonists in chronic pain has only recently been investigated. It is likely that they have no role in chronic pain, only in the treatment of the depression that it engenders¹²⁹.



INTRACTABLE LOW BACK PAIN

A few patients have persisting disabling pain despite the above management. Investigations, often including magnetic resonance imaging (MRI) have excluded systemic illness or demonstrated that the pathology is degenerative and not operable. The term "intractable" pain differentiates failed conservative management from previously untreated chronic pain, or "chronic pain syndrome", when changes in behaviour are apparent⁵⁸. Up to 10% of these patients may have an underlying medical problem¹³⁰.

At this stage both physical and psychological factors are likely to be significant^{49,131}, either predominating. Although they are described separately for clarity, both aspects must progress simultaneously.

Physical management

Clues that physical aspects are important lie in the pattern of pain which is often episodic and related to movement or posture (Table 13). Sleep patterns are important. After checking bedding^{49,57}, it must be established whether sleep loss is due to insomnia, pain or depression. Unvarying pain suggests that depression may be present. If pain disturbs sleep long-acting NSAIDs or analgesics are needed. If the pain is post-traumatic, flashbacks and nightmares need to be excluded^{132,133}.


There is limited evidence to support transcutaneous electrical nerve stimulation (TENS)¹³⁴ or acupuncture, although opinions vary^{135,136}. Serious adverse events have been reported following acupuncture, but seldom from orthodox medical practice¹³⁷.

Initially analgesics and / or NSAIDs are used separately or in combination to facilitate control of pain and the commencement of physical therapy (Table 11). If unsuccessful, tricyclic compounds may be added to the regime. Ultimately the objective is to wean of all regular analgesics and NSAIDs, using tricyclic compounds to facilitate this if needed. TCAs can be sometimes withdrawn over time. Patients with clinical depression and who do not tolerate TCAs will need SSRIs.

Injections may be given into the epidural space, facet joints, discs or the soft tissues. There is little evidence to support their use¹³⁸. They do not appear to get patients back to work¹³⁹. A recent study of sclerosants was unhelpful¹⁴⁰. Injections may establish the source of pain¹⁴¹ with psychological benefits and if transitory pain relief is achieved, intensive therapy may be effective.

Psychosocial management

A significant group of patients present with constant unchanging pain. Many suffer from the "pain or illness behaviour syndrome"^{50,142}, sometimes considered as "learned helplessness"¹⁴³. The effect on the spouse or "significant other" needs evaluation¹⁴³ particularly if over-protection or excessive sympathy has encouraged the adoption of a "sick/patient role". Patients may have problems consequent to loss of earnings or changing marital/sexual relationships^{49,57}. Often unrealistic fears exist - that long-standing pain must be caused by horrific disorders e.g. cancer or multiple



sclerosis⁵⁷; of long-term disability¹⁴⁴ or that activity will aggravate the pain¹⁴⁵. A disproportionate number appear to have post-traumatic pain. Issues relating to compensation, post-traumatic stress disorder and post-traumatic weight gain are discussed elsewhere^{132,133} and are often associated with depression.

Examination may reveal a non-anatomical pattern of pain¹⁴⁶ and is often accompanied by exaggerated gestures, particularly during the straight leg raise test¹⁴⁷. These patients used to be considered malingers, or to exhibit "functional overlay". It is now recognised that this pattern of behaviour reflects "distress" - an unconscious desire on the part of the patient to show the doctor just how much they hurt -often believing that their suffering has been trivialised by doctors and others¹⁴⁸.

Cognitive behavioural therapy (CBT) is the cornerstone of psychological pain management and is effective in chronic pain patients (about half suffering from back pain)⁷⁰. Groups help in boosting confidence through sharing experiences. Self-help may be facilitated by appropriate literature e.g. "Living with your pain"¹⁴⁹. Sometimes specialist counselling is needed.

Patients attitudes and beliefs about back pain, behaviours, compensation and emotional issues can all be noted early during the subacute period⁶³. Many of the psychological factors which may inhibit resolution of symptoms (Table 14) are amenable to psychological or psychiatric intervention.

Usually both physical and behavioural approaches are combined simultaneously^{30,96,110}. Pre-existing psychopathology is frequent in those requiring intensive rehabilitation¹⁵⁰ and does not militate against a successful outcome¹⁵¹.

"Back Schools" may mean any programme with an educational content varying from one outpatient session to an inpatient programme^{21,152-155}. A recent review¹⁵⁶ suggests they are effective, particularly in occupational settings.

Intensive rehabilitation programmes involve multi-professional physical, psycho-social, educational and vocational components which include learning to function in spite of pain, CBT, and developing physical tolerance in both fitness¹¹³ and vocational areas - e.g. the Helsinki^{155,157,158}, Texas¹⁵⁹ and Turku¹¹¹ programmes. Such programmes embrace the concepts of back school education, work hardening¹⁵⁹, sports injury approach^{97,98} and functional restoration¹⁶⁰⁻¹⁶². Results may be better if the programme is inpatient¹⁵⁵. The key features of facilitating a return to work are given in Table 15.

Advice on self care, living and coping with low back pain is explored elsewhere⁵⁷.



SERVICE IMPLICATIONS

Back pain and disability can be ameliorated and further episodes minimised by professional intervention at the initiation of the first episode through manipulation, physical training, a variety of exercise programmes and multi-professional rehabilitation. Minimal rest and a planned withdrawal of support are important developments with demonstrable benefit.

Delays in obtaining specialist advice contribute to non-resolution of acute/subacute back pain, increasing the pool of chronic pain patients and the risk of pain becoming intractable. These people may progress to pain behaviour, becoming more costly for employers and the state. Multi-professional integrated programs within 3 months can abort this process ⁹⁶.

Diagnostic and treatment protocols can decrease accidents and lost work days and be cost effective ⁶⁰ if costs to all sections of national life (state and industry) are considered. Good protocols exist for managing acute back pain but investment is needed to establish cost-effective intensive rehabilitation programmes in many countries.



11 SUMMARY

Although back pain appears prevalent throughout the world, back pain disability has now reached epidemic proportions in western (and probably most industrialised) societies. Few patients with back pain have serious medical pathology or direct involvement of nerve roots requiring surgery. Although the causes of back pain remain unclear, physical stress and its consequences on discs, facet joints and supporting soft tissues at work or leisure are thought to be mostly implicated.

Modern management emphasises the role of self-care which should begin in general practice at presentation of the first episode and be reinforced by all health professionals. Use of guidelines for the management of acute low back pain should greatly improve management of the acute episode. In the absence of root compression, bed rest should not exceed 48 hours. A positive approach is encouraged, acknowledging that returning to a normal life may require working through pain. Emphasis is on encouraging a rapid return to physical fitness and other activities, including employment. Medication facilitates this.

Studies demonstrate the ability to lessen the duration and severity of individual episodes of back pain and reduce recurrences and their cost in terms of suffering and lost work. No one should remain in pain longer than 6 weeks without being referred to a specialist service. This should include provision of information, access to manual therapists with postgraduate training, education on back strain prevention and physical fitness; clinical psychology pain services and consultant support. Comprehensive (and inpatient) pain and rehabilitation services should be available.

Future challenges lie in detecting those who are unlikely to respond to physical measures and back education so that developing coping strategies, reducing the inhibitory effects of fear and managing social issues may reduce those who require costly, intensive and probably inpatient rehabilitation.



Bibliography

Ref ID : 132

1 Clinical Standards Advisory Group - Chairman Prof M Rosen. Back Pain. 1994;1-89.

Ref ID : 1204

2 Agency for Health Care Policy and Research. Acute low back problems in adults. Clinical practice Guideline Number 14: Acute low back problems in adults: assessment and treatment. Quick reference guide for clinicians number 14: Understanding low back problems. Patient Guide. 1994;**AHCPR/PUB-95-0642**:1-200.

Ref ID : 816

3 Volinn E. The epidemiology of low back pain in the rest of the world. A review of surveys in low and middle income countries. *Spine* 1997;**22**:1747-1754.

Ref ID : 141

4 Armstrong D. What do patients want? *Br Med J* 1991;**303**:261-262.

Ref ID : 628

5 Indahl A, Velund L, Reikeraas O. Good prognosis for low back pain when left untampered: a randomized clinical trial. *Spine* 1995;**20**:473-477.

Ref ID : 703

6 Porter R. The nociceptive source. In: Porter R, ed. *Management of back pain*. Edinburgh: Churchill Livingstone, 1993;45-57.

Ref ID : 133

7 Spitzer W, LeBlanc F, Dupuis M. Scientific approach to the assessment and management of activity-related spinal disorders. A monograph for clinicians. *Spine* 1987;**12**:S1-S59.

Ref ID : 150

8 Deyo R. Conservative therapy for low back pain. Distinguishing useful from useless therapy. *J Am Med Ass* 1983;**250**:1057-1062.

Ref ID : 248

9 Koes B, Assendelft W, Heijden Gv, Bouter L, Knipschild P. Spinal manipulation and mobilization for back and neck pain: a blinded review. *Br Med J* 1991;**303**:1298-1303.

Ref ID : 508

10 Editorial. The back pain epidemic. *Acta Orthop Scand* 1989;**60**:633-634.

Ref ID : 1077

11 Underwood M. Crisis: what crisis? *Eur Spine J* 1998;**7**:2-5.

Ref ID : 1328

12 Maniadakis N, Gray A. The economic burden of back pain in the UK. *Pain* 2000;**84**:95-103.

Ref ID : 155

13 Lahad A, Malter A, Berg A, Deyo R. The effectiveness of four interventions for the prevention of low back pain. *J Am Med Ass* 1994;**272**:1286-1291.

Ref ID : 156

14 Frymoyer J. Introduction. In: Hochschuler S, Cotler H, Guyer R, eds. *Rehabilitation of the spine. Science and Practice*. St. Louis. Mosby, 1993;xix-xxi.

Ref ID : 158

15 Hanley E. The cost of surgical intervention for lumbar disc herniation. In: Weinstein J, ed. *Clinical efficacy and outcome in the diagnosis and treatment of low back pain*. New York: Raven Press, 1992;125-133.



Ref ID : 165

16 Deyo R, Tsui-Wu Y. Descriptive epidemiology of low back pain and its related medical care in the United States. *Spine* 1987;**12**:264-268.

Ref ID : 683

17 Papageorgiou A, Croft P, Ferry S, Jayson M, Silman A. Estimating the prevalence of low back pain in the general population. Evidence from the South Manchester Back Pain Survey. *Spine* 1995;**20**:1889-1894.

Ref ID : 401

18 Clinical Standards Advisory Group. Epidemiology Review: The epidemiology and cost of low back pain. 1994;1-72.

Ref ID : 188

19 Abenhaim L, Suissa S, Rossignol M. Risk of recurrence of occupational back pain over three year follow-up. *Br J Ind Med* 1988;**45**:829-833.

Ref ID : 194

20 Biering-Sorensen F, Thomsen C, Hilden J. Risk indicators for low back trouble. *Scand J Rehab Med* 1989;**21**:151-157.

Ref ID : 251

21 Stankovic R, Johnell O. Conservative treatment for acute low back pain. A prospective randomised trial: McKenzie method of treatment versus patient education in "Mini Back School". *Spine* 1990;**15**:120-123.

Ref ID : 434

22 Roland M, Morrell D, Morris R. Can general practitioners predict the outcome of episodes of back pain? *Br Med J* 1983;**286**:523-525.

Ref ID : 845

23 Stankovic R, Johnell O. Conservative treatment of acute low back pain. A five year follow-up study of two methods of treatment. *Spine* 1995;**20**:469-472.

Ref ID : 846

24 Watson P, Main C, Waddell G, Gales T, Purcell-Jones G. Medically certified work loss, recurrence and costs of compensation for back pain: a follow-up study of the working population of Jersey. *Br J Rheum* 1998;**37**:82-86.

Ref ID : 1065

25 Skargren E, Carlsson P, Oberg B. One year follow-up comparison of the cost and effectiveness of chiropractic and physiotherapy as primary management of back pain. *Spine* 1998;**23**:1875-1884.

Ref ID : 1197

26 Lonn J, Glomsrod B, Soukup M, Bo K, Larsen S. Active back school: prophylactic management for low back pain. *Spine* 1999;**24**:865-871.

Ref ID : 169

27 Walsh K, Cruddas M, Coggan D. Low back pain in eight areas of Britain. *J Epid and Community Health* 1992;**46**:227-230.

Ref ID : 176

28 Kelsey J, Githens P, O'Connor T, et al. Acute prolapsed lumbar intervertebral disc. An epidemiologic study with special reference to driving automobiles and cigarette smoking. *Spine* 1984;**9**:608-613.



Ref ID : 160

29 Davis H. Increasing rates of cervical and lumbar spine surgery in the United States 1979-90. *Spine* 1994;**19**:1117-1124.

Ref ID : 149

30 Fordyce W, Brockway J, Bergman J, Spengler D. Acute back pain: a control-group comparison of behavioural vs traditional management methods. *J of Behav Med* 1986;**9**:127-140.

Ref ID : 174

31 Svenson H, Andersson G, Hagstad A, Jansson P. The relationship of low back pain to pregnancy and gynecologic factors. *Spine* 1990;**15**:371-375.

Ref ID : 196

32 Editorial. Risk factors for back trouble. *Lancet* 1989;**1**:1305-1306.

Ref ID : 202

33 Magnusson M, Pope M, Wilder D, Areskoug B. Are occupational drivers at an increased risk for developing musculoskeletal disorders? *Spine* 1996;**21**:710-717.

Ref ID : 1043

34 Barnekow-Bergvist M, Hedberg G, Janlert U, Jansson E. Determinants of self-reported neck - shoulder and low back symptoms in a general population. *Spine* 1998;**23**:235-243.

Ref ID : 1336

35 Kumar A, Varghese M, Mohan D, Mahajan P, Gulati P, Kale S. Effect of whole body vibration on the low back: a study of tractor-driving farmers in North India. *Spine* 1999;**24**:2506-2515.

Ref ID : 175

36 Frymoyer J, Pope M, Clements J, Wilder D, Macpherson B, Ashikaga T. Risk factors in low back pain. *J Bone and Joint Surg* 1983;**65-A**:213-218.

Ref ID : 177

37 Behrens V, Seligman P, Cameron L, Mathias C, Fine L. The prevalence of back pain, hand discomfort, and dermatitis in the U.S. working population. *Am J Public Health* 1994;**84**:1780-1785.

Ref ID : 178

38 Andersson G. Epidemiological aspects on low back pain in industry. *Spine* 1981;**6**:53-60.

Ref ID : 182

39 Kaplan R, Deyo R. Back pain in hospital workers. In: Deyo R, ed. *SPINE: State of the Art reviews: occupational back pain*. Philadelphia: Hanley and Belsus, 1987;61-73.

Ref ID : 179

40 Bergenudd H, Nilsson B. The prevalence of locomotor complaints in middle age and their relationship to health and socioeconomic factors. *Clin Orth and Related Res* 1994;**308**:264-270.

Ref ID : 180

41 Bigos S, Battie M, Spengler D, et al. A prospective study of work perceptions and psychosocial factors affecting the report of back injury. *Spine* 1991;**16**:1-6.

Ref ID : 193

42 Battie M, Videman T, Gill K, et al. Smoking and lumbar intervertebral disc degeneration. An MRI study of identical twins. Volvo award in clinical sciences. *Spine* 1991;**16**:1015-1021.

Ref ID : 195



43 Dimberg L, Olafsson A, Stafansson E, et al. Sickness absenteeism in an engineering industry - an analysis with special reference to absence for neck and upper extremity symptoms. *Scand J Rehab Med* 1989;**17**:77-84.

Ref ID : 199

44 Kauppila L, Penttila A, Karhunen P, Lalu K, Hannikainen P. Lumbar disc degeneration and atherosclerosis of the abdominal aorta. *Spine* 1994;**19**:923-929.

Ref ID : 197

45 Walsh K, Cruddas M, Coggan D. Risk of low back pain in people admitted to hospital for traffic accidents and falls. *J Epid and Comm Health* 1992;**46**:231-233.

Ref ID : 775

46 Linton S. A population-based study of the relationship between sexual abuse and back pain: establishing a link. *Pain* 1997;**73**:47-53.

Ref ID : 173

47 Croft P, Rigby A. Socioeconomic influences on back problems in the community in Britain. *J Epid and Comm Health* 1994;**48**:166-170.

Ref ID : 198

48 Porter R, Oakshot G. Spinal Stenosis and health status. *Spine* 1994;**19**:901-903.

Ref ID : 72

49 Frank A, Hills J. Spinal pain. In: Frank A, Maguire G, eds. *Disabling Diseases. Physical, Environmental and Psychosocial Management*. Oxford: Heinemann Medical Books. 1989;41-78.

Ref ID : 206

50 Waddell G. A new clinical model for the treatment of low back pain. 1987 Volvo award in clinical sciences. *Spine* 1987;**12**:632-644.

Ref ID : 204

51 Bremner R, Simpson M. Management of chronic lumbosacral strain. *Lancet* 1959;**2**:949-950.

Ref ID : 205

52 Wilson R, Wilson S. A ten year follow-up of cases of low back pain. *The Practitioner* 1964;**192**:657-660.

Ref ID : 208

53 Von Korff M, Deyo R, Cherkin D, Barlow W. Back pain in primary care. Outcomes at one year. *Spine* 1993;**18**:855-862.

Ref ID : 940

54 Croft P, Macfarlane G, Papageorgiou A, Thomas E, Silman A. Outcome of low back pain in general practice: a prospective study. *Br Med J* 1998;**316**:1356-1359.

Ref ID : 135

55 Jayson M. Mechanisms underlying chronic back pain. Studies are looking at the brain and spinal cord. *Br Med J* 1994;**309**:681-682.

Ref ID : 667

56 Jayson M. Why does acute back pain become chronic? chronic back pain is not the same as acute back pain lasting longer. *Br Med J* 1997;**314**:1639-1640.

Ref ID : 800

57 Frank A. *Low back pain: diagnosis and management*. London: Current Medical Literature, 2000;1-80 - in press

Ref ID : 1



58 Frank A. Low back pain - Regular Review. *Br Med J* 1993;**306**:901-909.

Ref ID : 137

59 Waddell G. An approach to backache. *Br J Hosp Med* 1982;**September**:187-219.

Ref ID : 213

60 Gavin M, Wiesel S. Low back pain. *Curr Opinion Rheumatol* 1991;**3**:65-70.

Ref ID : 217

61 Bogduk N. The sources of low back pain. In: Jayson M, ed. *The Lumbar Spine and Back Pain*. Edinburgh: Churchill Livingstone, 1992;61-88.

Ref ID : 152

62 Porter R. The causes of back pain: aetiology. In: Porter R, ed. *Management of back pain*. Edinburgh: Churchill Livingstone, 1993;19-27.

Ref ID : 725

63 Kendall N, Linton S, Main C. *Guide to assessing psychosocial yellow flags in acute low back pain: risk factors for long-term disability and work loss*. Wellington, New Zealand: Accident Rehabilitation and Compensation Insurance Corporation of NZ and the National Health Committee, 1997;1-22

Ref ID : 360

64 Weiser S, Cedraschi C. Psychosocial issues in the prevention of chronic low back pain - a literature review. *Clin rheumatol* 1992;**6**:657-684.

Ref ID : 361

65 Brown R, Page G, McMahan P. Ergonomics in the US Railroad Industry. *Human Factors Society Bull* 1991;**34**:1-4.

Ref ID : 327

66 Shi L. A cost benefit analysis of a California county's back injury prevention program. *Pub Health Reports* 1993;**108**:204-211.

Ref ID : 759

67 Daltroy L, Iverson M, Larson M, et al. A controlled trial of an educational program to prevent low back injuries. *New Eng J Med* 1997;**337**:322-328.

Ref ID : 362

68 Daltroy L, Iverson M, Larson M, et al. Teaching and social support: effects on knowledge, attitudes, and behaviours to prevent low back injuries in industry. *Health Ed Quarterly* 1993;**20**:43-62.

Ref ID : 363

69 Feldstein A, Valanis B, Vollmer W, Stevens N, Overton C. The back injury prevention project pilot study: assessing the effectiveness of Back Attack, an injury prevention program among nurses, aides, and orderlies. *J Occupat Med* 1993;**35**:114-120.

Ref ID : 1168

70 Morley S, Eccleston C, Williams Ad. Systematic review and meta-analysis of randomized controlled trials of cognitive behavioural therapy and behaviour therapy for chronic pain in adults, excluding headache. *Pain* 1999;**80**:1-13.

Ref ID : 183

71 Ostgaard H, Zetherstrom G, Roos-Hansson E, Svanberg B. Reduction of back and posterior pelvic pain in pregnancy. *Spine* 1994;**19**:894-900.

Ref ID : 294

72 Chartered Society of Physiotherapy. Guide to the management of low back pain for General Practitioners. 1994



Ref ID : 520

73 Waddell G, Feder G, McIntosh A, Lewis M, Hutchinson A. Low back pain evidence review. 1996;1-35.

Ref ID : 972

74 Bogduk N. The National Musculoskeletal Health initiative: an algorithm for back pain management. *Proceedings of the 6th meeting of the Australasian Faculty of Rehabilitation Medicine* 1998;18-19.(Abstract)

Ref ID : 1161

75 Waddell G, McIntosh A, Hutchinson A, Feder G, Lewis M. Low back pain evidence review. 1999;1-34

Ref ID : 228

76 Roland M, Dixon M. Randomized controlled trial of an educational booklet for patients presenting with back pain in general practice. *J R Coll Gen Pract* 1989;**39**:244-246.

Ref ID : 1339

77 Burton A, Waddell G, Tillotson K, Summerton N. Information and advice to patients with back pain can have a positive effect: a randomized controlled trial of a novel educational booklet in primary care. *Spine* 1999;**24**:2484-2491.

Ref ID : 136

78 Roland M. Back pain in General Practice. Taking a positive view. *Musculoskeletal Medicine* 1994;**1**:2-4.

Ref ID : 1242

79 Bassols A, Bosch F, Campillo M, Canellas M, Banos J. An epidemiological comparison of pain complaints in the general population of Catalonia. *Pain* 1999;**83**:9-16.

Ref ID : 81

80 Frank A. Spinal Problems in Adults. In: Maddison P, Isenberg D, Woo P, Glass D, eds. *Oxford Textbook of Rheumatology*. Oxford: Oxford University Press, 1998;89-114.

Ref ID : 1216

81 Frank A. Don't neglect back pain. *Health and Aging* 1999;33-36.

Ref ID : 234

82 Szpalski M, Hayez J. Objective functional assessment of the efficacy of Tenoxicam in the treatment of acute low back pain. A double-blind placebo-controlled study. *Br J Rheum* 1994;**33**:74-78.

Ref ID : 704

83 Porter R, Ralston S. Pharmacological management of back pain syndromes. *Drugs* 1994;**48**:189-198.

Ref ID : 788

84 Deyo R. Drug therapy for back pain. Which drugs help patients? *Spine* 1996;**21**:2840-2850.

Ref ID : 1039

85 Cherkin D, Wheeler K, Barlow W, Deyo R. Medication use for low back pain in primary care. *Spine* 1998;**23**:607-614.

Ref ID : 1492

86 Innes G, Croskerry P, Worthington J, Beveridge R, Jones D. Ketorolac versus acetaminophen-codeine in the emergency department treatment of acute low back pain. *J Emerg Med* 1998;**16**:549-556.



Ref ID : 240

87 Manniche C, Lundberg E, Christensen I, Bentzen L, Hesselsoe G. Intensive dynamic back exercises for chronic low back pain: a clinical trial. *Pain* 1991;**47**:53-63.

Ref ID : 244

88 Overman S, Larson J, Dickstein D, Rockey P. Physical therapy care for low back pain: monitored program of first-contact nonphysician care. *Physical Therapy* 1988;**68**:199-207.

Ref ID : 2

89 Meade T, Dyer S, Browne W, Frank A. Randomised comparison of chiropractic and hospital outpatient management of low back pain: results from extended follow up. *Br Med J* 1995;**311**:349-351.

Ref ID : 8

90 Meade T, Dyer S, Browne W, Townsend J, Frank A. Low back pain of mechanical origin: randomised comparison of chiropractic and hospital outpatient treatment. *Br Med J* 1990;**300**:1431-1437.

Ref ID : 242

91 MacDonald R, Bell C. An open controlled assessment of osteopathic manipulation in nonspecific low back pain. *Spine* 1990;**15**:364-370.

Ref ID : 147

92 Koes B, Bouter L, van Mameren H, et al. Randomised clinical trial of manipulative therapy and physiotherapy for persistent back and neck complaints: results of one year follow-up. *Br Med J* 1992;**304**:601-605.

Ref ID : 239

93 Blomberg S, Svardsudd K, Mildenerger F. A controlled, multicentre trial of manual therapy in low back pain: initial status, sick leave and pain score during follow-up. *Scand J Prim Care* 1992;**10**:170-178.

Ref ID : 247

94 Shekelle P, Adams A, Chassin M, Hurwitz E, Brook R. Spinal manipulation for low back pain. A review. *Ann Int Med* 1992;**117**:590-598.

Ref ID : 283

95 Hadler N, Curtis P, Gillings D, Stinnett S. A benefit of spinal manipulation as adjunctive therapy for acute low back pain: a stratified controlled trial. *Spine* 1987;**12**:703-706.

Ref ID : 252

96 Lindstrom I, Ohlund C, Eek C, Wallin L, Peterson L, Nachemson A. Mobility, strength, and fitness after a graded activity program for patients with subacute low back pain. A randomised prospective clinical study with a behavioural therapy approach. *Spine* 1992;**17**:641-649.

Ref ID : 253

97 Mitchell R, Carmen G. The functional restoration approach to the treatment of chronic pain in patients with soft tissue and back injuries. *Spine* 1994;**19**:633-642.

Ref ID : 254

98 Mitchell R, Carmen G. Results of a multicenter trial using an intensive active exercise program for the treatment of acute soft tissue and back injuries. *Spine* 1990;**15**:514-521.

Ref ID : 250

99 Pope M, Phillips R, Haugh L, Hsieh C, MacDonald L, Haldeman S. A prospective randomised three-week trial of spinal manipulation, massage and corset in the treatment of subacute low back pain. *Spine* 1994;**19**:2571-2577.

Ref ID : 269



100 Jenkins E, Borenstein D. Exercise for the low back pain patient. *Clin rheumatol* 1994;**8**:191-197.

Ref ID : 241

101 Manniche C, Skall H, Braendholt L, et al. Clinical trial of postoperative dynamic back exercises after first lumbar discectomy. *Spine* 1993;**18**:92-97.

Ref ID : 243

102 Manniche C, Hesselsoe G, Bentzen L, Christensen I, Lundberg E. Clinical trial of intensive muscle training for chronic low back pain. *Lancet* 1988;**2**:1473-1476.

Ref ID : 270

103 Hansen F, Bendix T, Skov P, et al. Intensive, dynamic back-muscle exercises, conventional physiotherapy, or placebo-control treatment of low back pain. A randomised, observer-blind trial. *Spine* 1993;**18**:98-108.

Ref ID : 271

104 Risch S, Norvell N, Pollock M, et al. Lumbar strengthening in chronic low back pain patients. Physiological and psychological benefits. *Spine* 1993;**18**:232-238.

Ref ID : 272

105 Donchin M, Woolf O, Kaplan L, Floman Y. Secondary prevention of low back pain - a clinical trial. *Spine* 1990;**15**:1317-1320.

Ref ID : 273

106 Deyo R, Walsh N, Martin D, Schoenfeld L, Ramamurthy S. A controlled trial of transcutaneous electrical nerve stimulation (TENS) and exercise for chronic low back pain. *New Eng J Med* 1990;**322**:1627-1634.

Ref ID : 275

107 Zylbergold R, Piper M. Lumbar disc disease: comparative analysis of physical therapy treatments. *Arch Phys Med Rehabil* 1981;**62**:176-179.

Ref ID : 510

108 Hume Kendall P, Jenkins J. Exercises for backache: a double blind controlled trial. *Physiotherapy* 1968;**54**:154-157.

Ref ID : 276

109 DiMaggio A, Mooney V. The McKenzie program: exercise effective against back pain. *J of Musculoskeletal Med* 1987;**4**:63-74.

Ref ID : 277

110 Turner J, Clancy S, McQuade K, Cardenas D. Effectiveness of behavioural therapy for chronic low back pain: a component analysis. *J of Consulting and Clin Psychol* 1990;**58**:573-579.

Ref ID : 278

111 Alaranta H, Rytokoski U, Rissanen A, et al. Intensive physical and psychosocial training program for patients with chronic low back pain. A controlled clinical trial. *Spine* 1994;**19**:1339-1349.

Ref ID : 279

112 Elnaggar I, Nordin M, Sheikhzadeh A, Parnianpour M, Kahanovitz N. Effects of spinal flexion and extension exercises on low back pain and spinal mobility in chronic mechanical low back pain patients. *Spine* 1991;**16**:967-972.

Ref ID : 280

113 Frost H, Klaber Moffett J, Moser J, Fairbank J. Evaluation of a fitness programme for patients with chronic low back pain. A randomised controlled trial. *Br Med J* 1995;**310**:151-154.

Ref ID : 1214



114 Klaber Moffett J, Torgerson D, Bell-Syer S, et al. Randomised controlled trial of exercise for low back pain: clinical outcomes, costs, and preferences. *BMJ* 1999;**319**:279-283.

Ref ID : 281

115 Shekelle P. Spine Update - spinal manipulation. *Spine* 1994;**19**:858-861.

Ref ID : 1488

116 Assendelft W, Lankhorst G. Effectiveness of manipulative therapy in low back pain: systematic literature reviews and guidelines are inconclusive. *Nederlands Tijdschrift voor Geneeskunde* 1998;**142**

Ref ID : 290

117 Hsieh C, Phillips R, Adams A, Pope M. Functional outcomes of low back pain: comparison of four treatment groups in a randomised controlled trial. *J Manip and Physiolog Ther* 1992;**15**:4-9.

Ref ID : 1508

118 van der Heijden G, Beurskens A, Koes B, de Vet H, Assendelft W, Bouter L. The efficacy of traction for back and neck pain: a systematic, blinded review of randomized clinical trial methods. *Physical Therapy* 1995;**75**:93-104.

Ref ID : 1287

119 Vickers A, Zollman C. The manipulative therapies: osteopathy and chiropractic. *BMJ* 1999;**319**:1176-1179.

Ref ID : 291

120 Assendelft W, Koes B, Heijden Gv, Bouter L. The efficacy of chiropractic manipulation for back pain: blinded review of relevant randomized clinical trials. *J Manip and Physiolog Ther* 1992;**15**:487-493.

Ref ID : 1428

121 Anonymous. Why bogus therapies seem to work. *Bandolier* 2000;**7**:2-5.

Ref ID : 1439

122 Ernst E, White A. Acupuncture may be associated with serious adverse events. *BMJ* 2000;**320**:513-514.

Ref ID : 37

123 Frank A. Therapies for back pain. *Practitioner* 1990;**234**:113-115.

Ref ID : 1064

124 Nurmikko T, Nash T, Wiles J. Control of chronic pain. *Br Med J* 1998;**317**:1438-1441.

Ref ID : 1099

125 Jamison R, Raymond S, Slawsby E, Nedeljkovic S, Katz N. Opioid therapy for chronic noncancer back pain: a randomized prospective study. *Spine* 1998;**23**:2591-2600.

Ref ID : 292

126 Pheasant H, Bursk A, Goldfarb J, Azen S, Weiss J, Borelli L. Amitriptyline and chronic low back pain - a randomized double-blind crossover study. *Spine* 1983;**8**:552-557.

Ref ID : 293

127 Ward N. Tricyclic antidepressants for chronic low back pain - mechanisms of action and predictors of response. *Spine* 1986;**11**:661-665.

Ref ID : 605

128 McQuay H, Moore R. Antidepressants and chronic pain: effective analgesia in neuropathic pain and other syndromes. *Br Med J* 1997;**314**:763-764.



Ref ID : 731

129 Dickens C, Jayson M, Creed F. Treating depression in chronic low back pain sufferers. *Proc of 11 Congress of Europ Fed of AISP chapters* 1997;257-258.(Abstract)

Ref ID : 295

130 Wiesel S, Feffer H, Borenstein D. Evaluation and outcome of low back pain of unknown etiology. *Spine* 1988;**13**:679-680.

Ref ID : 296

131 Philips H, Grant L. The evolution of chronic back pain problems: a longitudinal study. *Behav Res and Therapy* 1991;**29**:435-441.

Ref ID : 29

132 Frank A. Psychiatric consequences of road traffic accidents - often disabling and unrecognised. *BMJ* 1993;**307**:1283

Ref ID : 399

133 Frank A. Understanding back injuries and back pain. In: Barnes M, Braithwaite B, Ward A, eds. *Medical Aspects of Personal Injury Litigation*. Oxford: Blackwell Science, 1997;164-209.

Ref ID : 1445

134 Gadsby J, Flowerdew M. *Transcutaneous Electrical Nerve Stimulation and acupuncture-like transcutaneous electrical nerve stimulation for chronic low back pain. (Cochrane review)*. Oxford: Update Software: The Cochrane Library, 1999

Ref ID : 554

135 Thomas M, Lundeberg T. Does acupuncture work? *Pain Clinical Update* 1996;**4**:1-4.(Abstract)

Ref ID : 1487

136 Tulder Mv, Cherkin D, Berman D, Lao L, Koes B. Acupuncture for low back pain (Cochrane review). In: AnonymousOxford: Update Software: The Cochrane Library, 1999;1-13.

Ref ID : 684

137 Ernst E, White A. Life-threatening adverse reactions after acupuncture? a systematic review. *Pain* 1997;**71**:123-126.

Ref ID : 1505

138 Nelemans P, Bie Rd, Vet Hd, Strumans F. *Injection therapy for subacute and chronic benign low back pain (Cochrane Review)*. Oxford: Update Software, 1999;1-15

Ref ID : 1443

139 Scheer S, Watanabe T, Radack K. Randomized controlled trials in industrial back pain relating to return to work: part 3 -subacute/chronic pain interventions. *Arch Phys Med Rehabil* 1997;**78**:414-423.

Ref ID : 1270

140 Dechow E, Davies R, Carr A, Thompson P. A randomized double-blind, placebo-controlled trial of sclerosant injections in patients with chronic low back pain. *Rheumatology* 1999;**38**:1255-1259.

Ref ID : 1067

141 Slosar P, White A, Wetzel F. Controversy: The use of selective nerve root blocks: diagnostic, therapeutic, or placebo. *Spine* 1998;**23**:2253-2256.

Ref ID : 314

142 Chapman S, Brena S. Learned helplessness and responses to nerve blocks in chronic low back pain patients. *Pain* 1982;**14**:355-364.

Ref ID : 315

143 Fordyce W. Psychological factors in the failed back. *Int Dis Studies* 1988;**10**:29-31.



Ref ID : 1500

144 Grogan E, Frank A, Keat A. Reassurance in rheumatology clinics. *BMJ* 2000;(in press)

Ref ID : 318

145 Waddell G, Newton M, Henderson I, Somerville D, Main C. A fear avoidance beliefs questionnaire (FABQ) and the role of fear avoidance beliefs in chronic low back pain and disability. *Pain* 1993;**52**:157-168.

Ref ID : 316

146 Ransford A, Cairns D, Mooney V. The pain drawing as an aid to the psychologic evaluation of patients with low back pain. *Spine* 1976;**1**:127-134.

Ref ID : 317

147 Waddell G, McCullough J, Kummel E, Venner R. Nonorganic physical signs in low back pain. *Spine* 1980;**5**:117-125.

Ref ID : 1068

148 Main C, Waddell G. Spine Update: Behavioral responses to examination: a reappraisal of the interpretation of "non-organic signs". *Spine* 1998;**23**:2367-2371.

Ref ID : 321

149 Broome A, Jellico H. *Living with your pain: a self-help guide to managing pain*. London: British Psychological Society, 1987;

Ref ID : 323

150 Polatin P, Kinney R, Gatchel R, Lillo E, Mayer T. Psychiatric illness and chronic low back pain. The mind and the spine - which goes first? *Spine* 1993;**18**:66-71.

Ref ID : 324

151 Gatchel R, Polatin P, Mayer T, Garcy P. Psychopathology and the rehabilitation of patients with chronic low back pain disability. *Arch Phys Med Rehabil* 1994;**75**:666-670.

Ref ID : 191

152 Bergquist-Ullman M. Acute low back pain in industry. A controlled prospective study with special reference to therapy and vocational factors. *Acta Orthop Scand* 1972;**Suppl 170**:1-117.

Ref ID : 232

153 Morrison G, Chase W, Young V, Roberts W. Back pain: treatment and prevention in a Community Hospital. *Arch Phys Med Rehabil* 1988;**69**:605-609.

Ref ID : 333

154 Klaber Moffett J, Chase S, Portek I, Ennis J. A controlled, prospective study to evaluate the effectiveness of a back school in the relief of chronic low back pain. *Spine* 1986;**11**:120-122.

Ref ID : 334

155 Harkapaa K, Mellin G, Jarvikoski A, Hurri H. A controlled study on the outcome of inpatient and outpatient treatment of low back pain. *Scand J Rehab Med* 1990;**22**:181-188.

Ref ID : 335

156 Koes B, van Tulder M, van der Windt D, Bouter L. The efficacy of back schools: a review of randomized clinical trials. *J Clin Epid* 1994;**47**:851-861.

Ref ID : 325

157 Harkapaa K, Jarvikoski A, Mellin G, Hurri H, Luoma J. Health locus of control beliefs and psychological distress as predictors for treatment outcome in low back pain patients: results of a 3 month follow-up of a controlled intervention study. *Pain* 1991;**46**:35-41.



Ref ID : 337

158 Mellin G, Harkapaa K, Hurri H, Jarvikoski A. A controlled study on the outcome of inpatient and outpatient treatment of low back pain. *Scand J Rehab Med* 1990;**22**:189-194.

Ref ID : 338

159 Mayer T, Gatchel R, Mayer H, Kishino N, Keeley J, Mooney V. A prospective two-year study of Functional Restoration in industrial low back injury. *JAMA* 1987;**258**:1763-1767.

Ref ID : 774

160 Pflingsten M, Hildebrandt J, Leibing E, Franz C, Saur P. Effectiveness of a multimodal treatment program for chronic low back pain. *Pain* 1997;**73**:77-85.

Ref ID : 1038

161 Mayer T, McMahon M, Gatchel R, Sparks B, Wright A, Peques P. Socioeconomic outcomes of combined spinal surgery and functional restoration in workers compensation spinal disorders with matched controls. *Spine* 1998;**23**:598-606.

Ref ID : 1504

162 Bendix A, Bendix T, Hastrup C, Busch E. A prospective randomised 5 year follow-up study of functional restoration in chronic low back pain patients. *Eur Spine J* 1998;**7**:111-119.

Ref ID : 94

163 Frank A. Spinal problems in adults. In: Maddison P, Isenberg D, Woo P, Glass D, eds. *Oxford Textbook of Rheumatology*. Oxford: Oxford University Press, 1993;41-54.

Ref ID : 563

164 Roland M, Waddell G, Klaber Moffett J, Burton A, Main C, Cantrell T. *The back book*. Norwich: The Stationary Office, 1996;1-23

Ref ID : 1484

165 International Association for the Study of Pain. *Classification of chronic pain: descriptions of chronic pain syndromes and definitions of pain terms*. Seattle: IASP Press, 1994;1-222

Ref ID : 1151

166 Waddell G. *The back pain revolution*. Edinburgh: Churchill Livingstone, 1998;1-438



Acknowledgments

We are grateful to Dr Martin Underwood for the figures on working days lost through incapacity and to the DHSS for permission to publish them.

AOF is indebted to Dr Laurie Allan (Pain service), Mr Ian Fyfe (Orthopaedic Surgeon), Dr Nan Mitchell (Musculo-skeletal radiologist) and many colleagues in physiotherapy and psychology from whom I have learned much.



Table 1

Common patterns of referral for spinal pain

Occiput to bifrontal areas (common cause of headaches)

Arms

Chest wall - common cause of non-cardiac chest pain

Loins - often mistaken for kidney pain

Groins

Femoral distribution

Sciatic distribution



TABLE 2

POTENTIAL FACTORS PREDISPOSING TO SPINAL PAIN

CONGENITAL

Shape and size of spinal canal

Predisposition to disc degeneration

Hypermobility

Abnormal bony segments:

- reduction in vertebrae
- fusion of all or part of segment
- abnormal segment eg spondylolysis
- pseudojoints

ACQUIRED

Excessive mechanical stress on normal structures

Normal stresses on weak / degenerate structures:

- soft tissues
- discs
- facet joints

Other spinal pathology

Modified from the Oxford Textbook of Rheumatology, Second Edition, 1998 ⁸⁰



Table 3
**ESTIMATED CONSEQUENCES OF THE BACK PAIN EPIDEMIC
IN THE UK**

Estimated total work incapacity (1997/8): 90 million working days (DHSS figures)

Lost output: Estimated loss (1997/8) £9,090 million ¹²

General practitioner consultations: estimated 8.2-12 million annually ^{12,18}

Hospital outpatient consultations: estimated 1.7-2.4 million annually ^{12,18}

Hospital inpatient episodes (1994/5): estimated 70,000 ¹²



TABLE 5

SOME CAUSES OF LOW BACK PAIN

Mechanical/ Degenerative Joints	Muscles and ligaments (eg hypermobility) Discs Nerve compression - root or cauda equina (spinal stenosis)
Inflammatory	Ankylosing Spondylitis Rheumatoid Arthritis (rare)
Infections	Bacterial osteomyelitis Tuberculous osteomyelitis Epidural abscess Brucellosis
Neoplasms	Multiple myeloma Lymphoma Secondary cancer Primary cancer (rare)
Bone disease	Osteoporosis Osteomalacia Paget's disease
Other	Gynaecological Neurological Renal Sickle cell disease Vascular "claudication"



TABLE 4
FEATURES OF NON-SPECIFIC “MECHANICAL” LOW BACK PAIN

Site (1 or more of the following)

Discomfort across lower back

Central pain, usually over L5

Leg pain and / or paraesthesiae within "sciatic" distribution

Unilateral or bilateral buttock or lateral back pain

Character

Episodic or cyclical pain in the middle years of life

Arises from L3-S1

Early morning stiffness/pain eases when up and about

Relationship to posture (often aggravated by sitting or standing still
and eased by walking normally)

NB Pain markedly aggravated by walking raises possibility of vascular
claudication, spinal or lateral canal stenosis.

Modified from the Oxford Textbook of Rheumatology, First Edition, 1993 ¹⁶³



TABLE 6

FEATURES SUGGESTIVE OF UNDERLYING ILLNESS

1. Absence of typical features - Table 4

2. Red flags

1

Constant unremitting pain in atypical or multiple sites

Pain unrelated to movement / posture
(non-mechanical pain)

Systemic / constitutional symptoms

Age less than 20 or over 55 with no previous similar episodes of pain

Widespread neurology

Structural deformity

Modified from the Oxford Textbook of Rheumatology, First Edition, 1993 ¹⁶³



TABLE 7

GUIDELINES FOR RADIOLOGICAL INVESTIGATION

First attack of pain

Age 25-55 pain resolving within 6 weeks - no X-Ray needed
static after 1 month - long lateral *

Age 0-25 pain resolving within 6 weeks - no X-Ray needed
static after 3 weeks - AP and lateral *

Age over 55 pain resolving within 3 weeks - no X-Ray needed
unresolved within 3 weeks - AP and lateral

Atypical back pain - well localised (1 or 2 adjacent levels)
CT scan or MRI if available

Atypical back pain - not well localised - Bone scan.
If positive, CT +/- biopsy under imaging control (or open).

Discogenic disease - well localised (1 or 2 adjacent levels) **
Surgery being considered/nerve compression CT scan or MRI **

Intractable mechanical low back pain - local anaesthetic +/- steroid
into one or more facet joints to localise source of pain ***

Second attack of pain

If similar to previous episode - no X-Rays needed

If different in character or level, as above

Indications for radiology for psycho-social management +

Plain films - for visual display of structural normality or abnormality
when explaining the mechanical nature of back pain

CT or MRI to define presence or absence of discogenic disease for prognostic reasons
e.g. timing return to work, to improve motivation to persist/embark on painful therapy

* If sacroiliitis suspected, dedicated views of sacroiliac joints (SIJ),
or CT scan may confirm if SIJ normal on AP film

** Rarely surgeons will require radiculography preoperatively particularly if non-invasive
investigations have suggested multi-level disease and MRI not available.

*** Discography seldom used now for such purposes

+ Not all radiologists understand the role of X-Rays in overcoming patient's fears

Modified from the Oxford Textbook of Rheumatology, First Edition, 1993 ¹⁶³

TABLE 8



Strategies for preventing back pain

Primary prevention

Elimination or minimisation of risks to health or well-being

Modify risk factors

- smoking
- obesity
- psychological profile
- level of physical fitness

Education - ergonomics in the work place
posture
stress management
lifting strategies
driving certain types of vehicles

Exercise - cardiovascular fitness (for example, aerobics)
spinal mobility / flexibility
muscle strength

Industrial back injury prevention programs

Secondary prevention

Preventing the acute attack from becoming chronic

- Use of treatment algorithms - rapid access to skilled advice
- Reorganisation of claims procedures
- Behavioural modification in back schools and intensive rehabilitation

Tertiary prevention

Rehabilitation - restoring to optimum function and accommodating residual disabilities

CBT and relaxation

Multi-disciplinary pain management programs (probably)

Behavioural modification in back schools and intensive rehabilitation



Table 9

Ergonomic approaches to minimising back injuries in the Railroad Industry ⁶⁵

Storing tools and materials off the ground - between knee and shoulder height

Heaviest items stored at knuckle height

Devising winches to lift and handle heavy equipment

Worktables, dollies and carts to handle heavy car parts and tools

NB Management took a participatory approach and encouraged workers to
configure their own tools and material handling devices



Table 12

**“Yellow flags” for risk of chronicity in
unresolved low back pain**

Presence of a belief that back pain is harmful or potentially severely disabling

Fear-avoidance behaviour (avoiding a movement or activity due to misplaced anticipation of pain) and reduced activity levels

Tendency to low mood and withdrawal from social interaction

An expectation that passive treatments rather than active participation will help

Suggested questions (phrased in treatment providers own words)

Have you had time off work in the past for low back pain?

What do you understand is the cause of your back pain?

How is your employer responding to your back pain? Your co-workers? your family?

What are you doing to cope with back pain?

Do you think you will return to work? when?

From Kendall NAS et al ⁶³



Table 10

The initial consultation for acute back pain

History

Listen to site and character of the pain - see Tables 5-6

Examination

Ask patient to show you the site - the characteristic hand movements
- particularly helpful if there are language barriers

Check for obvious spinal deformity, spasm or gross loss of movement

Check straight leg raise (SLR), reflexes and sensation if leg pain symptoms present

Management

Explanation and information - e.g The Back Book ¹⁶⁴

Minimise rest and encourage resumption of normal activity - including work - patient must take **active** part in recovery

Advise regular and not “as needed” analgesia - may enable activities to be maintained

Examine the reasons for the patient’s specific fears - reassure if appropriate:-

 Serious illness very unlikely

 X-Rays and blood tests unlikely to be needed

 Most severe back pain improves within days or a few weeks; milder symptoms may persist a few months

Recurrences are common

Offer early review if the pain is uncontrolled.

NB Ensure the patient feels comfortable with your actions and plan.



Table 11

Oral medication used in back pain

Standard analgesics Paracetamol, dextropropoxyphene, codeine & dihydrocodeine or in compound preparation formulation
(coproxamol - paracetamol and dextropropoxyphene)
(cocodamol - paracetamol and codeine)
(codydramol - paracetamol and dihydrocodeine)
long-acting compound preparations
e.g. Ibuprofen and codeine

Non-steroidal anti-inflammatory drugs (NSAIDs)

Mostly Cyclo-oxygenase 1 (COX-1) inhibition

Propionic acid derivatives - e.g. Ibuprofen, Naproxen

Others e.g. Soluble aspirin, Diclofenac

Some COX-2 inhibition

Nabumetone, Lodeine, Meloxicam

COX 2 inhibitors - Rofecoxib

Strong analgesics

Moderately potent:- Meptazinol, Nefopam, Pentazocine, Pethidine

Potent:- Buprenorphine, Tramadol

Very potent:- Diamorphine or Morphine
(? not appropriate for use in primary care)

Other Muscle relaxants or mild tranquillisers
e.g Diazepam for 5-7 days (acute back pain only)
Tricyclic antidepressant compounds (chronic pain only)



TABLE 13

PHYSICAL BARRIERS TO SUCCESSFUL OUTCOME

Repetitive poor posture at work / housework

leisure / or inappropriate activities

Repetitive lifting , particularly if involving twisting

Inappropriate bed or mattress

Sleeping in an inappropriate position

Getting in / out of bed incorrectly

Inappropriate exercise regime

Sitting > 1/2 hour (particularly when driving)

Recurrent bending (eg when brushing teeth)
(particularly if twisting)

Stopping exercise regime when "better"

Defaulting medication or corset

Lack of physical fitness



TABLE 14

PSYCHOLOGICAL BARRIERS TO SUCCESSFUL OUTCOME

1. Fear - of medication - drug dependency, side effects,
effectiveness will mask pain -
creating further damage
 - of physical activity - will increase pain particularly during recovery
 - of the cause - eg cancer, multiple sclerosis
 - of the consequences - loss of job (status and poverty)
 - disability (eg in a wheelchair)
 - of clicking in the back - frightened of damage to spine
2. Effects on the family - particularly on the partner - potential to develop learned helplessness or pain behaviour
3. Psychiatric consequences- Depression
Substance abuse eg alcohol
Post-traumatic stress disorder
- 4 Previous psychiatric history
5. Other - Job dissatisfaction, litigation / compensation



Table 15

Facilitating return to employment

The employers role

Take an interest in the patient - contact them when they are off sick - offer support

Offer assistance of the Occupational Health Service if available

Offer ergonomic advice at the work site

Allow a phased return to work

facillitate travel at non-peak times

minimise stress

accept that the patient may need less demanding work both physically and emotionally short term

The health professionals

Advise partner/spouse on safe activities, specify any risk factors and discourage excessive protection

Discuss the 'side-effects' of being 'off sick' and encourage a positive view of work

Liaise with employers about above recommendations

Encourage the patient to accept that returning to work is unlikely to aggravate the cause of the pain

Offer analgesia to facilitate getting fitter and to lessen pain at work

If physically demanding work, consider a surgical corset or a weight-lifters belt

Offer continued support for 1-2 weeks after return to work to ensure work patterns safely re-established if no occupational health service

From Frank AO ⁵⁷



Abbreviations

CBT	Cognitive behavioural therapy
COX	Cyclo-oxygenase
GP(s)	General practitioner(s)
MRI	Magnetic Resonance Imaging
NHS	National Health Service
NSAIDs	Non steroidal anti-inflammatory drugs
NSLBP	Non specific low back pain
TCAs	Tricyclic anti-depressant compounds
TENS	Transcutaneous electrical nerve stimulation
UK	United Kingdom
USA	United States of America



Glossary of terms

- Acute pain:** from 0-7 days duration (pain-free at onset)
- Acute on chronic pain:** significant exacerbation of pre-existing pain
- Allodynia:** pain stimulated by minor stimulation of the skin.
- Back school:** outpatient programme of treatment emphasising educational management
- Chronic pain:** duration of pain over 3 months
- Chronic pain syndrome:** psychological and social consequences of chronic pain influencing behaviour
- Hyperpathia:** A painful syndrome characterised by an abnormally painful reaction to a stimulus, especially if repetitive, as well as an increased threshold ¹⁶⁵.
- Intensive rehabilitation:** Multiprofessional combination of skills and therapeutic modalities to minimise pain and disability and improve function ⁵⁸
- Intractable pain:** failed conservative treatment of chronic pain ⁵⁸.
- Lumbar segment:** two adjacent vertebrae and their intervening soft tissues ⁵⁹. (the five lumbar vertebrae are linked by intervertebral discs anteriorly, two synovial facet joints posteriorly and supporting structures including ligaments and muscles).
- Nociceptive pain:** that arising from mechanical or other stimuli that could cause tissue damage These stimuli act on peripheral pain receptors to give activity in nerve fibres ¹⁶⁶.
- Pain:** An unpleasant sensory and emotional experience associated with actual or potential tissue damage, or described in terms of such damage ¹⁶⁵.
- Subacute pain:** from 7 days to 3 months duration {1}

