


Review: Strategies for enteric methane mitigation in cattle fed tropical forages

J. C. Ku-Vera¹, O. A. Castelán-Ortega², F. A. Galindo-Maldonado³, J. Arango⁴, N. Chirinda⁴, R. Jiménez-Ocampo^{1,5†} , S. S. Valencia-Salazar⁶, E. J. Flores-Santiago¹, M. D. Montoya-Flores⁷, I. C. Molina-Botero¹, A. T. Piñeiro-Vázquez⁸, J. I. Arceo-Castillo¹, C. F. Aguilar-Pérez¹, L. Ramírez-Avilés¹ and F. J. Solorio-Sánchez¹

¹Laboratory of Climate Change and Livestock Production, Department of Animal Nutrition, Faculty of Veterinary Medicine and Animal Science, University of Yucatan (UADY), Carretera Merida-Xmatkuil Km 15.5, C.P. 97100 Merida, Yucatan, Mexico; ²Laboratory of Livestock, Environment and Renewable Energies, Faculty of Veterinary Medicine and Animal Science, University of State of Mexico (UAEM), Instituto Literario 100, C.P. 50000 Toluca, Estado de Mexico, Mexico; ³Department of Ethology, Wildlife and Laboratory Animals, Faculty of Veterinary Medicine and Animal Science, National University of Mexico (UNAM), Ciudad Universitaria, C.P. 04510 Mexico City, Mexico; ⁴International Center for Tropical Agriculture (CIAT), Km 17, Recta Cali-Palmira, Palmira, C.P. 763537 Valle del Cauca, Colombia; ⁵National Institute of Research in Forestry, Agriculture and Livestock-INIFAP, Experimental Field Valle del Guadiana, Km 4.5 Carretera Durango – El Mezquital, C.P. 34170 Durango, Durango, Mexico; ⁶College of the Southern Border (ECOSUR), Livestock and Environment, Carretera Panamericana – Periferico Sur, C.P. 29290 San Cristobal de las Casas, Chiapas, Mexico; ⁷National Center for Disciplinary Research in Physiology and Animal Breeding-INIFAP, Km 1 Carretera a Colon, C.P. 76280 Ajuchitlan, Queretaro, Mexico; ⁸National Technologic of Mexico/I.T. Conkal, Avenida Tecnológico s/n, C.P. 97345 Conkal, Yucatan, Mexico

(Received 11 December 2019; Accepted 20 July 2020)

Methane (CH₄) is a greenhouse gas (GHG) produced and released by eructation to the atmosphere in large volumes by ruminants. Enteric CH₄ contributes significantly to global GHG emissions arising from animal agriculture. It has been contended that tropical grasses produce higher emissions of enteric CH₄ than temperate grasses, when they are fed to ruminants. A number of experiments have been performed in respiration chambers and head-boxes to assess the enteric CH₄ mitigation potential of foliage and pods of tropical plants, as well as nitrates (NO₃⁻) and vegetable oils in practical rations for cattle. On the basis of individual determinations of enteric CH₄ carried out in respiration chambers, the average CH₄ yield for cattle fed low-quality tropical grasses (>70% ration DM) was 17.0 g CH₄/kg DM intake. Results showed that when foliage and ground pods of tropical trees and shrubs were incorporated in cattle rations, methane yield (g CH₄/kg DM intake) was decreased by 10% to 25%, depending on plant species and level of intake of the ration. Incorporation of nitrates and vegetable oils in the ration decreased enteric CH₄ yield by ~6% to ~20%, respectively. Condensed tannins, saponins and starch contained in foliages, pods and seeds of tropical trees and shrubs, as well as nitrates and vegetable oils, can be fed to cattle to mitigate enteric CH₄ emissions under smallholder conditions. Strategies for enteric CH₄ mitigation in cattle grazing low-quality tropical forages can effectively increase productivity while decreasing enteric CH₄ emissions in absolute terms and per unit of product (e.g. meat, milk), thus reducing the contribution of ruminants to GHG emissions and therefore to climate change.

Keywords: tropical grasses, legumes, cattle, rumen fermentation, greenhouse gases

Implications

Feeding strategies designed to mitigate enteric CH₄ emissions in cattle have been tested by incorporating a range of tropical plants (foliages, pods and seeds), nitrates and vegetable oils to rations based on low-quality tropical grasses. These strategies have proved to be effective in decreasing enteric methane emissions by cattle. This could lead to an improvement in profitability of production by reducing the

incorporation of expensive feedstuffs in rations and thus to sustainable ruminant production systems for smallholders in the tropical regions of the world.

Introduction

Animal agriculture, in developing countries, will face tremendous challenges with respect to mitigation and adaptation to climate change in the following years as an increasing demand for meat and milk by the burgeoning population

† E-mail: rafax77@hotmail.com

is expected. This increase in demand is projected to result in an increase in livestock numbers in the near future. Considerable research efforts have been invested in accurately predicting enteric methane emissions from cattle (Escobar-Bahamondes *et al.*, 2016; Eugène *et al.*, 2019; van Lingen *et al.*, 2019) as well as in implementing appropriate enteric CH₄ mitigation strategies (Beauchemin *et al.*, 2020) in ruminant production systems.

During the dry season, most tropical grasses have a high NDF and low CP contents, resulting in limited fermentation of DM, long retention time of digesta and low absorption of volatile fatty acids (VFA) from the rumen, leading to modest daily weight gains in growing cattle and considerable emissions of enteric methane. More than 200 000 plant secondary compounds such as condensed tannins, saponins, flavonoids, organosulfur compounds and essential oils (EOs) have been identified as having potential to mitigate CH₄ emissions in grazing ruminants (Jafari *et al.*, 2019; Molina-Botero *et al.*, 2019a). Secondary compounds in plants should induce a reduction in CH₄ production (Figure 1) without negatively affecting rumen fermentation or being toxic to the animal, and should ideally improve productive performance, increase economic gains, be easy to handle and be of low cost to farmers. The effectiveness of these plant secondary compounds depends on their type, concentration and ingested amount. In the medium term, it is important to identify cost-effective secondary compounds which could improve rumen fermentation, VFA production and mitigate CH₄ emissions with a lasting effect. One effective methane mitigation strategy is the inclusion of low levels of oils in rations which induce biohydrogenation of unsaturated fatty acids, increase energy density and inhibit protozoa (Knapp *et al.*, 2014). A number of comprehensive reviews on methane mitigation in ruminants have been published recently (Berndt and Tomkins,

2013; Knapp *et al.*, 2014; Beauchemin *et al.*, 2020), but relatively few have emphasized the current situation in tropical cattle production systems in developing countries.

Ruminant production in developing countries (mostly in the tropics) faces enormous challenges to both increase productivity and decrease greenhouse gas (GHG) emissions (Ayantunde *et al.*, 2005). The achievement of CH₄ mitigation at the farm level can bring environmental benefits and increase profits for small farmers in tropical regions. Small-scale cattle farmers in the tropical regions of Latin America and the Caribbean, Asia and Africa confront formidable obstacles in achieving sustainable intensification of animal production (Ndung'u *et al.*, 2019). Poverty, poor literacy, scarcity of financial resources (credits), limited access to markets, lack of technical support and paradoxically, the ominous impacts of climate change (droughts, floods and fires) are just but a few of the present constraints.

The rationale behind this review is to describe and critically discuss the advantages of one of the simplest and most direct options for enteric CH₄ mitigation available to small-scale cattle farmers in the tropics, namely the incorporation of foliages, seeds and pods of widely distributed tropical plants, by-products and additives in practical rations.

Enteric methane production by cattle fed tropical grasses

There are many factors influencing CH₄ yield such as type of grass (C3 v. C4), chemical composition, forage : concentrate ratio, rate of passage of digesta through the rumen, physiological stage, breed, sex and it is assumed that all those are integrated within the single expression of DM intake (Archimède *et al.*, 2011 and 2018). Dry matter intake is




Figure 1 Metabolic and digestive effects of condensed tannins, essential oils and saponins on methane synthesis and animal production. (-): decrease; (+): increase. Modified from Valencia Salazar (2017).

the result of both grass and cattle interactions which affect rumen fermentation and, as such, represents a convenient basis for estimating enteric methane production. More work is required by research groups in tropical regions on the establishment of robust data sets from which accurate CH₄ yields could be derived for methane inventory purposes (Patra, 2017; Castelán Ortega *et al.*, 2019; Goopy *et al.*, 2020) and for evaluation of effective enteric CH₄ mitigation strategies.

A series of experiments involving methane emissions were carried out in Mexico with cattle housed in open-circuit respiration chambers and fed basal rations of low-quality tropical grass (>70% of ration DM) and a supplement (concentrate). Preliminary results showed a methane yield of 17.4 g CH₄/kg DMI in Nellore (*Bos indicus*) bulls that were fed chopped *Pennisetum purpureum* grass (Canul-Solís *et al.*, 2017), a value that compares reasonably well with the predicted methane yield of 18.0 g CH₄/kg DM intake ($n = 66$; $R^2 = 0.73$) in cattle fed low-quality tropical grass (~70% ration DM) reported by Ku-Vera *et al.* (2018) and with the methane yield reported by Noguera and Posada-Ochoa (2017) who reported 16.1 g CH₄/kg DMI in lactating cows fed *Pennisetum clandestinum* in Colombia. The values reported by the cited studies (Noguera and Posada-Ochoa, 2017; Canul-Solís *et al.*, 2017; Ku-Vera *et al.*, 2018) represent an approximation for cattle fed tropical grasses, and they are comparable to some extent to the data from Charmley *et al.* (2016) who reported a methane yield of 19.3 g CH₄/kg DMI ($n = 133$) for Zebu (Brahman) cattle fed tropical grasses (>70% of DM) as a basal ration in Australia. Although it has been claimed that tropical (C4) grasses contribute to enteric methane emissions to a greater extent than temperate (C3) grasses (Archimède *et al.*, 2018), the evidence hereby presented does not support that contention (Benaouda, 2018; Goopy *et al.*, 2018; Table 1). The overall enteric methane yield from experiments carried out in Mexico in

respiration chambers is 17.0 g CH₄/kg DMI ($n = 125$) for cattle consuming low-quality tropical grasses. This relatively low yield could be partially explained by the low rate and extent of fermentation of DM (~40%) in the rumen of cattle fed tropical grass during the dry season. Strategic grazing management (with a canopy with light interception of 95%) results in reduction in CH₄ emission intensity and greater milk yield in grazing cows, as well as 18% less CH₄/kg DM intake (Congio *et al.*, 2018). The so-called *rotatinoous* grazing management (Savian *et al.*, 2018) also results in a decrease in CH₄ emission intensity by carefully controlling grazing behavior of ruminants. Both approaches support the contention that it is through grazing management that CH₄ emission intensity in ruminants can be decreased without any other further intervention. Nonetheless, Berça *et al.* (2019) found no reductions in enteric methane emissions of crossbred dairy heifers grazing *Urochloa brizantha* cv. Marandu either fertilized or with the inclusion of the legume *Arachis pintoi* in Brazil. There are several possible strategies to mitigate methane emissions in ruminants that are fed low-quality tropical grasses. The selection criteria for a particular strategy must be based on sound economic grounds, as smallholders are limited economically in their capacity to buy expensive methane mitigating additives (e.g. monensin, 3-nitrooxypropanol). *Leucaena leucocephala*, a native legume shrub from Mexico and Central America, ranks first among the options available to develop sustainable beef cattle farming systems in tropical regions (Shelton and Dalzell, 2007; Harrison *et al.*, 2015; Ramírez-Avilés *et al.*, 2019). In Colombia, Molina *et al.* (2016) fed Lucerna heifers with a *Cynodon plectostachyus* (74% DM) and *Leucaena leucocephala* (26% DM) ration while cattle were housed in polytunnels for methane measurements and recorded a lower intensity of methane emissions compared to heifers fed only the tropical grass. Also in Colombia, Sierra-Montoya *et al.* (2017) found a reasonably good milk yield (~10 to 12 l cow/day) in recently

Table 1 Enteric methane emissions of cattle in the tropics of Asia, Africa, Oceania and Latin America as measured in respiration chambers

Cattle (breed, sex)	LW (kg)	Ration	DMI (kg/day)	CH ₄ (g/day)	CH ₄ (g/kg DMI)	Y _m (% GEI)	Country	References
Holstein cows	541	<i>Pennisetum clandestinum</i>	18.5	287	16.1	3.8	Colombia	Noguera and Posada-Ochoa (2017)
Holstein × Gyr, cows	–	TMR (maize silage, Tifton hay, concentrate)	20.2	382.7	18.9		Brazil	Machado <i>et al.</i> (2016)
Crossbred heifers	327	<i>Megathyrus maximus</i> hay + supplement	8.1	129.4	16.0	5.2	Mexico	Zavala-Escalante, L. unpublished
Boran steers	162.3	<i>Chloris gayana</i> hay	4.6	133.3	29.0	9.14	Kenya	Goopy <i>et al.</i> (2020)
Brahman steers	227	<i>H. contortus</i> C. <i>ciliaris</i>	4.0	99.3	19.6	6.3	Australia	Kennedy and Charmley (2012)
Brahman steers	309	<i>H. contortus</i> hay <i>ad libitum</i>	4.9	94.7	19.3	5.7	Australia	Charmley <i>et al.</i> (2016)
Brahman steers	266.4	<i>Chloris gayana</i> hay <i>ad libitum</i>	4.1	80 to 5	19.5	5.8	Australia	Perry <i>et al.</i> (2017)
Brahman steers	342.7	<i>Digitaria eriantha</i>	3.5	119.3	34.1	11.4	Thailand	Chaokaur <i>et al.</i> (2015)
Tropical cattle	277	Crop residues and by-products	4.6	123	26.7	8.2	Thailand	Kaewpila and Sommart (2016)

LW = live weight; DMI = DM intake; CH₄ = methane; TMR = total mixed ration; GEI = gross energy intake; Y_m = CH₄ loss expressed as percentage of GEI.

calved or early lactating Gyr × Holstein cows grazing a silvopastoral system consisting of *Cynodon plectostachyus* and *Leucaena leucocephala*, obtaining a good balance of metabolizable protein to metabolizable energy.

Enteric methane mitigation strategies in cattle under small-scale farming conditions

Before adoption or implementation at farm level, an enteric methane mitigation strategy must induce a significant extent of mitigation as demonstrated in various experiments, published equations applicable to various types of ruminants, feasibility of adoption by farmers, cost-effectiveness analysis and repeatability (Eugène *et al.*, 2019), as well as persistency. This review addresses three different options for mitigation: secondary compounds contained in plant species available in tropical regions that can be used in silvopastoral systems, high-energy compounds such as oils and finally a chemical compound exemplified by the feeding of additives such as nitrates. Nitrates require attention, particularly under the conditions of tropical ruminant production, since they supply N, an element which is usually deficient in C4 grasses during the long dry season.

Foliage and pods of tropical plants as a methane mitigation strategy

Foliages, seeds and pods of tropical trees and shrubs have been considered as important components of ruminant rations long ago (Topps, 1992; Shelton and Dalzell, 2007). Secondary compounds contained in such plant species exert different effects on the rumen microbial population (Patra and Saxena, 2009). Piñeiro-Vázquez *et al.* (2018) reported that foliage of the tropical legume *Leucaena leucocephala* induced a reduction in methane yield when it was included in ration DM in high proportions (40% to 80%) in crossbred heifers (*Bos indicus* × *Bos taurus*) housed in respiration chambers that were fed a basal ration of low-quality tropical grass (Table 2). However, the increase in CP intake from *L. leucocephala* induced higher N excretion (as urea) in the urine, probably leading to higher nitrous oxide emissions (Montoya-Flores *et al.*, 2020). This agrees with data reported by Harrison *et al.* (2015) who also found a reduction in enteric methane emissions, although nitrous oxide emissions increased in cattle fed *Leucaena leucocephala* in Australia. In a set of experiments involving the incorporation of foliages such as *Leucaena leucocephala* or *Gliricidia sepium* (Molina-Botero *et al.*, 2019a; Montoya-Flores *et al.*, 2020) or ground pods of *Enterolobium cyclocarpum* or *Samanea saman* (Lazos-Balbuena, 2015; Valencia Salazar *et al.*, 2018), a reduction in enteric methane yield of different magnitudes was recorded in cattle housed in respiration chambers (Table 2).

Pods of tropical legumes such as *Enterolobium cyclocarpum* and *Acacia pennatula* contain either condensed tannins or saponins, and when fed ground at 45% of ration DM, weight gains of up to 240 g/lamb per day were obtained

in small-scale sheep farms (Briceño-Poot *et al.*, 2012). Pods from those legumes can be hand collected by smallholders, stored in bags, ground and directly fed to cattle and sheep up to 50% of ration DM (Briceño-Poot *et al.*, 2012). Pods are palatable and they are eagerly consumed by cattle and sheep after falling from the trees during the dry season. Rumen DM degradability of ground pods is high: 87% for *Enterolobium cyclocarpum* (Piñeiro-Vázquez *et al.*, 2013) and 80% for *Samanea saman* (Valencia Salazar *et al.*, 2018); they provide (through anaerobic fermentation) the necessary ATP for the maintenance and growth of bacteria in the rumen. The high level of rumen DM degradation of tropical pods suggests that they contain a high concentration of readily available carbohydrates (i.e. pectins, starch), which may change the pattern of rumen fermentation (Valencia Salazar *et al.*, 2018) and decrease methane production per kg of DM consumed. Table 2 shows the magnitude of reduction in enteric methane yield when cattle were fed foliage or pods from tropical trees. Both options are simple and straightforward strategies for reducing methane emissions in farms under smallholder conditions in the tropics.

The proportion of gross energy intake (GEI) lost as methane gas: Y_m (Table 2) ranged from 5.0% to 9.5% throughout the experiments, and it generally decreased as the levels of mitigating compounds (i.e. foliage, pods) were incorporated at increasing levels in the rations. Those authors found that Y_m fluctuated between 4.8% and 13.7% and that IPCC (2006) procedures underestimated Y_m by ~26.1%. Therefore, the derivation of precise Y_m values remains an important challenge for estimating CH₄ emissions from *Bos indicus* type of cattle fed low-quality forages in tropical developing countries. In Thailand, Chaokaur *et al.* (2015) working with Brahman cattle fed Pangola (*Digitaria eriantha*) grass as basal ration found that Y_m ranged between 7.3% and 11.5% (above the IPCC (2006) default value of 6.5%) and that enteric methane emissions were linearly increased from the maintenance level of feeding up to *ad libitum* intake.

There are several possible explanations for the reduction of enteric methane emissions as a result of feeding foliage and pods of tropical plant species to cattle and sheep (see Figure 1). Condensed tannins contained in such plants may reduce CH₄ in two different ways, first, by directly inhibiting methanogenic archaea, and second by decreasing H₂ availability in the rumen and the apparent digestibility of DM and protozoal population (Patra *et al.*, 2017; Piñeiro-Vázquez *et al.*, 2018), Aboagye (2018 and 2019) reported a reduction in fiber digestibility, rather than a decrease in the amount of structural carbohydrates. Nonetheless, Canul-Solis *et al.* (2014) found no effect of saponins (6 g/day) of *Yucca schidigera* on methane emissions of hair sheep fed low-quality tropical grass; this could be associated with the lack of effect on methanogenic archaea or the relatively low levels of incorporation of the mitigating commercial compound.

Saponins in pods such as those in *Samanea saman* may form complexes with sterols in the protozoal membrane

Table 2 Methane production, methane yield and methane conversion factor (Y_m) in cattle fed tropical plants for enteric methane mitigation as measured in respiration chambers

Source	Species	Level of incorporation (% DM)	DMI (kg/day)	CH ₄ (g/day)	CH ₄ (g/kg DMI)	Y _m (% GEI)	Design and statistical analysis	Authors	
<i>Enterolobium cyclocarpum</i> (ground dry pods)	Cattle	0	8.6 ^a	108.20	12.86	–	4 × 4 crossover LSD	Lazos-Balbuena (2015) (Thesis)	
		12	9.7 ^{ab}	97.80	10.02	–			
		24	10.0 ^b	100.02	10.00	–	Tukey's test		
		36	10.2 ^b	101.68	9.89	–			
		SE		0.26	6.90	1.01	–		
P-value		0.018	ns	ns	–				
<i>Leucaena leucocephala</i> (fresh forage)	Cattle	0	7.03	104.51	15.30	5.42	5 × 5 crossover LSD	Piñeiro-Vázquez et al. (2018)	
		20	7.15	77.03	11.18	4.00			
		40	7.07	66.52	9.21	3.44	Surface response analysis		
		60	7.00	57.01	7.99	2.95			
		80	7.00	40.72	5.86	2.11			
SE		0.60	11.26	1.59	0.58				
Linear contrast		ns	**	**	**				
<i>Samanea saman</i> (ground dry pods)	Cattle	0	6.26	87.14	13.73	4.70	4 × 4 crossover LSD	Valencia-Salazar et al. (2018)	
		10	6.44	64.64	10.06	3.38			
		20	6.16	51.94	9.12	2.82	Surface response analysis		
		30	6.49	42.76	5.90	2.15			
		SE		0.28	6.77	0.78			0.31
Linear contrast		ns	**	**	**				
<i>Leucaena leucocephala</i> (dry leaves)	Cattle	0	8.36	174.2	20.8	6.5	4 × 4 crossover LSD	Montoya-Flores et al. (2020)	
		12	8.32	162.9	19.7	6.07			
		24	8.63	154.8	18.0	5.54	Surface response analysis		
		36	8.54	140.0	16.5	5.05			
		SE		0.13	3.03	0.41			0.13
Linear contrast		0.09	**	**	**				
<i>E. cyclocarpum</i> (ground dry pods) 50% + <i>Gliricidia sepium</i> (dry foliage) 50%	Cattle	0	5.18	144.8	28.50	9.57	4 × 4 crossover LSD	Molina-Botero et al. (2019a)	
		15	5.23	140.1	26.80	9.09			
		30	5.57	141.3	25.56	8.57	Surface response analysis		
		45	5.57	143.3	25.89	8.80			
		SE		0.33	4.88	1.50			0.51
Linear contrast		0.989	0.788	0.050	0.051				
Quadratic contrast		0.258	0.218	0.299	0.281				
Cubic contrast		0.500	0.655	0.689	0.683				
<i>E. cyclocarpum</i> (ground dry pods) + <i>G. sepium</i> (dry foliage)	Cattle	E + G : 0	6.11	145.99	23.95 ^a	8.15 ^a	Repeated measures design	Molina-Botero et al. (2019b)	
		E + G : 15	6.62	147.78	23.32 ^b	7.59 ^b			
		SE		0.20	3.02	0.33	0.11		
		P-value diet		0.598	0.631	0.016	0.016		Tukey's test
		P-value time		0.018	0.262	0.885	0.886		
P-value (diet × time)		0.383	0.440	0.682	0.682				

CH₄ = methane; DMI = DM intake; Y_m = CH₄ loss expressed as percentage of gross energy intake (GEI); LSD = Latin square design; ns = non-significant ($P > 0.05$).

^{a,b}Means in the same column with different superscript are statistically different according to Tukey's test ($P > 0.05$) (for experiments that used Tukey's test).

** $P < 0.01$.

surface and induce lysis of such membrane (Anantasook et al., 2014), thus reducing the population of methanogenic archaea which are usually associated with protozoa. Incorporation of foliage and pods of plants containing condensed tannins and steroidal saponins in ruminant rations can increase the molar proportion of propionic acid in rumen liquor and decrease the availability of metabolic H which is the specific substrate for carbon dioxide reduction to methane (Valencia-Salazar et al., 2018). Feeding of starch-containing ingredients in the ration such as in the pods of the legume tree *Samanea saman* (rain tree) may also drive a

reduction in methane emissions by changing the pattern of rumen fermentation, increasing the molar proportion of propionic acid and lowering that of acetic acid in the rumen (Valencia-Salazar et al., 2018). Anantasook et al. (2014) similarly reported an increase in propionic acid concentration in the rumen and a reduction in methane production, when they fed pods of *Samanea saman* to dairy steers fed urea-treated rice straw as basal ration. Ground pods of *Samanea saman* have been successfully used for feeding dairy cows (Anantasook et al., 2014) in Thailand. Changes in the molar proportions of VFA in the rumen (i.e. increasing

propionic acid relative to acetic acid) may improve the efficiency of metabolizable energy utilization by decreasing heat increment and increasing energy retention as protein and fat synthesis (weight gain) in the body (NASEM, 2016) or in milk as fat and lactose. Thus, this kind of mitigation strategy may be more acceptable to smallholders, since they would see their effort compensated in both productivity and probably in financial revenues.

An increase in animal productivity has been observed under production conditions in cattle and sheep in small farms where both leucaena and pod legume incorporation in practical rations were implemented. Growing crossbred bulls showed a daily weight gain of 770 g, while grazing in a silvopastoral system with *Leucaena leucocephala* (Mayo-Eusebio, 2013) and confined hair sheep gained 240 g/day when pods of *Enterolobium cyclocarpum* were incorporated at 50% (DM basis) of their ration (Briceño-Poot et al., 2012) under smallholder farming conditions. These results agree with those of Harrison et al. (2015) with grazing cows on leucaena pastures in Australia and Moscoso et al. (1995) with confined sheep fed ground pods of *Enterolobium cyclocarpum* (36% of ration DM) in Honduras (Central America). We do not fully understand the lack of changes in terms of the rumen microbial population when using tropical feedstuffs containing secondary compounds. So far, the work carried out on DNA (qPCR) techniques for counting the rumen microbial population has been inconclusive with regard to changes in the composition of the microbial population. No changes have been observed so far in the structure of the rumen population when cattle were fed either *Leucaena leucocephala* or *Enterolobium cyclocarpum*. Nonetheless, the length of time of feeding apparently does not lead to an adaptation of the rumen microbial population to secondary compounds (such as the saponins contained in pods of *Enterolobium cyclocarpum*) (Molina-Botero et al., 2019b). Disposal of H₂ in the rumen through methanogenesis is aimed at maintaining the partial rumen pressure of H₂ low so that fermentation can proceed efficiently. So far, there is still not a plausible explanation for the fate of the excess metabolic H as a result of methane mitigation in ruminants fed tropical C4 grasses (4 mol H₂ remains unused per mol CH₄ decreased; Hegarty and Gerdes, 1999). No alternative H₂ sink has been proposed as yet, although it is recognized that there may be an excess H₂ in the rumen when methane mitigating compounds are fed (up to 35.9 g/day in heifers; Romero-Perez et al. (2015)). Some of the excess H₂ in the treatments involving incorporation of mitigating compounds may have been incorporated into rumen microbes (Kennedy and Charmley, 2012). Nevertheless, more work has to be done on this subject. Rumen pH of cattle fed foliages and pods of tropical plants where substantial reductions in methane emissions have been recorded showed no apparent changes 4 h after feeding (Valencia-Salazar et al., 2018; Montoya-Flores et al., 2020). Since rumination is normally carried out in such conditions, rumen pH generally remains within the physiological range (6 to 7), as well as the osmotic pressure (i.e. 250 to 300 mosmol/kg).

Essential oils are secondary compounds (alcohol, ester or aldehyde derivatives of phenylpropanoids and terpenoids) present in some tropical plants. They are volatile, aromatic substances. Essential oils display antimicrobial actions related to processes in the bacterial cell membrane involving electron transport, ion gradients, phosphorylation and protein translocation (Benchaar et al., 2008). Essential oils exert a high affinity for lipids in cell membranes of bacteria which results from their hydrophobic nature and their antibacterial actions are related to their lipophilic character; thus, they can disrupt the cytoplasmic membrane either directly or by damaging proteins in the membrane, resulting in increased permeability of the membrane and then leakage of cytoplasmic constituents thus affecting the proton motive force (Hart et al., 2008). Scientific evidence on the potential of EOs to reduce enteric methane in ruminants *in vivo* in tropical regions is still scarce (Wang et al., 2018). Effective doses, mode of action, voluntary intake and the cost-benefit ratio must be assessed. Most of the studies involving EOs have been conducted under *in vitro* conditions (Ye et al., 2018) and those results must be validated *in vivo* at the farm level under smallholder conditions. Recent work, carried out at the University of Yucatan in Mexico, with crossbred heifers held in respiration chambers (Jiménez-Ocampo, R. *unpublished* results) has shown that EO from *Citrus sinensis* (0.5% DM) reduced methane yield in heifers fed low-quality tropical grasses by up to ~18% (compared to a control ration without oil), which agrees with experiments carried out in Hu sheep supplemented with EO from grapefruit peels (Wu et al., 2018). Figure 1 shows the mechanisms of action of some secondary compounds for mitigation of enteric methane in ruminants. The mechanisms are rather different: condensed tannins may both affect feed digestibility and inhibit archaea, saponins may disrupt protozoa membranes and reduce the population of methanogenic archaea, while EOs may affect protozoa and methanogens, depending on the type and concentration. In the tropics, there is a wide range of plants which may induce such mechanisms in the rumen of cattle under practical feeding conditions.

Nitrate supplementation as a methane mitigation strategy
Nitrate (NO₃⁻) has been proposed as a replacement for urea (non-protein N) as a source of rumen degradable N for the rumen microbes (Lee and Beauchemin, 2014). Therefore, nitrate could work well under dry season cattle farming conditions in the tropics when grasses are characterized by deficient N concentration for optimal microbial growth supplying N essential for adequate bacterial protein synthesis (Leng, 1990). Nitrate is an alternative H sink which competes with methane synthesis; the stoichiometric potential of 1 mol of nitrate can reduce methane synthesis by 1 mol (Leng, 2008) and nitrate can decrease the number of rumen protozoa resulting in CH₄ reduction. Nitrate reduction in the rumen to nitrite and then to NH₃ is energetically more favorable than carbon dioxide reduction for methane synthesis, which implies that 4 mol H₂ is redirected toward nitrate reduction. Therefore, methane production can be mitigated by 1 mol per

each mole of nitrate reduced, which is equivalent to a CH₄ emission reduction of 25.8/100 g nitrate fed. However, high levels of nitrate in the rumen may induce toxicity, as nitrate is transformed into nitrite and an excess may induce an increase in methemoglobin (**MetHb**) in the bloodstream, leading to hypoxia, dyspnea and even death if the animal is not treated immediately (Callaghan *et al.*, 2014). Nitrate toxicity and its potentially hazardous consequences can be avoided by supplying less than 9 g NO₃⁻/kg DM intake in cattle, but the concentrations of basal NO₃⁻ in the consumed grass must also be accounted for.

Concentrations of NO₃⁻ are related to plant species, phenological stage and other plant stressors (e.g. drought, fertilization). According to Callaghan *et al.* (2014), in Northern Australia they recommended ~15 g/day of NO₃⁻ in breeding beef cows grazing dry season pastures. Nonetheless, there is a risk in the control of the intake of the supplement and the heterogeneous structure of herds of grazing cattle with different feed intakes. An experiment was carried out to evaluate the effect of dietary nitrate on enteric methane emissions in dairy cows grazing pastures during the summer and it was found that 8 g NO₃⁻/kg DM in grazing dairy cows led to a decrease in CH₄ emissions (van Wyngaard *et al.*, 2018). Hulshof *et al.* (2012) using sugarcane-based rations reported that 22 g NO₃⁻/kg DM reduced enteric methane emission by 32%.

In the dry season, tropical grasses usually contain low concentrations of CP and nitrates could supply the N required for improving rumen fermentation of low-quality basal rations. A gradual adaptation period of 15 days and the use of slow-release NO₃⁻ in the ration are alternative ways of reducing the risk of toxicity (Alemu *et al.*, 2019). El-Zaiat *et al.* (2014) showed that encapsulated nitrate was not toxic to lambs (4.51% encapsulated nitrate product (60.83% nitrate in the product DM)). At the same time, those authors identified higher risks of toxicity in rations with low levels of readily fermented carbohydrates. For that reason, a practice that reduces feeding problems is to supply non-structural carbohydrates in the ration, as grains (i.e. starch) which appear to exert some prophylactic effect. Nitrates can be offered twice a day or mixed with molasses in roller drums to reduce MetHb synthesis (Benu *et al.*, 2016). However, those practices are complicated in pastoral systems in the tropics, since forage quality varies with season. Villar *et al.* (2020) found that mature steers fed a basal ration of lucerne chaff and supplemented with a mixture of NO₃⁻ and canola oil reduced daily methane production by 26%, which suggested a synergistic effect of the two mitigating compounds, when combined.

Recent work, conducted at the University of Yucatan-Mexico, with crossbred heifers (*Bos indicus* × *Bos taurus*) housed in open-circuit respiration chambers showed that supplementation with increasing levels of NO₃⁻ (0.1%, 0.2% and 0.3% DM) to a basal ration of low-quality chopped tropical grass (*Pennisetum purpureum*) resulted in a substantial reduction in enteric CH₄ emissions from 135.5 g/day (0% NO₃⁻) to 93.3 g/day (with 0.3% NO₃⁻ DM), although a reduction in DM intake was also noticed. Nitrate seems to

be a viable methane mitigating additive which simultaneously supplies N to the rumen microorganisms during the dry season when N concentration in tropical grasses is deficient for optimal microbial protein synthesis in the rumen.

Lipid supplementation as a methane mitigation strategy

Lipid sources as a methane mitigation strategy under tropical conditions (rations with high NDF and low digestibility and intake) can play an important role in the improvement of cattle performance and therefore in farm productivity. In their comprehensive review, Grainger and Beauchemin (2011) suggested that the mitigation potential of fats is 1 g CH₄/kg DM intake for each 10 g fat/kg DM in the ration and this has shown persistent results, although no effect of the chain length of the fatty acids on the magnitude of mitigation has been detected. Feeding sunflower oil can effectively reduce enteric methane emissions of dairy cows (Bayat *et al.*, 2017). Similar effects of vegetable oils on reducing methane emissions have been reported under *in vitro* conditions (Vargas *et al.*, 2020). Recent work carried out at the University of Yucatan, Mexico (Flores-Santiago, E. unpublished results), has shown that by feeding palm oil (6% of ration DM) to hair Pelibuey sheep housed in respiration chambers, which were fed a low-quality basal ration of tropical C4 grass (*Brachiaria brizantha*) hay, methane yield was reduced by ~14%, compared to a control treatment without palm oil. Palm oil contains saturated fatty acids (50%; palmitic acid: 44% and stearic acid: 5%) and unsaturated fatty acids (oleic acid: 40% and polyunsaturated linoleic and α-linoleic acid: 10%). Palm oil is available for ruminant feeding in several Latin American and other tropical countries.

A number of experiments have been carried out in Brazil with grazing cattle supplemented with diverse sources of lipids. Mata e Silva *et al.* (2017) found that sunflower oil (383 g/day) given to lactating dairy cows grazing a tropical pasture (*Urochloa brizantha*) decreased enteric CH₄ by ~23%. Neto *et al.* (2015) reported that grazing Nelore bulls supplemented with a concentrate containing oil was an effective means for reducing enteric methane emissions as a percentage of GEI by 12.8%. Similarly, Carvalho *et al.* (2016) supplemented Nelore steers grazing *Brachiaria brizantha* with linseed oil and found a reduction in enteric methane emissions. In Nelore bulls grazing *Brachiaria brizantha* supplemented with starch, the use of oil (~198 g soybean grain/kg DM) reduced enteric CH₄ emissions (Neto *et al.*, 2019).

Practical implications of methane mitigation in tropical cattle systems

There is a shortage of information on enteric methane emissions from dairy and beef cattle production systems in the tropics (Bartl *et al.*, 2011; Niu *et al.*, 2018). Enteric CH₄ mitigation in small-scale cattle farms may be a rather

straightforward practice, from a purely technical standpoint. For instance, if it is assumed that crossbred (*Bos indicus* × *Bos taurus*) dual-purpose cows grazing on silvopastoral systems in the tropics consume ~30% of total daily DM intake (measured with the *n*-alkane technique) as *Leucaena leucocephala* (Bottini-Luzardo *et al.*, 2016), and by using the results obtained by Montoya-Flores *et al.* (2020) who found a reduction in CH₄ yield of ~20% in crossbred cows fed *Leucaena leucocephala* (36% DM; measured in respiration chambers). If those data are combined and extrapolated as a first approximation, it follows that the actual mitigation of enteric CH₄ emissions in silvopastoral systems with leucaena may possibly range from ~15% to ~20%, compared to an only-grass, monoculture system. In dual-purpose cattle production systems in the tropics, average milk yield per cow usually ranges between 7 and 10 litres/day, and there is an urgent need to increase productivity. Under those conditions, financial feasibility improves in the silvopastoral systems because the farmer relies less on imported (expensive) grains (i.e. concentrates) when using locally available high-quality forages such as leucaena (Bottini-Luzardo *et al.*, 2016; Kumari *et al.*, 2020). Reviewed data clearly show the range of methane yields (16.1 to 34.1 g CH₄/kg DM intake) which can be expected from cattle fed tropical grasses, with values at the lower end, probably reflecting the slow rate and low extent of fermentation of organic matter in the rumen (i.e. dry season grasses). Suybeng *et al.* (2019) have advanced the potential of *Desmanthus leptophyllus* as a promising legume for methane mitigation and improvement of cattle productivity under tropical grazing conditions in Australia.

Strategies based on the incorporation of natural resources such as vegetable oils, foliages, pods and seeds of tropical plants in the ration represent an approach for enteric CH₄ mitigation in ruminants in the vast areas of land covered with low-quality tropical grasses. Recently, Ali *et al.* (2019b) observed that when Holstein × Boran heifers were fed a basal ration of low-quality Rhodes hay and supplemented with sweet potato vine silage or a urea-molasses block, enteric CH₄ emissions were lowered while cattle were kept in respiration chambers in Kenya. In developing countries, it is important to work more intensively on technology transfer (extension) and farmer organization (cooperatives) in order to realize actual changes in cattle production, by introducing technologies with low environmental impact through the mitigation of GHG emissions such as CH₄ and N₂O. It is likely that in the near future it will be necessary to reach decreases in net methane emissions of cattle (g CH₄/head per day), to comply with international agreements (United Nations Framework Convention on Climate Change) where commitments towards absolute reductions in national GHG inventories have been signed by governments (Beauchemin *et al.*, 2020).

At low levels of nutrition of cattle, a situation which is common in the tropics during the dry season, it has been claimed that the slow rate of passage of solid digesta through the rumen may play a role in methane emissions (Ali *et al.*, 2019a), implying that if solid digesta is retained for a long

time in the rumen, methane emissions may increase. These data have obvious implications for methane inventories from tropical, developing countries, if this is substantiated further. In the event of that information being validated by further research, it would be rather easy to increase passage rate of digesta by feeding highly rumen fermentable forages (i.e. *Brosimum alicastrum*, *Leucaena leucocephala*, *Gliricidia sepium* and *Desmanthus* spp.) which have been proven to stimulate DM intake and thus passage rate in ruminants. In developing tropical countries, reduction of methane emissions in cattle by decreasing retention time of digesta in the rumen seems an achievable goal in the short-term, simply by supplementing highly rumen fermentable forages.

Productivity is expected to increase to some extent as a result of rechanneling energy (H₂) saved from CH₄ synthesis into protein and fat stored in the body or secreted in milk (van Zijderveld *et al.*, 2011). Indeed, some tropical plants (i.e. pods of *Samanea saman*; Valencia Salazar *et al.*, 2018) with mitigation potential may be particularly useful for significantly reducing enteric CH₄ production in cattle fed basal rations of low-quality grasses. Greening *et al.* (2019) have advanced the concept that the increase in animal productivity while mitigating CH₄ emissions relies on the control of H₂ use by methanogenic archaea. It seems then that redirection of H₂ flux by manipulating rumen fermentation can increase animal production by recovering energy that would be otherwise lost through methane synthesis in the rumen. This is a subject which requires further attention by animal scientists in the tropics, along with work on kinetics of digesta passage through the rumen (Ali *et al.*, 2019a), since mean retention time of solid digesta in the rumen is expected to contribute in a given extent to animal variation in feed efficiency (Cantalapiedra-Hijar *et al.*, 2018).


Conclusions

Foliages, pods and seeds of tropical trees and shrubs containing either condensed tannins, EOs, saponins or starch, as well as other compounds such as nitrates and by-products such as vegetable oils can be fed to cattle to mitigate enteric methane emissions under smallholder conditions. Methane yields can be reduced by approximately 10% to 25% by adopting those practices in ruminant production systems. Silvopastoral systems with high-quality legumes (i.e. leucaena, desmanthus), intensification of grazing management (i.e. *rotatinoous*) and supplementation with nutritional blocks containing CH₄ depressors (pods of *Enterolobium cyclocarpum* or *Samanea saman* or seeds of *Brosimum alicastrum*) are practical options available to smallholders for mitigating enteric methane emissions in cattle raised under tropical conditions.

Acknowledgements

A preliminary version of this paper was published in an abstract form at the 7th Greenhouse Gas and Animal

Agriculture Conference 2019, held at Iguassu Falls, Brazil. We are indebted to CONACYT-Mexico for granting fellowships to students to carry out MSc and PhD studies, as well as postdoctoral research at the University of Yucatan, Mexico. We thank the State Government of Yucatan (SIIDETEY)-Mexico for financial support to keep the respiration chambers running. We are grateful to CONACYT-Mexico for support (CB: A1-S-23910; Convocatoria 2018-1) to carry out experiments on energy metabolism of cattle. This work was implemented as part of the CGIAR Research Programs on Livestock and Climate Change, Agriculture and Food Security (CCAFS), which are carried out with support from CGIAR Fund Donors and through bilateral funding agreements. For details please visit <https://ccafs.cgiar.org/donors>. The views expressed in this document cannot be taken to reflect the official opinions of these organizations. We thank all donors that globally support our work through their contributions to the CGIAR system. We are indebted to the anonymous referees of this paper, for their kind suggestions towards its improvement.

 R. Jiménez-Ocampo 0000-0003-2171-1357

Declaration of interest

The authors declare no conflict of interest.

Ethics statement

The authors did not use any live animal to conduct this review.

Software and data repository resources

None of the data were deposited in an official repository.

References

- Aboagye IA, Oba M, Castillo AR, Koenig KM, Iwaasa AD and Beauchemin KA 2018. Effects of hydrolyzable tannin with or without condensed tannin on methane emissions, nitrogen use, and performance of beef cattle fed a high-forage diet. *Journal of Animal Science* 96, 5276–5286.
- Aboagye IA, Oba M, Koenig KM, Zhao GY and Beauchemin KA 2019. Use of gallic acid and hydrolyzable tannins to reduce methane emission and nitrogen excretion in beef cattle fed a diet containing alfalfa silage. *Journal of Animal Science* 97, 2230–2244.
- Albores-Moreno S, Alayón-Gamboa JA, Ayala-Burgos AJ, Solorio-Sánchez FJ, Aguilar-Pérez CF, Olivera-Castillo L and Ku-Vera JC 2017. Effects of feeding ground pods of *Enterolobium cyclocarpum* Jacq. Griseb on dry matter intake, rumen fermentation, and enteric methane production by Pelibuey sheep fed tropical grass. *Tropical Animal Health and Production* 49, 857–866.
- Alemu A, Romero-Pérez A, Araujo R and Beauchemin K 2019. Effect of encapsulated nitrate and microencapsulated blend of essential oils on growth performance and methane emissions from beef steers fed backgrounding diets. *Animals* 9, 21.
- Ali AIM, Wassie SE, Korir D, Goopy JP, Merbold L, Butterbach-Bahl K, Dickhoefer U and Schlecht E 2019a. Digesta passage and nutrient digestibility in Boran steers at low feed intake levels. *Journal of Animal Physiology and Animal Nutrition* 103, 1325–1337.
- Ali AIM, Wassie SE, Korir D, Merbold L, Goopy JP, Butterbach-Bahl K, Dickhoefer U and Schlecht E 2019b. Supplementing tropical cattle for improved nutrient utilization and reduced enteric methane emissions. *Animals* 9, 210.
- Anantasook N, Wanapat M and Cherdthong A 2014. Manipulation of ruminal fermentation and methane production by supplementation of rain tree pod meal containing tannins and saponins in growing dairy steers. *Journal of Animal Physiology and Animal Nutrition* 98, 50–55.
- Archimède H, Eugène M, Marie Magdeleine C, Boval M, Martin C, Morgavi DP, Lecomte P and Doreau M 2011. Comparison of methane production between C3 and C4 grasses and legumes. *Animal Feed Science and Technology* 166, 59–64.
- Archimède H, Rira M, Eugène M, Fleury J, Lastel ML, Periacarpan F, Silou-Etienne T, Morgavi DP and Doreau M 2018. Intake, total-tract digestibility and methane emissions of Texel and Blackbelly sheep fed C4 and C3 grasses tested simultaneously in a temperate and a tropical area. *Journal of Cleaner Production* 185, 455–463.
- Ayantunde AA, Fernández-Rivera S, McCrabb G and International Livestock Research Institute 2005. Coping with feed scarcity in smallholder livestock systems in developing countries. International Livestock Research Institute, Nairobi, Kenya.
- Bartl K, Gómez CA and Nemecek T 2011. Life cycle assessment of milk produced in two smallholder dairy systems in the highlands and the coast of Peru. *Journal of Cleaner Production* 19, 1494–1505.
- Bayat AR, Ventto L, Kairenius P, Stefański T, Leskinen H, Tapio I, Negussie E, Viikki J and Shingfield KJ 2017. Dietary forage to concentrate ratio and sunflower oil supplement alter rumen fermentation, ruminal methane emissions, and nutrient utilization in lactating cows. *Translational Animal Science* 1, 277–286.
- Beauchemin KA, Ungerfeld EM, Eckard RJ and Wang M 2020. Review: fifty years of research on rumen methanogenesis: lessons learned and future challenges for mitigation. *Animal* 14, s2–s16.
- Benaouda M 2018. Efecto de la calidad de la dieta y el nivel de la fibra en la cinética de digestión y la producción de metano en bovino lechero. Universidad Autónoma del Estado de México, Toluca, Estado de México.
- Benchaar C, Calsamiglia S, Chaves AV, Fraser GR, Colombatto D, McAllister TA and Beauchemin KA 2008. A review of plant-derived essential oils in ruminant nutrition and production. *Animal Feed Science and Technology* 145, 209–228.
- Benu I, Callaghan MJ, Tomkins N, Hepworth G, Fitzpatrick LA and Parker AJ 2016. The effect of feeding frequency and dose rate of nitrate supplements on blood haemoglobin fractions in *Bos indicus* cattle fed Flinders grass (*Iseilemia* spp.) hay. *Animal Production Science* 56, 1605–1611.
- Berça AS, Cardoso ADS, Longhini VZ, Tedeschi LO, Boddey RM, Berndt A, Reis RA and Ruggieri AC 2019. Methane production and nitrogen balance of dairy heifers grazing palisade grass cv. Marandu alone or with forage peanut. *Journal of Animal Science* 97, 4625–4634.
- Berndt A and Tomkins NW 2013. Measurement and mitigation of methane emissions from beef cattle in tropical grazing systems: a perspective from Australia and Brazil. *Animal* 7, 363–372.
- Bottini-Luzardo MB, Aguilar-Pérez CF, Centurión-Castro FG, Solorio-Sánchez FJ and Ku-Vera JC 2016. Milk yield and blood urea nitrogen in crossbred cows grazing *Leucaena leucocephala* in a silvopastoral system in the Mexican tropics. *Tropical Grasslands-Forrajeras Tropicales* 4, 159–167.
- Briceño-Poot EG, Ruiz-González A, Chay-Canul AJ, Ayala-Burgos AJ, Aguilar-Pérez CF, Solorio-Sánchez FJ and Ku-Vera JC 2012. Voluntary intake, apparent digestibility and prediction of methane production by rumen stoichiometry in sheep fed pods of tropical legumes. *Animal Feed Science and Technology* 176, 117–122.
- Callaghan MJ, Tomkins NW, Benu I and Parker AJ 2014. How feasible is it to replace urea with nitrates to mitigate greenhouse gas emissions from extensively managed beef cattle? *Animal Production Science* 54, 1300–1304.
- Cantalapiedra-Hijar G, Abo-Ismael M, Carstens GE, Guan LL, Hegarty R, Kenny DA, McGee M, Plastow G, Relling A and Ortigues-Marty I 2018. Review: biological determinants of between-animal variation in feed efficiency of growing beef cattle. *Animal* 12, s321–s335.
- Canul-Solis JR, Piñero-Vázquez AT, Arceo-Castillo JI, Alayón-Gamboa JA, Ayala-Burgos AJ, Aguilar-Pérez CF, Solorio-Sánchez FJ, Castelan-Ortega OA, Lachica-López M, Quintana-Owen P and Ku-Vera JC 2017. Design and construction of low-cost respiration chambers for ruminal methane measurements in ruminants. *Revista Mexicana de Ciencias Pecuarias* 8, 185–191.
- Canul-Solis JR, Piñero-Vázquez AT, Briceño-Poot EG, Chay-Canul AJ, Alayón-Gamboa JA, Ayala-Burgos AJ, Aguilar-Pérez CF, Solorio-Sánchez FJ, Castelan-Ortega OA and Ku-Vera JC 2014. Effect of supplementation with saponins from *Yucca schidigera* on ruminal methane production by Pelibuey sheep fed *Pennisetum purpureum* grass. *Animal Production Science* 54, 1834–1837.
- de Carvalho IPC, Fiorentini G, Berndt A, Castagnino PdS, Messana JD, Frighetto RTS, Reis RA, Berchielli T 2016. Performance and methane emissions of Nelore steers grazing tropical pasture supplemented with lipid sources. *Revista Brasileira de Zootecnia* 45, 760–767.

- Castelán Ortega OA, Ku Vera JC, Ángeles-Hernández JC, Hernández-Pineda GS, Benaouda M, Molina LT, Ramírez-Cancino L, Castelán-Jaime SV, Praga-Ayala AR, Lazos-Balbuena F, Montelongo-Pérez H, González-Ronquillo M, Vázquez-Carrillo MF, Cardoso-Gutiérrez E, Aranda-Aguirre E, Villegas-Estrada D, Guadarrama-López AP and Apodaca-Martínez G 2019. Ganadería. In Estado del Ciclo del Carbono: agenda Azul y Verde, Programa Mexicano del Carbono, pp. 492–528. Texcoco, Estado de México, México.
- Chaokaur A, Nishida T, Phaowphaisal I and Sommart K 2015. Effects of feeding level on methane emissions and energy utilization of Brahman cattle in the tropics. *Agriculture, Ecosystems & Environment* 199, 225–230.
- Charmley E, Williams SRO, Moate PJ, Hegarty RS, Herd RM, Oddy VH, Reyenga P, Staunton KM, Anderson A and Hannah MC 2016. A universal equation to predict methane production of forage-fed cattle in Australia. *Animal Production Science* 56, 169–180.
- Congio GFS, Batalha CDA, Chiavegato MB, Berndt A, Oliveira PPA, Frighetto RTS, Maxwell TMR, Gregorini P and Da Silva SC 2018. Strategic grazing management towards sustainable intensification at tropical pasture-based dairy systems. *Science of the Total Environment* 636, 872–880.
- El-Zaiat HM, Araujo RC, Soltan YA, Morsy AS, Louvandini H, Pires AV, Patino HO, Correa PS and Abdalla AL 2014. Encapsulated nitrate and cashew nut shell liquid on blood and rumen constituents, methane emission, and growth performance of lambs. *Journal of Animal Science* 92, 2214–2224.
- Escobar-Bahamondes P, Oba M and Beauchemin KA 2016. Universally applicable methane prediction equations for beef cattle fed high- or low-forage diets. *Canadian Journal of Animal Science* 97, 83–94.
- Eugène M, Sauvant D, Nozière P, Viillard D, Oueslati K, Lherm M, Mathias E and Doreau M 2019. A new Tier 3 method to calculate methane emission inventory for ruminants. *Journal of Environmental Management* 231, 982–988.
- Goopy JP, Korir D, Pelster D, Ali AIM, Wassie SE, Schlecht E, Dickhoefer U, Merbold L and Butterbach-Bahl K 2020. Severe below-maintenance feed intake increases methane yield from enteric fermentation in cattle. *British Journal of Nutrition* 123, 1239–1246.
- Goopy JP, Onyango AA, Dickhoefer U and Butterbach-Bahl K 2018. A new approach for improving emission factors for enteric methane emissions of cattle in smallholder systems of East Africa - Results for Nyando, Western Kenya. *Agricultural Systems* 161, 72–80.
- Grainger C and Beauchemin KA 2011. Can enteric methane emissions from ruminants be lowered without lowering their production? *Animal Feed Science and Technology* 166–167, 308–320.
- Greening C, Geier R, Wang C, Woods LC, Morales SE, McDonald MJ, Rushton-Green R, Morgan XC, Koike S, Leahy SC, Kelly WJ, Cann I, Attwood GT, Cook GM and Mackie RI 2019. Diverse hydrogen production and consumption pathways influence methane production in ruminants. *The ISME Journal* 13, 2617–2632.
- Harrison MT, McSweeney C, Tomkins NW and Eckard RJ 2015. Improving greenhouse gas emissions intensities of subtropical and tropical beef farming systems using *Leucaena leucocephala*. *Agricultural Systems* 136, 138–146.
- Hart KJ, Yañez-Ruiz DR, Duval SM, McEwan NR and Newbold CJ 2008. Plant extracts to manipulate rumen fermentation. *Animal Feed Science and Technology* 147, 8–35.
- Hegarty RS and Gerdes R 1999. Hydrogen production and transfer in the rumen. *Recent Advances in Animal Nutrition in Australia* 12, 37–44.
- Hulshof RBA, Berndt A, Gerrits WJJ, Dijkstra J, van Zijderveld SM, Newbold JR and Perdok HB 2012. Dietary nitrate supplementation reduces methane emission in beef cattle fed sugarcane-based diets. *Journal of Animal Science* 90, 2317–2323.
- IPCC 2006. IPCC Guidelines for National Greenhouse Gas Inventories.
- Jafari S, Ebrahimi M, Goh YM, Rajion MA, Jahromi MF and Al-Jumaili WS 2019. Manipulation of rumen fermentation and methane gas production by plant secondary metabolites (saponin, tannin and essential oil) – a review of ten-year studies. *Annals of Animal Science* 19, 3–29.
- Kaewpila C and Sommart K 2016. Development of methane conversion factor models for Zebu beef cattle fed low-quality crop residues and by-products in tropical regions. *Ecology and Evolution* 6, 7422–7432.
- Kennedy PM and Charmley E 2012. Methane yields from Brahman cattle fed tropical grasses and legumes. *Animal Production Science* 52, 225–239.
- Knapp JR, Laur GL, Vadas PA, Weiss WP and Tricarico JM 2014. Invited review: enteric methane in dairy cattle production: quantifying the opportunities and impact of reducing emissions. *Journal of Dairy Science* 97, 3231–3261.
- Kumari S, Fagodiya RK, Hiloidhari M, Dahiya RP and Kumar A 2020. Methane production and estimation from livestock husbandry: A mechanistic understanding and emerging mitigation options. *Science of the Total Environment* 709, 136135.
- Ku-Vera JC, Valencia-Salazar SS, Piñero-Vázquez AT, Molina-Botero IC, Arroyave-Jaramillo J, Montoya-Flores MD, Lazos-Balbuena FJ, Canul-Solis JR, Arceo-Castillo JI, Ramírez-Cancino L, Escobar-Restrepo CS, Alayón-Gamboa JA, Jiménez-Ferrer G, Zavala-Escalante LM, Castelán-Ortega OA, Quintana-Owen P, Ayala-Burgos AJ, Aguilar-Pérez CF and Solorio-Sánchez FJ 2018. Determination of methane yield in cattle fed tropical grasses as measured in open-circuit respiration chambers. *Agricultural and Forest Meteorology* 258, 3–7.
- Lazos Balbuena FJ 2015. Uso del fruto de *Enterolobium cyclocarpum* como fuente de saponinas esteroidales para reducir la producción de metano entérico en bovinos. Tesis de Maestría thesis. Universidad Autónoma de Yucatán, Mérida, Yucatán.
- Lee C and Beauchemin KA 2014. A review of feeding supplementary nitrate to ruminant animals: nitrate toxicity, methane emissions, and production performance. *Canadian Journal of Animal Science* 94, 557–570.
- Leng RA 1990. Factors affecting the utilization of 'poor-quality' forages by ruminants particularly under tropical conditions. *Nutrition Research Reviews* 3, 277–303.
- Leng RA 2008. The potential of feeding nitrate to reduce enteric methane production in ruminants. A Report to the Department of Climate Change, Commonwealth Government of Australia, Canberra, Australia.
- van Lingen HJ, Niu M, Kebreab E, Valadares Filho SC, Rooke JA, Duthie C-A, Schwarm A, Kreuzer M, Hynd PI, Caetano M, Eugène M, Martin C, McGee M, O'Kiely P, Hünerberg M, McAllister TA, Berchielli TT, Messana JD, Peiren N, Chaves AV, Charmley E, Cole NA, Hales KE, Lee S-S, Berndt A, Reynolds CK, Crompton LA, Bayat A-R, Yañez-Ruiz DR, Yu Z, Bannink A, Dijkstra J, Casper DP and Hristov AN 2019. Prediction of enteric methane production, yield and intensity of beef cattle using an intercontinental database. *Agriculture, Ecosystems & Environment* 283, 106575.
- Machado FS, Tomich TR, Ferreira AL, Cavalcanti LFL, Campos MM, Paiva CAV, Ribas MN and Pereira LGR 2016. Technical note: A facility for respiration measurements in cattle. *Journal of Dairy Science* 99, 4899–4906.
- Mata e Silva BC, Lopes FCF, Pereira LGR, Tomich TR, Morenz MJF, Martins CE, Gomide CAM, Paciullo DSC, Maurício RM and Chaves AV 2017. Effect of sunflower oil supplementation on methane emissions of dairy cows grazing *Urochloa brizantha* cv. Marandu. *Animal Production Science* 57, 1431–1436.
- Mayo-Eusebio R 2013. Ganancia de peso de bovino en crecimiento en un sistema silvopastoril intensivo y en un sistema en confinamiento en Michoacán, México. MSc thesis in agricultural science thesis. University of Yucatan, Yucatan, Mexico.
- Molina IC, Angarita EA, Mayorga OL, Chará JE and Barahona-Rosales R 2016. Effect of *Leucaena leucocephala* on methane production of Lucerna heifers fed a diet based on *Cynodon plectostachyus*. *Livestock Science* 185, 24–29.
- Molina-Botero IC, Arroyave-Jaramillo J, Valencia-Salazar S, Barahona-Rosales R, Aguilar-Pérez CF, Ayala Burgos A, Arango J and Ku-Vera JC 2019a. Effects of tannins and saponins contained in foliage of *Gliricidia sepium* and pods of *Enterolobium cyclocarpum* on fermentation, methane emissions and rumen microbial population in crossbred heifers. *Animal Feed Science and Technology* 251, 1–11.
- Molina-Botero IC, Montoya-Flores MD, Zavala-Escalante LM, Barahona-Rosales R, Arango J and Ku-Vera JC 2019b. Effects of long-term diet supplementation with *Gliricidia sepium* foliage mixed with *Enterolobium cyclocarpum* pods on enteric methane, apparent digestibility, and rumen microbial population in crossbred heifers. *Journal of Animal Science* 97, 1619–1633.
- Montoya-Flores MD, Molina-Botero IC, Arango J, Romano-Muñoz JL, Solorio-Sánchez FJ, Aguilar-Pérez CF and Ku-Vera JC 2020. Effect of dried leaves of *Leucaena leucocephala* on rumen fermentation, rumen microbial population, and enteric methane production in crossbred heifers. *Animals* 10, 300.
- Moscoco C, Vélez M, Flores A and Agudelo N 1995. Effects of guanacaste tree (*Enterolobium cyclocarpum* Jacq. Griseb.) fruit as replacement for sorghum grain and cotton-seed meal in lamb diets. *Small Ruminant Research* 18, 121–124.
- National Academies of Sciences, Engineering, and Medicine (NASEM) 2016. Nutrient requirements of beef cattle, 8th revised edition. The National Academies Press, Washington, DC, USA. <https://doi.org/10-17226/19014>
- Ndung'u PW, Bebe BO, Ondiek JO, Butterbach-Bahl K, Merbold L and Goopy JP 2019. Improved region-specific emission factors for enteric methane emissions from cattle in smallholder mixed crop: livestock systems of Nandi County, Kenya. *Animal Production Science* 59, 1136.

- Neto AJ, Messana JD, Ribeiro AF, Vito ES, Rossi LG and Berchielli TT 2015. Effect of starch-based supplementation level combined with oil on intake, performance, and methane emissions of growing Nelore bulls on pasture. *Journal of Animal Science* 93, 2275–2284.
- Neto JA, Messana JD, Rossi LG, Carvalho IPC and Berchielli TT 2019. Methane emissions from Nelore bulls on pasture fed two levels of starch-based supplement with or without a source of oil. *Animal Production Science* 59, 654–663.
- Niu M, Kebreab E, Hristov AN, Oh J, Arndt C, Bannink A, Bayat AR, Brito AF, Boland T, Casper D, Crompton LA, Dijkstra J, Eugène MA, Garnsworthy PC, Haque MN, Hellwing ALF, Huhtanen P, Kreuzer M, Kuhla B, Lund P, Madsen J, Martin C, McClelland SC, McGee M, Moate PJ, Muetzel S, Muñoz C, O’Kiely P, Peiren N, Reynolds CK, Schwarm A, Shingfield KJ, Storlien TM, Weisbjerg MR, Yáñez-Ruiz DR and Yu Z 2018. Prediction of enteric methane production, yield, and intensity in dairy cattle using an intercontinental database. *Global Change Biology* 24, 3368–3389.
- Noguera RR and Posada-Ochoa SL 2017. Factor de emisión de metano entérico para vacas Holstein lactantes en la zona norte de Antioquia - Colombia. *Livestock Research for Rural Development* 29, 6.
- Patra A 2017. Prediction of enteric methane emission from cattle using linear and non-linear statistical models in tropical production systems. *Mitigation and Adaptation Strategies for Global Change* 22, 629–650.
- Patra A, Park T, Kim M and Yu Z 2017. Rumen methanogens and mitigation of methane emission by anti-methanogenic compounds and substances. *Journal of Animal Science and Biotechnology* 8, 13.
- Patra A and Saxena J 2009. Dietary phytochemicals as rumen modifiers: a review of the effects on microbial populations. *Antonie van Leeuwenhoek* 96, 363–375.
- Perry LA, Al Jassim R, Gaughan JB and Tomkins NW 2017. Effect of feeding forage characteristic of wet- or dry-season tropical C4 grass in northern Australia, on methane production, intake and rumen outflow rates in *Bos indicus* steers. *Animal Production Science* 57, 2033.
- Piñeiro-Vázquez AT, Ayala-Burgos AJ, Chay-Canul AJ and Ku-Vera JC 2013. Dry matter intake and digestibility of rations replacing concentrates with graded levels of *Enterolobium cyclocarpum* in Pelibuey lambs. *Tropical Animal Health and Production* 45, 577–583.
- Piñeiro-Vázquez AT, Canul-Solis JR, Jiménez-Ferrer GO, Alayón-Gamboa JA, Chay-Canul AJ, Ayala-Burgos AJ, Aguilar-Pérez CF and Ku-Vera JC 2018. Effect of condensed tannins from *Leucaena leucocephala* on rumen fermentation, methane production and population of rumen protozoa in heifers fed low-quality forage. *Asian-Australasian Journal of Animal Sciences* 31, 1738–1746.
- Ramírez-Avilés L, Solorio-Sánchez FJ, Aguilar-Pérez CF, Ayala-Burgos AJ and Ku-Vera JC 2019. *Leucaena leucocephala* feeding systems for cattle production in Mexico. *Tropical Grasslands-Forrajes Tropicales* 7, 375–380.
- Romero-Perez A, Okine EK, McGinn SM, Guan LL, Oba M, Duval SM, Kindermann M and Beauchemin KA 2015. Sustained reduction in methane production from long-term addition of 3-nitrooxypropanol to a beef cattle diet. *Journal of Animal Science* 93, 1780–1791.
- Savian JV, Schons RMT, Marchi DE, de Freitas TS, da Silva Neto GF, Mezzalana JC, Berndt A, Bayer C and Carvalho PCdF 2018. Rotatinuous stocking: a grazing management innovation that has high potential to mitigate methane emissions by sheep. *Journal of Cleaner Production* 186, 602–608.
- Shelton M and Dalzell S 2007. Production, economic and environmental benefits of leucaena pastures. *Tropical Grasslands* 41, 174–190.
- Sierra-Montoya E, Chará JE and Barahona-Rosales R 2017. The nutritional balance of early lactation dairy cows grazing in intensive silvopastoral systems. *Ciencia Animal Brasileira* 18, 1–12.
- Suybeng B, Charmley E, Gardiner CP, Malau-Aduli BS and Malau-Aduli AEO 2019. Methane emissions and the use of desmanthus in beef cattle production in Northern Australia. *Animals* 9, 542.
- Topps JH 1992. Potential, composition and use of legume shrubs and trees as fodders for livestock in the tropics. *The Journal of Agricultural Science* 118, 1–8.
- Valencia Salazar SS 2017. Efecto del consumo de frutos de *Samanea saman* sobre la digestibilidad, patrón de fermentación y producción de metano entérico en novillas. Master in science thesis, Universidad Autónoma de Yucatán, Mérida, Yucatán.
- Valencia Salazar SS, Piñeiro Vázquez AT, Molina Botero IC, Lazos Balbuena FJ, Uuh Narváez JJ, Segura Campos MR, Ramírez Avilés L, Solorio Sánchez FJ and Ku Vera JC 2018. Potential of *Samanea saman* pod meal for enteric methane mitigation in crossbred heifers fed low-quality tropical grass. *Agricultural and Forest Meteorology* 258, 108–116.
- Vargas JE, Andrés S, López-Ferreras L, Snelling TJ, Yáñez-Ruiz DR, García-Estrada C and López S 2020. Dietary supplemental plant oils reduce methanogenesis from anaerobic microbial fermentation in the rumen. *Scientific Reports* 10, 1613.
- Villar ML, Hegarty RS, Nolan JV, Godwin IR and McPhee M 2020. The effect of dietary nitrate and canola oil alone or in combination on fermentation, digesta kinetics and methane emissions from cattle. *Animal Feed Science and Technology* 259, 114294.
- Wang C, Liu Q, Guo G, Huo WJ, Ma L, Zhang YL, Pei CX, Zhang SL and Wang H 2018. Effects of dietary soybean oil and coated folic acid on ruminal digestion kinetics, fermentation, microbial enzyme activity and bacterial abundance in Jinnan beef steers. *Livestock Science* 217, 92–98.
- Wu L, Koerkamp PWGG and Ogink N 2018. Uncertainty assessment of the breath methane concentration method to determine methane production of dairy cows. *Journal of Dairy Science* 101, 1554–1564.
- van Wyngaard JDV, Meeske R and Erasmus LJ 2018. Effect of dietary nitrate on enteric methane emissions, production performance and rumen fermentation of dairy cows grazing kikuyu-dominant pasture during summer. *Animal Feed Science and Technology* 244, 76–87.
- Ye D, Karnati SKR, Wagner B, Firkins JL, Eastridge ML and Aldrich JM 2018. Essential oil and monensin affect ruminal fermentation and the protozoal population in continuous culture. *Journal of Dairy Science* 101, 5069–5081.
- van Zijderveld SM, Gerrits WJJ, Dijkstra J, Newbold JR, Hulshof RBA and Perdok HB 2011. Persistency of methane mitigation by dietary nitrate supplementation in dairy cows. *Journal of Dairy Science* 94, 4028–4038.