

1 **TITLE PAGE**

2

3 **Translation, Cross-Cultural Adaptation and Preliminary Validation of the Transsexual Voice**
4 **Questionnaire for Male-to-Female Transsexuals (I-TVQ^{MtF}) into Italian**

5 Carlo Robotti¹, Francesco Mozzanica^{2,3}, Chiara Atzori⁴, Andrea Cavalot⁵, Diego Cossu⁵, Adi
6 Primov-Fever⁶, Marco Benazzo¹, Luca Negri⁷, Antonio Schindler⁸

7

8 1. Department of Otolaryngology - Head and Neck Surgery, University of Pavia, IRCCS Policlinico
9 San Matteo Foundation, Pavia, Italy

10 2. Department of Clinical Sciences and Community Health, University of Milan, Milan, Italy

11 3. Department of Otorhinolaryngology, San Giuseppe Hospital, IRCCS Multimedica, Milan, Italy

12 4. Department of Infectious Diseases, ASST Fatebenefratelli Sacco, Luigi Sacco University
13 Hospital, Milan, Italy

14 5. Otorhinolaryngology Division, Santa Croce Hospital, Moncalieri, Italy

15 6. Department of Otolaryngology and Head and Neck Surgery, Chaim Sheba Medical Center,
16 Ramat-Gan, Israel; Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel.

17 7. Department of Pathophysiology and Transplantation, University of Milan, Milan, Italy.

18 8. Department of Biochemical and Clinical science "L. Sacco", Luigi Sacco University Hospital,
19 University of Milan, Milan, Italy

20

21 **Author for correspondence**

22 Francesco Mozzanica, MD, PhD. Department of Biochemical and Clinical science "Luigi Sacco"

23 Via G.B. Grassi 74, Milan, 20157, Italy;

24 Phone: +39 340 4662852; Email: francesco.mozzanica@gmail.com

25

26 Declarations of interest: none

1 **Abstract**

2 **Objective.** To perform a cross-cultural adaptation into Italian and to analyse reliability and validity
3 of the Transsexual Voice Questionnaire for male-to-female transsexuals (I-TVQ^{MtF}).

4 **Study Design.** Cross-sectional non-randomized survey study.

5 **Methods.** For item-generation, a cross-cultural adaptation and translation process was performed
6 following standard guidelines. Transgender women were consecutively recruited and asked to fill
7 out the I-TVQ^{MtF} and a form on social, demographic and transition-related variables. Firstly, data
8 collected from participants were used to perform confirmatory factor analysis (CFA), and to
9 evaluate internal consistency and test-retest reliability. Subsequently, convergent validity was
10 evaluated comparing I-TVQ^{MtF} total scores with the two extra items addressing self-perception
11 (SPVF) and aspiration (AVF) of voice femininity. To further evaluate convergent validity, scores of
12 the Italian version of the Voice Handicap Index (I-VHI) were considered for comparisons. A
13 correlation analysis was performed to verify potential association between I-TVQ^{MtF} scores and
14 social, demographic and transition-related variables.

15 **Results.** CFA demonstrated that a two-factor model fits data better than the unidimensional one.
16 Both internal consistency and test retest reliability of the I-TVQ^{MtF} were satisfactory. Negative
17 correlations were highlighted between I-TVQ^{MtF} scores on one side and SPVF and AVF on the
18 other. Positive correlations between I-TVQ^{MtF} and I-VHI scores were also found. Finally, negative
19 correlations were demonstrated between I-TVQ^{MtF} scores and time spent living in the female role.

20 **Conclusion.** The I-TVQ^{MtF} appears to be a reliable and valid instrument for the assessment of
21 voice-related quality of life in transgender women.

22 **Keywords:** transgender; voice; quality of life; cross-cultural adaptation; dysphonia

23

1 **1. Introduction**

2 **Transgender (or trans) is an umbrella term used to describe people whose gender identities**
3 **(or gender roles) differ from those typically associated with the sex they were assigned at**
4 **birth. Gender identity is an individual's inner sense of being a man or a woman. For some**
5 **individuals, their gender identity does not fit neatly into those two choices. For transgender**
6 **individuals, the sex they were assigned at birth and their own internal gender identity do not**
7 **match. [1-4]. This cross-gender identification is usually accompanied by a persistent sense of**
8 **inappropriateness of the anatomic sex, which may determine subjective discomfort, feelings of**
9 **inadequacy, and negative societal attitudes in terms of activity limitations and participation**
10 **restrictions [5]. Therefore, in order to reach an acceptable grade of gender conformity, transgender**
11 **individuals frequently seek medical and surgical treatments. In the search of adequacy between**
12 **physical appearance and gender identity, voice can represent a critical obstacle, because it acts as a**
13 **salient cue to the listener's perception of a speaker's gender [6-8]. Interestingly, the magnitude of**
14 **that obstacle is peculiarly different between transgender men and transgender women. As far**
15 **as transgender men are concerned, voice changes are achieved effortlessly in most cases. In**
16 **fact, hormone replacement therapy (HRT) with androgens (testosterone) induces hypertrophy**
17 **of the thyroarytenoid muscles, thus increasing the mass of the vocal folds and lowering the**
18 **fundamental frequency (F0) towards the male range [9-11]. Therefore, transgender men may**
19 **reach the desired low-pitched voice with HRT alone, with little or no need for further**
20 **improvements through voice therapy [12]. On the other hand, transgender women usually**
21 **struggle to achieve a gender-congruent voice, as HRT with antiandrogens (estrogens)**
22 **feminizes the body but has no effects on their voices [13]. This is the reason why voice**
23 **modification is requested especially by trans women, whose voice characteristics are often male-**
24 **like and thus strongly contrasting with their identity and presentation as women [14, 15]. Failing**
25 **to reach a gender congruent voice might determine important negative effects on these**
26 **subjects. For example, Dacakis and collaborators reported that psychosocial issues were the**

1 **most frequently reported voice-related impacts in transgender women. Specifically, their**
2 **results highlighted how the anxiety and the frustration caused by the misalignment between**
3 **physical appearance and voice may raise feelings of inadequacy and self-consciousness**
4 **towards strangers in transgender women [16]. Conversely, transgender subjects whose voices**
5 **sound more congruent with the experienced gender, report greater well-being (greater life**
6 **satisfaction, better QOL and better self-esteem, plus lower levels of anxiety and depression)**
7 **than the ones with less gender congruent voices [17, 18]. In other words, trans individuals**
8 **whose vocal gender better corresponds to their experienced gender enjoy a higher level of**
9 **well-being along a wide-ranging array of measures, not limited to simple satisfaction with**
10 **their own voices [18].** Because of its potential psychosocial effect, the management of voice
11 change in **transgender** women requires a more comprehensive evaluation, including not only
12 acoustic analysis and auditory-perceptual evaluation, but also the evaluation of the individual's
13 voice-related difficulties and their impacts on everyday life. The importance of evaluating a
14 **transgender** woman's perception of her voice is highlighted by the fact that voice feminization
15 goes way beyond mere pitch elevation, since it has been demonstrated that a higher pitch does not
16 necessarily result in listeners perceiving a voice as female [19]. As a matter of fact, F0 – which is
17 the acoustic correlate of pitch – seems to account for just 10% of the variance in happiness with
18 voice in **trans** women [20]. Likewise, clinicians' perceptions of voices of **trans women** do not
19 consistently correlate with the clients' satisfaction regarding their own voice [21, 22]. Transgender
20 individuals are highly heterogeneous in terms of both current voice perception and future voice
21 expectations. Since each transgender individual experiences different challenges, self-evaluation
22 should represent a crucial element from the very beginning of clinical evaluation. Moreover,
23 objective and subjective voice assessments are not necessarily always correlated [23, 24].
24 Therefore, given the multitude of factors playing a role in determining voice-related QOL in
25 transgender individuals, clinicians should understand the uniqueness of each trans patient's
26 perspective, in order to tailor client-centered goals.

1 Only a few instruments specifically designed to investigate the self-perception of voice limitations
2 and the impact of voice on quality of life (QOL) in **transgender women** are available so far [16,
3 25]. The Transgender Self-Evaluation Questionnaire (TSEQ) was the first patient-reported outcome
4 (PRO) instrument developed for the assessment of voice-related QOL in **transgender** subjects. It
5 was based on the well-known Voice Handicap Index (VHI), which was partially modified to
6 capture relevant voice-related features in **trans** patients [25]. Subsequently, an extensive reviewing
7 and refining process of the TSEQ, conducted by Dacakis and colleagues, led to the Transsexual
8 Voice Questionnaire for Male-to-Female Transsexuals (TVQ^{MtF}) [16]. The TVQ^{MtF} is a self-report
9 measure of vocal functioning and voice-related impact on **transgender** women's QOL. It contains
10 30 items, each rated on a 4-point Likert scale (1 = "never or rarely"; 2 = "sometimes"; 3 = "often";
11 4 = "usually or always"), for a total score ranging from 30 to 120; a lower score represents a more
12 desirable outcome. **As far as the name of the original questionnaire is concerned (TVQ^{MtF}), the
13 terminology related to this populations significantly changed during the last few years, and
14 terms such as "transsexual" and "male-to-female" may now be considered as offensive by
15 members of the population. Therefore, the authors are willing to change the original title, and
16 a new name for this outcome instrument will be released in the upcoming months [3].
17 Therefore, the title of the original questionnaire will be used in the present paper solely for
18 the purpose of referring to the instrument itself.** Through principal component analysis, Dacakis
19 demonstrated the two-component structure of the TVQ^{MtF}, accounting for almost 58% of the
20 variance in the questionnaire: (a) "vocal functioning" (VF), fourteen items, dealing with voice
21 production and its relationship with gender identity; (b) "social participation" (SP), twelve items,
22 relating to the impact of an individual's voice on participation in everyday life. Four items were not
23 allocated to either domain for loading on both factors. The two-domain structure enhances the
24 ability of the questionnaire to provide significant information regarding the nature of the
25 impairment and to tailor strategies and goals of treatment [15]. Moreover, the TVQ^{MtF} comes with
26 two extra items – "Currently my voice is" and "My ideal voice would sound" – dealing with self-

1 perception (SPVF) and **aspiration (AVF)** of voice femininity, both rated on a 5-point Likert scale
2 (1 = “very female”; 2 = “somewhat female”; 3 = “gender neutral”; 4 = “somewhat male”; 5 = “very
3 male”) [20, 26]. Several studies have already assessed the reliability and the validity of the TVQ^{MtF}
4 [15, 16, 25]. Moreover, the TVQ^{MtF} has been already adapted and validated for different cultural
5 and linguistic contexts [8, 27-30]. To date, however, a validated Italian version of this instrument is
6 not yet available. **In addition, to the best of our knowledge, no studies have been conducted so
7 far to investigate voice-related QOL in Italian transgender women.** Therefore, this study has
8 been structured to evaluate the reliability and the validity of the Italian version of the TVQ^{MtF} (I-
9 TVQ^{MtF}). We surmise that the TVQ^{MtF} can be adapted to the Italian language and that both validity
10 and reliability of the I-TVQ^{MtF} are satisfactory. A validated Italian version of this instrument will
11 allow clinicians to thoroughly assess voice-related impairments and needs in **transgender** women.
12 Additionally, the Italian validated translation of this outcome instrument will be useful in designing
13 cross-cultural and international outcome studies addressing voice-related issues in **transgender**
14 women.

15

16

1 **2. Materials and Methods**

2 This non-randomized cross-sectional survey study was carried out according to the principles stated
3 in the Declaration of Helsinki, after being approved by the Institutional Review Board of our
4 institution (**Luigi Sacco University Hospital, University of Milan, Milan, Italy**). Authors of the
5 original English version of the TVQ^{MtF} were informed about our proposal in conducting the present
6 study, aimed at producing the validated Italian version of this PRO instrument. The study was
7 structured as six different phases: I-TVQ^{MtF} item generation (phase 1); participants recruitment
8 (phase 2); **I-TVQ^{MtF} confirmatory factor analysis (phase 3)**; I-TVQ^{MtF} reliability analysis (phase
9 4); I-TVQ^{MtF} validity analysis (phase 5); I-TVQ^{MtF} correlation analysis (phase 6). In order to
10 guarantee appropriate conclusions about the psychometric properties of the I-TVQ^{MtF}, the
11 Consensus-based Standards for the Selection of Health Measurement Instruments (COSMIN)
12 checklist [31] was followed.

13 14 *2.1 Phase 1: I-TVQ^{MtF} item generation*

15 A cross-cultural adaptation process was performed following standard techniques [32, 33] and also
16 in accordance with the World Health Organization (WHO) recommendations for the process of
17 translation and adaptation of instruments [34]. Items of the original TVQ^{MtF} questionnaire were first
18 translated into Italian by two independent bilingual speech and language pathologists (SLPs)
19 familiar with **transgender** patients care and with experience in translation; moreover, both SLPs
20 were instructed to produce a conceptual translation rather than a literal one (*stage 1: forward*
21 *translation*). Subsequently, a bilingual expert panel (two independent phoniatricians familiar with
22 the process of instrument validation and the SLPs involved in the first stage) examined idiomatic,
23 semantic and conceptual issues of the two translations, in order to further refine them. An Italian
24 final-consensus version was therefore obtained (*stage 2: synthesis*) and given to two independent
25 professional translators, who had no knowledge of the questionnaire, to have it translated back into
26 English (*stage 3: back translation*). Once this task was completed, the two translators and an expert

1 committee reviewed all reports in order to assess conceptual and cultural equivalence and to
2 produce a pre-final version of the instrument (*stage 4: expert committee review*). Finally, twenty-
3 five **transgender women** (mean age 45.4 years, SD \pm 9.81 years, age range 19-63 years, **who had**
4 **been at least 1 year in transition**) were enrolled in a pilot study (*stage 5: pretesting*). **Specifically,**
5 **participants of the pretesting stage were consecutively recruited at Otolaryngology and**
6 **Infectious Diseases outpatient clinics of our institution.** Each **trans woman** autonomously filled
7 out the pre-final version of the Italian TVQ^{MtF} and then discussed meaning, relevance and clarity of
8 each item together with the two phoniatricians and the two SLPs. Afterwards, in order to improve
9 the readability of the questionnaire, the wording of each item of the pre-final version of the Italian
10 TVQ^{MtF} was refined according to the suggestions given by the participants of the pretesting stage.
11 This revision process led to the final version of the Italian TVQ^{MtF} (I-TVQ^{MtF}; see Appendix).

12

13 2.2 Phase 2: Participants recruitment

14 For the present study, 153 **transgender women** (mean age 39.59 years; standard deviation [SD] \pm
15 10.50 years; age range 18-64 years) were consecutively recruited between October 2018 and July
16 2019. **All participants came from regions in the north of Italy. Moreover, none of these**
17 **subjects participated in the pre-testing phase.** Inclusion criteria were as follows: age of 18 years
18 or older; male sex at birth; living in the female role sometimes, often or always; good understanding
19 of Italian; preserved reading skills. Exclusion criteria included: female sex at birth; history of
20 pathologies of the larynx in the previous 6 months; history of head and neck malignancies.
21 Recruitment strategies included: (a) direct recruitment of **transgender women** referred to our
22 institution (71 subjects, 46.41%); (b) advertisement of the study in Facebook groups of Italian
23 lesbian, gay, bisexual and transgender (LGBT) associations, publishing the link to the web version
24 of the study form (82 subjects, 53.59%). **Moreover, transgender women** who heard about the
25 ongoing project from other participants were also allowed to volunteer for this research. All
26 participants were informed about the objectives of the study and gave their consent in written form

1 or online, depending on the recruitment modality. First, all subjects were asked to fill in the I-
2 TVQ^{MtF}. Moreover, social and demographic data were collected for each participant through a
3 dedicated series of questions assessing the following variables: age, education, occupation,
4 relationship status, children. Additionally, the following data regarding the transition process were
5 collected: psychological counselling, frequency and total duration of life in the female role,
6 hormonal therapy, gender-affirming surgery (GAS), voice feminization procedures (speech therapy,
7 surgery). The relevance of the abovementioned questions was discussed together with a team of
8 national and international clinicians with at least a decade of experience in **transgender** patients
9 care.

10

11 ***2.3 Phase 3: I-TVQ^{MtF} Confirmatory Factor Analysis (CFA)***

12 **To be consistent with the results from Dacakis and colleagues' principal component analysis**
13 **[35], item 22, 26, 27, and 28 were excluded. Two competitive models were evaluated: (1) a**
14 **unidimensional model with the 26 retained items loading on a single global factor; (2) a two-**
15 **factor (VF and SP) model [35]. The following indices were employed to evaluate each model's**
16 **goodness of fit: Satorra–Bentler scaled chi-squared statistic ($SB\chi^2$), root mean square error of**
17 **approximation (RMSEA), comparative fit index (CFI), and standardized root mean square**
18 **residual (SRMR). Model acceptability was evaluated through the following cutoff criteria:**
19 **RMSEA < 0.08, CFI > 0.95 and SRMR < 0.08 [36]. In order to choose between competing**
20 **models, Akaike's information criterion was employed, with lower values indicating the model**
21 **to be preferred; a scaled difference chi-square test [37] further compared nested models' fit.**

22

23 ***2.4 Phase 4: I-TVQ^{MtF} reproducibility analysis***

24 This phase of the study was aimed at evaluating both internal consistency and test-retest reliability
25 of the I-TVQ^{MtF}. **For this purpose, the I-TVQ^{MtF} scores obtained from the enrolled 153**
26 **transgender women were analysed.** Internal consistency assesses the extent to which each item in

1 a factor measures the same underlying construct. It was assessed using Cronbach's alpha (α)
2 coefficient, which measures how much the items of a questionnaire are interrelated as a group.
3 Values of this coefficient may range between 0.0 and 1.0: the higher the value, the stronger the
4 internal consistency of the instrument is. Therefore, Cronbach's alpha estimates between 0.7 and
5 0.9 were taken to indicate acceptable internal consistency [38]. **Out of the 71 transgender women**
6 **directly recruited at our institution, 50 participants (mean age 57.3 years; SD \pm 11.47; age**
7 **range 34–77 years) were randomly selected for test–retest reliability analysis. Specifically, this**
8 **analysis was limited to participants physically enrolled at our institution in order to ensure**
9 **that those who were going to complete the questionnaire a second time didn't keep a copy of**
10 **their first completed questionnaire, to which they could later refer to.** All subjects filled out the
11 I-TVQ^{MtF} twice within a three-week interval. Variations of five days before or after the requested
12 interval between trials were considered tolerable, in compliance with the subjects' needs. This
13 three-week interval was selected since no significant changes were expected to take place within
14 this period. **All participants involved in this step of the study did not undergo any intervention**
15 **between the two assessments, nor access to previous responses was allowed when filling out**
16 **the I-TVQ^{MtF} for the second time. Moreover, during the retest evaluation, items of the I-**
17 **TVQ^{MtF} were presented in a different order, with the purpose of preventing participants from**
18 **recalling previous responses.** Test–retest reliability was assessed through two-way random
19 internal consistency coefficient (ICC), which gives a measure of the temporal stability of answers to
20 the items of an outcome instrument. Its value can range between 0.0 and 1.0: the higher the value,
21 the stronger the temporal stability of the answers is. Values between 0.4 and 0.75 indicate a good
22 correlation, while values above 0.75 indicate an excellent one [39].

23

24 *2.5 Phase 5: I-TVQ^{MtF} validity analysis*

25 The aim of this phase of the study was to assess the degree to which the I-TVQ^{MtF} measures the
26 construct it purports to measure [38]. In particular, we analysed convergent validity. Convergent

1 validity, which is a subtype of construct validity, measures the degree to which two measures of
2 constructs that should be theoretically related, are in fact related. Therefore, similarly to the study
3 conducted by Dacakis and colleagues [15], I-TVQ^{MtF} total scores were correlated with the results of
4 the two extra items assessing self-perception and **aspiration** of voice femininity (SPVF, AVF).
5 These correlations were assessed using Spearman's test. In order to further analyse convergent
6 validity, a subgroup of 40 **transgender women** (mean age 51.6 years; SD ± 12.09 years; age range
7 24-73 years) completed the I-TVQ^{MtF} and the Italian version of the Voice Handicap Index (I-VHI)
8 [40, 41] at the same time. **Specifically, 17 participants (42.5%) were selected from the**
9 **transgender women who were directly recruited at our institution, while 23 participants**
10 **(47.5%) were selected from the subjects who were recruited online.** The I-VHI consists of 30
11 items, each one scoring from 0 to 4 (0 = "no disability"; 1 = "mild disability"; 2 = "moderate
12 disability"; 3 = "severe disability"; 4 = "complete disability"), divided into three subscales: (a)
13 *functional*, relating the impact of dysphonia on common daily activities; (b) *emotional*, evaluating
14 the psychological impact of dysphonia; (c) *physical*, dealing with the self-perception of laryngeal
15 and vocal discomfort. **The VHI was the base to develop the TVQ^{MtF}, but it was properly**
16 **modified to capture relevant situations and experiences of trans-persons; previous authors**
17 **also used the VHI to analyse convergent validity [28, 42], showing strong correlation between**
18 **these two measures. Although the VHI and the TVQ^{MtF} do not have an overlapping theoretical**
19 **construct, the VHI was selected as no other voice-related QOL assessment tools specifically**
20 **developed and validated for transgender subjects are available to date.** The correlation between
21 I-TVQ^{MtF} and I-VHI scores was evaluated using Spearman's test.

22

23 2.6 Phase 6: I-TVQ^{MtF} Correlation analysis

24 In order to assess potential relationships between I-TVQ^{MtF} scores on one side and social,
25 demographic and transition-related variables on the other, a correlation analysis was conducted.
26 Point-biserial correlation was used for binary coded (yes/no) variables (partner, psychological

1 counselling). Spearman's test was used instead for all other variables (age, educational level, job,
2 children, time living as a woman, frequency of living as a woman, hormonal therapy, GAS, name
3 change, civil status change, speech therapy for voice feminization, voice feminization surgery).
4 Correlational strength was considered strong for values above 0.5, moderate for values ranging
5 between 0.3 and 0.5, and weak for values below 0.3 [43].

6

7 *2.7 Statistical analysis*

8 **CFA was conducted using Lisrel ver. 8 [44], while** other statistical tests were performed using the
9 SPSS ver. 24.0 statistical software (SPSS, Inc., Chicago, IL). Kolmogorov–Smirnov test was used
10 to test the normality of the distribution of I-TVQ^{MIF} scores among **transgender women**. Since this
11 test demonstrated that the distribution of the scores among the participants was not normal, non-
12 parametric tests were used when requested. For all statistical comparisons an $\alpha = 0.05$ and a power
13 of 0.80 were used [45].

14

15

1 **3. Results**

2 *3.1 Phase 1: I-TVQ^{MtF} item generation*

3 The cross-cultural adaptation process for the adaptation of the TVQ^{MtF} questionnaire into Italian
4 was conducted, leading to the final version of the I-TVQ^{MtF} (see Appendix).

6 *3.2 Phase 2: Participants recruitment*

7 The time required to fill in the questionnaire and the additional form never exceeded 10 minutes in
8 the group of participants directly recruited in our institution. For the group of participants recruited
9 online, this parameter could not be verified. Moreover, given the peculiar recruitment strategies of
10 this survey study (direct recruitment and online advertisement), the exact number of potential
11 participants who received notification of the study was not available; therefore, the response rate
12 could not be calculated. Social, demographic and transition-related data of the participants are
13 reported in Table 1, Table 2 and Table 3. Eighty-six participants (87.58%, $n = 134$) indicated that
14 they had been living full-time in the female role. Specifically, one hundred and fifteen subjects
15 (75.16%) had been living as women for more than 5 years. Of all participants, one hundred and
16 seventeen (76.47%) were taking hormones, thirty (19.61%) had undergone GAS, twelve (7.84%)
17 had chosen speech-therapy for voice feminization and six (3.92%) had undergone voice
18 feminization surgery. Mean scores for all the items of the questionnaire are reported in Table 4. The
19 mean total score for the I-TVQ was 58.43 ($SD \pm 23.89$; range 30-114). Regarding the two extra
20 items dealing with voice femininity, SPVF had a mean score of 2.96 ($SD \pm 1.15$), while a mean
21 score of 1.63 ($SD \pm 0.82$) was calculated for AVF.

23 *3.3 Phase 3: I-TVQ^{MtF} Confirmatory Factor Analysis*

24 **Satorra-Bentler scaled chi square was employed to account for non-normal data distribution**
25 **resulting from Mardia's multivariate omnibus test ($\chi^2 = 1300.13$; $p < 0.001$). The goodness of**
26 **fit values of the unidimensional and the two-factor models are reported in Table 5. Both**

1 **models showed acceptable to good goodness-of-fit indices, RMSEA score associated with the**
2 **unidimensional model was the sole index slightly below acceptance threshold. When models**
3 **were compared, the two-factor model showed a lower AIC value ($\Delta AIC = 24.24$); further, the**
4 **scaled difference chi-square test was found to be significant ($\Delta SB\chi^2_{(1)} = 89.89$; $p < 0.001$). Both**
5 **these results suggest that the two-factor model fits data better than the unidimensional one.**
6 **The standardized loading estimates for the two-factor model are reported in Table 4.**
7 **Loadings ranged from 0.39 (item 1) to 0.85 (item 6); VF and SP factors were very highly**
8 **correlated ($\rho = 0.96$).**

10 *3.4 Phase 4: I-TVQ^{MtF} reproducibility analysis*

11 The internal consistency of the I-TVQ^{MtF}, assessed using Cronbach's alpha, was satisfactory ($\alpha =$
12 0.912). Also test-retest reliability analysis for the I-TVQ^{MtF} yielded satisfactory results (ICC =
13 0.849, 95% confidence interval, range 0.791-0.898).

15 *3.5 Phase 5: I-TVQ^{MtF} validity analysis*

16 As far as convergent validity of the I-TVQ^{MtF} is concerned, total scores recorded for all participants
17 were compared with the scores of the two extra items assessing voice femininity (SPVF, AVF),
18 This analysis, conducted using Spearman's test, revealed a strong negative correlation between the
19 I-TVQ^{MtF} total score and SPVF ($r = -0.612$; $p = 0.001$), while a weak negative correlation was
20 found between the I-TVQ^{MtF} total score and AVF ($r = -0.311$; $p = 0.001$). For convergent validity
21 analysis, the correlations between I-TVQ^{MtF} and I-VHI total scores in a subgroup of 40 participants
22 were analyzed using Spearman's test. The mean I-VHI total score was 34.19 (SD \pm 30.30, range 0-
23 107), while the mean I-TVQ^{MtF} score was 62.62 (SD \pm 21.77, range 30-113). A strong positive
24 correlation between I-TVQ^{MtF} and I-VHI total scores was observed ($r = 0.862$; $p = 0.001$).

26 *3.6 Phase 6: I-TVQ^{MtF} correlation analysis*

1 A correlation analysis was conducted to assess potential association between social, demographic
2 and transition-related variables on one side and I-TVQ^{MtF} total scores on the other. No relevant
3 correlations were found for the following variables: age, educational level, job, partner, children,
4 psychological counselling, hormonal therapy, GAS, name change, civil status change. Interestingly,
5 a weak but significant negative correlation between I-TVQ^{MtF} total score on one side and time spent
6 living in the female role on the other was found ($r = -0.266, p < 0.01$). A weak negative correlation
7 was also highlighted between I-TVQ^{MtF} total score and frequency of living in the female role ($r = -$
8 $0.189, p < 0.05$). Finally, moderate positive correlations were revealed between I-TVQ^{MtF} total
9 score on one side and voice therapy ($r = 0.388, p < 0.01$) and vocal folds surgery ($r = 0.348, p <$
10 0.01) for voice feminization on the other.

11

12

1 **4. Discussion**

2 The present study was designed to produce and validate the Italian version of the TVQ^{MtF}. In order
3 to guarantee the cross-cultural equivalence of the questionnaire and to allow comparisons of results
4 between populations divided by language, the five-step items generation method suggested by
5 Beaton and colleagues [34] and the WHO recommendations for the process of translation and
6 adaptation of instruments [34] were applied thoroughly. Both the experts and the bilingual
7 translators involved in the item generation phase judged the cross-cultural equivalence between the
8 English version of the TVQ^{MtF} and the Italian one as satisfactory, suggesting that the items of the
9 latter had retained the meaning of the original instrument. All study participants directly recruited at
10 our institution ($n = 71$, 46.40%) managed to fill out autonomously the questionnaire and the form
11 assessing social, demographic and transition-related variables in less than 10 minutes. Therefore,
12 even though data regarding the time required to complete the task was not available for participants
13 who were enrolled online, it might be speculated that the I-TVQ^{MtF} can be easily administered with
14 no major comprehension difficulties.

15 The mean I-TVQ^{MtF} total score recorded for the enrolled 153 **transgender women** was 58.43 (SD \pm
16 23.89, range 30-114). In previous reports, the mean baseline total score ranged between 51.55 (SD
17 \pm 18.90, range 30-97) [15] and 71.23 (SD \pm 22.27, range 34-107) [16]. Interestingly, as shown in
18 Table 5, items of the “vocal function” (VF) domain recorded on average higher scores than the
19 items of the “social participation” (SP) domain. A similar trend was also highlighted for previous
20 validation studies, for which mean items scores were published and thus available for comparison
21 (Table 6). This trend suggests that voice impairments in **transgender women** may affect QOL
22 mostly by reason of the intimate connections between vocal functioning and gender identity, rather
23 than the potential impact of those impairments on participation in everyday life. **As far as voice**
24 **feminization treatments are concerned, very few participants underwent voice therapy ($n =$**
25 **12; 7.84%) and voice surgery ($n = 6$; 3.92%). Therefore, as this subpopulation is too small, no**
26 **assumptions can be made regarding treatment satisfaction. Future studies should investigate**

1 **patient satisfaction with voice feminization therapies, in order to identify the most effective**
2 **strategies and to better select the right option for each transgender patient, in a perspective of**
3 **“patient-tailored” gender-affirming voice therapy (either towards feminization or**
4 **masculinization).**

5 The psychometric properties of the I-TVQ^{MtF} were analyzed following the COSMIN checklist. The
6 results revealed good internal consistency, test-retest reliability and validity, supporting the
7 adoption of the I-TVQ^{MtF} for the self-assessment of voice-related QOL in **transgender women,**
8 **both in clinical practice and in outcome research. In terms of CFA, both the unidimensional and**
9 **two-factor models were found to adequately fit the data. Results from model comparison,**
10 **however, showed that the two-factor model – the one including more freely estimated**
11 **parameters – fits data better than the unidimensional model. All items were found to load on**
12 **the expected factor [35] with standardized loadings well above the 0.40 threshold, apart for**
13 **item number 1 (“People have difficulty hearing me in a noisy room”). The correlation**
14 **between factors was found to be positive and very high; hence, future studies should further**
15 **focus on the level of association between VF and SP to determine the clinical benefit in**
16 **considering these two constructs as unique or as separated components.**

17 The internal consistency of the I-TVQ^{MtF} appeared to be excellent, with an overall Cronbach’s α
18 coefficient value of 0.912 in 153 **transgender women.** This result is similar to the ones obtained by
19 the English ($\alpha = 0.964$) [16], Portuguese ($\alpha = 0.911$) [8], Spanish ($\alpha = 0.976$) [29], French ($\alpha =$
20 0.97) [30] and German ($\alpha = 0.97$) [28] versions of the instrument. As far as the reliability of the I-
21 TVQ^{MtF} is concerned, the results of test-retest analysis, with an ICC of 0.849, support the high
22 stability and reproducibility over time of the Italian version of this PRO instrument. Similar results
23 were highlighted for the English (ICC = 0.979) [16], Spanish (ICC = 0.885) [29] and Portuguese
24 (ICC = 0.957) [8] validated versions. Convergent validity analysis, as done in a previous study by
25 Dacakis and colleagues [15], was conducted comparing I-TVQ^{MtF} total scores with the scores of the
26 two extra items assessing self-perception and **aspiration** of voice femininity (SPVF, AVF).

1 Significant negative correlations were highlighted, in accordance with the findings by the
2 abovementioned research, demonstrating that a lower self-rating of voice femininity was associated
3 with a more negative impact of voice impairments on **transgender women's** QOL. Pasricha [5]
4 explained how **transgender women** may be assisted in being perceived as women in diverse
5 contexts by feeling proud of and comfortable with who they are. Consequently, a more solid self-
6 perception of voice femininity should induce a more confident approach to communication tasks,
7 with a higher chance to be perceived as women and, therefore, reducing the impact of voice
8 impairments on QOL [16]. **Basing on these evidences, it can be surmised that psychological**
9 **counseling might be helpful for transgender women in reducing the negative impact of vocal**
10 **impairments on QOL. Centering counseling on what the patient perceives as voice femininity**
11 **and improving patient's self-awareness – both internally (how the patient sees her own values,**
12 **passions, aspirations, fit with her environment) and externally (how the patient understands**
13 **how other people view her in terms of the same factors) – might strengthen the results of voice**
14 **feminization therapies.** However, further studies are necessary to assess the potential beneficial
15 effect of psychotherapy on voice-related QOL during and after the transition process. **Convergent**
16 **validity was assessed comparing I-TVQ^{MtF} and I-VHI total scores in a subgroup of 40**
17 **participants. A significant strong correlation was highlighted ($r = 0.862$), in accordance with**
18 **the findings by the research group led by Salm ($r = 0.88$) [28], which tested this correlation in**
19 **a group of 108 German transgender women. However, it must be stressed that the VHI has**
20 **been previously described as inappropriate for the evaluation of vocal impairments in**
21 **transgender women [46], not having being specifically designed to address the very specific**
22 **vocal needs of transgender subjects. Moreover, Hancock [42], Salm [28] and the present study**
23 **revealed that the generic VHI might over-rate or under-rate voice-related QOL of**
24 **transgender women. Therefore, except for the purposes of cross-cultural validation studies,**
25 **the recommendation of not including the VHI in voice-related QOL assessment protocols for**
26 **transgender women can be further emphasized.**

1 Findings of the correlation analysis, conducted to verify potential associations between the I-
2 TVQ^{MtF} total score and all the screened variables (social, demographic and transition-related), are
3 noteworthy. In particular, weak but significant negative correlations were demonstrated for the
4 variables “time living in the female role” ($r = -0.266$) and “frequency of living in the female role” (r
5 $= -0.189$). A significant correlation between the total TVQ^{MtF} score and time spent in the female
6 role was also demonstrated by a previous report [28], suggesting that the lower (and the less
7 frequent) the time spent living in the female role, the higher is the subjective voice-related
8 impairment on overall QOL. Remarkably, moderate positive correlations were found for the
9 variables “speech therapy for voice feminization” ($r = 0.388$) and “voice feminization surgery” ($r =$
10 0.348). It might be hypothesized that, since these treatment options are not mandatory, only
11 **transgender** women experiencing significant vocal impairments are prone to resort to speech
12 therapy and surgery for voice feminization.

13 **In conclusion, the I-TVQ^{MtF} appears to be a reliable and valid instrument for the assessment**
14 **of voice-related QOL in transgender women. Research aimed at assessing and improving**
15 **QOL in transgender subject is a compelling need and it should be implemented. In fact, a**
16 **recent systematic review and meta-analysis demonstrated that transgender people generally**
17 **display poor QOL, regardless of the domain investigated [47]. Moreover, transgender people**
18 **appear to report poorer mental health compared to the general population [47]. Specifically, a**
19 **study conducted on a sample of the US transgender population revealed how respondents had**
20 **a high prevalence of clinical depression (44.1%), anxiety (33.2%), and somatization (27.5%);**
21 **moreover, social stigma was positively associated with psychological distress [48]. Therefore,**
22 **the introduction of the I-TVQ^{MtF} as an additional instrument to assess and possibly improve**
23 **QOL in transgender women is highly recommended in everyday clinical practice as well as in**
24 **research settings.**

25 The present study has several limitations. First of all, the number of enrolled subjects is quite small
26 even if in line with previous reports. Thus, the results reported in this paper should be considered as

1 preliminary. Second, a selection bias might be postulated since only 46.4% of individuals were
2 directly recruited at our institution, while the remaining participants were recruited through
3 advertisements in Facebook groups of LGBT associations. For the latter individuals, study
4 participation was possible only if they had access to the Internet and were connected with peers via
5 social media and networks. These aspects may have had an impact on the representativeness of the
6 sample. In addition, it is also possible that the majority of the **transgender women** recruited online
7 who decided to participate in the study were those more concerned about their voice [28]. Third, a
8 responsiveness analysis was not performed and, consequently, no information regarding the
9 sensitivity to changes (for example after voice therapy or surgery) of the I-TVQ^{MtF} is available.
10 Further studies analyzing this aspect are therefore needed. **Fourth, reliability analysis was**
11 **performed using the results obtained from individuals directly recruited at our institution and**
12 **no information regarding the reliability of the online-administered questionnaire is available.**

13

14

15

1 **5. Conclusions**

2 The findings of this preliminary validation study support the reliability and the validity of the I-
3 TVQ^{MtF} for the assessment of voice-related QOL in Italian **transgender women**. The application of
4 this patient-reported outcome instrument in everyday clinical practice and in outcome research is
5 therefore recommended, as it could represent a more rigorous assessment strategy of voice-related
6 needs and impairments in MtF patients. Finally, given the cross-cultural equivalence of the I-
7 TVQ^{MtF}, cross-country and multi-center studies are advisable in order to test the responsiveness of
8 this voice-related PRO instrument.

9

10

1 **6. Acknowledgments**

2 We would like to thank all the participants of the study and all the Italian LGBT associations who
3 kindly supported and advertised the present research.

4

1 **7. References**

- 2 1. Human Rights Campaign, Transgender FAQ. <https://www.hrc.org/resources/transgender-faq#1/>, 2020 (accessed 28 June 2020).
- 3
- 4 2. Gay and Lesbian Alliance Against Defamation (GLAAD), Transgender FAQ. <https://www.glaad.org/transgender/transfaq/>, 2020 (accessed 28 June 2020).
- 5
- 6 3. Shelagh Davies, Transsexual Voice Questionnaire (Male to Female). <http://www.shelaghdavies.com/questionnaire/>, 2020 (accessed 28 June 2020).
- 7
- 8 4. American Psychological Association (APA). A glossary: Defining transgender terms. <https://www.apa.org/monitor/2018/09/ce-corner-glossary/>, 2020 (accessed 7 July 2020).
- 9
- 10 5. N. Pasricha, G. Dacakis, J. Oates. Communicative satisfaction of male-to-female transsexuals. *Logoped. Phoniatr. Vocol.* 33 (2008) 25–34.
- 11
- 12 6. M.P. Gelfer, V.A. Mikos. The relative contributions of speaking fundamental frequency and formant frequencies to sex identification based on isolated vowels. *J. Voice* 19 (2005) 544–554.
- 13
- 14 7. R. King, G. Brown, C. McCrea. Voice parameters that result in identification or misidentification of biological gender in male-to-female transgender veterans. *Int. J. Transgend.* 13 (2011) 117–130.
- 15
- 16
- 17 8. H.H. Santos, A.G. Aguiar, H.E. Baeck, J. Van Borsel. Translation and preliminary evaluation of the Brazilian Portuguese version of the Transgender Voice Questionnaire for male-to-female transsexuals. *Codas* 27 (2015) 89-96.
- 18
- 19
- 20 9. J. Abitbol, P. Abitbol, B. Abitbol. Sex hormones and the female voice. *J. Voice* (1999) 424–446.
- 21
- 22 10. M. Cosyns, J. Van Borsel, K. Wierckx, et al. Voice in female-to-male transsexual persons after long-term androgen therapy. *Laryngoscope* 124 (2014) 1409–1414.
- 23
- 24 11. M. Talaat, AM. Talaat, I. Kelada, et al. Histologic and histochemical study of effects of anabolic steroids on the female larynx. *Ann. Otol. Rhinol. Laryngol.* 96 (1987) 468–471.
- 25

- 1 12. D.P. Buckley, K.L. Dahl, G.C. Cler, C.E. Stepp. Transmasculine Voice Modification: A Case
2 Study. *J. Voice* (2019) S0892-1997(19)30116-X [Online ahead of print].
- 3 13. H. Asscheman, L.J. Gooren. Hormone treatment in transsexuals. In: W.O. Bocking, E.
4 Coleman, eds. *Gender Dysphoria: Interdisciplinary Approaches in Clinical Management*. New
5 York, NY: Haworth Press; 1992.
- 6 14. J. Van Borsel, G. De Cuyper, H. Van de Berghe. Physical appearance and voice in male-to-
7 female transsexuals. *J. Voice* 15 (2001) 570-575.
- 8 15. G. Dacakis, J. Oates, J. Douglas. Associations between the Transsexual Voice Questionnaire
9 (TVQ^{MtF}) and self-report of voice femininity and acoustic voice measures. *Int. J. Lang.*
10 *Commun. Disord.* 52 (2017) 831-838.
- 11 16. G. Dacakis, S. Davies, J.M. Oates, J.M. Douglas, J.R. Johnston. Development and Preliminary
12 Evaluation of the Transsexual Voice Questionnaire for Male-to-Female Transsexuals. *J. Voice*
13 27 (2013) 312-320.
- 14 17. A.B. Hancock, J. Krissinger, K. Owen. Voice Perceptions and Quality of Life of Transgender
15 People. *J. Voice* 25 (2011) 553-558.
- 16 18. S.O. Watt, K.O. Tskhay, N.O. Rule. Masculine Voices Predict Well-Being in Female-to-Male
17 Transgender Individuals. *Arch. Sex. Behav.* 47 (2018) 963-972.
- 18 19. K.H. Mount, S.J. Salmon. Changing the vocal characteristics of a postoperative transgender
19 patient: a longitudinal study. *J. Commun. Disord.* 21 (1988) 229-238.
- 20 20. E. McNeill, J. Wilson, S. Clark, J. Deakin. Perception of voice in the transgender client. *J.*
21 *Voice* 22 (2008) 727-733.
- 22 21. R.C. Bralley, G.L. Bull, C.E. Gove, M.J. Edgerton. Evaluation of the vocal pitch in male
23 transsexuals. *J. Commun. Disord.* 11 (1978) 443-449.
- 24 22. G. Dacakis. Long-term maintenance of fundamental frequency increases in male-to female
25 transsexuals. *J. Voice* 14 (2000) 549-556.

- 1 23. S. Sirin, A. Polat, F. Alioglu. Psychometric Evaluation of Adapted Transsexual Voice
2 Questionnaire for Turkish Trans Male Individuals. *J. Voice* 20 (2020) S0892-1997(19)30601-0.
3 Online ahead of print.
- 4 24. S. Sirin, A. Polat. The association between subjective and objective masculine vocal quality in
5 hormone-naïve trans-male individuals. *ENT Updates* 9 (2019) 219-226.
- 6 25. S. Davies, J.M. Goldberg. Clinical aspects of transgender speech feminization and
7 masculinization. *Int. J. Transgend.* 9 (2006) 167–196.
- 8 26. L. Carew, G. Dacakis, J.M. Oates. The effectiveness of oral resonance therapy on the
9 perception of femininity of voice in male-to-female transsexuals. *J. Voice* 21 (2017) 591–603.
- 10 27. K. Schwarz, A.M.V. Fontanari, A. Mueller, et al. Transsexual Voice Questionnaire for Male-
11 to-female Brazilian Transsexual People. *J. Voice* 31 (2017) 120.e15-120.e20.
- 12 28. S. Salm, K. Hower, S. Neumann, L. Ansmann. Validation of the German Version of the
13 Transsexual Voice Questionnaire for male-to-Female Transsexuals. *J. Voice* 34 (2020) 68-77.
- 14 29. E. Mora, A. Carrillo, A. Giribet, A. Becerra, M.J. Lucio, I. Cobeta. Translation, Cultural
15 Adaptation, and Preliminary Evaluation of the Spanish Version of the Transgender Voice
16 Questionnaire for Male-to-Female Transsexuals (TVQ^{MtF}). *J. Voice* 32 (2018) 514.e1-514.e6.
- 17 30. D. Morsomme, J. Revis, E. Thomas. Translation, Adaptation, and Preliminary Validation of
18 Dacakis and Davies’ “Transsexual Voice Questionnaire (Male to Female)” in French. *J. Voice*
19 33 (2019) 807.e13-807.e24.
- 20 31. L.B. Mokkink, C.B. Terwee, D.L. Patrick. International consensus on taxonomy, terminology,
21 and definitions of measurement properties for health-related patient-reported outcomes: results
22 of the COSMIN study. *J. Clin. Epidemiol.* 63 (2010) 737–745.
- 23 32. D.E. Beaton, C. Bombardier, F. Guillemin, et al. Guidelines for the process of cross-cultural
24 adaptation of self-report measures. *Spine* 25 (2000) 3186– 3191.
- 25 33. P.M. Boynton, T. Greenhalgh. Selecting, designing, and developing your questionnaire. *BMJ*
26 328 (2004) 1312–1315.

- 1 34. World Health Organization (WHO). Process of translation and adaptation of instruments. 2019.
2 Available at http://www.who.int/substance_abuse/research_tools/translation/en/. Accessed on
3 November 4, 2019.
- 4 35. G. Dacakis, J.M. Oates, J.M. Douglas. Further Evidence of the Construct Validity of the
5 Transsexual Voice Questionnaire (TVQMtF) Using Principal Components Analysis. *J Voice*.
6 31 (2017) 142-148.
- 7 36. L. Hu, P.M. Bentler. Evaluating model fit. In: Hoyle RH (ed) *Structural equation modeling:*
8 *concepts, issues, and applications*. Sage Publications, Inc, Thousand Oaks (1995) pp 76–99.
- 9 37. F.B. Bryant, A. Satorra. Principles and practice of scaled difference chi-square testing.
10 *Structural Equation Modeling* 19 (2012) 372-398.
- 11 38. C.B. Terwee, S.D.M. BotM, M.R. de Boer, et al. Quality criteria were proposed for
12 measurement properties of health status questionnaires. *J. Clin. Epidemiol.* 60 (2007) 34–42.
- 13 39. L. Prieto, R. Lamarca, A. Casado. Assessment of the reliability of clinical findings: the
14 intraclass correlation coefficient. *Med. Clin. (Barc)* 110 (1998) 142–145.
- 15 40. B.H. Jacobson, A. Johnson, C. Grywalski, et al. The Voice Handicap Index (VHI):
16 development and validation. *Am. J. Speech. Lang. Pathol.* 6 (1997) 66–70.
- 17 41. A. Schindler, F. Ottaviani, F. Mozzanica, et al. Cross-cultural adaptation and validation of the
18 Voice Handicap Index into Italian. *J. Voice* 24 (2010) 708–714.
- 19 42. A.B. Hancock. An ICF Perspective on Voice-related Quality of Life of American Transgender
20 Women. *J. Voice* 31 (2017) 115.e1-115.e8.
- 21 43. E. Vittinghoff, D.V. Glidden, S.C. Shiboski, et al. *Regression Methods in Biostatistics: Linear,*
22 *Logistic, Survival, and Repeated Measures Models*. Springer, New York, 2005.
- 23 44. K.G. Jöreskog, D. Sörbom. *LISREL 8: User's reference guide*. Lincolnwood: Scientific
24 Software International (2001).
- 25 45. G.M. Sullivan, R. Feinn. Using effect size—or why the P value is not enough. *J. Grad. Med.*
26 *Educ.* 24 (2012) 279–282.

- 1 46. G. T'Sjoen, M. Moerman, J. Van Borsel, et al. Impact of voice in transsexuals. *Int. J.*
2 *Transgender* 9 (2006) 1–7.
- 3 47. A. Nobili, C. Glazebrook, J. Arcelus. Quality of life of treatment-seeking transgender adults: A
4 systematic review and meta-analysis. *Rev. Endocr. Metab. Disord.* 19 (2018) 199–220.
- 5 48. W.O. Bockting, M.H. Miner, R.E. Swinburne Romine, A. Hamilton, E. Coleman. Stigma,
6 Mental Health, and Resilience in an Online Sample of the US Transgender Population. *Am. J.*
7 *Public Health* 103 (2013) 943–951.

8
9
10
11
12
13

- 1 **Conflict of interest:** none
- 2 **Source funding:** nothing to declare
- 3

1 **Table 1.** General demographic characteristics of the enrolled transgender women.

Parameter	Value (N = 153)
Age ± SD (years)	39.59 ± 10.14
Age range (years)	18 – 64
Education (n, %)	No school 5 (3.27%) Elementary school 13 (8.50%) Middle school 23 (15.03%) High school 72 (47.06%) University 33 (21.57%) Master/PhD 7 (4.58%)
Job (n, %)	No job 50 (32.68%) Part-time job 49 (32.03%) Full-time job 54 (35.29%)
Partner (n, %)	Yes 63 (41.18%) No 90 (58.82%)
Children (n, %)	No 136 (88.89%) Yes, but we're not in touch 0 (0.0%) Yes, and we're in touch 17 (11.11%)
Psychological counselling (n, %)	Yes 23 (15.03%) No 130 (84.97%)

2
3
4

1
2

Table 2. Transition-related characteristics of the enrolled transgender women.

Parameter	Value (N = 153)	
Time living in the female role (n, %)	Less than 1 year	12 (7.84%)
	Between 1 and 5 years	26 (16.99%)
	More than 5 years	115 (75.16%)
Frequency of living in the female role (n, %)	Rarely	0 (0.00%)
	Sometimes	0 (0.00%)
	Often	19 (12.42%)
	Always	134 (87.58%)
Hormonal therapy (n, %)	No	36 (23.53%)
	Yes, less than 1 year	10 (6.54%)
	Yes, between 1 and 5 years	22 (14.38%)
GAS ¹ (n, %)	Yes, more than 5 years	85 (55.56%)
	No, not planned	65 (42.48%)
	No, but considering it	44 (28.76%)
	No, but planned	14 (9.15%)
Name change (n, %)	Yes	30 (19.61%)
	No	68 (44.44%)
	Yes, but not legally	40 (26.14%)
Civil status change (n, %)	Yes, legally	45 (29.41%)
	No	105 (68.63%)
	Yes, but not legally	9 (5.88%)
	Yes, legally	39 (25.49%)

¹GAS = Gender-affirming surgery

3
4
5
6

1
2

Table 3. History of voice feminization treatments (speech and language therapy, surgery).

Parameter	Value (<i>N</i> = 153)	
Speech and language therapy for voice feminization (<i>n</i> , %) (<i>N</i> = 153)	No, not planned	70 (45.75%)
	No, but considering it	54 (35.29%)
	No, but planned	17 (11.11%)
	Yes	12 (7.84%)
Speech and language therapy for voice feminization: satisfaction (<i>n</i> , %) (<i>N</i> = 12)	Not satisfied	2 (16.67%)
	Little satisfied	7 (58.33%)
	Very satisfied	1 (8.33%)
	Extremely satisfied	2 (16.67%)
Voice feminization surgery (<i>n</i> , %) (<i>N</i> = 153)	No, not planned	87 (58.86%)
	No, but considering it	53 (34.64%)
	No, but planned	7 (4.58%)
	Yes	6 (3.92%)
Voice feminization surgery: satisfaction (<i>n</i> , %) (<i>N</i> = 6)	Not satisfied	0 (0.0%)
	Little satisfied	3 (50.0%)
	Very satisfied	0 (0.0%)
	Extremely satisfied	3 (50.0%)

3
4

1
2
3
4

Table 4. I-TVQ^{MtF} and voice femininity mean values and standard deviations and Confirmatory Factor Analysis standardized factor loadings (two-factor model). I-TVQ^{MtF} items are listed in descending order of mean score.

I-TVQ ^{MtF}					
N°	VF ¹	SP ²	Item	Mean	SD
3	0.78		My voice makes me feel less feminine than I would like	2.32	1.06
20	0.78		My voice doesn't match my physical appearance	2.32	1.18
24	0.82		I feel my voice does not reflect the "true me"	2.24	1.25
18	0.74		When I am not paying attention, my pitch goes down	2.22	1.11
28			It distresses me when I'm perceived as a man because of my voice	2.20	1.23
15	0.65		I have to concentrate to make my voice sound the way I want it to sound	2.14	1.13
26			I feel self-conscious about how strangers perceive my voice	2.12	1.09
4	0.78		The pitch of my speaking voice is too low	2.10	1.09
5	0.57		The pitch of my voice is unreliable	2.10	0.94
6		0.85	My voice gets in the way of me living as a woman	2.06	1.13
2		0.73	I feel anxious when I know I have to use my voice	2.01	1.02
19	0.76		When I laugh I sound like a man	2.01	1.10
16	0.81		I feel frustrated with trying to change my voice	1.99	1.10
11	0.72		When I speak the pitch of my voice does not vary enough	1.94	0.89
22			My voice gets tired quickly	1.94	1.09
29	0.66		The pitch range of my speaking voice is restricted	1.92	1.01
8		0.79	I'm tense when talking to others because of my voice	1.91	0.99
10	0.82		My voice makes it hard for me to be identified as a woman	1.90	1.02
25		0.82	I am less outgoing because of my voice	1.87	1.06
9	0.72		My voice gets croaky, hoarse or husky when I try to speak in a female voice	1.83	1.05
13		0.80	I avoid speaking in public because of my voice	1.83	1.08
21	0.74		I use a great deal of effort to produce my voice	1.83	0.98
12		0.81	I feel uncomfortable talking to friends neighbors and relatives because of my voice	1.80	1.04
17		0.79	My voice difficulties restrict my social life	1.77	1.04
30		0.82	I feel discriminated against because of my voice	1.74	1.05
7		0.70	I avoid using the phone because of my voice	1.71	0.97
23		0.77	My voice restricts the sort of work I do	1.69	1.08
1		0.39	People have difficulty hearing me in a noisy room	1.67	0.83
14		0.77	My voice sounds artificial	1.65	0.91
27			My voice "gives out" in the middle of speaking	1.58	0.89
I-TVQ^{MtF} VF score				28.87	11.55
I-TVQ^{MtF} SP score				21.71	9.69
I-TVQ^{MtF} Total Score				58.43	23.89
Voice Femininity					
Item				Mean	SD
SPVF ³ Currently, my voice is				2.96	1.15
AVF ⁴ My ideal voice would sound				1.63	0.82

¹VF = Vocal Functioning domain; ²SP = Social participation domain; ³SPVF = Self-perception of voice femininity; ⁴AVF = Aspiration of voice femininity

5
6
7

1 **Table 5.** Confirmatory Factor Analysis goodness of fit indices ($N = 153$).

Model	SB χ^2	df	RMSEA	CFI	SRMR	AIC
Unidimensional	599.98*	299	.081	.98	.058	703.98
Two Factors	573.74*	298	.078	.98	.057	679.74

2 Note. SB χ^2 = Satorra-Bentler Scaled Chi-Square; df = Degree of Freedom; RMSEA = Root Mean Square Error of
3 Approximation; CFI = Comparative Fit Index; SRMR = Standard Root Mean Square Residual; AIC = Akaike's
4 Information Criterion. * $p < .001$.

5

1
2
3
4

Table 6. Comparison of mean items scores and standard deviations between different validated versions of the TVQ^{MtF} (Italian, French, English). TVQ^{MtF} items are listed in descending order of mean score.

Italian TVQ ^{MtF} (Robotti et al, 2020)					French TVQ ^{MtF} (Morsomme et al, 2019)					English (original) TVQ ^{MtF} (Dacakis et al, 2013)				
N°	VF ¹	SP ²	Mean	SD	N°	VF ¹	SP ²	Mean	SD	N°	VF ¹	SP ²	Mean	SD
3	■		2.32	1.06	15	■		2.89	1.14	28			3.09	1.20
20	■		2.32	1.18	4	■		2.72	0.88	15	■		3.09	1.17
24	■		2.24	1.25	18	■		2.67	1.01	4	■		2.98	1.03
18	■		2.22	1.11	28			2.67	1.22	26			2.89	1.11
28			2.20	1.23	20	■		2.53	1.13	24	■		2.87	1.25
15	■		2.14	1.13	3	■		2.50	1.13	3	■		2.83	1.07
26			2.12	1.09	11	■		2.44	0.97	16	■		2.64	1.10
4	■		2.10	1.09	24	■		2.42	1.11	20	■		2.71	1.18
5	■		2.10	0.94	19	■		2.31	1.12	18	■		2.86	1.00
6		■	2.06	1.13	1		■	2.28	0.88	29	■		2.80	1.02
2		■	2.01	1.02	29	■		2.28	0.88	21	■		2.54	1.12
19	■		2.01	1.10	5	■		2.25	0.77	10	■		2.51	1.10
16	■		1.99	1.10	21	■		2.22	1.10	11	■		2.49	0.95
11	■		1.94	0.89	2		■	2.19	1.14	19	■		2.49	1.17
22			1.94	1.09	10	■		2.19	0.98	5	■		2.46	0.92
29	■		1.92	1.01	26			2.19	1.12	1		■	2.29	0.96
8		■	1.91	0.99	7		■	2.17	1.13	22			2.29	1.07
10	■		1.90	1.02	22			2.17	1.03	2		■	2.17	0.99
25		■	1.87	1.06	16	■		2.14	1.15	13		■	2.09	1.07
9	■		1.83	1.05	27			2.06	0.86	14		■	2.09	0.98
13		■	1.83	1.08	8		■	2.03	0.94	8		■	1.91	0.92
21	■		1.83	0.98	13		■	2.03	1.16	27			1.85	0.81
12		■	1.80	1.04	9	■		2.00	0.86	25		■	2.26	1.22
17		■	1.77	1.04	6		■	1.89	1.09	6		■	2.09	1.31
30		■	1.74	1.05	23		■	1.89	1.19	17		■	2.00	1.11
7		■	1.71	0.97	17		■	1.86	1.10	30		■	1.97	1.18
23		■	1.69	1.08	25		■	1.81	1.09	9	■		1.94	1.00
1		■	1.67	0.83	30	■		1.72	1.03	23	■	■	1.79	0.96
14		■	1.65	0.91	12		■	1.47	0.81	12		■	1.63	0.88
27			1.58	0.89	14		■	1.47	0.77	7		■	1.63	0.77

5
6

¹VF = Vocal Functioning domain; ²SP = Social participation domain

1 Appendix

2

3 **Italian Transsexual Voice Questionnaire for male-to-female Transsexuals (I-TVQ^{MtF})**

4

5 In base alla tua esperienza di vita nel ruolo femminile, rispondi alle seguenti domande segnando la
6 risposta che descrive al meglio la tua condizione attuale. Per favore, fornisci una risposta per ciascuna
7 delle domande riportate. Considera inoltre la seguente legenda:

8

9 1 – “mai” o “molto raramente”

10 2 – “a volte”

11 3 – “spesso”

12 4 – “frequentemente” o “sempre”

	1	2	3
	4		
1. Le persone hanno difficoltà a sentire la mia voce in un ambiente rumoroso.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
2. Provo ansia quando so di dover utilizzare la mia voce.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
3. La mia voce mi fa sentire meno femminile di quanto vorrei.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
4. Il tono della mia voce parlata è troppo grave.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
5. Il tono della mia voce è imprevedibile.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
6. La mia voce rappresenta un ostacolo per vivere come donna.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
7. Evito di utilizzare il telefono a causa della mia voce.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
8. Mi sento tesa nel parlare con altre persone a causa della mia voce.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
9. La mia voce è gracchiante o rauca quando provo a parlare con una voce femminile.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
10. La mia voce mi crea difficoltà nel farmi identificare come donna.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
11. Quando parlo il tono della mia voce non varia abbastanza.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
12. Mi sento a disagio quando parlo con amici, vicini e familiari a causa della mia voce.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
13. Evito di parlare in pubblico a causa della mia voce.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
14. La mia voce suona come se fosse artificiale.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
15. Devo concentrarmi per far sì che la mia voce suoni come vorrei.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
16. Mi sento frustrata quando cerco di cambiare la mia voce.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
17. Le difficoltà connesse alla mia voce limitano la mia vita sociale.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
18. Quando non presto attenzione, il tono della mia voce diventa più grave.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
19. Quando rido, la mia voce suona come quella di un uomo.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		
20. La mia voce non rispecchia il mio aspetto fisico.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>		

21.	Devo utilizzare un grande sforzo per produrre la mia voce.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
22.	La mia voce si affatica rapidamente.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
23.	La mia voce limita la tipologia di lavoro che posso svolgere.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
24.	Credo che la mia voce non rifletta il mio "vero io".	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
25.	Sono meno estroversa a causa della mia voce.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
26.	Mi sento a disagio per come gli estranei percepiscono la mia voce.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
27.	La mia voce si esaurisce a metà della conversazione.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
28.	Provo angoscia quando vengo percepita come uomo a causa della mia voce.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
29.	L'estensione vocale della mia voce parlata è limitata.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
30.	Mi sento discriminata a causa della mia voce.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18

Per favore, formula ora un giudizio globale sulla tua voce:

(SPVF) Attualmente la mia voce è:

molto in parte neutra in parte

molto

femminile femminile maschile

maschile

(AVF) La mia voce ideale sarebbe:

molto in parte neutra in parte

molto

femminile femminile maschile

maschile

Vitae

- **Carlo Robotti.** Born in Voghera (Italy) in 1987. Medical Degree in 2014. Specialization in Otorhinolaryngology in 2019. Research Fellow in Otorhinolaryngology at University of Pavia (Pavia, Italy) since 2020.
- **Francesco Mozzanica.** Born in Milan (Italy) in 1981. Medical Degree in 2007. Specialization in Otorhinolaryngology in 2014. PhD in Nutritional Sciences in 2018. Assistant Professor of Otorhinolaryngology at University of Milan (Milan, Italy) since 2019.
- **Chiara Atzori.** Born in Milan (Italy) in 1964. Medical Degree in 1989. Specialization in Infectious Diseases in 1993.

- 1 • **Andrea Cavalot.** Born in Turin (Italy) in 1960. Medical Degree in 1986. Specialization in
2 Otorhinolaryngology in 1990.
- 3 • **Diego Cossu.** Born in Genova (Italy) in 1961. Medical Degree and Specialization in
4 Audiology and Phoniatics.
- 5 • **Adi Primov-Fever.** Born in Tel Aviv (Israel) in 1968. Medical Degree in 1995.
6 Specialization in Otorhinolaryngology in 2005. Laryngology Fellowship in 2006. Lecturer
7 of Otorhinolaryngology at Sackler Faculty of Medicine, Tel Aviv University (Tel Aviv,
8 Israel) since 2014.
- 9 • **Marco Benazzo.** Born in Pavia (Italy) in 1959. Medical Degree in 1984. Specialization in
10 Otorhinolaryngology in 1987. Specialization in Audiology in 1991. Associate Professor of
11 Otorhinolaryngology at University of Pavia (Pavia, Italy) from 2000 to 2010. Full Professor
12 of Otorhinolaryngology at University of Pavia (Pavia, Italy) since 2011.
- 13 • **Luca Negri.** Born in Chiaravalle (Italy) in 1979. Psychology degree in 2007. Ph.D in
14 Neuropsychobiology in 2013. Specialization in Cognitive Psychotherapy in 2013. Post-doc
15 Research Fellow at University of Milan. (Milan, Italy)
- 16 • **Antonio Schindler.** Born in Turin (Italy) in 1972. Medical degree in 1997. Specialization in
17 Audiology and Phoniatics in 2001. Specialization Otorhinolaryngology in 2005. Associate
18 Professor of Audiology at University of Milan (Milan, Italy) since 2015.