

## **Bread-making performance of durum wheat as affected by sprouting**

Gaetano Cardone, Silvia Grassi, Anna Scipioni, Alessandra Marti\*

Department of Food, Environmental, and Nutritional Sciences (DeFENS). Università degli Studi di

Milano, via G. Celoria 2, 20133 Milan, Italy

\*Corresponding author:

E-mail address: [alessandra.marti@unimi.it](mailto:alessandra.marti@unimi.it)

address: via G. Celoria 2, 20133 Milan, Italy.

## **Abstract**

The effects of sprouting duration (24 h, 38 h, 48 h, and 62 h) were assessed on durum wheat kernel characteristics (hardness, test weight), semolina chemical composition, pasting and gluten aggregation properties, and leavening and bread-making performance (bread volume and crumb porosity). Sprouting decreased both kernel hardness (~29 %) and test weight (~19 %). Starch gelatinization and retrogradation capability, as well as the gluten aggregation properties, decreased as sprouting duration increased. The 62 h sample showed the worst aggregation properties leading to a bread with the lowest specific volume (2.69 mL/g). The best results in terms of bread specific volume (3.08 mL/g) and crumb porosity distribution were obtained using semolina from sprouted wheat up to 38 h. A multivariate approach by Principal Component Analysis and clustering confirmed the relationships between all the considered variables and allowed to assess three sprouting levels: 24-38 h with improved bread-making performance; 48 h with decreased overall quality; 62 h with the worst quality. In conclusion, the sprouting of durum wheat up to 38 h could improve its bread-making attitude.

**Keywords:** semolina; germination; pasting properties; gluten functionality; bread

**Abbreviations:**  $A_0$ , radial area of the dough at the beginning of the leavening; A-am,  $\alpha$ -amylase activity; AgEn, Aggregation Energy;  $A_t$ , radial area of the dough at time t; BD, Breakdown index; CTRL: unsprouted durum wheat; DS, Damaged Starch; FV, Final Viscosity; Glu, D-glucose; GPE, GlutoPeak Equivalent; GPU, GlutoPeak Unit; Mal, Maltose; MT, Maximum Torque; PCA, Principal Component Analysis; PMT, Peak Maximum Time; Prot, Protein; PV, Peak Viscosity; SpV, Specific Volume; Suc, Sucrose; TS, Total Starch; V, bread volume.

## 1 **1. Introduction**

2 Durum wheat (*Triticum turgidum* subsp. *durum*) is characterized by a peculiar hard and vitreous  
3 endosperm which influences its milling behavior, e.g., milling energy, yield and the starch damage  
4 (Turnbull & Rahman, 2002). The strength and poor extensibility of its gluten network makes durum  
5 wheat the ideal raw material for pasta-making but unsuitable for baked-goods (Ammar, Kronstad, &  
6 Morris, 2000). Despite the enhanced nutritional traits thanks to the carotenoids (Pasqualone, Caponio,  
7 & Simeone, 2004), using durum wheat in bread-making results in low loaf volume and dense crumb  
8 structure (Sissons, 2008). However, dough extensibility and bread volume improved using sourdough  
9 fermentation, since the combination of acidity and hydrolytic activity of both lactic acid bacteria and  
10 yeasts positively affect durum wheat gluten functionality (Barber, Ortolá, Barber, & Fernández,  
11 1992). Considering the above, this study investigated the exploitation of the enzymatic pattern  
12 developed throughout sprouting to improve the bread-making performance of durum wheat.  
13 Although, an excessive accumulation of enzymes in wheat has always represented a negative event  
14 from a technological standpoint, recently it has been reported that sprouting improved the bread-  
15 making performance of common wheat (Cardone, D’Incecco, Pagani, & Marti, 2020a; Marti,  
16 Cardone, Nicolodi, Quaglia, & Pagani, 2017; Marti, Cardone, Pagani, & Casiraghi, 2018). In the case  
17 of durum wheat, the sprouting process have been recently investigated in relation to bioactive  
18 compounds (Jribi et al., 2019a) and functional properties (Jribi, Sahagùn, Debbabi, & Gomez, 2019b)  
19 of wholemeal semolina. To the best of our knowledge, no study has focused yet on the relationship  
20 between sprouting and bread-making performance of durum wheat. Since the understanding of flour  
21 functionality is a key element in the production of cereal-based products, the aim of this study was to  
22 evaluate the effects of sprouting duration on durum wheat kernel characteristics, starch and gluten  
23 behavior, and their relationship with the bread characteristics also from a multivariate point of view,  
24 thus applying Principal Component Analysis and clustering.

## 25 **2. Materials and methods**

## 26 2.1. Sample preparation

27 Five aliquots (1 kg each) of durum wheat (*Triticum durum* Desf.), supplied by Molino Quaglia S.p.A.  
28 (Vighizzolo d'Este, Italy), were sprouted at 20° C for 24 h, 38 h, 48 h and 62 h and dried at 50° C for  
29 9 h, as previously reported by Grassi et al. (2018). Unsprouted durum wheat was used as control  
30 (CTRL). Unsprouted and sprouted samples were conditioned until they reached 165 g/kg of water  
31 content and milled into refined semolina using a laboratory mill (RM1300, Erkaya, Ankara, Turkey),  
32 equipped with a 250 µm sieve.

## 33 2.2. Kernel hardness and test weight

34 Kernel hardness was assessed by NIR (6500, Foss, Hilleroed, Denmark) following the AACC method  
35 39-70.02 (AACCI 2011). Test weight was determined with a Grain Analysis Computer (2100b,  
36 DICKEY-john, Auburn, USA).

## 37 2.3. Chemical composition and $\alpha$ -amylase activity

38 Total and damaged starch content were evaluated according to AACC methods (76-13.01 and 76-  
39 31.01, respectively; AACCI 2001). Simple sugars were quantified by means of the  
40 Maltose/Sucrose/D-Glucose Assay kit commercialized by Megazyme (Wicklow, Ireland). Protein  
41 content was quantified by following the ISO method 20483:2006 (ISO, 2006).  $\alpha$ -amylase activity  
42 was determined according to the AACC method 22-02.01 (AACCI 2001). All the measurements were  
43 carried out in triplicate.

## 44 2.4. Pasting properties

45 Starch pasting properties were evaluated in duplicate by using the Rapid Viscoanalyzer® (4500,  
46 Perten Instrument, Stockholm, Sweden) according to the AACC method 76–21.01 (AACCI 2001) in  
47 presence of either water or silver nitrate (AgNO<sub>3</sub>; 0.001mol/L) as enzymatic inhibitor.

## 48 2.5. Gluten aggregation properties

49 Gluten aggregation kinetic was assessed in triplicate by using the GlutoPeak® (Brabender  
50 GmbH&Co., Duisburg, Germany) device, according to Suárez-Estrella et al. (2020).

## 51 2.6. Dough preparation and leavening properties

52 Semolina was kneaded with fresh yeast (30 g/kg semolina; Carrefour, Milan, Italy) and salt (15 g/kg  
53 semolina; Candor®, Com-Sal s.r.l., Pesaro, Italy) in an automatic mixer equipped with a spiral hook  
54 (KitchenAid 5KSM125EER, Whirlpool, St. Paul, USA) for 6 min, until a smooth and non-sticky  
55 dough was obtained. The amount of water used in the formulations has been added on the basis on  
56 preliminary farinographic tests. Specifically, 645 g/kg of water was added to CTRL and 24 h sample,  
57 605 g/kg of water for 38 h and 48 h samples and, finally, 585 g/kg of water for 62 h sample. Three  
58 portions (5 g each) of the resulted doughs were molded in a spherical shape and then placed in three  
59 Petri dishes, and subjected to leavening at 30° C. The Petri dishes were scanned at 300 dpi with a  
60 flatbed scanner (Epson Perfection 550 Photo, Seiko-Epson, Suwa, Japan) at the beginning of the test,  
61 and after 15 min, 30 min, 45 min, 60 min, 90 min, 120 min and 180 min. The radial increase of the  
62 dough area (mm<sup>2</sup>) was determined by image analysis using the Image Pro Plus software v. 6.0 (Media  
63 Cybernetics, Inc., Rockville, USA) and it was used to determine the relative increase of dough surface  
64 ( $A_t/A_{t0}$ ), through the ratio between the area at time t ( $A_t$ ) and the area of the dough at the  
65 beginning of the test ( $A_{t0}$ ), according to (Caramanico et al., 2018).

## 66 2.7. Micro-baking test

67 Dough samples were obtained as reported in the previous paragraph. Samples were shaped, left to  
68 rise (90 min at 30° C) and baked (20 min at 200° C) as reported by Cardone et al. (2020a). The  
69 obtained loaves were characterized 2 h after baking. One baking test was performed for each sample  
70 and two loaves were obtained.

## 71 2.8. Bread properties

72 Each loaf was characterized for specific volume (SpV) through the ratio between the bread volume,  
73 evaluated by seed replacement method (AACC 10-05.01; AACCI, 2001) and the bread weight.

74 Crumb porosity was assessed on three slices from each loaf as described by Marti et al. (2017) with  
75 some modifications about pore dimensional classes (i.e. < 0.09 mm<sup>2</sup>; 0.10-0.99 mm<sup>2</sup>; 1.00-2.99 mm<sup>2</sup>;  
76 3.00-9.99 mm<sup>2</sup>; > 10.00 mm<sup>2</sup>). Crumb yellowness was evaluated on three points of three central slices  
77 from each loaf by means of digital colorimeter (Digital Color Meter, Apple Inc., Cupertino, USA).

## 78 2.9. Statistical analysis

79 Data were elaborated by a paired t-Test ( $\alpha=0.05$ ) through the software StatPlus:mac (v.7.3.31,  
80 (Analystsoft, Inc., Walnut, USA), to compare differences between mean for unsprouted (CTRL) and  
81 each sprouted sample for different duration for each parameter. Moreover, a type of homoscedastic  
82 or heteroscedastic t-Test was selected according to whether the variance of the pair of the tested  
83 samples was equal or different, respectively. In order to provide the precision of the measurements,  
84 for the parameters in which the variance of the samples was comparable, the pooled SD (i.e. the  
85 square-root of a pooled variance estimator) was calculated. Data were also explored by Principal  
86 Component Analysis (PCA) after data mean centering by means of Matlab software (v. 2016a,  
87 Mathworks, Inc., Natick, USA). Samples grouping was confirmed by K-Nearest Neighbor cluster  
88 analysis (PLS toolbox, v. 8.5, Eigenvector Research, Inc., Manson, USA).

## 89 3. Results

### 90 3.1. Kernel characteristics

91 The sprouting process caused a significant decrease in both kernel hardness (from 112 to 78 after 24  
92 h of sprouting) and test weight (from 80 kg/hL to 69 kg/hL after 24 h of sprouting).

### 93 3.2 Chemical composition and $\alpha$ -amylase activity

94 Sprouting did not affect the starch content, instead the damaged starch fraction statistically  
95 ( $p=9.01*10^{-5}$ ) increased after 38 h of sprouting (Table 1). As the damaged starch increased also  
96 simple sugars statistically increased; in particular, maltose increased ( $p=4.47*10^{-4}$ ) after 24 h, instead  
97 sucrose ( $p=3.44*10^{-2}$ ) after 38 h, and glucose ( $p=4.61*10^{-2}$ ) after 48 h of sprouting (Table 1).  $\alpha$ -

98 amylase activity statistically ( $p=1.60*10^{-4}$ ) increased by about 260 folds, already after 24 h of  
99 sprouting (Table 1).

100 Sprouting duration did not strongly affect the protein content of semolina, which decreased by about  
101 6% (Table 1).

### 102 3.2. Pasting properties

103 Regardless the sprouting duration, in presence of water, sprouted samples showed low viscosity  
104 values ( $< 0.1 \text{ Pa}\cdot\text{s}$ ), in both heating and cooling stages (data not shown). Inhibiting the amylase  
105 activity with a solution of silver nitrate ( $\text{AgNO}_3$ ;  $0.001 \text{ mol/L}$ ) all samples showed a higher viscosity,  
106 indicating that the pasting and gelation properties of sprouted samples were not drastically affected  
107 by sprouting (Figure 1a). Specifically, the peak viscosity ( $1.866\pm 0.008 \text{ Pa}\cdot\text{s}$  for CTRL,  $1.58\pm 0.02$ ,  
108  $1.34\pm 0.04$ ,  $1.1755\pm 0.0007$  and  $1.156\pm 0.008 \text{ Pa}\cdot\text{s}$  for 24 h, 38 h, 48 h and 62 h, respectively) and the  
109 breakdown index (i.e. resistance of the gel to mechanical stress) ( $0.44\pm 0.01 \text{ Pa}\cdot\text{s}$  for CTRL,  
110  $0.39\pm 0.03$ ,  $0.33\pm 0.04$ ,  $0.275\pm 0.006$  and  $0.36\pm 0.04 \text{ Pa}\cdot\text{s}$  for 24 h, 38 h, 48 h and 62 h, respectively)  
111 significantly decreased after 24 h ( $p=3.46*10^{-2}$ ) and 48 h ( $p=3.38*10^{-2}$ ) of sprouting, respectively.  
112 Moreover, the final viscosity and the setback index (i.e. the tendency of starch to retrograde)  
113 statistically ( $p=4.34*10^{-2}$ ) decreased as sprouting duration increased, starting from 24 h of sprouting  
114 ( $2.92\pm 0.04 \text{ Pa}\cdot\text{s}$  for CTRL,  $2.471\pm 0.008$ ,  $2.163\pm 0.002$ ,  $1.95\pm 0.01$  and  $1.71\pm 0.08 \text{ Pa}\cdot\text{s}$  for 24 h, 38 h,  
115 48 h and 62 h, respectively).

### 116 3.3. Gluten aggregation properties

117 As regards changes in gluten aggregation kinetics (Figure 1b), sprouting led to a significant  
118 ( $p=1.21*10^{-3}$ ) increase in the peak maximum time starting from 38 h of sprouting ( $60\pm 2 \text{ s}$  for CTRL,  
119  $62\pm 3$ ,  $83\pm 2$ ,  $77\pm 2$  and  $98\pm 6 \text{ s}$  for 24 h, 38 h, 48 h and 62 h, respectively), and a significant decrease  
120 in both maximum torque ( $p=3.34*10^{-4}$ ) ( $47.0\pm 0.8 \text{ GPU}$  for CTRL,  $31.8\pm 0.9$ ,  $26.4\pm 0.1$ ,  $24.2\pm 0.9$  and  
121  $20.5\pm 0.7 \text{ GPU}$  for 24 h, 38 h, 48 h and 62 h, respectively) and aggregation energy ( $p=4.48*10^{-2}$ ) (i.e.  
122 energy required for gluten aggregation;  $1239\pm 47 \text{ GPE}$  for CTRL,  $887\pm 15$ ,  $758\pm 9$ ,  $694\pm 21$  and

123 592±15 GPE for 24 h, 38 h, 48 h and 62 h, respectively), already after 24 h and 38 h of sprouting,  
124 respectively.

### 125 3.4. Dough leavening properties

126 Dough leavening properties were evaluated by monitoring changes in radial area. CTRL reached the  
127 maximum development in 45 min ( $A_{t45}/A_{t0}=2.3$ ) and no longer increased up to 120 min of  
128 leavening; after that, it decreased ( $A_{t180}/A_{t0}=2.0$ ) (Figure 2). In contrast, the radial area of  
129 sprouted wheat dough constantly increased until the end of the test period ( $A_{t180}/A_{t0}=2.7$ ) (Figure  
130 2). The fastest area expansion was observed after 24 h and 36 h of sprouting, subsequent to leavening  
131 for 15 min.

### 132 3.5. Bread-making properties

133 Using sprouted wheat did not lead to a drastic worsening of bread properties, in terms of volume, not  
134 even after 62 h of sprouting (178±4, 173±4, 180±1, 180±1 and 178±4 mL for CTRL, 24 h, 38 h, 48  
135 h and 62 h, respectively). Samples from 38 h sprouted wheat showed the best bread-making  
136 performances, in terms of specific volume (Figure 3). Instead, 62 h sprouted sample showed the worst  
137 crumb structure, that appeared sticky and irregular (Figure 3). As regards crumb yellowness, loaves  
138 from sprouted wheat showed a more intense yellowness (Figure 3).

139 No significant differences ( $p>0.05$ ) were observed among the samples in terms of number of cells  
140 (data not shown). Unlike that, differences were observed in cell area (Table 2). Specifically, CTRL  
141 bread showed a crumb characterized by about 70 % of small cells (< 1 mm<sup>2</sup>), instead this pore class  
142 represented about 50 % of the total in loaves from sprouted wheat. Moreover, large pores (> 10 mm<sup>2</sup>)  
143 were found only in bread from sprouted wheat whose area accounted for the 10 % of the total for 24  
144 h bread, instead about 5 % for 38 h and 48 h loaves.

### 145 3.6 PCA and cluster analysis

146 PCA results showed sample distribution according to chemical composition,  $\alpha$ -amylase activity,  
147 dough leavening properties and bread-making properties. The scores plot defined by the first PCs



148 described almost the 83 % of the data variability (PC1=55.87 %; PC2=27.11 %) and showed a clear  
149 separation of CTRL samples from sprouted samples (Figure 4a). Indeed, CTRL samples assumed  
150 highly positive PC1 and PC2 values, being in the I quadrant of the plot. 24 h sprouted sample is  
151 located in the IV quarter, assuming the lowest PC2 value; 38 h sprouted samples was well separated  
152 in the III quarter; finally, 48 h and 62 h samples were grouped in the II quarter. Scores vs time  
153 representation (Figure 4b) enabled to highlight that PC1 described an unique process as the scores  
154 values decreased with time progress, whereas PC2 trajectory was characterized by a sudden decrease  
155 in the first 24 h followed by an increment of the scores after 38 h and a consecutive decrement in the  
156 last sampling time. In order to uncover the variables responsible for sample grouping the loadings  
157 plot was presented (Figure 4c). Most of the chemical indexes and  $\alpha$ -amylase drove the separation of  
158 CTRL sample from sprouted samples along PC1, together with gluten aggregation properties;  
159 whereas leavening properties and bread characteristics resulted relevant in the discrimination among  
160 samples subjected to different sprouting duration (24 h, 38 h, 48 h and 62 h).

161 The explorative data analysis showed a sample distribution according to the sprouting duration  
162 (Figure 4c), envisioning the possibility of defining sprouting classes according to the considered  
163 parameter. However, the confirmation of sample grouping according to sprouting duration needed  
164 more solid bases, thus a cluster analysis was performed. The cluster analysis based on K-Nearest  
165 Neighbor algorithm identified four clusters based on the whole results collected. From the  
166 dendrogram (Figure 4d), the first cluster, i.e. the group that differed the most from the others, was the  
167 one formed by CTRL which resulted highly different (distance = 7) from the sprouted samples, no  
168 matter the sprouting duration. By reducing the distance to 5, the analysis individuated three sprouting  
169 levels: a cluster consisting of 24 h and 38 h sprouted samples and other two separated clusters for 48  
170 h and 62 h sprouted samples.

#### 171 **4. Discussion**

172 Compared to common wheat, durum wheat is characterized by high kernel hardness, high gluten  
173 tenacity and intensive yellowness – due to its high carotenoid content. All these characteristics are  
174 used to evaluate the grain quality on the market. As regards the kernel characteristics, sprouting  
175 process led to a significant decrease in hardness (Figure S1), with the greatest changes occurring at  
176 48 h sprouting duration (Figure S1). The decrease in kernel hardness might positively affect the  
177 milling behavior. Indeed, hard kernels, such as durum wheat, require more energy to be milled than  
178 both soft and hard kernels (Różyło et al., 2003). Specifically, the decrease in kernel hardness might  
179 be attributed to the decrease in starch-protein matrix density in the endosperm. This hypothesis has  
180 been confirmed by the decrease in test weight (i.e. index related to the kernel density; Figure S1) due  
181 to the high  $\alpha$ -amylase activity associated with sprouting (Table 1). The effect of enzymatic activity  
182 on decreasing the endosperm density as a consequence of sprouting has been recently shown in  
183 sprouted common wheat (Cardone, D’Incecco, Casiraghi, & Marti, 2020b). Moreover, the decrease  
184 in kernel hardness and test weight were in line with previous study carried out on sprouted common  
185 wheat (Miś & Grundas, 2002; Różyło, Laskowski, & Grundas, 2003). However, both the indices  
186 seemed not to be affected by the sprouting duration (Figure S1).

187 In addition to milling energy, hardness also affects the milling yield and the damaged starch content  
188 of flours (Turnbull & Rahman, 2002). In this study, the milling yield did not appear to be affected by  
189 the sprouting duration within 48 h, ranging from 49 g/100 g for CTRL, to 48, 46, 47 and 38 g/100 g  
190 for 24 h, 38 h, 48 h and 62 h, respectively (data not shown). The low yield ratio obtained could be  
191 due to the use of a laboratory mill that allowed to extract mainly the innermost regions of the  
192 endosperm, at the expenses of the yield. The decrease in milling yield might be related to the decrease  
193 in test weight (Figure S1), with evidence at prolonged sprouting durations. Indeed, after 62 h the  
194 rootlet was quite evident (Figure S1), suggesting an intense hydrolysis of the storage macromolecules,  
195 as confirmed by the increased  $\alpha$ -amylase activity. It is generally recognized that the sprouting process  
196 is considered concluded when the rootlet reached the kernel length, in order to avoid strongly negative  
197 effects on the kernel properties and flour functionality (Marti, Cardone, & Pagani, 2020). During

198 sprouting, high levels of hydrolytic enzymes – specifically  $\alpha$ -amylases – are released and create some  
199 holes on the surface of the starch granules (Cardone et al., 2020a; Faltermaier, Zarnkow, Becker,  
200 Gastl, & Arendt, 2015), making them more accessible to a further enzymatic action. Thus, the level  
201 of damaged starch (which is defined as the amount of starch readily accessible to  $\alpha$ -amylase) might  
202 provide information about the intensity of the sprouting process. In general, high damaged starch  
203 content adversely affects the dough handling (e.g. greater water absorption and dough stickiness) and  
204 the bread characteristics (e.g. lower development in volume and darker crust color) (Sapirstein,  
205 David, Preston, & Dexter, 2007). Under the condition applied in this study, the damaged starch  
206 content increased as the sprouting duration increased too (Table 1), as an effect of the increased  $\alpha$ -  
207 amylase activity (Table 1), rather than exclusively as mechanical damage of the starch granules  
208 during milling. These findings were confirmed by the multivariate exploration by PCA, indeed  
209 damaged starch and  $\alpha$ -amylase activity were close to each other and located in the II quarter of the  
210 loadings plot (Figure 4c) affecting the separation of samples sprouted 48 h and 62 h from lower  
211 germination exposure (24 h and 38 h) and CTRL (Figure 4a), thus driving the separation of these  
212 samples along PC1 according to sprouting duration (Figure 4b)

213 Sprouting resulted in lower pasting and gelation properties (Figure 1a), because of the lower  
214 gelatinization and retrogradation ability of the smaller starch polymers accumulated in sprouted  
215 samples than CTRL. These changes were in line with other studies on sprouted durum (Jribi et al.,  
216 2019b) and common (Cardone et al., 2020a; Grassi et al., 2018) wheat and also remarked by the PCA  
217 loadings plot (Figure 4c), in which the pasting and gelation indexes calculated from the analysis  
218 performed in presence of water or silver nitrate assumed positive PC1 scores, thus separating the  
219 CTRL from the sprouted samples (Figure 4a). Furthermore, the lower ability to retrograde of the  
220 sprouted samples might have led to obtain a fresh bread with a softer crumb, compared to the CTRL  
221 one, as shown in common wheat (Cardone et al., 2020a,b) and quinoa-enriched bread (Suárez-Estrella  
222 et al., 2020).

223 As regards the proteins, the decrease (Table 1) might be attributable to their hydrolysis into soluble  
224 peptides due to the proteolytic activity (Mbithi-Mwikya, Ooghe, Van Camp, Ngundi, & Huyghebaert,  
225 2000). On the other hand, it is reported that changes in protein content less than 10 % indicates that  
226 the sprouting process did not significantly affect the protein content of grains (Lemmens et al., 2019).  
227 Similar changes are reported in previous studies on sprouted durum (Jribi et al., 2019a) and common  
228 (Cardone et al., 2020a; Grassi et al., 2018; Koehler, Hartmann, Wieser, & Rychlik, 2007; Marti et al.,  
229 2017) wheat.

230 Moving to gluten properties, the sprouting duration negatively affected the aggregation properties of  
231 the gluten-forming proteins (Figure 1b), in terms of peak maximum time (increased by ~63 % after  
232 62 h of sprouting), maximum torque (decreased by ~56 % after 62 h of sprouting) and aggregation  
233 energy (decreased by ~52 % after 62 h of sprouting), suggesting a weakening of the gluten network  
234 (Grassi et al., 2018; Marti, Augst, Cox, & Koehler, 2015a), as a consequence of the proteolytic  
235 activity. In general, flour with good bread-making performance are characterized by a faster  
236 aggregation (i.e., low peak maximum time) and higher maximum torque compared to those with poor  
237 bread-making attitude (Quayson, Atwell, Morris, & Marti, 2016). Actually, the aggregation  
238 properties of the gluten-forming proteins resulted the ones most affecting the separation between  
239 CTRL and the highly sprouted samples along the PC1 of the PCA scores plot (Figure 4a), being the  
240 peak maximum time highly negative and maximum torque and aggregation energy highly positive.  
241 A possible explanation of the maximum torque and the peak maximum time shifts is that sprouting  
242 induced changes in the profile of gluten proteins (i.e. gliadin and glutenin fractions) (Koehler et al.,  
243 2007). Indeed, Marti et al. (2015b) found a positive correlation between maximum torque and gliadin  
244 content and between energy and glutenin with high molecular weight. In particular, it is already  
245 reported that sprouting caused a significant degradation of glutenins, already after 48 h of sprouting,  
246 instead longer duration was required for degrading gliadins, about 102 h (Koehler et al., 2007).  
247 Although the sprouted samples showed a different gluten aggregation profiles that would suggest  
248 gluten weakening, they were still able to aggregate and form a gluten network with good performance

249 in bread-making (Figure 1b), confirming previous studies on common wheat (Cardone et al., 2020a;  
250 Marti et al., 2018). The only exception was the 62 h sample that lost its ability to form gluten (Figure  
251 1b), likely due to the stronger intensity of the sprouting process (Figure S1; Table 1).

252 In comparison with common wheat, durum wheat is characterized by a very stiff and not very  
253 extensible gluten, making it suitable for the pasta-making but unsuitable for leavened baked-goods  
254 (Ammar et al. 2000). Indeed, the resulting bread will be characterized by a high density and a hard  
255 texture (Sissons, 2008). The interest in durum wheat bread lies in the fact that this raw material is  
256 richer in carotenoids (i.e. antioxidant compounds) compared to common wheat. Generally, to  
257 overcome the negative technological properties (i.e., low volume and high crumb density) of durum  
258 wheat bread, sourdough fermentation is used as leavening agent. Indeed, the low pH and the  
259 enzymatic activities of lactic bacteria and yeasts enhanced bread-making performance, in terms of  
260 bread volume (Barber et al., 1992; Pagani, Lucisano & Mariotti, 2014). In this context, the increased  
261 enzymatic activity developed during sprouting process might represent a good strategy to improve  
262 the bread-making attitude of durum wheat.

263 Thanks to the correlations between dough tenacity and strength and maximum torque and aggregation  
264 energy (Marti et al., 2015b; Rakita, Dokić, Dapčević Hadnađev, Hadnađev, & Torbica, 2018), it is  
265 possible to hypothesize that sprouting could represent a good way to decrease dough tenacity and  
266 consequently improve its bread-making performance. Despite the gluten weakening (Figure 1b), the  
267 dough from sprouted durum wheat was able to withstand the leavening stresses, expanding itself  
268 without collapsing (Figure 2). The increased CO<sub>2</sub> production during leavening - thanks to the  
269 increased amount of fermentable sugars by yeasts, resulting from the  $\alpha$ -amylase activity (Table 1) –  
270 increased loaf specific volume, mainly for the 38 h sample (Figure 3 and Figure 4a). Similar results  
271 are reported for common wheat (Cardone et al., 2020a; Marti et al., 2018). The worsening of crumb  
272 structure in bread from 62 h sprouted wheat (Figure 3) agreed with the excessive gluten weakening  
273 (Figure 1b). Indeed, the poor gluten aggregation properties and its gas retention capacity resulted in  
274 the lowest specific volume (Figure 3). As regards the pore distribution, large pores (>10 mm<sup>2</sup>) were

275 found only in bread from sprouted wheat, probably due to the coalescence of the gas cells, favored  
276 by  $\alpha$ -amylase activity (Lagrain, Leman, Goesart, & Delcour, 2008). In addition, bread from sprouted  
277 wheat resulted in a higher crumb yellowness (Figure 3), following a similar trend of the yellow index  
278 of semolina (from  $19\pm 1$  for CTRL to  $25.4\pm 0.8$  after 62h; data not shown). Yang et al. (2001) report  
279 that the  $\beta$ -carotene content increased upon sprouting and the color intensity of the carotenoid extract  
280 increased as the sprouting duration increased too. Although this aspect needs to be further  
281 investigated, finding suggests that sprouting process might have a positive effect on the carotenoid  
282 content in bread from sprouted durum wheat.

283 All the considered chemical composition,  $\alpha$ -amylase activity, dough leavening properties and bread-  
284 making properties do not act separately but are interconnected and correlated. Thus, the multivariate  
285 approach led us to confirm the relationships between all the considered variables and to define which  
286 of them contributed most in the sample distribution, i.e. in assessing the sprouting influence in the  
287 final product, as Grassi et al. (2018) speculate. Indeed, the dendrogram obtained by the cluster  
288 analysis (Figure 4d) confirmed that samples sprouted up to 38 h had similar and improved bread-  
289 making performance. The two distinct clusters for 48 h and 62 h sprouted samples (Figure 4d)  
290 indicated a progressive and significant decrease of the overall quality.

## 291 **5. Conclusions**

292 Changes induced by sprouting strongly depended on the process duration. Specifically, sprouting  
293 under controlled conditions (i.e., up to 48 h) did not strongly compromise the functional properties  
294 of starch (i.e., gelatinization and retrogradation phenomena). As regards proteins, despite the  
295 sprouting process weakened the gluten network, gluten proteins were still able to aggregate and retain  
296 gas during leavening, resulting in bread with improved volume. Specifically, the best bread-making  
297 performance were achieved using durum wheat that was sprouted for 38 h.

298 Overall results suggest that sprouting carried out under controlled conditions could improve the  
299 bread-making attitude of durum wheat and produce a more attractive product (i.e. improved bread

300 volume and crumb porosity) for the consumer and with high carotenoid content compared to common  
301 bread. However, the effects of sprouting process on gliadin and glutenin fractions need to be studied  
302 in depth, as well as the potential application of the process on various durum wheat varieties.

303 **Acknowledgments.** The authors wish to thank M.A. Pagani (formerly Università degli Studi di  
304 Milano, Italy) for the critical discussion during the manuscript preparation.

305 **Declarations of interest:** none

## 306 **References**

307 AACC Approved Methods of Analysis, 11th Ed. Cereals & Grains Association, St. Paul, MN, USA.

308 Ammar, K., Kronstad, W. E., & Morris, C. F. (2000). Breadmaking quality of selected durum wheat  
309 genotypes and its relationship with high molecular weight glutenin subunits allelic variation and  
310 gluten protein polymeric composition. *Cereal Chemistry*, 77, 230–236.

311 Barber, B., Ortolá, C., Barber, S., & Fernández, F. (1992). III. Effects of sour dough and addition of  
312 acids on bread characteristics. *Zeitschrift Für Lebensmittel-Untersuchung Und - For - Schung*,  
313 194, 442–449.

314 Caramanico, R., Marti, A., Vaccino, P., Bottega, G., Cappa, C., Lucisano, M., & Pagani, M. A.  
315 (2018). Rheological properties and baking performance of new waxy lines: Strengths and  
316 weaknesses. *LWT - Food Science and Technology*, 88, 159-164.

317 Cardone, G., D’Incecco, P., Casiraghi, M. C., & Marti, A. (2020b). Exploiting milling by-products  
318 in bread-making: the case of sprouted wheat. *Foods*, 9, 260.

319 Cardone, G., D’Incecco, P., Pagani, M. A., & Marti, A. (2020a). Sprouting improves the bread-  
320 making performance of whole wheat flour (*Triticum aestivum* L.). *Journal of the Science of Food*  
321 *and Agriculture*, 1000, 2453-2459.

322 Faltermaier, A., Zarnkow, M., Becker, T., Gastl, M., & Arendt, E. K. (2015). Common wheat  
323 (*Triticum aestivum* L.): evaluating microstructural changes during the malting process by using

324 confocal laser scanning microscopy and scanning electron microscopy. *European Food Research*  
325 *and Technology*, 241, 239-252.

326 Grassi, S., Cardone, G., Bigagnoli, D., & Marti, A. (2018). Monitoring the sprouting process of wheat  
327 by non-conventional approaches. *Journal of Cereal Science*, 83, 180–187.

328 ISO 20483. (2006). Cereals and pulses–Determination of the nitrogen content and calculation of the  
329 crude protein content–Kjeldahl method. *International Organization for Standardization*,  
330 Geneva, Switzerland.

331 Jribi, S., Molnar, H., Adanyi, N., Marzougui, S., Naar, Z. & Debbabi, H. (2019b). Effect of sprouting  
332 temperature on durum wheat (*Triticum Durum*) sprouts nutritional properties and bioactive  
333 compounds. *International Journal of Innovative Approaches in Agricultural Research*, 3, 87-95.

334 Jribi, S., Sahagùn, M., Debbabi, H., & Gomez, M. (2019a). Evolution of functional, thermal and  
335 pasting properties of sprouted whole durum wheat flour with sprouting time. *International*  
336 *Journal of Food Science & Technology*, 54, 2718–2724.

337 Koehler, P., Hartmann, G., Wieser, H., & Rychlik, M. (2007). Changes of folates, dietary fiber, and  
338 proteins in wheat as affected by germination. *Journal of Agricultural and Food Chemistry*, 55,  
339 4678–4683.

340 Lagrain, B., Leman, P., Goesart, H., & Delcour, J. A. (2008). Impact of thermostable amylases  
341 during bread making on wheat bread crumb structure and texture. *Food Research International*,  
342 41, 819–827.

343 Lemmens, E., Moroni, A. V., Pagand, J., Heirbaut, P., Ritala, A., Karlen, Y., ..., & Delcour, J. A.  
344 (2019). Impact of cereal seed sprouting on its nutritional and technological properties: a critical  
345 review. *Comprehensive Reviews in Food Science and Food Safety*, 18, 305–328.

346 Marti, A., Augst, E., Cox, S., & Koehler, P. (2015a). Correlations between gluten aggregation  
347 properties defined by the GlutoPeak test and content of quality-related protein fractions of winter  
348 wheat flour. *Journal of Cereal Science*, 66, 89–95.



- 349 Marti, A., Cardone, G., Nicolodi, A., Quaglia, L., & Pagani, M. A. (2017). Sprouted wheat as an  
350 alternative to conventional flour improvers in bread-making. *LWT - Food Science and*  
351 *Technology*, 80, 230–236.
- 352 Marti, A., Cardone, G., & Pagani, M. A. (2020). Sprouted cereal grains and products. In *Innovative*  
353 *Processing Technologies for Healthy Grains*. Submitted
- 354 Marti, A., Cardone, G., Pagani, M. A., & Casiraghi, M. C. (2018). Flour from sprouted wheat as a  
355 new ingredient in bread-making. *LWT - Food Science and Technology*, 89, 237–243.
- 356 Marti, A., Ulrici, A., Foca, G., Quaglia, L., & Pagani, M. A. (2015b). Characterization of common  
357 wheat flours (*Triticum aestivum* L.) through multivariate analysis of conventional rheological  
358 parameters and gluten peak test indices. *LWT - Food Science and Technology*, 64, 95–103.
- 359 Mbithi-Mwikya, S., Ooghe, W., Van Camp, J., Ngundi, D., & Huyghebaert, A. (2000). Amino acid  
360 profiles after sprouting, autoclaving, and lactic acid fermentation of finger millet (*Eleusine*  
361 *coracana*) and kidney beans (*Phaseolus vulgaris* L.). *Journal of Agricultural and Food Chemistry*,  
362 48, 3081-3085.
- 363 Miś, A., & Grundas, S. (2002). Wheat grain hardness modified by the laboratory sprouting test.  
364 *International Agrophysics*, 16, 283–288.
- 365 Pagani, M.A., Lucisano, M., & Mariotti, M. (2014). Italian bakery products. In W. Zhou, Y. H. Hui,  
366 I. De Leyn, M. A. Pagani, C. M. Rosell, J. D. Selman, & N. Therdthai (Eds.), *Bakery Products*  
367 *Science and Technology* (pp. 685-721). Hoboken: John Wiley & Sons, Ltd.
- 368 Pasqualone, A., Caponio, F., & Simeone, R. (2004). Quality evaluation of re-milled durum wheat  
369 semolinas used for bread-making in Southern Italy. *European Food Research and Technology*,  
370 219, 630–634.
- 371 Quayson, E. T., Atwell, W., Morris, C. F., & Marti, A. (2016). Empirical rheology and pasting  
372 properties of soft-textured durum wheat (*Triticum turgidum* ssp. *durum*) and hard-textured  
373 common wheat (*T. aestivum*). *Journal of Cereal Science*, 69, 252–258.

- 374 Rakita, S., Dokić, L., Dapčević Hadnađev, T., Hadnađev, M., & Torbica, A. (2018). Predicting  
375 rheological behavior and baking quality of wheat flour using a GlutoPeak test. *Journal of Texture*  
376 *Studies*, 49, 339–347.
- 377 Różyło, R., Laskowski, J., & Grundas, S. (2003). Study of how protein content in wheat grain relates  
378 to hardness index and grinding energy. *Acta Agrophysica*, 2, 173–178.
- 379 Sapirstein, H. D., David, P., Preston, K. R., & Dexter, J. E. (2007). Durum wheat breadmaking  
380 quality: effects of gluten strength, protein composition, semolina particle size and fermentation  
381 time. *Journal of Cereal Science*, 45, 150–161.
- 382 Sissons, M. (2008). Role of durum wheat composition on the quality of pasta and bread. *Food*, 2,  
383 75–90.
- 384 Suárez-Estrella, D., Cardone, G., Buratti, S., Pagani, M. A., & Marti, A. (2020). Sprouting as a pre-  
385 processing for producing quinoa-enriched bread. *Submitted*
- 386 Yang, T. K., Basu, B., & Ooraikul, F. (2001). Studies on germination conditions and antioxidant  
387 contents of wheat grain. *International Journal of Food Sciences and Nutrition*, 52, 319-330.

## Figure captions

Figure 1. Rapid Viscoanalyzer (in presence of silver nitrate - AgNO<sub>3</sub>; 0.001 mol/L) (a) and GlutoPeak (b) profiles of semolina from unsprouted (CTRL) and sprouted durum wheat. CTRL: solid line; 24 h: dotted line; 38 h: short dash line; 48 h: dash-dot-dot line; 62 h: long dash line. 24 h, 38 h, 48 h, 62 h: sprouting duration; CTRL: unsprouted durum wheat; GPU: GlutoPeak Units.

Figure 2. Increasing the radial area ( $A_t/A_{t0}$ ) of the dough during leavening. CTRL: dash-line; 24 h: black square; 38 h: grey circle; 48 h: black triangle; 62 h: grey diamond.

Asterisk indicates a significant difference between CTRL and each sample from sprouted wheat (paired t-Test;  $\alpha=0.05$ ;  $n=3$ ). n.s.: not significant differences. 24 h, 38 h, 48 h, 62 h: sprouting duration;  $A_{t0}$ , radial area of the dough at the beginning of the leavening;  $A_t$ , radial area of the dough at time t; CTRL: unsprouted durum wheat.

Figure 3. Pictures of the bread loaves, crumb yellowness and specific volume (SpV) of bread prepared from semolina from unsprouted (CTRL) and sprouted durum wheat. Asterisk indicates a significant difference between CTRL and each bread sample from sprouted wheat (paired t-Test;  $\alpha=0.05$ ;  $n=5$  for crumb yellowness;  $n=2$  for specific volume). Scale bar is 1 cm. 24 h, 38 h, 48 h, 62 h: sprouting duration; CTRL: unsprouted durum wheat.

Figure 4. Multivariate data analysis on data collected for chemical composition,  $\alpha$ -amylase activity, dough leavening properties and bread-making properties: scores plot for Principal Component Analysis (a), scores vs sprouting duration plot (b), loadings plot (c), dendrogram for cluster analysis by K-Nearest Neighbor (d)

A-am,  $\alpha$ -amylase activity; TS, Total Starch; DS, Damaged Starch; Mal, Maltose; Suc, Sucrose; Glu, D-glucose; Prot, Protein. Pasting properties: PV, Peak Viscosity; BD, Breakdown index; FV, Final Viscosity. Gluten aggregation properties: PMT, Peak Maximum Time; MT, Maximum Torque; AgEn, Aggregation Energy. Leavening properties: relative increase of dough surface at 15 min

(A\_t15), 30 min (A\_t30), 45 min (A\_t45), 60 min (A\_t60), 90 min (A\_t90), 120 min (A\_t120) and 180 min (A\_t180). Bread characteristics: SpV, Specific Volume; V, Bread Volume.

Figure S1. Kernel hardness and test weight of durum wheat kernels during sprouting process, from 24 h to 62 h




Figure 1.




Figure 2.




Figure 3.




Figure 4.



Table 1. Chemical characteristics (starch, simple sugar and protein contents) and  $\alpha$ -amylase activity of semolina from unsprouted (CTRL) and sprouted durum wheat at different sprouting duration (24 h, 38 h, 48 h and 62 h).

	<b>CTRL</b>	<b>24 h</b>	<b>38 h</b>	<b>48 h</b>	<b>62 h</b>	<b>Pooled SD</b>
Total starch	71	71 <sub>ns</sub>	72 <sub>ns</sub>	71 <sub>ns</sub>	69 <sub>ns</sub>	1
Damaged starch	10.3	9.7*	13.2*	13.4*	15.9*	0.3
Maltose	0.3	2.1*	4.7*	5.4*	6.6*	0.4
Sucrose	1.5	1.9 <sub>ns</sub>	2.0*	2.0*	2.1*	0.2
D-glucose	0.20	0.16 <sub>ns</sub>	0.3 <sub>ns</sub>	0.41*	0.42*	0.2
Protein	14.18	14.11 <sub>ns</sub>	13.80 <sub>ns</sub>	13.88*	13.32*	0.03
$\alpha$ -amylase activity	0.089±0.004	3.8±0.3*	9.9±0.5*	21.6±0.9*	24.3±0.2*	-

Chemical data are expressed as g/100 g sample (dry basis). Damaged starch is expressed as g/100 g of total starch (dry basis).  $\alpha$ -amylase activity is expressed as Ceralpha Units/g flour (dry basis). Asterisk indicates a significant difference between CTRL and each sprouted sample (paired t-Test;  $\alpha=0.05$ ; n=3). CTRL: unsprouted durum wheat; 24 h, 38 h, 48 h, 62 h: sprouting duration; ns: not significant difference.

Table 2. Area occupied by each pore dimensional class of the bread crumb (%).

<b>Dimensional classes (mm<sup>2</sup>)</b>	<b>CTRL</b>	<b>24 h</b>	<b>38 h</b>	<b>48 h</b>	<b>Pooled SD</b>
< 0.09	8.9	8.7 <sub>ns</sub>	9.3 <sub>ns</sub>	7.8 <sub>ns</sub>	0.7
0.10 – 0.99	59	42*	47*	43*	2
1.00 – 2.99	26	25 <sub>ns</sub>	23 <sub>ns</sub>	28 <sub>ns</sub>	4
3.00 – 9.99	8	14 <sub>ns</sub>	14 <sub>ns</sub>	17 <sub>ns</sub>	3
> 10.00	-	7	10	4	2

Asterisk indicates a significant difference between CTRL and each sprouted sample (paired t-Test;  $\alpha=0.05$ ;  $n=3$ ). CTRL: unsprouted durum wheat; 24 h, 38 h, 48 h: sprouting duration; ns: not significant difference.




Figure S1. Kernel hardness and hectolitre weight of durum wheat kernels during sprouting process, from 24 h to 62 h.

Asterisk indicates a significant difference between CTRL and each bread sample from sprouted wheat (paired t-Test;  $\alpha=0.05$ ;  $n=2$ ).

**Highlights:**

- Sprouting of durum wheat decreased the kernel hardness and hectolitre weight
- Starch and gluten properties were not strongly affected up to 48 h of sprouting
- Sprouting process improved dough leavening attitude of durum wheat
- Using sprouted durum wheat up to 38 h addressed to the highest bread specific volume