



Cardioprotective effect of succinate dehydrogenase inhibition in rat hearts and human myocardium with and without diabetes mellitus

Jespersen, Nichlas Riise; Hjortbak, Marie Vognstoff; Lassen, Thomas Ravn; Støttrup, Nicolaj Brejnholt; Johnsen, Jacob; Tonnesen, Pernille Tilma; Larsen, Steen; Kimose, Hans Henrik; Bøtker, Hans Erik

Published in:
Scientific Reports

DOI:
[10.1038/s41598-020-67247-4](https://doi.org/10.1038/s41598-020-67247-4)

Publication date:
2020

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY](https://creativecommons.org/licenses/by/4.0/)

Citation for published version (APA):
Jespersen, N. R., Hjortbak, M. V., Lassen, T. R., Støttrup, N. B., Johnsen, J., Tonnesen, P. T., ... Bøtker, H. E. (2020). Cardioprotective effect of succinate dehydrogenase inhibition in rat hearts and human myocardium with and without diabetes mellitus. *Scientific Reports*, 10(1), [10344]. <https://doi.org/10.1038/s41598-020-67247-4>



OPEN

Cardioprotective effect of succinate dehydrogenase inhibition in rat hearts and human myocardium with and without diabetes mellitus

Nichlas Riise Jespersen¹✉, Marie Vognstoft Hjortbak¹, Thomas Ravn Lassen¹, Nicolaj Brejnholt Støttrup¹, Jacob Johnsen¹, Pernille Tilma Tonnesen¹, Steen Larsen^{2,3}, Hans-Henrik Kimose⁴ & Hans Erik Bøtker¹

Ischemia reperfusion (IR) injury may be attenuated through succinate dehydrogenase (SDH) inhibition by dimethyl malonate (DiMAL). Whether SDH inhibition yields protection in diabetic individuals and translates into human cardiac tissue remain unknown. In isolated perfused hearts from 24 weeks old male Zucker diabetic fatty (ZDF) and age matched non-diabetic control rats and atrial trabeculae from patients with and without diabetes, we compared infarct size, contractile force recovery and mitochondrial function. The cardioprotective effect of a 10 minutes DiMAL administration prior to global ischemia and ischemic preconditioning (IPC) was evaluated. In non-diabetic hearts exposed to IR, DiMAL 0.1 mM reduced infarct size compared to IR ($55 \pm 7\%$ vs. $69 \pm 6\%$, $p < 0.05$). Mitochondrial respiration was reduced by DiMAL 0.6 mM compared to sham and DiMAL 0.1 mM ($p < 0.05$). In diabetic hearts an increased concentration of DiMAL (0.6 mM) was required for protection compared to IR ($64 \pm 13\%$ vs. $79 \pm 8\%$, $p < 0.05$). Mitochondrial function remained unchanged. In trabeculae from humans without diabetes, IPC and DiMAL improved contractile force recovery compared to IR ($43 \pm 12\%$ and $43 \pm 13\%$ vs. $23 \pm 13\%$, $p < 0.05$) but in patients with diabetes only IPC provided protection compared to IR ($51 \pm 15\%$ vs. $21 \pm 8\%$, $p < 0.05$). Neither IPC nor DiMAL modulated mitochondrial respiration in patients. Cardioprotection by SDH inhibition is possible in human tissue, but depends on diabetes status. The narrow therapeutic range and discrepancy in respiration between experimental and human studies may limit clinical translation.

Attenuation of ischemia reperfusion (IR) injury is fundamental for reducing mortality and morbidity in patients suffering from a myocardial infarction. Ischemic and pharmacological conditioning strategies, as a supplement to revascularization, may reduce IR injury¹ by ultimately modulating mitochondrial function². While experimental and clinical conditioning studies have shown compelling evidence of infarct size reduction^{3,4}, translation into a clinical benefit has been challenging^{5,6}. Clinical outcome in patients with ST-elevation myocardial infarction (STEMI) undergoing immediate revascularization has improved significantly⁷. However, the development seems to level out signifying a continuous need for new cardioprotective strategies that go beyond current conditioning modalities in a translational perspective⁸.

During ischemia, hypoxia leads to suppression of the mitochondrial oxidative phosphorylation (OXPHOS)⁹. Revascularization causes a rapid re-establishment of oxygen and consequently mitochondrial OXPHOS. This results in a burst of reactive oxygen species (ROS) emission originating predominantly from complex I in the electron transport chain^{10,11}. Succinate increases the complex II (the succinate dehydrogenase (SDH)) linked respiration, which seems to overload the capacity of the electron transport system (ETS), and induce reverse electron transport to complex I and increased ROS production¹¹. Increased level of ROS, in combination with

¹Department of Cardiology, Aarhus University Hospital, Aarhus, Denmark. ²Lab, Center for Healthy Aging, Department of Biomedical Sciences, University of Copenhagen, Copenhagen, Denmark. ³Clinical Research Centre, Medical University of Bialystok, Bialystok, Poland. ⁴Department of Cardiothoracic Surgery, Aarhus University Hospital, Aarhus, Denmark. ✉e-mail: n.riise.jespersen@clin.au.dk

mitochondrial damage and dysfunction, initiates a vicious circle of increased mitochondrial dysfunction and further ROS production ultimately causing myocyte death and irreversible myocardial damage¹².

Mitochondrial metabolism can be modulated to reduce excessive ROS production and limit the myocardial injury^{13–15}. Inhibition of the malate aspartate shuttle (MAS) (transport of redox compounds from the cytosol into the mitochondria) protects the myocardium through reduced succinate accumulation and attenuated ROS mediated mitochondrial dysfunction¹⁶. As SDH is the key enzyme responsible for succinate build-up during ischemia, inhibition of the SDH represents a more targeted approach to reduce succinate levels during ischemia and reperfusion than inhibition of the MAS. Inhibition of SDH protects against IR injury in cardiomyocytes¹¹. Studies examining the cardioprotective capacities of SDH inhibition have been conducted in young healthy, non-diabetic animals, so the clinical perspectives of SDH inhibition remain limited.

In patients with diabetes, the outcome following an acute myocardial infarction is impaired compared to patients without diabetes¹⁷. Compromised outcome in diabetes may be due to metabolic disarrays in the cardiac cells beyond the vascular disease burden^{18,19}. Diabetes is associated with mitochondrial dysfunction²⁰, even when compared to overweight non-diabetic controls²¹. Additionally, susceptibility to ischemia is impacted by progression of the disease with increased injury in mature diabetes²² and the stimulus required for protection increases with duration and severity of the disease²³. We hypothesized that hearts from mature diabetic individuals have alterations in mitochondrial metabolism leading to mitochondrial dysfunction and an inability to achieve protection by inhibition of the SDH.

We evaluated cardioprotection by pre-ischemic inhibition of the SDH by dimethyl malonate (DiMAL) in animals with or without type 2 diabetes. We transferred these findings into human cardiac tissue by examining the effect of pre-ischemic SDH inhibition by DiMAL on contractile force recovery and mitochondrial function in trabeculae from patients with or without type 2 diabetes.

Methods

Ethical approval. The isolated rat heart study conformed to the Danish law for animal research (Act. No. 1306 of 23/11/2007, Danish Ministry of Justice) and the guidelines from *Guide for the Care and Use of Laboratory Animals* published by the US National Institutes of Health (NIH Publication No. 85–23, revised 1996). The Danish Animal Experimental Inspectorate approved the experimental work (Authorization No. 2018-15-0201-01446). The human study conformed to the Danish law for clinical studies and the study was approved by the Danish health research ethical committee (Authorization No. 1-10-72-361-15) and registered on clinicaltrials.gov (registration number: NCT02993484). Informed consent was obtained from all participating patients prior to enrollment in the study.

Biological material. We used 24 weeks old male Zucker diabetic fatty (ZDF) rats (homozygote (fa/fa), n = 57, approximately 400 g, Charles River Laboratories, USA) and age matched non-diabetic controls (heterozygote (fa/+), n = 59, approximately 400 g, Charles River Laboratories, USA) as a model of mature type 2 diabetes following guidelines for rigor and reproducibility in preclinical and clinical studies on cardioprotection^{24,25}. Due to excessive food intake the animals developed mature diabetes at 24 weeks of age. Animals were kept at a constant temperature of 23 °C with a 12 hours light-dark cycle and allowed unlimited access to enriched food (Purina 5008; recommended by supplier) and water. No anti-diabetic treatment was given. Rats were fasted 10–12 hours prior to the experiments in order to allow correct measurement of fasting blood glucose levels²².

We obtained human cardiac atrial appendage tissue from patients undergoing elective coronary artery bypass grafting (CABG) or valve replacement surgery with the use of extracorporeal circulation. The right atrial appendage was removed to allow insertion of the venous tube into the heart. Following excision of the appendage it was immediately immersed in oxygenated KH buffer (pH 7.35–7.45, room temperature) and transported to the laboratory within 5 minutes.

Study design. In the isolated rat heart study, isolated perfused hearts were divided into 4 groups based on the type of interventions (n = 7–9 in each group): (I) Sham hearts (Sham group), (II) IR-injured hearts (IR group) and IR-injured hearts co-perfused with (III) 0.1 mM DiMAL or (IV) 0.6 mM DiMAL for 10 minutes prior to global no-flow ischemia to mimic preconditioning (Fig. 1). All interventions were evaluated in both diabetic and non-diabetic hearts.

We conducted a dose-response correlation and two experimental series: an infarct size study and a mitochondrial respiratory and function study. We did not evaluate infarct size in the Sham groups as it is known to be negligible²⁶ and without importance for the conclusions of the study.

Following isolation, hearts were allowed to stabilize for 30–40 minutes depending on the type of intervention. Subsequently, they received 40 minutes global no-flow ischemia followed by 30 minutes (experimental series II) or 120 minutes (experimental series I) of reperfusion. After 30 minutes of reperfusion in experimental series II, the left ventricular muscle was quickly removed and divided into three parts. One part was immediately stored in an ice-cold relaxing solution (BIOPS, composition in mM: 2.77 CaK₂, EGTA, 7.23 EGTA, 20 taurine, 6.56 MgCl₂, 5.77 ATP, 15 phosphocreatine, 0.5 dithiothreitol and 50 4-morpholineethanesulphonic acid; pH 7.1) for the measurements of mitochondrial respiratory capacity and mitochondrial fatty acid oxidation. The remaining parts were quickly frozen in liquid nitrogen and stored at –80 °C for later analyses, including mitochondrial enzymatic activities and protein content.

In the human study, patients were recruited after project information and giving written consent. The following criteria led to exclusion: age >85 years, atrial fibrillation, ejection fraction <30% or CKMB or Troponin T elevation within 2 weeks. Following excision of the right atrial appendage it was immediately immersed in oxygenated Krebs-Henseleit buffer (pH 7.35–7.45, room temperature) and transported to the laboratory. After suspension in the superperfused atrial strip myograph, each working trabecula was randomized to one of the





Figure 1. Study design of the isolated rat heart study. An overview of the two experimental series of the isolated rat heart study including subgroups and perfusion protocols. IR: Ischemia reperfusion, DiMAL: Dimethyl Malonate, VF: Ventricular fibrillation.

following groups: (I) IR-injured trabeculae (IR group), (II) IR-injured trabeculae treated with IPC by 10 minutes of anoxia followed by 10 minutes of reperfusion prior to sustained anoxia (IPC group) and (III) IR-injured trabeculae co-perfused with 5 mM DiMAL for 20 minutes prior to sustained anoxia (DiMAL group) (Fig. 2).

Analysis of blood glucose and plasma metabolites. Prior to isolation of the heart pre-anesthetic fasting blood samples were taken by tail-vein bleeding for measurements of fasting blood glucose (FreeStyle Precision, Abbott Diabetes Care, Copenhagen, Denmark) and plasma insulin as previously described²². A fasting blood glucose measurement above 7 mmol/L was used to define the diabetic status. Samples for insulin measurements were collected in heparinized tubes (approximately 400 μ L) and centrifuged (5000 RPM for 1 minute at ambient temperature). The supernatant was stored at -80°C before analysis with a hypersensitive rat insulin ELISA kit (DRG instruments, Marburg, Germany). Serum total-cholesterol, triglycerides and free fatty acids were measured in 1 ml blood samples drawn from the abdominal aorta immediately before excision of the heart. Preparations included centrifugation (3500 RPM for 10 minutes at -4°C), collection of the supernatant and storage at -80°C until analysis on a Cobas Integra Analyzer (Roche Diagnostics, Rotkreuz, Switzerland).

Isolated heart preparation. Isolated perfused rat hearts (Langendorff perfusion) were prepared as previously described²⁷. Rats were anesthetized by a subcutaneous injection with a mixture of Dormicum (midazolam, 0.5 mg (kg body weight⁻¹); Matrix Pharmaceuticals, Herlev, Denmark) and Hypnorm (fentanyl citrate, 0.158 mg (kg body weight⁻¹ and fluanisone, 0.5 mg (kg body weight⁻¹)).

A tracheotomy was performed and the rat was connected to a rodent ventilator (Ugo Basile 7025 rodent ventilator, Comerio, Italy) and ventilated at 60 breaths/minute with a tidal volume of 3 mL. Subsequently, a laparotomy



1) Human study



Figure 2. Study design of human study. An overview of the experimental series of the study including subgroups and perfusion protocols. IR: Ischemia reperfusion, IPC: Ischemic preconditioning, DiMAL: Dimethyl Malonate, AFLI: Atrial fibrillation.

and thoracotomy were performed and a bolus of 1000 IU/kg heparin (Leo Pharma, Ballerup, Denmark) was administered through the femoral vein. The ascending aorta was cannulated *in situ* and retrograde perfusion of the heart was commenced at a constant pressure of 80 mmHg with an oxygenated (95% O₂ and 5% CO₂) Krebs Henseleit buffer (KHB: composition in mM: NaCl 118.5, KCl 4.7, NaHCO₃ 25.0, glucosemonohydrate 11.0, MgSO₄·7H₂O 1.2, CaCl₂ 2.4 and KH₂PO₄ 1.2). The heart was excised under continuous perfusion and mounted in an isolated perfused heart system (IH-SR type844/1; HSE, March-Hugstetten, Germany) where temperature is kept constant at 37 °C. An intraventricular balloon (Size 7, HSE, March-Hugstetten, Germany) was inserted in the left ventricle after removing the left atrial appendage and the intraventricular balloon volume was adjusted to a left ventricular end-diastolic pressure of 4–8 mmHg to simulate preload. Coronary flow was monitored continuously using an inline flow probe (Type 2.5SB, Transonic System Inc., Ithaca, NY, USA). Data was digitally converted (DT9804; Data Translation, Marlboro, MA, USA) and stored using Notocord Hem software (version 2.0, Notocord systems, Croissy sur Seine, France).

Atrial strip model. Isolated atrial trabeculae were prepared as previously described²⁸. Patients undergoing CABG or valve-replacement surgery were enrolled in the study. Within 20 minutes of excision of the atrial appendage, atrial trabeculae were tied by a suture (4.0 silk) in each end, excised from the remaining atrial appendage and mounted in the atrial strip myograph. Sutures are connected to a force transducer and a fixed hook.

The atrial trabeculae (length ranging between 5 and 10 mm and diameter ranging between 400–600 μm) were submerged into 37 °C Krebs-Henseleit buffer oxygenated with 95% O₂ and 5% CO₂ to maintain pH between 7.35 and 7.45. The isolated trabeculae were stimulated with 1 Hz by platinum electrodes connected to stimulators (Harvard Apparatus, amplitude 30 V, duration 1 ms) during stabilization and reoxygenation. Simulated ischemia was established by increased pacing (3 Hz) and changing to a hypoxic buffer (composition in mM: NaCl 118.5, KCl 4.7, NaHCO₃ 25.0, choline chloride 14.0, MgSO₄·7H₂O 1.2, CaCl₂ 2.4 and KH₂PO₄ 1.2), which was deoxygenated with 95% N₂ and 5% CO₂. After mounting the trabeculae in the organ baths, all trabeculae were stretched

passively to a pretension of 0.5 g and left to stabilize for 30 minutes. Subsequently pretension was increased to 0.75 g for additional 45 minutes of stabilization.

Infarct size. At the end of perfusion in experimental series I of the isolated rat heart study, hearts were frozen at -80°C , sliced (1.5 mm thick) and vital stained with 1% 2,3,5-triphenyltetrazoliumchloride (Merck, Darmstadt, Germany) for 3 minutes at 37°C to delineate areas of infarction. The stained heart slices were preserved in formaldehyde (10%) for 24–48 hours and then scanned using a flatbed scanner (Epson Perfection V600, Epson, Suwa, Japan) and weighed (Mettler-Toledo level balances, Mettler-Toledo, USA)¹⁶. Area-at-risk and area-of-infarction (infarct size) were assessed using image analysis software (ImageJ, National Institute of Health, USA) and weighted with the wet weight of the individual slices. Finally, infarct size/area-at-risk (IS/AAR) ratio was calculated. An observer blinded to treatment and disease groups analyzed the data.

Contractile force assessment. Contractile force was measured using a force transducer and data was digitally converted (DT9804; Data Translation, Marlboro, MA, USA) and stored using Notocord Hem software (version 2.0) (Notocord systems, Croissy sur Seine, France). Contractile force recovery was calculated as a ratio of post-ischemic contractile force divided by pre-ischemic contractile force and served as a surrogate marker of myocardial IR injury²⁸.

Mitochondrial respiratory capacity. At the end of the perfusion protocol in experimental series II of the isolated rat heart study, the left ventricle was quickly divided in two samples during continued perfusion for subsequent measurement of mitochondrial respiration and various tissue analyses. In the human study the trabeculae were removed and the areas affected by the ligature were discarded. The tissue was stored temporarily in a cold ($0-4^{\circ}\text{C}$) relaxation buffer (BIOPS; composition in mM: CaK_2EGTA 2.77, K_2EGTA 7.23, Na_2ATP 5.77, $\text{MgCl}_2\cdot 6\text{H}_2\text{O}$ 6.56, Taurine 20, $\text{Na}_2\text{Phosphocreatine}$ 15, Imidazole 20, Dithiothreitol 0.5 and MES 50; pH 7.1; kept between $0-4^{\circ}\text{C}$) until isolation of the fibers.

Mitochondrial respiration was measured in permeabilized cardiac muscle fibers as previously described¹⁶. Muscle fibers were dissected free of connective tissue in ice-cold BIOPS buffer using sharp forceps. After dissection, fibers were placed in ice-cold BIOPS-buffer supplemented with $50\ \mu\text{g}\cdot\text{mL}^{-1}$ Saponin for 30 minutes to ensure permeabilization. Fibers were subsequently washed by agitation in ice-cold MiR05-buffer (composition in mM): EGTA 0.5, $\text{MgCl}_2\cdot 6\text{H}_2\text{O}$ 3.0, K-lactobionate 60, Taurine 20, KH_2PO_4 10, HEPES 20, Sucrose 110 and BSA $1\ \text{g}\cdot\text{L}^{-1}$; pH 7.1) for two times 10 minutes. Muscle fibers were then weighed and transferred to an oxygraph (Oxygraph-2k; Oroboros, Innsbruck, Austria) for measurement of glucose supported metabolism (protocol 1) and fatty acid supported metabolism (protocol 2). All measurements were performed as duplicates in hyperoxygenation to avoid potential oxygen limitations to respiration. Due to limited tissue available for high-resolution respirometry only protocol 1 was applied in the human study.

Protocol 1 (complex I + II-linked respiration): Malate (2 mM) and glutamate (10 mM) were added to evaluate state 2 respiration (GM). After addition of ADP (5 mM) state 3 respiration was achieved (GM3). Respiratory control ratio (RCR) was calculated as state 3/state 2. Cytochrome *c* ($10\ \mu\text{M}$) was added to evaluate outer mitochondrial membrane integrity (more than 10% increases in respiration lead to exclusion). Subsequently, succinate (20 mM) was added to activate complex II during state 3 (GMS3) followed by Oligomycin ($2.5\ \mu\text{M}$) to achieve state 4 respiration (4o). Carbonyl cyanide-p-trifluoromethoxyphenylhydrazone (FCCP) was added to evaluate maximal respiratory capacity in a non-coupled state (E). Finally, Rotenone ($0.5\ \mu\text{M}$) and Antimycin A ($2.5\ \mu\text{M}$) were added to estimate residual oxygen consumption (ROX).

Protocol 2 (fatty acid oxidation): Malate (2 mM) was added to give fundamental stimulation of complex I (state 2 respiration) followed by multiple titrations of Octanoyl-1-carnitine to achieve maximal β -oxidation and complex II respiration (MOc). State 3 respiration is achieved by adding ADP (5 mM, MOC3) and finally cytochrome *c* ($10\ \mu\text{M}$) was added to examine outer membrane integrity.

Mitochondrial enzyme activity. Citrate synthase (CS) activity, a key enzyme in the glucose oxidation and marker of mitochondrial content, and β -hydroxyacyl-CoA dehydrogenase (HAD) activity, a key enzyme in the fatty acid oxidation, were measured in cardiac tissue homogenate by spectrophotometry as previously described²⁹. In short, 2–3 mg wet weight of cardiac muscle were homogenized in a TissueLyser (Qiagen, Venlo, Limburg, The Netherlands). For CS, the homogenate was diluted 50 times and the conversion of 5,5-dithiobis-(2-nitrobenzoic acid) to 5-thiobis-(2-nitrobenzoic acid) was measured at 37°C spectrophotometrically at 415 nm on an automatic analyzer (Cobas 6000, C 501, Roche Diagnostics, Mannheim, Germany). For HAD, the homogenate was diluted 70 times and changes in NADH at 37°C were measured spectrophotometrically at 340 nm on an automatic analyzer (Cobas 6000, C 501, Roche Diagnostics).

Mitochondrial protein content of ETC complexes. Mitochondrial protein content was measured as previously described³⁰. In short, the protein homogenates were diluted in Laemmli buffer and $15\ \mu\text{g}$ of total protein from each sample was separated on precast criterion gels (Bio-Rad, Copenhagen, Denmark). After electrophoresis, the gel was electrophoretically transferred to a polyvinylidene fluoride (PVDF) membrane. The membranes were blocked over night at 4°C with skim milk diluted in phosphate-buffered saline. The Total OXPHOS antibody cocktail (Abcam, ab110413) was used to determine the proteins of interest and set to incubate at room temperature for 1 hour followed by incubation with horseradish peroxidase-conjugated secondary antibodies (Dako, Glostrup, Denmark). Visualization of antibody-specific labeling bands were revealed with activation with enhanced chemiluminescence Western blotting detection system (ECL, GE Healthcare, Little Chalfont, UK) and a CCD camera (LAS 4000, GE Healthcare). We quantified intensities of each specific band obtained by chemiluminescence using Image Quant TL software (GE Healthcare, USA) and normalized to the entire UV intensity of the corresponding sample representing the total protein content on the membrane.

Type		Non-diabetic	Non-diabetic	Non-diabetic		Diabetic	Diabetic	Diabetic
Group		IR (n = 8)	DiMAL 0.1 mM (n = 8)	DiMAL 0.6 mM (n = 8)		IR (n = 9)	DiMAL 0.1 mM (n = 8)	DiMAL 0.6 mM (n = 8)
Characteristics of the ZDF rat model								
Infarct size study								
Body weight (BW), g		402 ± 22	405 ± 30	404 ± 25		403 ± 37	407 ± 28	387 ± 21
Heart weight (HW), g		1.18 ± 0.15	1.20 ± 0.10	1.23 ± 0.12		1.23 ± 0.11	1.18 ± 0.07	1.19 ± 0.05
HW/BW ratio		0.29 ± 0.03	0.30 ± 0.02	0.30 ± 0.02		0.31 ± 0.03	0.29 ± 0.02	0.31 ± 0.02
B-glucose, mmol/L		4.7 ± 0.4	5.5 ± 0.7	4.7 ± 0.3		15.0 ± 4.1 F [#]	24.7 ± 1.9 F	12.8 ± 3.4 F [#]
P-insulin, pmol/L		44.2 ± 18.1	47.9 ± 9.7	41.2 ± 15.4		88.0 ± 27.6 F	89.4 ± 21.7 F	82.7 ± 14.3 F
S-total cholesterol, mmol/L		2.17 ± 0.20	2.46 ± 0.27	2.29 ± 0.40		4.11 ± 0.72 F [#]	5.18 ± 0.77 F	4.00 ± 0.46 F [#]
S-triglyceride, mmol/L		0.88 ± 0.52	0.78 ± 0.32	0.97 ± 0.39		9.81 ± 1.50 F	8.87 ± 0.53 F	9.90 ± 0.56 F
S-free fatty acids, mmol/L		1.70 ± 1.34	1.80 ± 0.78	1.71 ± 0.58		6.14 ± 0.56 F	5.83 ± 1.49 F	5.21 ± 1.77 F
Mitochondrial function study								
Type	Non-diabetic	Non-diabetic	Non-diabetic	Non-diabetic	Diabetic	Diabetic	Diabetic	Diabetic
Group	Sham (n = 8)	IR (n = 8)	DiMAL 0.1 mM (n = 8)	DiMAL 0.6 mM (n = 7)	Sham (n = 8)	IR (n = 9)	DiMAL 0.1 mM (n = 8)	DiMAL 0.6 mM (n = 8)
Body weight (BW), g	413 ± 26	397 ± 18	400 ± 34	423 ± 31	408 ± 46	370 ± 38	390 ± 19	392 ± 42
B-glucose, mmol/L	4.8 ± 0.2	5.0 ± 0.4	5.1 ± 0.5	4.7 ± 0.3	12.5 ± 3.1 F [#]	13.0 ± 2.7 F [#]	24.2 ± 4.0 F	11.9 ± 1.8 F [#]
P-insulin, pmol/L	47.2 ± 16.2	36.4 ± 12.9	49.4 ± 10.1	46.3 ± 12.1	82.0 ± 24.8 F	75.3 ± 18.0 F	74.3 ± 18.0	83.1 ± 32.8 F
S-total cholesterol, mmol/L	2.19 ± 0.15	2.23 ± 0.18	2.09 ± 0.57	2.43 ± 0.14	4.32 ± 0.32 F	3.83 ± 0.75 F [#]	4.95 ± 0.95 F	4.04 ± 0.27 F [#]
S-triglyceride, mmol/L	0.94 ± 0.44	0.94 ± 0.37	0.87 ± 0.21	1.18 ± 0.42	9.97 ± 0.77 F [#]	10.14 ± 0.53 F [#]	8.37 ± 0.96 F	9.99 ± 0.81 F [#]
S-free fatty acids, mmol/L	1.73 ± 0.72	1.50 ± 0.90	1.83 ± 0.51	2.03 ± 1.14	5.92 ± 0.78 F	5.18 ± 2.07 F	4.71 ± 1.45 F	5.063 ± 1.51 F

Table 1. Characteristics of the ZDF rat model. Data are mean ± SD. F P < 0.05 grouped diabetic rats (fa/fa) compared to their grouped age-matched controls. [#]P < 0.05 diabetic group vs. diabetic DiMAL 0.1 mM.

Complex II activity. Frozen tissue was homogenized in a PBS buffer supplemented with protease inhibitors and prepared for complex II activity measurements using a Complex II Enzyme activity assay kit (Abcam, ab109908). Colorimetric measurements were achieved using PheraStar FS (BMG Labtech, Aylesbury, UK).

Myocardial interstitial concentrations of succinate. Myocardial interstitial concentration of succinate was measured in the 0.6 mM DiMAL group in non-diabetic and diabetic rats by microdialysis as previously described²⁷. A microdialysis probe (membrane length 4 mm, cut-off 6 Da; AgnTho's, Lidingö, Sweden) was inserted into the free wall of the left ventricle. The microdialysis probe was continuously perfused at a flow speed of 1 µL/minutes. with Krebs-Henseleit buffer solution deoxygenated with 95% N₂ and 5% CO₂. Following the implantation, a 20 minutes stabilization period was allowed for the metabolites to reach equilibrium in the perturbed tissue. Samples were collected at 10 minutes intervals and later analyzed by liquid chromatography and mass spectrometry.

Statistical analysis. All results are expressed as mean ± SD. Non-diabetic and diabetic groups were compared by two-way ANOVA. Comparison between baseline characteristics of patients with and without diabetes was done by t-test. Between groups comparison was performed using one-way ANOVA followed by a Bonferroni post-hoc analysis when appropriate. We used Prism, version 6.0 (GraphPad Software, La Jolla, CA, USA) for all statistical analyses. P < 0.05 was considered statistically significant.

Results

Exclusions. *Isolated rat heart study.* A total of 4 rats (all non-diabetic) were excluded prior to enrolment in the study (Fig. 1). Three rats were excluded due to ventricular fibrillation in the stabilization period and one rat due to unsuccessful surgical procedure. Complete datasets were achieved from the remaining rats.

Human study. In the human study 5 patients were excluded after giving written consent due to small atrial tissue sample sizes (Fig. 2). 10 patients were allocated to protocol optimization and dose/response correlation. 33 patients were enrolled in the study and 12 patients were excluded because the fibers did not reach sufficient hemodynamic performance of 0.5 g of contractility at the end of stabilization²⁸.

Baseline characteristics. *Isolated rat heart study.* Bodyweight, heart weight, fasting blood glucose levels and biochemical characteristics are summarized in Table 1. Body and heart weight did not differ between diabetic and non-diabetic rats. Fasting blood glucose was significantly higher in diabetic groups compared to the respective non-diabetic groups. Diabetic rats in the DiMAL 0.1 mM groups had significantly higher glucose levels than the other diabetic groups. We found no significant differences between all other diabetic and non-diabetic disease groups.

The diabetic groups had significant elevation in the biochemical variables, including insulin, cholesterol, triglycerides and free fatty acids (Table 1), compared to their respective non-diabetic groups. S-cholesterol was

Patients characteristics			
Type	Patient without diabetes	Patient with diabetes	p value
	(n = 12)	(n = 9)	
Age, year (mean ± SD)	64 ± 9	63 ± 7	0.75
Male gender, n (%)	11 (92)	8 (77)	0.78
Smoker, n (%)	1 (8)	1 (11)	0.78
Former smoker, n (%)	7 (58)	6 (67)	0.47
BMI, kg/m ² n (mean ± SD)	27.1 ± 4.4	30.0 ± 4.0	0.15
HbA1c, mmol/L (mean ± SD)	6.2 ± 0.6	8.7 ± 1.0	<0.05
Total cholesterol, mmol/L (mean ± SD)	4.0 ± 1.0	4.3 ± 1.4	0.63
LDL, mmol/L (mean ± SD)	2.2 ± 0.7	2.5 ± 1.1	0.41
HDL, mmol/L (mean ± SD)	1.1 ± 0.3	0.9 ± 0.2	0.30
Triglyceride, mmol/L (mean ± SD)	1.7 ± 1.0	1.8 ± 0.5	0.70
Statin therapy, n (%)	3 (25)	9 (100)	<0.05
Antihypertensive therapy, n (%)	7 (58)	9 (100)	<0.05

Table 2. Patient characteristics. Data are mean ± SD.




Figure 3. Isolated rat heart study – Dose-response relationship and infarct size. Dose-response relationship showing the effect of DiMAL on infarct size (IS) as a percentage of area at risk (AAR) in non-diabetic rat hearts (a) and IS/AAR of different intervention groups in non-diabetic and diabetic ZDF rat hearts (b). IR: Ischemia reperfusion, DiMAL: Dimethyl Malonate. Data are the mean ± SD. * $P < 0.05$ between indicated groups.

significantly elevated in DiMAL 0.1 mM compared to other diabetic groups. Between all other groups within the diabetic and non-diabetic disease groups, we found no significant differences.

Human study. We found no differences between patients with and without diabetes on age, BMI, total cholesterol, LDL-, HDL-cholesterol or triglyceride (Table 2). We found significantly higher HbA1c levels in patients with diabetes compared to patients without diabetes (8.7 ± 1.0 vs. 6.2 ± 0.6 mmol/L of HbA1c, $P < 0.05$). All patients with diabetes received antihypertensive and statin treatment, compared to only 58% and 25% in patients without diabetes ($p < 0.05$).

Infarct size. Isolated rat heart study - Dose-response relationship. The dose-response relationship of DiMAL treatment in non-diabetic rat hearts revealed a u-shaped relationship between dose and infarct size. Infarct size was reduced in the 0.1 mM DiMAL group compared to IR alone ($54 \pm 6\%$ vs. $69 \pm 6\%$ of LV, $p < 0.05$) (Fig. 3). DiMAL concentrations of 0.05 mM, 0.3 mM and 0.6 mM did not provide reductions in infarct size compared to IR ($64 \pm 14\%$, $63 \pm 16\%$ and $73 \pm 13\%$ vs. $69 \pm 6\%$).

Isolated rat heart study - infarct size. Infarct size following IR did not differ statistically significantly in diabetic compared to non-diabetic rats ($79 \pm 8\%$ and $69 \pm 6\%$, $p = 0.07$) (Fig. 3). In non-diabetic animals, DiMAL 0.1 mM reduced infarct size compared to the IR group ($55 \pm 7\%$ vs. $69 \pm 6\%$, $p < 0.05$). DiMAL 0.6 mM yielded no protection ($73 \pm 13\%$ and $69 \pm 6\%$, $p = 0.71$) (Fig. 3). In diabetic hearts, DiMAL 0.1 mM treatment did not reduce infarct size compared to the IR group ($68 \pm 8\%$ and $79 \pm 8\%$, $p = 0.14$) whereas DiMAL 0.6 mM yielded significant protection ($64 \pm 13\%$ vs. $79 \pm 8\%$, $p < 0.05$).

Contractile force recovery. Human study - Dose-response relationship. The dose-response relationship of DiMAL treatment in non-diabetic human atrial trabeculae showed that a concentration of 5 mM yielded the largest improvement in contractile force recovery compared to IR alone ($54 \pm 34\%$ vs. $24 \pm 6\%$ of contractile force recovery, $p = 0.67$) (Fig. 4). Reduction of DiMAL concentration to 3 mM or an increment to 10 mM attenuated or




Figure 4. Human study – Dose-response relationship and contractile force recovery. Dose-response relationship between DiMAL and contractile force recovery in non-diabetic human atrial trabeculae (a) and contractile force recovery of non-diabetic and diabetic human atrial trabeculae given as a ratio of contractile force in the reperfusion compared to the pre-ischemic value (b). IR: Ischemia reperfusion, DiMAL: Dimethyl Malonate, IPC: Ischemic preconditioning. Data are mean \pm SD. * $P < 0.05$ between indicated groups. P-values at the top indicate overall effect of diabetes and interventions by two-way ANOVA, respectively.

abolished contractile force recovery improvements, respectively, compared to IR alone ($35 \pm 1\%$, $22 \pm 10\%$ and $24 \pm 6\%$ of contractile force recovery, $p > 0.99$).

Human study. Contractile force did not differ between patients without and with diabetes (ANOVA $p > 0.99$). IPC improved contractile force recovery in trabeculae from patients without diabetes compared to trabeculae undergoing IR alone ($43 \pm 12\%$ vs. $23 \pm 13\%$, $p < 0.05$) (Fig. 4). DiMAL 5 mM yielded the same improvement in non-diabetic trabeculae ($43 \pm 13\%$ vs. $23 \pm 13\%$, $p < 0.05$). IPC in trabeculae from patients with diabetes improved contractile force recovery compared to IR and DiMAL ($51 \pm 15\%$ vs. $21 \pm 8\%$ and $29 \pm 16\%$, $p < 0.05$). In contrast, DiMAL did not significantly affect contractile force recovery compared to IR in diabetic trabeculae ($29 \pm 16\%$ and $21 \pm 8\%$, $p = 0.83$).

Mitochondrial respiratory capacity. Isolated rat heart study. Diabetes reduced mitochondrial respiration capacity compared to non-diabetic hearts as evaluated by GM, GM3, GMS3 and RCR (Fig. 5, ANOVA: $p = 0.02$, $p = 0.002$, $p = 0.004$ and $p = 0.04$, respectively). DiMAL 0.1 mM increased State 2 respiration (GM) compared to DiMAL 0.6 mM but not compared to IR in non-diabetic hearts (41.6 ± 8.7 vs. 28.5 ± 5.3 and 33.9 ± 7.9 pmol $O_2/(s \cdot mg)$, $p < 0.05$ and $p = 0.25$). We observed no other differences in state 2 respiration in either the non-diabetic or the diabetic group. State 3 complex I-linked respiration capacity (GM3) was higher in both the Sham and DiMAL 0.1 mM groups than in the DiMAL 0.6 mM group in non-diabetic hearts (157.4 ± 26.5 and 140.0 ± 35.3 vs. 81.8 ± 42.4 pmol $O_2/(s \cdot mg)$, $p < 0.01$ and $p < 0.05$). IR alone numerically reduced the respiration compared to Sham (157.4 ± 26.5 vs. 112.3 ± 27.0 pmol $O_2/(s \cdot mg)$, $p = 0.09$). State 3 complex I- and II-linked respiration (GMS3) was higher in non-diabetic Sham hearts than in non-diabetic hearts exposed to IR, DiMAL 0.1 mM, DiMAL 0.6 mM as well as in the diabetic Sham group (283.5 ± 53.6 vs. 207.6 ± 42.6 , 211.0 ± 44.1 , 213.3 ± 48.0 and 198.8 ± 50.0 pmol $O_2/(s \cdot mg)$) ($p < 0.05$), respectively. The RCR was higher in the Sham groups compared to any of the intervention groups regardless of the presence of diabetes. No differences were observed between intervention groups in state 4o (Fig. 5), non-coupled state E with FCCP (Fig. 5) or ROX (data not shown).

Human study. State 2 respiration (GM) and state 3 complex I-linked respiration (GM3) did not differ between trabeculae from patients with and without diabetes and was not affected by IPC or DiMAL (Fig. 6). Also, State 3 complex I + II linked respiration (GMS3) further increased the level of respiration in all groups but with no differences between trabeculae from patients with and without diabetes or between interventions. Consistent with this finding, we observed no change in RCR by any intervention. State 4o, non-coupled state E with FCCP and ROX were similar in all intervention groups and between trabeculae from patients with and without diabetes (ROX data not shown).

Mitochondrial fatty acid oxidation. Isolated rat heart study. Diabetes reduced mitochondrial respiration capacity compared to non-diabetic hearts as evaluated by MOc3 and RCR (Fig. 7, ANOVA: $p = 0.007$ and $p = 0.02$). All interventions reduced mitochondrial fatty acid oxidation compared to the non-diabetic sham group. (MOc: Sham 50.3 ± 6.7 vs. IR 35.4 ± 5.2 vs. DiMAL 0.1 mM 37.7 ± 10.2 vs. DiMAL 0.6 mM 32.2 ± 10.6 pmol $O_2/(s \cdot mg)$) ($p < 0.05$); MOc3: Sham 106.6 ± 14.0 vs. IR 75.5 ± 12.0 vs. DiMAL 0.1 mM 58.2 ± 21.6 vs. DiMAL 0.6 mM 73.6 ± 30.2 pmol $O_2/(s \cdot mg)$) ($p < 0.05$) (Fig. 7). Pre-ischemic inhibition by DiMAL 0.1 mM reduced state 3 respiration (MOc3) in diabetic hearts compared to the Sham group (Sham 80.3 ± 14.4 vs. 50.1 ± 5.7 pmol $O_2/(s \cdot mg)$). RCR was reduced in the DiMAL 0.1 mM group in both non-diabetic (DiMAL 0.1 mM 1.52 ± 0.21 vs. Sham 2.15 ± 0.34 vs. IR 2.03 ± 0.23 vs. DiMAL 0.6 mM 2.23 ± 0.27 , ($p < 0.05$)) and diabetic hearts (DiMAL 0.1 mM 1.52 ± 0.23 vs. Sham 2.03 ± 0.14 vs. IR 1.88 ± 0.23 vs. DiMAL 0.6 mM 1.95 ± 0.16 , ($p < 0.05$)). Finally, diabetes was associated with an overall decrease in respiration as observed by state 3 respiration and RCR (ANOVA: $p = 0.007$ and 0.02).




Figure 5. Isolated rat heart study - Mitochondrial respiratory capacity. Representative graph of mitochondrial respiratory capacity from a Sham heart. Below mitochondrial respiratory capacity. GM, state 2 respiration with glutamate + malate (a); GMS, state 3 respiration with glutamate and malate (b); GMS3, state 3 respiration with glutamate, malate and succinate (c); 4o, state 4 respiration with oligomycin (d); FCCP, non-coupled state by FCCP (e); RCR with complex I-linked substrates (f). Data are mean \pm SD. * $P < 0.05$ between indicated groups. P-values at the top indicate overall effect of diabetes and interventions by two-way ANOVA, respectively.

Mitochondrial enzyme activity. *Isolated rat heart study.* No overall difference in CS activity was seen between the diabetic and non-diabetic group. Citrate synthase (CS) activity did not differ between groups in the non-diabetic hearts (Sham 196 ± 61 and IR 182 ± 25 and DiMAL 0.1 mM 201 ± 61 and DiMAL 0.6 mM 212 ± 85 $\mu\text{mol}/\text{min}/\text{g}$ ($p = 0.82$)) or the diabetic hearts (Sham 198 ± 44 and IR 196 ± 66 and DiMAL 0.1 mM 201 ± 44 and DiMAL 0.6 mM 194 ± 69 $\mu\text{mol}/\text{min}/\text{g}$ ($p = 0.99$)) (Fig. 8). β -Hydroxyacyl-CoA dehydrogenase (HAD) activity was not different between intervention groups in either the non-diabetic (Sham 187 ± 55 and IR 168 ± 26 and DiMAL 0.1 mM 190 ± 65 and DiMAL 0.6 mM 201 ± 80 $\mu\text{mol}/\text{min}/\text{g}$) or diabetic hearts (Sham 230 ± 64 and IR 240 ± 81 and DiMAL 0.1 mM 239 ± 63 and DiMAL 0.6 mM 236 ± 83 $\mu\text{mol}/\text{min}/\text{g}$). Diabetic hearts had a higher HAD activity than non-diabetic hearts (ANOVA $p < 0.05$).




Figure 6. Human study - Mitochondrial respiratory capacity. Representative graph of mitochondrial respiratory capacity from an IR heart trabecula. Below mitochondrial respiratory capacity. GM, state 2 respiration with glutamate + malate (a); GM3, state 3 respiration with glutamate and malate (b); GMS3, state 3 respiration with glutamate, malate and succinate (c); 4o, state 4 respiration with oligomycin (d); FCCP, non-coupled state by FCCP (e); RCR with complex I-linked substrates (f). Data are mean \pm SD. * $P < 0.05$ between indicated groups. P-values at the top indicate overall effect of diabetes and interventions by two-way ANOVA, respectively.

Mitochondrial protein content of ETC complexes and complex II activity. The protein content of the mitochondrial ETC complexes I-V in cardiac muscle did not differ between groups (Fig. 8) or between non-diabetic and diabetic cardiac tissue. Complex II activity in cardiac muscle was similar in all groups (Fig. 8).

Interstitial succinate concentration. Diabetic IR hearts had statistically non-significantly lower succinate concentration than non-diabetic IR hearts (58 ± 14 vs. 90 ± 35 $\mu\text{mol/L}$, $p = 0.30$) (Fig. 8). After 60 minutes of ischemia DiMAL reduced succinate concentrations in the non-diabetic group (90 ± 35 vs. 33 ± 42 $\mu\text{mol/L}$, $p < 0.05$). DiMAL also reduced interstitial succinate concentration in diabetic hearts but the difference was not statistically significant (43 ± 18 vs. 58 ± 14 $\mu\text{mol/L}$, $p > 0.99$).




Figure 7. Isolated rat heart study - Mitochondrial fatty acid respiratory capacity. Summarized data of mitochondrial fatty acid respiratory capacity in each respiratory state. MOC, state 2 respiration with malate and octanoyl-L-carnitine; MOC3, state 3 respiration with malate and octanoyl-L-carnitine; RCR with fatty acids. Data are the mean \pm SD. * $P < 0.05$ between indicated groups. P-values at the top indicate overall effect of diabetes and interventions by two-way ANOVA, respectively.

Discussion

The main finding of the present study is that inhibition of the SDH by administration of DiMAL protects the heart from IR injury in non-diabetic and diabetic animals. However, diabetes seems to reduce the sensitivity to DiMAL such that an increased dose is required to elicit a cardioprotective response. The DiMAL induced cardioprotection was associated with an improvement of post-ischemic mitochondrial function in non-diabetic animals but not in diabetic animals. Diabetes *per se* compromised mitochondrial respiration capacity, reflecting that mitochondrial dysfunction may explain reduced sensitivity to cardioprotective strategies in experimental models of diabetes. At a dose optimized through a dose-response experiment, a cardioprotective effect of DiMAL was present in human cardiac tissue from patients without diabetes, while cardiac tissue from patients with diabetes was not protected by the administered doses of DiMAL. In contrast to our experimental findings, mitochondrial function was not compromised in human cardiac tissue from diabetes patients and the absent effect of DiMAL was not associated with mitochondrial respiratory capacity.

Our results support previous findings of a cardioprotective effect by DiMAL^{11,31,32}. SDH inhibition has become an attractive target in pharmacologic conditioning because it may interact with central mechanisms underlying IR injury¹¹. Complex I is the major source of mitochondrial ROS production. During ischemia, the purine nucleotide cycle (PNC)¹¹ and the MAS²⁷ supply fumarate to Complex II, which acts in reverse to reduce fumarate to succinate. At reperfusion, oxygen supply is restored, and Complex II rapidly metabolizes the excess succinate in its forward direction to fumarate. However, a delay in the subsequent steps prevents further Complex III-V metabolism, when the mitochondrial membrane potential increases. As a consequence, electrons are forced in a reverse direction through Complex I generating large amounts of ROS. Excessive ROS production ultimately cause opening of the mitochondrial permeability pore complex and cell death⁹.

DiMAL is a competitive inhibitor of the SDH that binds to the succinate site of the enzyme¹⁵. DiMAL can be administered prior to index ischemia to reduce succinate accumulation¹¹ or during early reperfusion to inhibit succinate oxidation thereby limiting reverse electron transport and ROS production³². Our data demonstrate that pre-ischemic administration of DiMAL, mimicking preconditioning, is cardioprotective both in rats and in human cardiac tissue with an efficacy comparable to IPC. Remote IPC, postconditioning, hypothermia and cyclosporine A can induce the same level of cardioprotection as DiMAL^{33,34}. The additive effect of hypothermia and DiMAL on infarct size was protective by distinct mechanisms. It is unknown if remote IPC, postconditioning or cyclosporine A modifies ROS production by attenuation of succinate production. Our data showed that the succinate/ROS pathway does not seem to be an inherent part of IPC, since IPC does not inhibit succinate accumulation during ischemia^{16,27,35} or succinate oxidation during early reperfusion^{27,36}. These findings suggest that a




Figure 8. Isolated rat heart study - Mitochondrial enzyme activity. Enzyme activity of citrate synthase (CS) and β -hydroxyacyl-CoA dehydrogenase (HAD) (**a,b**). Below protein content of individual electron transport chain complex I-V in non-diabetic (**c**) and diabetic (**d**) heart tissue. At the bottom Complex II activity in non-diabetic and diabetic tissue (**e**) and interstitial succinate concentrations (**f**). IR: Ischemia reperfusion, DiMAL: Dimethyl malonate. Data are the mean \pm SD. * $P < 0.05$ between indicated groups. P-values at the top indicate overall effect of diabetes and interventions by two-way ANOVA, respectively.

combination of different therapeutic approaches could further protect against IR injury during acute myocardial infarction.

We found that an increased concentration of DiMAL was needed to reduce infarct size in diabetic compared to non-diabetic animals. This might be caused by changes in the amount or activity of SDH, since DiMAL is a competitive inhibitor of SDH. We did not observe any change in protein content of any of the complexes of the ETC or differences in complex II activity between groups. The finding indicates that the need for higher concentration of DiMAL may be caused by factors other than protein content and complex II activity, e.g. metabolic rate control or substrate selection beyond the mechanisms traditionally claimed to explain the glucose-fatty-acid cycle³⁷. Using dosages optimized to non-diabetic trabeculae, we found no protection by DiMAL in patients with diabetes. The dosage used in human tissue was chosen from the dose-response relationship in non-diabetic tissue and not customized to the diabetic phenotype. The optimized dosage for tissue from patients without diabetes was 10 fold higher than in the experimental setting. This is due to difference in drug delivery. In the isolated hearts DiMAL was delivered directly through the coronary vessels, whereas the atrial trabeculae were superperfused, demanding a higher concentration to achieve effect. Taking the experimental data into consideration, it seems plausible that the dosage was insufficient because diabetic animals needed a higher dose than non-diabetic animals to achieve protection. Due to low inclusion rates of patients with diabetes, it was not possible to evaluate the effect of a higher dosage in the current study, so we cannot exclude that a higher threshold might be present in humans as previously demonstrated in the diabetic rats³⁸. Our data suggest that the therapeutic window for SDH inhibition is narrow and this may be a translational challenge. We found that even small changes in

concentrations of DiMAL could abrogate the protection, supporting previous observations in mice that revealed a toxic effect of DiMAL on hemodynamic performance under normoxic conditions³¹. The respiratory capacity of complex I was severely compromised by the high DiMAL dosage in non-diabetic rats (DiMAL 0.6 mM) reflecting the narrow therapeutic window of DiMAL. The deleterious effect may be explained by an irreversible, detrimental effect of combining excessive SDH blockade with IR injury, which may inhibit restoration of normal respiration during reperfusion. No improvement in mitochondrial respiration was observed in human trabeculae treated with DiMAL or IPC indicating that cardioprotection may be achievable without measurable modulation of post-ischemic mitochondrial function.

Preserved post-ischemic mitochondrial function is important for cellular function and survival^{9,39}. Our data confirmed that diabetes has an inherent deleterious effect on overall mitochondrial respiration in our experimental setting, which adds to the impaired respiratory capacity following ischemia. We found that administration of 0.1 mM DiMAL to non-diabetic hearts tended to improve complex I-linked respiration reflecting reduced injury to the complex. This improvement in complex I-linked respiration correlates with the current hypothesis that DiMAL limits succinate accumulation, reverses electron transport to complex I and ROS-mediated injuries¹¹.

DiMAL reduced accumulation of interstitial succinate during ischemia in non-diabetic and diabetic rat hearts. Diabetic hearts seemed to have an inherent ability to reduce succinate accumulation compared to non-diabetic hearts. Reduced succinate accumulation during an ischemic event has previously been reported to provide cardioprotection¹⁶. We found no correlation between reduced succinate accumulation and cardioprotection since IR exposed diabetic hearts and DiMAL treated non-diabetic hearts did not have lower infarct sizes compared to non-diabetic IR hearts. The missing correlation between succinate buildup during ischemia and cardioprotection has previously been reported in the setting of ischemic preconditioning^{40,41}, so the role of succinate in IR injury remains unresolved.

Diabetic tissue from rat hearts showed elevated levels of HAD activity corresponding to an elevated fatty acid metabolism possibly caused by a reduced ability to metabolize glucose. However, *ex vivo* rat heart fatty acid oxidation was reduced indicating that other steps in the fatty acid metabolism may be impaired in the diabetic hearts. IR reduced fatty acid oxidation and treatment with DiMAL did not improve the respiratory capacity in spite of its cardioprotective effect. This finding may simply reflect that the isolated rat hearts only received glucose during the perfusion protocol or that fatty acid metabolism may be less important compared to glucose metabolism in the protection against IR injury.

CS activity did not differ between non-diabetic and diabetic hearts indicating a similar level of mitochondrial content. Diabetes can both up- and down-regulate mitochondrial content over the course of disease progression⁴². Our findings may suggest that the diabetic disease model had not yet developed major changes in mitochondrial content and support that the respiratory findings are mainly driven by changed respiratory capacity rather than the number of mitochondria.

The fact that co-morbidities such as diabetes and co-medication including glucose lowering therapy modify the sensitivity to cardioprotection and consequently changes the concentration required to elicit cardioprotection increases the complexity of applying the concept into clinical use. This challenge is reflected by the discrepant findings between our experimental and human studies. This is probably not a specific characteristic for DiMAL but for all modulators of the mitochondrial function. Future studies should focus on safe delivery of ETC modifiers to the target area in the optimal concentration with absent systemic effect. A recent study addressed this problem and demonstrated the potential of intracoronary administration of DiMAL in pigs³². While intracoronary administration increases the therapeutic value of the drug, it will most likely be a delicate balance to achieve optimal effect without introducing side effects.

An alternative and safer way to modulate mitochondria may be utilization of a physiological inhibitor rather than a synthetic inhibitor, which cannot be metabolized in the cell. Oxaloacetate and malate are tricarboxylic acid cycle intermediates with strong inhibitory effect on SDH¹⁵, similar to that of DiMAL. It may be possible to achieve a protective effect by increasing the levels of e.g. oxaloacetate. Protection by oxaloacetate has been tested in rats in a focal brain ischemic model with evidence suggesting a neuroprotective effect⁴³. The concept is further supported by the possibility to induce cardioprotection through MAS inhibition by aminooxyacetate (AOA) during late ischemia¹⁶. AOA inhibits the aspartate amino transferase, which uses oxaloacetate as a substrate and it is possible that the protection works through upregulation of oxaloacetate¹⁶. Finally, short cycles of IPC have been shown to increase malate, the precursor to oxaloacetate, indicating a possible involvement of these tricarboxylic acid cycle intermediates in the protection of IPC¹⁶.

The current study has limitations. First, we used a ZDF rat model to examine the impact of mature diabetes on myocardial IR injury. The ZDF rat is mutated in the leptin receptor resulting in a diabetic phenotype similar to the human. Because the leptin receptor is defect, the animals have high circulating levels of leptin, which may activate protective pathways⁴⁴. However, the ZDF model is considered a valid model for type 2 diabetes because of the similarities with the human phenotype. Second, the baseline characteristics of the animals were consistent with a diabetic phenotype similar to humans with elevated fasting blood glucose, insulin, cholesterol, triglycerides and FFA. However, compared to previous studies, the animals in the diabetic groups had lower levels of circulating glucose concentrations²². Only animals randomized to the DiMAL 0.1 mM groups had blood glucose levels similar to previous observations²². However, all animals in the diabetic groups had elevated blood glucose and insulin levels and a phenotype consistent with a metabolic syndrome. Third, we used an isolated rat heart model to evaluate the cardioprotective effects. Isolation of the heart excludes the influence of systemic and humoral factors during myocardial IR. Our circulation buffer did not contain insulin or FFA and does not resemble normal physiology. Fourth, we used permeabilized myocardial fibers for evaluation of mitochondrial respiration. This is a validated method to investigate the mitochondria *in situ*, and seems to be more physiologically correct than isolated mitochondria⁴⁵. Even so, it does not reflect physiological conditions, which may limit the translation into the clinical setting. Fifth, tissue from the right atrial trabeculae may not reflect the function, physiology

and metabolism of the left ventricle. Nevertheless, similarities between atrial and ventricular tissue have been demonstrated^{46,47} and the right atrial appendage remains the most ethically applicable method to examine human myocardium. Tissue was transported from the operating theater to our laboratory in an oxygenated buffer with a transport time below 5 minutes. It cannot be excluded that partial hypoxia might affect the tissue prior to suspension in the atrial strip model. Finally, we acknowledge the lack of insight into ROS production and injury, which limits the mechanistic understanding and interpretation of our data.

In summary, pre-ischemic inhibition of the SDH by DiMAL protects the non-diabetic and diabetic rat heart from IR injury. The protection can be translated into human cardiac tissue from patients without diabetes and is as effective as IPC. In our animal model, mature diabetes alters the metabolic phenotype that resulted in higher resistance to SDH inhibition and requirement for a higher dosage to induce cardioprotection. Accordingly, DiMAL treatment with an optimized dosage through dose-response correlation did not protect the diabetic human myocardium. The protection is associated with improved post-ischemic mitochondrial function in our isolated heart model, but not in the human atrial trabeculae. Conceptually, SDH inhibition may provide a pharmacological target to counteract IR injury. However, synthetic agents have narrow therapeutic range and efficacy may be influenced by diabetes.

Received: 6 April 2020; Accepted: 2 June 2020;

Published online: 25 June 2020

References

1. Botker, H. E., Lassen, T. R. & Jespersen, N. R. Clinical translation of myocardial conditioning. *Am. J. Physiol. Heart Circ. Physiol.* **314**, H1225–H1252 (2018).
2. Di Lisa, F. & Bernardi, P. Mitochondria and ischemia-reperfusion injury of the heart: fixing a hole. *Cardiovasc. Res.* **70**, 191–199 (2006).
3. Hausenloy, D. J. *et al.* Ischaemic conditioning and targeting reperfusion injury: a 30 year voyage of discovery. *Basic Res. Cardiol.* **111**, 70-016-0588-8. Epub 2016 Oct 20 (2016).
4. Heusch, G., Botker, H. E., Przyklenk, K., Redington, A. & Yellon, D. Remote ischemic conditioning. *J. Am. Coll. Cardiol.* **65**, 177–195 (2015).
5. Heusch, G. Critical Issues for the Translation of Cardioprotection. *Circ. Res.* **120**, 1477–1486 (2017).
6. Hausenloy, D. J. *et al.* Effect of remote ischaemic conditioning on clinical outcomes in patients with acute myocardial infarction (CONDI-2/ERIC-PPCI): a single-blind randomised controlled trial. *Lancet* **394**, 1415–1424 (2019).
7. Szummer, K. *et al.* Improved outcomes in patients with ST-elevation myocardial infarction during the last 20 years are related to implementation of evidence-based treatments: experiences from the SWEDEHEART registry 1995–2014. *Eur. Heart J.* **38**, 3056–3065 (2017).
8. Botker, H. E., Cabrera-Fuentes, H. A., Ruiz-Meana, M., Heusch, G. & Ovize, M. Translational issues for mitoprotective agents as adjunct to reperfusion therapy in patients with ST-segment elevation myocardial infarction. *J. Cell. Mol. Med.* **24**, 2717–2729 (2020).
9. Halestrap, A. P. A pore way to die: the role of mitochondria in reperfusion injury and cardioprotection. *Biochem. Soc. Trans.* **38**, 841–860 (2010).
10. Burwell, L. S., Nadochiy, S. M. & Brookes, P. S. Cardioprotection by metabolic shut-down and gradual wake-up. *J. Mol. Cell. Cardiol.* **46**, 804–810 (2009).
11. Chouchani, E. T. *et al.* Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS. *Nature* **515**, 431–435 (2014).
12. Zweier, J. L. & Talukder, M. A. The role of oxidants and free radicals in reperfusion injury. *Cardiovasc. Res.* **70**, 181–190 (2006).
13. Lesnefsky, E. J. *et al.* Blockade of electron transport during ischemia protects cardiac mitochondria. *J. Biol. Chem.* **279**, 47961–47967 (2004).
14. Elrod, J. W. *et al.* Hydrogen sulfide attenuates myocardial ischemia-reperfusion injury by preservation of mitochondrial function. *Proc. Natl. Acad. Sci. USA* **104**, 15560–15565 (2007).
15. Drose, S., Bleier, L. & Brandt, U. A common mechanism links differently acting complex II inhibitors to cardioprotection: modulation of mitochondrial reactive oxygen species production. *Mol. Pharmacol.* **79**, 814–822 (2011).
16. Jespersen, N. R. *et al.* Pre-ischaemic mitochondrial substrate constraint by inhibition of malate-aspartate shuttle preserves mitochondrial function after ischaemia-reperfusion. *J. Physiol.* **595**(12):3765–3780. (Jun 15, 2017).
17. Haffner, S. M., Lehto, S., Ronnema, T., Pyorala, K. & Laakso, M. Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. *N. Engl. J. Med.* **339**, 229–234 (1998).
18. Lehto, S. *et al.* Myocardial infarct size and mortality in patients with non-insulin-dependent diabetes mellitus. *J. Intern. Med.* **236**, 291–297 (1994).
19. Kleinbongard, P., Botker, H. E., Ovize, M., Hausenloy, D. J. & Heusch, G. Co-morbidities and co-medications as confounders of cardioprotection-Does it matter in the clinical setting? *Br. J. Pharmacol.* (2019).
20. Bugger, H. & Abel, E. D. Mitochondria in the diabetic heart. *Cardiovasc. Res.* **88**, 229–240 (2010).
21. Montaigne, D. *et al.* Myocardial Contractile Dysfunction is Associated with Impaired Mitochondrial Function and Dynamics in Type 2 Diabetic but not in Obese Patients. *Circulation* **130**(7):554–64. doi: 10.1161/CIRCULATIONAHA.113.008476. Epub 2014 Jun 13. (2014).
22. Povlsen, J. A. *et al.* Protection against myocardial ischemia-reperfusion injury at onset of type 2 diabetes in Zucker diabetic fatty rats is associated with altered glucose oxidation. *PLoS One* **8**, e64093 (2013).
23. Ferdinandy, P., Hausenloy, D. J., Heusch, G., Baxter, G. F. & Schulz, R. Interaction of risk factors, comorbidities, and comedications with ischemia/reperfusion injury and cardioprotection by preconditioning, postconditioning, and remote conditioning. *Pharmacol. Rev.* **66**, 1142–1174 (2014).
24. Lindsey, M. L. *et al.* Guidelines for experimental models of myocardial ischemia and infarction. *Am. J. Physiol. Heart Circ. Physiol.* **314**, H812–H838 (2018).
25. Botker, H. E. *et al.* Practical guidelines for rigor and reproducibility in preclinical and clinical studies on cardioprotection. *Basic Res. Cardiol.* **113**, 39-018-0696-8 (2018).
26. Jespersen, N., Lassen, T., Hjortbak, M., Støttrup, N. & Botker, H. Sodium Glucose Transporter 2 (SGLT2) Inhibition does not Protect the Myocardium from Acute Ischemic Reperfusion Injury but Modulates Post- Ischemic Mitochondrial Function. *Cardiovasc. Pharm Open Access* **6**, 2 (2017).
27. Støttrup, N. B. *et al.* Inhibition of the malate-aspartate shuttle by pre-ischaemic aminoxyacetate loading of the heart induces cardioprotection. *Cardiovasc. Res.* **88**, 257–266 (2010).
28. Jensen, R. V. *et al.* Impact of O-GlcNAc on cardioprotection by remote ischaemic preconditioning in non-diabetic and diabetic patients. *Cardiovasc. Res.* **97**, 369–378 (2013).

29. Christiansen, L. B. *et al.* Impaired cardiac mitochondrial oxidative phosphorylation and enhanced mitochondrial oxidative stress in feline hypertrophic cardiomyopathy. *Am. J. Physiol. Heart Circ. Physiol.* **308**, H1237–47 (2015).
30. Vigelso, A. *et al.* GAPDH and beta-actin protein decreases with aging, making Stain-Free technology a superior loading control in Western blotting of human skeletal muscle. *J. Appl. Physiol.* (1985) **118**, 386–394 (2015).
31. Valls-Lacalle, L. *et al.* Succinate dehydrogenase inhibition with malonate during reperfusion reduces infarct size by preventing mitochondrial permeability transition. *Cardiovasc. Res.* **109**, 374–384 (2016).
32. Valls-Lacalle, L. *et al.* Selective Inhibition of Succinate Dehydrogenase in Reperfused Myocardium with Intracoronary Malonate Reduces Infarct Size. *Sci. Rep.* **8**, 2442–018–20866–4 (2018).
33. Gedik, N. *et al.* Cardiomyocyte mitochondria as targets of humoral factors released by remote ischemic preconditioning. *Arch. Med. Sci.* **13**, 448–458 (2017).
34. Kleinbongard, P. *et al.* Mitochondrial and Contractile Function of Human Right Atrial Tissue in Response to Remote Ischemic Conditioning. *J Am Heart Assoc.* **7**, e009540, <https://doi.org/10.1161/JAHA.118.009540>. (2018).
35. Andrienko, T., Pasdois, P., Rossbach, A. & Halestrap, A. P. Real-Time Fluorescence Measurements of ROS and [Ca²⁺] in Ischemic/Reperfused Rat Hearts: Detectable Increases Occur only after Mitochondrial Pore Opening and Are Attenuated by Ischemic Preconditioning. *PLoS One* **11**, e0167300 (2016).
36. Pell, V. R. *et al.* Ischemic preconditioning protects against cardiac ischemia reperfusion injury without affecting succinate accumulation or oxidation. *J. Mol. Cell. Cardiol.* **123**, 88–91 (2018).
37. Cortassa, S., Aon, M. A. & Sollott, S. J. Control and Regulation of Substrate Selection in Cytoplasmic and Mitochondrial Catabolic Networks. A Systems Biology Analysis. *Front. Physiol.* **10**, 201 (2019).
38. Tsang, A., Hausenloy, D. J., Mocanu, M. M., Carr, R. D. & Yellon, D. M. Preconditioning the diabetic heart: the importance of Akt phosphorylation. *Diabetes* **54**, 2360–2364 (2005).
39. Hori, M. & Nishida, K. Oxidative stress and left ventricular remodelling after myocardial infarction. *Cardiovasc. Res.* **81**, 457–464 (2009).
40. Jose Alburquerque-Bejar, J. *et al.* Remote ischemic conditioning provides humoral cross-species cardioprotection through glycine receptor activation. *Cardiovasc. Res.* **113**, 52–60 (2017).
41. Hjortbak, M. V. *et al.* Influence of diabetes mellitus duration on the efficacy of ischemic preconditioning in a Zucker diabetic fatty rat model. *PLoS One* **13**, e0192981 (2018).
42. Duncan, J. G. Mitochondrial dysfunction in diabetic cardiomyopathy. *Biochim. Biophys. Acta* **1813**, 1351–1359 (2011).
43. Knapp, L. *et al.* Neuroprotective effect of oxaloacetate in a focal brain ischemic model in the rat. *Cell. Mol. Neurobiol.* **35**, 17–22 (2015).
44. Smith, C. C. *et al.* Leptin-induced cardioprotection involves JAK/STAT signaling that may be linked to the mitochondrial permeability transition pore. *Am. J. Physiol. Heart Circ. Physiol.* **299**, H1265–70 (2010).
45. Pesta, D. & Gnaiger, E. High-resolution respirometry: OXPHOS protocols for human cells and permeabilized fibers from small biopsies of human muscle. *Methods Mol. Biol.* **810**, 25–58 (2012).
46. Bohm, M., Pieske, B., Ungerer, M. & Erdmann, E. Characterization of A1 adenosine receptors in atrial and ventricular myocardium from diseased human hearts. *Circ. Res.* **65**, 1201–1211 (1989).
47. Heidebuchel, H., Vereecke, J. & Carmeliet, E. Three different potassium channels in human atrium. Contribution to the basal potassium conductance. *Circ. Res.* **66**, 1277–1286 (1990).

Acknowledgements

We sincerely thank Casper Carlsen Elkjær, Hans Pedersen, Regitze Kraunsøe and Christina Hansen for technical assistance. The present study was financially supported by grants from Kirsten Antonius foundation, Aarhus University; Novo Nordisk foundation, AP Møller foundation, The Danish Council for Independent Research (11-108354), the Danish Council for Strategic Research (11-115818).

Author contributions

N.R.J., N.B.S., and H.E.B. contributed to the conception and design of this study. N.R.J., T.R.L., M.V.H. and P.T.T. performed the experiments and analyzed the data. H.H.K. and N.R.J. obtained atrial appendage tissue from patients. N.R.J., T.R.L., M.V.H., N.B.S., J.J., S.L. and H.E.B. interpreted the results of the experiments. N.R.J., N.B.S., J.J. and H.E.B. wrote the manuscript. N.R.J., T.R.L., M.V.H., J.J., N.B.S., S.L., H.H.K. and H.E.B. edited and revised the manuscript. All authors approved the final version of the manuscript submitted for publication.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to N.R.J.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020