



## The equation of state at high temperatures from lattice QCD

---

**G. Endrődi<sup>1</sup>, Z. Fodor<sup>1,2</sup>, S.D. Katz<sup>1,2</sup>, K.K. Szabó<sup>\*2</sup>**

<sup>1</sup>*Institute for Theoretical Physics, Eötvös University, H-1117 Budapest, Hungary.*

<sup>2</sup>*Department of Physics, University of Wuppertal, D-42097 Wuppertal, Germany.*

*E-mail: [szaboka@general.elte.hu](mailto:szaboka@general.elte.hu)*

We present results for the equation of state upto previously unreachable, high temperatures. Since the temperature range is quite large, a comparison with perturbation theory can be done directly.

*The XXV International Symposium on Lattice Field Theory*

*July 30 - August 4 2007*

*Regensburg, Germany*

---

\*Speaker.

## 1. Introduction

Quantum chromodynamics (QCD) is the theory describing the strong interactions. According to QCD, as the temperature increases, hadronic matter undergoes a transition to quark gluon plasma. The function, which describes the equilibrium in this system for different temperatures, is the equation of state. For the equation of state, practically one has to measure only the pressure, which then will unambiguously determine the energy density as well. A difficulty emerging about the equation of state is, that perturbation theory does not seem to fit lattice results even at higher temperatures (smaller couplings). While available lattice results both for pure gauge theory [1, 2] and for full QCD [3, 4, 5] (and see also [6]) end at around  $5 \cdot T_C$ , standard perturbation theory converges only at extremely high temperatures. To create a link between these two methods, we present lattice results on the pressure at temperatures, which were previously unreachable. It becomes possible to compare our data with perturbation theory formulae. Our results are obtained using two different approaches: we present a new way to renormalize the pressure (for this we have results for  $N_t = 4$ ), and a direct method to measure the pressure (results for  $N_t = 4, 6$  and  $8$ ).

## 2. Renormalization of the pressure

For illustration, we present here the technique and the results for pure SU(3) gauge theory. The extension to full QCD is straightforward.

In order to obtain results which can be extrapolated to the continuum, one has to renormalize the pressure. This is usually done using the standard integral method [7]:

$$p_{\text{ren}}(T) = p(T) - p(0) = \int d\beta (\langle \text{Pl} \rangle_T - \langle \text{Pl} \rangle_0) \quad (2.1)$$

This way one has to carry out simulations on finite and zero temperature lattices. However, exactly zero temperature cannot be realized. Together with the fact, that divergences that are removed by renormalization are independent of the temperature, rises the question: why not use finite  $T$  lattices for renormalization? In order to do so, let us introduce the following quantity:

$$\bar{p}(T) = p(T) - p(T/2) = \int d\beta (\langle \text{Pl} \rangle_{N_t} - \langle \text{Pl} \rangle_{2 \cdot N_t}) \quad (2.2)$$

Here, instead of using  $1/2$  for the subtraction temperature, one can use any factor, which is smaller than 1. Now, we may build up  $p_{\text{ren}}$  as a sum of differences, so it can be expressed with  $\bar{p}$  as:

$$p_{\text{ren}}(T) = p(T) - p(T/2) + p(T/2) - p(T/4) + \dots = \bar{p}(T) + \bar{p}(T/2) + \dots \quad (2.3)$$

In fact, we usually measure the dimensionless pressure, which can be obtained by including the  $T^4$  factors:

$$\frac{p_{\text{ren}}}{T^4} = \left. \frac{\bar{p}}{T^4} \right|_T + \frac{1}{16} \cdot \left. \frac{\bar{p}}{T^4} \right|_{T/2} + \frac{1}{256} \cdot \dots \quad (2.4)$$

Due to the increasing powers of  $1/16$  in the forthcoming terms, one practically needs only a few of them. Applying this scheme, we may reach arbitrary high temperatures using lattices with only  $N_t$

and  $2 \cdot N_t$  temporal extent. It is worth mentioning, that a similar formula can be constructed for the case of the normalized interaction measure  $I \equiv (\varepsilon - 3 \cdot p)/T^4$ :

$$I_{\text{ren}}(T) = \bar{I}(T) + \frac{1}{16} \bar{I}(T/2) + \dots \quad (2.5)$$

where


$$\bar{I}(T) = N_t^4 \cdot d\beta/d\log a \cdot (\langle \text{PI} \rangle_{N_t} - \langle \text{PI} \rangle_{2 \cdot N_t})$$

The above method can also be easily generalized for the case of dynamical fermions.

We can demonstrate the applicability of this technique by using only the finite temperature  $N_t = 6$  and  $N_t = 8$  data of [1] to reproduce their results obtained by the standard method (which uses the more expensive  $T = 0$  data). Here we will use the ratio  $x = 6/8$  instead of the previously shown  $1/2$ . We keep only four terms in the sum (2.3), since the forthcoming terms are practically negligible:

$$\frac{p_{\text{ren}}}{T^4} = \bar{p}(T) + x^4 \cdot \bar{p}(xT) + \overbrace{x^8 \cdot \bar{p}(x^2T)}^{p'} + x^{12} \cdot \bar{p}(x^3T) \quad (2.6)$$

On figure 1. we show the results obtained by the standard renormalization procedure, and also those by our method.




**Figure 1:** The dimensionless pressure as a function of the temperature. By summing up the intermediate terms (indicated by solid blue lines, denoted by  $p'$ ) we arrive at the total renormalized pressure (solid red line), which agrees completely with results obtained by the usual integral method.

### 3. Setting the scale

Our new method is not only capable of reconstructing results of the standard procedure, but it also has a clear advantage over that, since  $T = 0$  lattices are not needed for the subtraction, only for scale determination. So we need large statistics only for  $T > 0$  simulations, which are less demanding in terms of computer resources (memory, CPU time). If we wanted to reach very high temperatures, we still needed large, and therefore expensive zero temperature lattices to set the scale. Alternatively, in the asymptotic scaling region, one might determine the scale using improved


perturbation theory. To verify that we are indeed in the asymptotic scaling region, we compare results of the Sommer parameter from [8], and a 3-loop improved perturbation theory formula [9] inserted into the expression of the lattice spacing  $a = \Lambda^{-1}(a \rightarrow 0)f_{\text{impPT}}(\beta)$ . This comparison on figure 2 shows that the last few lattice simulation points are in the asymptotic scaling regime, which means that an extrapolation with the perturbative formula might be trusted.



**Figure 2:** Asymptotic scaling seems to be realized, as the improved perturbation theory formula fits the lattice results of the Sommer parameter for larger values of  $\beta$ .

#### 4. Results I.


For our simulations we used tree level improved Symanzik action, and an overrelaxation-heatbath algorithm. The simulations were carried out on lattices with temporal extent  $N_t = 4$ ; we performed the subtraction on  $N_t = 8$  lattices. In order to account for decreasing screening masses, we used a rather large aspect ratio of  $N_s/N_t = 8$ . On figure 3. we show our results on the pressure.



**Figure 3:** We show the two, non-negligible terms (denoted by dotted lines) of the normalized pressure, and their sum (solid line).  $P_{SB}$  denotes the pressure of the non-interacting gluon gas.

It is worth mentioning here, that independently of the renormalization procedure, a problem emerges within the integral method framework. Since strictly speaking, the pressure is only exactly





**Figure 4:** Cancellation effect in calculating the pressure by our new method. These results were obtained on  $N_t = 4$  lattices at  $\beta = 50$  (which roughly corresponds to the Planck temperature).

## 6. Results II.

The method discussed above gives us the possibility to measure the pressure at very high temperatures. This was carried out using lattices with temporal extension  $N_t = 4$ ,  $N_t = 6$  and  $N_t = 8$ . The temperature interval ranged from  $4 \cdot T_C$  upto  $3 \cdot 10^7 \cdot T_C$ . Our results are shown on figure 5. A comparison can be done with the standard method, namely the result of [1] for smaller temperatures. Here we present the results which are closest to the continuum limit, thus  $N_t = 8$ . These results nicely follow the perturbative predictions (see [10, 11, 12], and also [13, 14]). However, more statistics is needed to determine the whole applicability region of the perturbative approach. It is important to point out, that at the highest temperature point, the pressure (within its statistical uncertainty) is already consistent with the Stefan-Boltzmann limit.


## 7. Conclusion

We presented two new methods to determine the equation of state for QCD. Compared to previous techniques, these methods allow to extend the temperature range by orders of magnitude. We developed a renormalization procedure of the pressure, for which no  $T = 0$  simulations are needed, and thus is less expensive than the methods, which were used previously. We also presented a new approach, which enables us to measure the pressure directly. Using this approach we gain a factor of  $N_t$  in CPU time with respect to the standard method. Our approach can also be used to set the integration constant, which provides a reference point for the pressure. Based on these new methods we presented first results on the pressure of the pure gauge theory at temperatures that could not have been reached before.

**Acknowledgements** This research was supported by the EU under grant I3HP and DFG under grant FO 502/1. We would like to acknowledge helpful discussions with M. Laine and Y. Schröder.

## References

- [1] G. Boyd, et al., *Thermodynamics of SU(3) Lattice Gauge Theory*, *Nucl. Phys.* **B469** (1996) 419-444 [hep-lat/9602007].



**Figure 5:** The pressure, normalized to its Stefan-Boltzmann value, as a function of the temperature obtained by our new technique. Results with smaller discretization errors ( $N_t = 8$ , blue circles), seem to fit improved perturbation theory, and also reproduce results obtained by the standard method at lower temperatures. At the highest temperature,  $3 \cdot 10^7 \cdot T_C$ , the pressure (within its statistical uncertainty) is consistent with the Stefan-Boltzmann limit.

- [2] **CP-PACS** collaboration, M. Okamoto et al., *Equation of state for pure SU(3) gauge theory with renormalization group improved action*, *Phys. Rev.* **D60** (1999) 094510 [hep-lat/9905005].
- [3] F. Karsch, E. Laermann and A. Peikert, *The pressure in 2, 2+1 and 3 flavour QCD*, *Phys. Lett.* **B478** (2000) 447-455 [hep-lat/0002003].
- [4] **MILC** collaboration, C. Bernard et al., *QCD equation of state with 2+1 flavors of improved staggered quarks*, *Phys. Rev.* **D75** (2007) 094505 [hep-lat/0611031].
- [5] Y. Aoki, Z. Fodor, S.D. Katz and K.K. Szabo, *The equation of state in lattice QCD: With physical quark masses towards the continuum limit*, *JHEP* **01** (2006) 089 [hep-lat/0510084].
- [6] F. Karsch, *Recent lattice results on finite temperature and density QCD*, In proceedings of XXV *International Symposium on Lattice Field Theory* (2007) PoS(LATTICE 2007) 015.
- [7] J. Engels, J. Fingberg, F. Karsch, D. Miller and M. Weber, *Nonperturbative thermodynamics of SU(N) gauge theories*, *Phys. Lett.* **B252** (1990) 625-630.
- [8] S. Necco and R. Sommer, *The  $N(f) = 0$  heavy quark potential from short to intermediate distances*, *Nucl. Phys.* **B622** (2002) 328-346 [hep-lat/0108008].
- [9] M. Gockeler et al., *A determination of the Lambda parameter from full lattice QCD*, *Phys. Rev.* **D73** (2006) 014513 [hep-ph/0502212].
- [10] E. Braaten A. Nieto, *Free Energy of QCD at High Temperature*, *Phys. Rev.* **D53** (1996) 3421-3437 [hep-ph/9510408].
- [11] K. Kajantie, M. Laine, K. Rummukainen and Y. Schroder, *The pressure of hot QCD up to  $g^{**6} \ln(1/g)$* , *Phys. Rev.* **D67** (2003) 105008 [hep-ph/0211321].
- [12] M. Laine and Y. Schroder, *Quark mass thresholds in QCD thermodynamics*, *Phys. Rev.* **D73** (2006) 085009 [hep-ph/0603048].
- [13] J. Blaizot, E. Iancu and A. Rebhan, *Thermodynamics of the high-temperature quark gluon plasma* (2003) [hep-ph/0303185].
- [14] J.O. Andersen, E. Braaten, E. Petitgirard and M. Strickland, *HTL perturbation theory to two loops*, *Phys. Rev.* **D66** (2002) 085016 [hep-ph/0205085].