

DIVISORS ON GRAPHS, CONNECTED FLAGS, AND SYZYGIES

FATEMEH MOHAMMADI AND FARBOD SHOKRIEH

ABSTRACT. We study the binomial and monomial ideals arising from linear equivalence of divisors on graphs from the point of view of Gröbner theory. We give an explicit description of a minimal Gröbner bases for each higher syzygy module. In each case the given minimal Gröbner bases is also a minimal generating set. The Betti numbers of the binomial ideal and its natural initial ideal coincide and they correspond to the number of “connected flags” in the graph. In particular the Betti numbers are independent of the characteristic of the base field. For complete graphs the problem was previously studied by Postnikov and Shapiro ([27]) and by Manjunath and Sturmfels ([19]). The case of a general graph was stated as an open problem.

1. INTRODUCTION

The theory of divisors on finite graphs can be viewed as a discrete version of the analogous theory on Riemann surfaces. This notion arises in different fields of research including the study of “abelian sandpiles” ([8, 12]), the study of component groups of Néron models of Jacobians of algebraic curves ([28, 16]), and the theory of chip-firing games on graphs ([5]). Riemann-Roch theory for finite graphs (and generalizations to tropical curves) is developed in this setting ([1, 13, 20]).

We are interested in the linear equivalence of divisors on graphs from the point of view of commutative algebra. Associated to every graph G there is a canonical binomial ideal I_G which encodes the linear equivalences of divisors on G . Let R denote the polynomial ring with one variable associated to each vertex. For any two effective divisors $D_1 \sim D_2$ one can write a binomial $\mathbf{x}^{D_1} - \mathbf{x}^{D_2}$. The ideal $I_G \subset R$ is generated by all such binomials. Two effective divisors are linearly equivalent if and only if their associated monomials are equal in R/I_G . This ideal is implicitly defined in Dhar’s seminal statistical physics paper [8]; R/I_G is the “operator algebra” defined there. To our knowledge, this ideal (more precisely, an affine piece of it) was first introduced in [7] to address computational questions in chip-firing dynamics using Gröbner bases. From a purely computational point of view, there are now much more efficient methods available (see, e.g., [2] and references therein). However, this ideal seems to encode a lot of interesting information about G and its linear systems. Some of the algebraic properties of I_G (and its generalization for directed graphs) are studied in [26]. Manjunath and Sturmfels [19] relate Riemann-Roch theory for finite graphs to Alexander duality in commutative algebra using this ideal.

In this paper, we study the syzygies and free resolutions of the ideals I_G and $\text{in}(I_G)$ from the point of view of Gröbner theory. Here $\text{in}(I_G)$ denotes the initial ideal with respect to a natural term order which is defined after distinguishing a vertex q (see Definition 2.4). When G is a complete graph, the syzygies and Betti numbers of the ideal $\text{in}(I_G)$ are studied

2010 *Mathematics Subject Classification.* 05C40, 13D02, 05E40, 13A02, 13P10.

Key words and phrases. Graphs, divisors, chip-firing, syzygy modules, Gröbner bases, free resolution, Betti numbers, connected flags.

by Postnikov and Shapiro [27]. Again for complete graphs, Manjunath and Sturmfels [19] study the ideal I_G and show that the Betti numbers coincide with the Betti numbers of $\text{in}(I_G)$. Finding minimal free resolutions for a general graph G was stated as an open problem in both [27] and [19] (also in [31, 26], where a conjecture is formulated). In particular, it was not known whether the Betti numbers for a general graph depend on the characteristic of the base field or not.

We explicitly construct free resolutions for both $\text{in}(I_G)$ and I_G for a general graph G using Schreyer’s algorithm. We remark that Schreyer’s algorithm “almost never” gives a *minimal* free resolution. However in our situation we are able to carefully order our combinatorial objects to enforce the minimality. As a result we describe, combinatorially, the minimal Gröbner bases for all higher syzygy modules of I_G and $\text{in}(I_G)$. In each case the minimal Gröbner bases is also a minimal generating set and the given resolution is minimal. This is shown by explicitly describing the differential maps in the constructed resolutions. In particular, the Betti numbers of $\text{in}(I_G)$ and I_G coincide. In other words, we have a positive answer to [6, Question 1.1] for I_G (see [22] and references therein, for other such examples). For a complete graph the minimal free resolution for $\text{in}(I_G)$ is nicely structured by a Scarf complex. The resolution for I_G when G is a tree is given by a Koszul complex since I_G is a complete intersection.

The description of the generating sets and the Betti numbers is in terms of the “connected flags” of G . Fix a vertex $q \in V(G)$ and an integer k . A *connected k -flag* of G (based at q) is a strictly increasing sequence $U_1 \subsetneq U_2 \subsetneq \cdots \subsetneq U_k = V(G)$ such that $q \in U_1$ and all induced subgraphs on vertex sets U_i and $U_{i+1} \setminus U_i$ are connected. Associated to any connected k -flag one can assign a “partial orientation” on G (Definition 4.3). Two connected k -flags are considered equivalent if the associated partially oriented graphs coincide. The Betti numbers correspond to the numbers of connected flags up to this equivalence. We give a bijective map between the connected flags of G and minimal Gröbner bases for higher syzygy modules of I_G and $\text{in}(I_G)$. For a complete graph all flags are connected and all distinct flags are inequivalent. So in this case the Betti numbers are simply the face numbers of the order complex of the poset of those subsets of $V(G)$ that contain q (ordered by inclusion). These numbers can be described using classical Stirling numbers (see Example 7.6). Hence our results directly generalize the analogous results in [27] and [19].

The paper is structured as follows. In §2 we fix our notation and provide the necessary background from the theory of divisors on graphs. We also define the ideal I_G and the natural $\text{Pic}(G)$ -grading and a term order $<$ on the polynomial ring relevant to our setting. In §3 we quickly recall some basic notions from commutative algebra. Our main goal is to fix our notation for Schreyer’s algorithm for computing higher syzygies, which is slightly different from what appears in the existing literature but is more convenient for our application. Also, to our knowledge Theorem 3.11, which gives a general sufficient criterion for an ideal to have the same graded Betti numbers as its initial ideal, has not appeared in the literature. In §4 we define connected flags and their equivalence relation. Basic properties of connected flags (up to equivalence) are studied in §4.4. In §5 we study the free resolution and higher syzygies of our ideals from the point of view of Gröbner theory, and in §6 we show that the constructed free resolutions are minimal. As a corollary we give our description of the graded Betti numbers in §7 and we describe some connections with the theory of reduced divisors.

Analogous results were obtained simultaneously (and independently) by Manjunath, Schreyer, and Wilmes in [18] using different techniques. Mania [17] gives an alternate proof for the expression of the first Betti number (see Remark 7.13(ii)). The constructed minimal free resolutions in this paper are in fact supported on certain cellular complexes. In [23] we describe this geometric picture for both I_G and $\text{in}(I_G)$, making precise connections with Lawrence and oriented matroid ideals of [3, 24]. Dochtermann and Sanyal have recently (independently) worked out this geometric picture in [9] for the monomial ideal $\text{in}(I_G)$.

Acknowledgements. The first author acknowledges support from the Mathematical Sciences Research Institute and the Alexander von Humboldt Foundation. She is grateful to H el ene Barcelo and Volkmar Welker for numerous comments and many helpful conversations. She would like to express her gratitude to Bernd Sturmfels for introducing her to this concept. The second author’s work was partially supported by NSF grants DMS-0901487. Part of this work was done while the second author was visiting UC Berkeley, which he would like to thank for the hospitality. We thank the referees for their very helpful suggestions and remarks. We would like to thank Lukas Katth an for pointing out the reference [15].

2. DEFINITIONS AND BACKGROUND

2.1. Graphs and divisors. Throughout this paper, a *graph* means a finite, connected, unweighted multigraph with no loops. As usual, the set of vertices and edges of a graph G are denoted by $V(G)$ and $E(G)$. We set $n = |V(G)|$. A *pointed graph* (G, q) is a graph together with a choice of a distinguished vertex $q \in V(G)$.

For a subset $S \subseteq V(G)$, we denote by $G[S]$ the induced subgraph of G with the vertex set S ; the edges of $G[S]$ are exactly the edges that appear in G over the set S . We use “ S is connected” and “ $G[S]$ is connected” interchangeably.

Let $\text{Div}(G)$ be the free abelian group generated by $V(G)$. An element of $\text{Div}(G)$ is written as $\sum_{v \in V(G)} a_v(v)$ and is called a *divisor* on G . The coefficient a_v in D is also denoted by $D(v)$. A divisor D is called *effective* if $D(v) \geq 0$ for all $v \in V(G)$. The set of effective divisors is denoted by $\text{Div}_+(G)$. We write $D \leq E$ if $E - D \in \text{Div}_+(G)$. For $D \in \text{Div}(G)$, let $\text{deg}(D) = \sum_{v \in V(G)} D(v)$. For $D_1, D_2 \in \text{Div}(G)$, the divisor $E = \max(D_1, D_2)$ is defined by $E(v) = \max(D_1(v), D_2(v))$ for $v \in V(G)$.

We denote by $\mathcal{M}(G)$ the group of integer-valued functions on the vertices. For $A \subseteq V(G)$, $\chi_A \in \mathcal{M}(G)$ denotes the $\{0, 1\}$ -valued characteristic function of A . The *Laplacian operator* $\Delta : \mathcal{M}(G) \rightarrow \text{Div}(G)$ is defined by

$$\Delta(f) = \sum_{v \in V(G)} \sum_{\{v, w\} \in E(G)} (f(v) - f(w))(v) .$$

The group of *principal divisors* is defined as the image of the Laplacian operator and is denoted by $\text{Prin}(G)$. It is easy to check that $\text{Prin}(G) \subseteq \text{Div}^0(G)$, where $\text{Div}^0(G)$ denotes the subgroup consisting of divisors of degree zero. The quotient $\text{Pic}^0(G) = \text{Div}^0(G)/\text{Prin}(G)$ is a finite group whose cardinality is the number of spanning trees of G (see, e.g., [2] and references therein). The full *Picard group* of G is defined as

$$\text{Pic}(G) = \text{Div}(G)/\text{Prin}(G)$$

which is isomorphic to $\mathbb{Z} \oplus \text{Pic}^0(G)$. Since principal divisors have degree zero, the map $\text{deg} : \text{Div}(G) \rightarrow \mathbb{Z}$ descends to a well-defined map $\text{deg} : \text{Pic}(G) \rightarrow \mathbb{Z}$. Two divisors D_1

and D_2 are called *linearly equivalent* if they become equal in $\text{Pic}(G)$. In this case we write $D_1 \sim D_2$. The *linear system* $|D|$ of D is defined as the set of effective divisors that are linearly equivalent to D .

To an ordered pair of *disjoint* subsets $A, B \subseteq V(G)$ we assign an effective divisor

$$(2.1) \quad D(A, B) = \sum_{v \in A} |\{w \in B : \{v, w\} \in E(G)\}|(v) .$$

In other words, the support of $D(A, B)$ is a subset of A and for $v \in A$ the coefficient of (v) in $D(A, B)$ is the number of edges between v and B . We define

$$d(A, B) = \deg(D(A, B))$$

which is the number of edges of G with one end in A and the other end in B . Although in general $D(A, B) \neq D(B, A)$ we always have $d(A, B) = d(B, A)$.

2.2. Divisors on graphs and the polynomial ring. Let K be a field and let $R = K[\mathbf{x}]$ be the polynomial ring in the n variables $\{x_v : v \in V(G)\}$. Any effective divisor D gives rise to a monomial

$$\mathbf{x}^D := \prod_{v \in V(G)} x_v^{D(v)} .$$

2.2.1. Gradings. For an abelian group \mathbf{A} , the polynomial ring R is said to be \mathbf{A} -graded (or graded by \mathbf{A}) if it is endowed with an \mathbf{A} -valued degree homomorphism $\deg_{\mathbf{A}} : \text{Div}(G) \rightarrow \mathbf{A}$. This is equivalent to fixing a semigroup homomorphism $\deg_{\mathbf{A}} : \text{Div}_+(G) \rightarrow \mathbf{A}$. Let $\deg_{\mathbf{A}}(\mathbf{x}^D) = \deg_{\mathbf{A}}(D)$. For $\mathbf{a} \in \mathbf{A}$ let $R_{\mathbf{a}}$ denote the K -vector space of homogeneous polynomials of degree \mathbf{a} . If there is no homogeneous polynomial of degree \mathbf{a} we let $R_{\mathbf{a}} = 0$.

There are three natural gradings of R in our setting:

- (i) $\mathbf{A} = \mathbb{Z}$ and $\deg_{\mathbf{A}}(\mathbf{x}^D) = \deg(D)$. This is the coarse \mathbb{Z} -grading of R .
- (ii) $\mathbf{A} = \text{Div}(G)$ and $\deg_{\mathbf{A}}(\mathbf{x}^D) = D$. This is the fine \mathbb{Z}^n -grading of R .
- (iii) $\mathbf{A} = \text{Pic}(G)$ and $\deg_{\mathbf{A}}(\mathbf{x}^D) = [D]$, where $[D]$ denotes the equivalence class of D in $\text{Pic}(G)$.

Gradings (i) and (ii) are, of course, well known. For the grading in (iii) (which is finer than the grading in (i) and is coarser than the grading in (ii)) we have the following lemma.

Lemma 2.1. *Let $\mathbf{A} = \text{Pic}(G)$ and $\deg_{\mathbf{A}}(\mathbf{x}^D) = [D]$ as above. Then $R_0 = K$ and, for each $\mathbf{a} \in \text{Pic}(G)$, the graded piece $R_{\mathbf{a}}$ is finite-dimensional.*

We remark that by [21, Theorem 8.6] the two conclusions in this lemma are, in general, equivalent.

Proof. For each $[D] \in \text{Pic}(G)$, the graded piece $R_{[D]}$ is spanned (as a K -vector space) by $\{\mathbf{x}^E : E \in |D|\}$ which is a finite set. This is because if $E \in |D|$, then, in particular, $\deg(E) = \deg(D)$. If $D \sim 0$ then $\deg(D) = 0$. So if D is effective we get $D = 0$. This means $R_0 = K$. \square

Let $R = \bigoplus_{\mathbf{a} \in \text{Pic}(G)} R_{\mathbf{a}}$ and $\mathfrak{m} = \bigoplus_{0 \neq \mathbf{a} \in \text{Pic}(G)} R_{\mathbf{a}}$. It follows from Lemma 2.1 that \mathfrak{m} is a maximal ideal of R . Consider the map $u := \deg \circ \deg_{\mathbf{A}} : \text{Div}(G) \rightarrow \mathbb{Z}$ sending D to $\deg([D]) = \deg(D)$. It is clear that u takes every nonzero element of $\text{Div}_+(G)$ to a strictly positive integer. Equivalently $u(r) \geq 1$ for every nonzero monomial. We can use this observation to prove that Nakayama's lemma holds for R with respect to the $\text{Pic}(G)$ -grading (see, e.g., [15, Proposition 1.4]).

Lemma 2.2. *Let R and \mathfrak{m} be as above. Then for every finitely generated $\text{Pic}(G)$ -graded module M such that $\mathfrak{m}M = M$ we have $M = 0$.*

Proof. Suppose $M \neq 0$. Write $M = \bigoplus_{\mathfrak{a} \in \text{Pic}(G)} M_{\mathfrak{a}}$. For any graded piece $M_{\mathfrak{a}}$ let $u(M_{\mathfrak{a}})$ denote the integer $u(m_{\mathfrak{a}})$ for any $m_{\mathfrak{a}} \in M_{\mathfrak{a}}$. Let $u(M) = \min_{\mathfrak{a} \in \text{Pic}(G)} u(M_{\mathfrak{a}})$. Since M is assumed to be finitely generated $u(M) > -\infty$. Since $u(r) \geq 1$ for all $r \in \mathfrak{m}$ we have $u(\mathfrak{m}M) > u(M)$ and therefore $\mathfrak{m}M \neq M$. \square

Because of Lemma 2.2 the notions of minimal generating set for (finitely generated) modules, minimal free resolution, and graded Betti numbers all make sense for the $\text{Pic}(G)$ -grading. Note that we need Lemma 2.2 since $\text{Pic}(G)$ is *not* in general torsion-free, and the $\text{Pic}(G)$ -grading of R is not a “positive multigrading” in the sense of [21, Definition 8.7].

2.2.2. The binomial ideal I_G . Associated to every graph G there is a canonical ideal which encodes the linear equivalences of divisors on G . This ideal is implicitly defined in Dhar’s seminal paper [8]. The ideal was introduced in [7] to address computational questions in chip-firing dynamics using Gröbner bases.

Definition 2.3.

$$\begin{aligned} I_G &:= \langle \mathbf{x}^{D_1} - \mathbf{x}^{D_2} : D_1 \sim D_2 \text{ both effective divisors} \rangle \\ &= \text{span}_K \{ \mathbf{x}^{D_1} - \mathbf{x}^{D_2} : D_1 \sim D_2 \text{ both effective divisors} \} . \end{aligned}$$

Clearly this ideal is graded (or homogeneous) with respect to the \mathbb{Z} and $\text{Pic}(G)$ gradings described in §2.2.1 ((i) and (iii)). It follows that the quotient R/I_G is both \mathbb{Z} -graded and $\text{Pic}(G)$ -graded as an R -module.

2.2.3. A natural term order. Once we fix a vertex q , there is a family of natural monomial orders that gives rise to a particularly nice Gröbner bases for I_G . This term order was first introduced in [7].

Fix a pointed graph (G, q) . Consider a total ordering of the set of variables $\{x_v : v \in V(G)\}$ compatible with the distances of vertices from q in G :

$$(2.2) \quad \text{dist}(w, q) < \text{dist}(v, q) \Rightarrow x_w < x_v .$$

Here, the distance between two vertices in a graph is the number of edges in a shortest path connecting them. The above ordering can be thought of an ordering on vertices induced by running the breadth-first search algorithm starting at the root vertex q .

Definition 2.4. We denote by $<$ the degree reverse lexicographic ordering on $R = K[\mathbf{x}]$ induced by the total ordering on the variables given in (2.2).

We remark that the choice of the vertex q is implicit in this notation.

Remark 2.5. The “total potential” functional $b_q(\cdot)$ from [2] is in the Gröbner cone of $<$. In fact it corresponds to the barycenter of this cone (see [23, §3.3 and §3.4]).

Throughout this paper $\text{in}(I_G)$ denotes the initial ideal of I_G with respect to this term order. Note that $\text{in}(I_G)$ is denoted by M_G in [27].

3. COMMUTATIVE ALGEBRA: SYZYGIES AND BETTI NUMBERS

In this section, we quickly recall some basic notions from commutative algebra. Our main goal is to fix our notation. A secondary goal is to keep the paper self-contained. Most of the material here is well known and we refer to standard books (e.g. [10, 14]) for proofs and more details. To our knowledge Theorem 3.11, which gives a general sufficient criterion for an ideal to have the same Betti numbers as its initial ideal, has not appeared in the literature.

Let K be any field and let $R = K[\mathbf{x}]$ be the polynomial ring in n variables graded by an abelian group A . The degree map will be denoted by \deg . Whenever we talk about notions like minimal generating sets, minimal free resolutions, or graded Betti numbers, we further assume that the grading is “nice” in the sense that Nakayama’s lemma holds and these notions are well defined. In this case we let \mathfrak{m} denote the corresponding maximal ideal of R consisting of nonunit elements. Examples of such nice gradings are all the gradings in §2.2.1 as well as “positive multigradings” in the sense of [21, Definition 8.7] (which generalizes the \mathbb{Z} and $\text{Div}(G)$ gradings, but not the $\text{Pic}(G)$ -grading).

3.1. Syzygies. Let F_{-1} be the free R -module generated by a finite set E . Elements of F_{-1} will be written as formal sums (with coefficients in R) of symbols $[e]$ (one symbol $[e]$ for each $e \in E$). Fix a *module ordering* $<_0$ on F_{-1} extending the monomial ordering $<$ on R . Recall that a module ordering on F_{-1} is a total ordering on the set of “monomials” $\mathbf{x}^\alpha[e]$ (for $\alpha \in \mathbb{N}^n$ and $e \in E$) extending a monomial ordering on R and compatible with the R -module structure. As usual LM will denote the *leading monomial* with respect to the associated ordering on monomials.

Let M be a graded submodule of F_{-1} . Assume that the finite totally ordered set (\mathbf{G}, \prec) forms a Gröbner bases for $(M, <_0)$ consisting of homogeneous elements. Let F_0 be the free module generated by \mathbf{G} . For $g \in \mathbf{G}$ we let the formal symbol $[g]$ denote the corresponding generator for F_0 ; each element of F_0 can be written as a sum of these formal symbols with coefficients in R .

There is a natural surjective homomorphism

$$\varphi_0 : F_0 \longrightarrow M \subseteq F_{-1}$$

sending $[g]$ to g for each $g \in \mathbf{G}$. Moreover, we force this homomorphism to be graded (or homogeneous of degree 0) by defining

$$\deg([g]) := \deg(g) \quad \text{for all } g \in \mathbf{G} .$$

By definition the syzygy module of M with respect to \mathbf{G} , denoted by $\text{syz}(\mathbf{G})$, is the kernel of this map. Let $\text{syz}_0(\mathbf{G}) := M$ and $\text{syz}_1(\mathbf{G}) := \text{syz}(\mathbf{G})$. For $i > 1$ the higher syzygy modules are defined as $\text{syz}_i(\mathbf{G}) := \text{syz}(\text{syz}_{i-1}(\mathbf{G}))$.

Remark 3.1. Since R is a graded ring, if \mathbf{G} is a *minimal* set of homogeneous generators of M then $\text{syz}(M) := \text{syz}(\mathbf{G})$ is well defined (i.e. independent of the choice of the generating set \mathbf{G}) up to a graded isomorphism.

3.2. Gröbner bases for syzygy modules. We now discuss a method to compute a Gröbner bases for $\text{syz}(\mathbf{G})$.

One can “pull back” the module ordering $<_0$ from F_{-1} along φ_0 to get a compatible module ordering $<_1$ on F_0 ; for $f, h \in \mathbf{G}$ define

$$(3.1) \quad \mathbf{x}^\beta[h] <_1 \mathbf{x}^\alpha[f] \Leftrightarrow \begin{cases} \text{LM}(\mathbf{x}^\beta h) <_0 \text{LM}(\mathbf{x}^\alpha f) \\ \text{or} \\ \text{LM}(\mathbf{x}^\beta h) = \text{LM}(\mathbf{x}^\alpha f) \quad \text{and} \quad f \prec h. \end{cases}$$

Note that both $<_0$ and $<_1$ extend the same monomial ordering $<$ on R . Also, the module ordering $<_1$ on F_0 depends on both $<_0$ on F_{-1} and on the totally ordered set (\mathbf{G}, \prec) .

To simplify the notation we assume the leading coefficients of all elements of \mathbf{G} are 1. Suppose we are given a pair of elements $f \prec h$ of \mathbf{G} such that

$$\text{LM}(f) = \mathbf{x}^{\alpha(f)}[e] \quad \text{and} \quad \text{LM}(h) = \mathbf{x}^{\alpha(h)}[e]$$

for some $e \in E$. Since \mathbf{G} is a Gröbner bases, setting $\gamma(f, h) := \max(\alpha(f), \alpha(h))$ (the entry-wise maximum), we have the “standard representation”:

$$(3.2) \quad \text{spoly}(f, h) = \mathbf{x}^{\gamma(f, h) - \alpha(f)} f - \mathbf{x}^{\gamma(f, h) - \alpha(h)} h = \sum_{g \in \mathbf{G}} a_g^{(f, h)} g$$

for some polynomials $a_g^{(f, h)} \in R$. We set

$$(3.3) \quad s(f, h) = \mathbf{x}^{\gamma(f, h) - \alpha(f)} [f] - \mathbf{x}^{\gamma(f, h) - \alpha(h)} [h] - \sum_{g \in \mathbf{G}} a_g^{(f, h)} [g] \in F_0 .$$

Since $f, h \in \mathbf{G}$ are by assumption homogeneous

$$\begin{aligned} \deg([f]) &= \deg(f) = \deg(\text{LM}(f)) = \deg(\alpha(f)) + \deg([e]) , \\ \deg([h]) &= \deg(h) = \deg(\text{LM}(h)) = \deg(\alpha(h)) + \deg([e]) . \end{aligned}$$

It follows that $s(f, h)$ is also homogeneous and its degree is equal to $\deg(\gamma(f, h)) + \deg([e])$. Also, by definition $s(f, h) \in \text{syz}(\mathbf{G})$. More is true:

Theorem 3.2 (Schreyer [29]). *The set*

$\mathcal{S}(\mathbf{G}) = \{s(f, h) : f, h \in \mathbf{G}, f \prec h, \text{LM}(f) = \mathbf{x}^{\alpha(f)}[e], \text{LM}(h) = \mathbf{x}^{\alpha(h)}[e] \text{ for some } e \in E\}$
forms a homogeneous Gröbner bases for $(\text{syz}(\mathbf{G}), <_1)$.

Both the module ordering $<_1$ and the Gröbner bases $\mathcal{S}(\mathbf{G})$ depend on $<_0$ and on (\mathbf{G}, \prec) .

Lemma 3.3. *With respect to $<_1$ we have $\text{LM}(s(f, h)) = \mathbf{x}^{\gamma(f, h) - \alpha(f)} [f]$.*

Proof. From the “standard representation” (3.2) we know

$$\text{LM}\left(\sum_{g \in \mathbf{G}} a_g^{(f, h)} g\right) = \text{LM}(\mathbf{x}^{\gamma(f, h) - \alpha(f)} f - \mathbf{x}^{\gamma(f, h) - \alpha(h)} h) <_0 \text{LM}(\mathbf{x}^{\gamma(f, h) - \alpha(f)} f)$$

and therefore from (3.1) we obtain

$$\text{LM}\left(\sum_{g \in \mathbf{G}} a_g^{(f, h)} [g]\right) <_1 \mathbf{x}^{\gamma(f, h) - \alpha(f)} [f] .$$

Moreover we have

$$\text{LM}(\mathbf{x}^{\gamma(f, h) - \alpha(h)} h) = \text{LM}(\mathbf{x}^{\gamma(f, h) - \alpha(f)} f) \quad \text{and} \quad f \prec h .$$

Again (3.1) implies

$$\mathbf{x}^{\gamma(f, h) - \alpha(h)} [h] <_1 \mathbf{x}^{\gamma(f, h) - \alpha(f)} [f] . \quad \square$$

To read the Betti numbers for M one needs to find a *minimal generating set* for the syzygy modules. In general the set $\mathcal{S}(\mathbf{G})$ is far from being even a *minimal Gröbner bases*. One criterion to detect some of the redundant bases elements is given in the following lemma.

Lemma 3.4. *Let $\mathcal{S}(\mathbf{G})$ be as in Theorem 3.2. Let $f_1 \prec f_2$ and $f_1 \prec f_3$ and $\text{LM}(f_i) = \mathbf{x}^{\alpha(f_i)}[e]$ for some $e \in E$ and $1 \leq i \leq 3$. If $\alpha(f_2) \leq \gamma(f_1, f_3)$ then $\mathcal{S}(\mathbf{G}) \setminus \{s(f_1, f_3)\}$ is already a Gröbner bases for $(\text{syz}(\mathbf{G}), <_1)$.*

Proof. The inequality implies $\gamma(f_1, f_2) - \alpha(f_1) \leq \gamma(f_1, f_3) - \alpha(f_1)$ which means

$$\text{LM}(s(f_1, f_2)) \mid \text{LM}(s(f_1, f_3)) . \quad \square$$

Remark 3.5. By repeatedly applying Lemma 3.4 we can find a subset $\mathcal{S}_{\min}(\mathbf{G})$ of $\mathcal{S}(\mathbf{G})$ which has the following properties:

- (1) $\mathcal{S}_{\min}(\mathbf{G})$ forms a Gröbner bases for $(\text{syz}(\mathbf{G}), <_1)$,
- (2) there are no pair of elements $s(f, h), s(f, g) \in \mathcal{S}_{\min}(\mathbf{G})$ such that

$$\text{LM}(s(f, h)) \mid \text{LM}(s(f, g)) .$$

In other words (see Lemma 3.3) $\mathcal{S}_{\min}(\mathbf{G})$ is a *minimal Gröbner bases* for $(\text{syz}(\mathbf{G}), <_1)$.

3.3. Free resolutions from Gröbner theory. One can use Theorem 3.2 to construct a graded free resolution of M by induction on the homological degree. We summarize this procedure in Algorithm 1 which is due to Schreyer [29] (also Spear [30], see, e.g., [10]).

It follows immediately from definitions that the output sequence is exact and that we obtain a free resolution in this way. We note that the constructed free resolution is in general not minimal.

Remark 3.6. The map $\varphi_{i+1} : F_{i+1} \rightarrow F_i$ can be described more explicitly. Since $u \in \mathbf{G}_{i+1} = \mathcal{S}_{\min}(\mathbf{G}_i)$ there are two elements $f \prec h$ in \mathbf{G}_i such that

$$u = s(f, h) = \mathbf{x}^{\gamma(f, h) - \alpha(f)}[f] - \mathbf{x}^{\gamma(f, h) - \alpha(h)}[h] - \sum_{g \in \mathbf{G}_i} a_g^{(f, h)}[g] .$$

In other words

$$(3.4) \quad \varphi_{i+1}([u]) = \mathbf{x}^{\gamma(f, h) - \alpha(f)}[f] - \mathbf{x}^{\gamma(f, h) - \alpha(h)}[h] - \sum_{g \in \mathbf{G}_i} a_g^{(f, h)}[g] .$$

Since $\{[u] : u \in \mathbf{G}_{i+1}\}$ is the set of bases elements for F_{i+1} and $\{[g] : g \in \mathbf{G}_i\}$ is the set of bases elements for F_i , the set of equalities in (3.4), as u runs through the set \mathbf{G}_i , determines the map φ_{i+1} completely. If we fix a labeling for the elements of $(\mathbf{G}_{i+1}, \prec_{i+1})$ and (\mathbf{G}_i, \prec_i) we can write down the corresponding matrix for φ_{i+1} from this data.

Remark 3.7. Although any total ordering \prec_i on the sets \mathbf{G}_i would work in Algorithm 1, it follows from Lemma 3.3 and Lemma 3.4 that the “quality of output” very much depends on the choice of these total orderings; how far the free resolution produced by the algorithm is from being minimal depends on the choice of the total ordering in a crucial way.

Input:

Graded polynomial ring $R = K[\mathbf{x}]$,
 Monomial ordering $<$ on R ,
 Free R -module F_{-1} generated by formal symbols $\{[e]\}_{e \in E}$,
 Graded R -submodule M of F_{-1} ,
 Module ordering $<_0$ on F_{-1} extending the monomial ordering $<$,
 Finite set \mathbf{G} forming a homogeneous Gröbner bases for $(M, <_0)$.

Output:

A graded free resolution: $\cdots \rightarrow F_i \xrightarrow{\varphi_i} F_{i-1} \rightarrow \cdots \rightarrow F_0 \xrightarrow{\varphi_0} M \rightarrow 0$.

Initialization:

$\mathbf{G}_0 := \mathbf{G}$;

$F_0 :=$ free R -module generated by formal symbols $\{[g]\}_{g \in \mathbf{G}_0}$; **Output** F_0 ;

$\varphi_0 : F_0 \rightarrow M \subseteq F_{-1}$ defined by $[g] \mapsto g$ for each $g \in \mathbf{G}_0$; **Output** φ_0 ;

$i = 0$;

while $F_i \neq 0$ **do**

$<_i$: arbitrary total ordering on \mathbf{G}_i ;

$<_{i+1}$: module ordering on F_i obtained from $<_i$ on F_{i-1} (as in (3.1)) ;

$\mathbf{G}_{i+1} := \mathcal{S}_{\min}(\mathbf{G}_i) \subset F_i$, a minimal Gröbner bases of $(\text{syz}_{i+1}(\mathbf{G}), <_{i+1})$ (as in Theorem 3.2 and Remark 3.5) ;

$F_{i+1} :=$ free R -module generated by formal symbols $\{[u]\}_{u \in \mathbf{G}_{i+1}}$; **Output** F_{i+1} ;

$\varphi_{i+1} : F_{i+1} \rightarrow F_i$ defined by $[u] \mapsto u$ for each $u \in \mathbf{G}_{i+1}$; **Output** φ_{i+1} ;

$i \leftarrow i + 1$;

end

Algorithm 1: Algorithm for computing a free resolution of M (Schreyer's algorithm)

3.4. Minimal free resolutions and Betti numbers. Let R be a graded ring and M be a graded R -module. Assume that

$$\mathcal{F} : 0 \rightarrow \cdots \rightarrow F_i \xrightarrow{\varphi_i} F_{i-1} \rightarrow \cdots \rightarrow F_0 \xrightarrow{\varphi_0} M \rightarrow 0$$

is a *minimal* graded free resolution (i.e., a graded free resolution such that $\varphi_{i+1}(F_{i+1}) \subseteq \mathfrak{m}F_i$ for all $i \geq 0$). The i -th *Betti number* $\beta_i(M)$ of M is by definition the rank of F_i . The i -th *graded Betti number* in degree $j \in \mathbf{A}$, denoted by $\beta_{i,j}(M)$, is the rank of the degree j part of F_i . It is a consequence of Nakayama's lemma for graded rings that any finitely generated graded R -module has minimal free resolution, and that the numbers $\beta_{i,j}(M)$ and $\beta_i(M)$ are independent of the choice of the minimal resolution.

If \mathbf{G} is a minimal set of homogeneous generators of M then $\text{syz}(M) := \text{syz}(\mathbf{G})$ is well defined up to graded isomorphism. Similarly, setting $\text{syz}_0(M) := M$, the i -th syzygy modules $\text{syz}_i(M) := \text{syz}(\text{syz}_{i-1}(M))$ are well defined for all $i \geq 0$. In this case the i -th Betti number $\beta_i(M)$ is also the minimal number of generators of $\text{syz}_i(M)$ and the graded Betti number $\beta_{i,j}(M)$ is the minimal number of generators of the i -th syzygy module $\text{syz}_i(M)$ in degree j .

Theorem 3.8. *Suppose that \mathcal{F} is a minimal graded free resolution of M . Fix an $i \geq 0$. Let E_i denote a bases for the free module F_i . Then $\{\varphi_i(f) : f \in E_i\}$ is a minimal system of homogeneous generators of $\text{syz}_i(M)$.*

For a proof see, for example, [25, Theorem 10.2].

Remark 3.9. It follows from Theorem 3.8 that if for all $i \geq 1$ and all $u \in \mathbf{G}_i$ the coefficients appearing in the expression (3.4) of $\varphi_i([u])$ in terms of $\{[g] : g \in \mathbf{G}_{i-1}\}$ are all nonunit elements of R (i.e. if they belong to the ideal \mathfrak{m}), then the resolution is a *minimal* free resolution of M . In this case it follows from Theorem 3.8 that the sets \mathbf{G}_i are *minimal generating sets* of $\text{syz}_i(M)$ for all $i \geq 0$.

3.5. Betti numbers of M and $\text{in}(M)$. For a module ordering $<_0$ on F_{-1} let $\text{in}(M)$ denote the “leading module” (i.e. the module generated by leading monomials) of M with respect to $<_0$. The following theorem is well known and is a consequence of the fact that passing to $\text{in}(M)$ is a flat deformation (see, e.g., [21, Theorem 8.29]).

Theorem 3.10 (Upper-semicontinuity). $\beta_{i,j}(M) \leq \beta_{i,j}(\text{in}(M))$ for all $i \geq 0$ and $j \in \mathbf{A}$.

In the setting of Algorithm 1, there is a general sufficient condition for equality to hold. The following result gives a general criterion guaranteeing that, if it is satisfied, then the answer of [6, Question 1.1] is positive.

Theorem 3.11. *If the output of Algorithm 1 is a minimal graded free resolution then $\beta_{i,j}(M) = \beta_{i,j}(\text{in}(M))$ for all $i \geq 0$ and $j \in \mathbf{A}$.*

Proof. Let $\mathbf{G}' = \{\text{LM}(g) : g \in \mathbf{G}\} \subset \text{in}(M)$. Since \mathbf{G} forms a minimal Gröbner bases of $(M, <_0)$ the map

$$\pi_0 : \mathbf{G} \rightarrow \mathbf{G}' \subset \text{in}(M) \quad \text{with} \quad \pi_0(g) := \text{LM}(g) \quad \text{for} \quad g \in \mathbf{G}$$

is a bijection between \mathbf{G} and \mathbf{G}' . The proof for $i \geq 0$ is by induction on i . We show that for each $i \geq 0$ there is a free module F'_i with bases elements

$$[s(\pi_{i-1}(f), \pi_{i-1}(h))]$$

corresponding to bases elements $[s(f, h)]$ of \mathbf{G}_i . This extends to a natural bijective map from F_i to F'_i . Moreover, this bijection induces maps

$$\pi_i : \mathbf{G}_i \rightarrow \text{syz}_i(\mathbf{G}') \quad \text{with} \quad \pi_i(s(f, h)) := s(\pi_{i-1}(f), \pi_{i-1}(h)) \quad \text{for} \quad f \prec h \text{ in } \mathbf{G}_{i-1}$$

such that

- (1) $<_i$ is a term order on F'_i .
- (2) $\text{LM}(\pi_i(s(f, h))) = \mathbf{x}^\theta[\pi_{i-1}(f)]$, where $\text{LM}(s(f, h)) = \mathbf{x}^\theta[f]$.
- (3) $\pi_i(\mathbf{G}_i)$ forms a minimal Gröbner bases of $(\text{syz}_i(\mathbf{G}'), <_i)$.

We have already shown the case $i = 0$. Now assume that $i > 0$ and the result holds for $i - 1$. Note that by the induction hypothesis π_{i-1} is injective. This together with (2) and (3) for $i - 1$ implies that the elements $[s(\pi_{i-1}(f), \pi_{i-1}(h))]$ are pairwise distinct, since their leading terms are pairwise distinct. Assume that $<_{i-1}$ is the term order on F_{i-1} . By the induction hypothesis we have the total order \prec'_{i-1} on the elements $\pi_{i-1}(\mathbf{G}_i)$ such that

$$(3.5) \quad \pi_{i-1}(f) \prec'_{i-1} \pi_{i-1}(h) \quad \text{if and only if} \quad f \prec_{i-1} h .$$

Since (2) and (3) hold for $i - 1$, choosing the total order of (3.5) on the elements $\pi_{i-1}(\mathbf{G}_i)$ we will get the term order $<_i$ on F'_i by (3.1). To prove (2) for i , let $s(f, h)$ be an element of \mathbf{G}_i with

$$\text{LM}(f) = \mathbf{x}^{\alpha(f)}[u] , \quad \text{LM}(h) = \mathbf{x}^{\alpha(h)}[u] \quad \text{and} \quad \gamma(f, h) = \max(\alpha(f), \alpha(h))$$

for some bases element $[u] \in \mathbf{G}_{i-2}$. Then by the induction hypothesis we have

$$\text{LM}(\pi_{i-1}(f)) = \mathbf{x}^{\alpha(f)}[\pi_{i-2}(u)] \quad \text{and} \quad \text{LM}(\pi_{i-1}(h)) = \mathbf{x}^{\alpha(h)}[\pi_{i-2}(u)] .$$

Therefore Lemma 3.3 together with induction hypothesis implies

$$\text{LM}(\pi_i(s(f, h))) = \text{LM}(s(\pi_{i-1}(f), \pi_{i-1}(h))) = \mathbf{x}^{\gamma(f, h) - \alpha(f)}[\pi_{i-1}(f)]$$

which is (2). Now it follows that π_i is injective; assume that $\pi_i(s(f, h)) = \pi_i(s(p, q))$. Therefore $\text{LM}(s(f, h)) = \text{LM}(s(p, q)) = \mathbf{x}^{\gamma(f, h) - \alpha(f)}[\pi_{i-1}(f)]$. Now (2) implies that

$$\text{LM}(s(f, h)) = \text{LM}(s(p, q)) = \mathbf{x}^{\gamma(f, h) - \alpha(f)}[f] ,$$

which is a contradiction by our assumption on $\mathbf{G}_i = \mathcal{S}_{\min}(\mathbf{G}_{i-1})$. The fact that π_i is injective implies that its extension from F_i to F'_i is a bijective map, as desired.

Now we show that $\pi_i(\mathbf{G}_i)$ forms a Gröbner bases for $(\text{syz}_i(\mathbf{G}'), <_i)$. For the sake of contradiction, assume that $\pi_i(\mathbf{G}_i)$ does not form a Gröbner bases for $(\text{syz}_i(\mathbf{G}'), <_i)$. Then our induction hypothesis that $\pi_{i-1}(\mathbf{G}_{i-1})$ forms a Gröbner bases for $(\text{syz}_{i-1}(\mathbf{G}'), <_{i-1})$, implies that there exist elements f and h of \mathbf{G}_{i-1} such that $f \prec_{i-1} h$ and $\text{LM}(s(\pi_{i-1}(f), \pi_{i-1}(h)))$ is not divisible by the leading monomial of any element of $\pi_i(\mathbf{G}_i)$. On the other hand, our assumption on \mathbf{G}_i and the fact that $\text{LM}(s(\pi_{i-1}(f), \pi_{i-1}(h))) = \text{LM}(s(f, h))$ imply that there exists an element g in \mathbf{G}_{i-1} such that $f \prec_{i-1} g$ and $\text{LM}(s(f, g))$ divides $\text{LM}(s(f, h))$. By the induction hypothesis $\pi_{i-1}(f)$ and $\pi_{i-1}(g)$ belong to $\text{syz}_{i-1}(\mathbf{G}')$. Moreover by (2) $\text{LM}(s(\pi_{i-1}(f), \pi_{i-1}(g))) = \text{LM}(s(f, g))$ divides $\text{LM}(s(\pi_{i-1}(f), \pi_{i-1}(h)))$ which is a contradiction. Thus $\pi_i(\mathbf{G}_i)$ forms a Gröbner bases for $(\text{syz}_i(\mathbf{G}'), <_i)$.

Note that π_i is a graded map of degree zero which preserves the degree of the elements $s(f, h)$ of \mathbf{G}_i . This together with the fact that $\pi_i(\mathbf{G}_i)$ forms a minimal Gröbner bases of $(\text{syz}_i(\mathbf{G}'), <_i)$ implies that $\beta_{i,j}(\text{in}(M)) \leq \beta_{i,j}(M)$. Now Theorem 3.10 completes the proof. \square

4. CONNECTED FLAGS ON GRAPHS

4.1. Connected flags, partial orientations, and divisors. From now on we fix a pointed graph (G, q) and we let $n = |V(G)|$. Consider the poset

$$\mathfrak{C}(G, q) := \{U \subseteq V(G) : q \in U\}$$

ordered by inclusion. The following special chains of this poset arise naturally in our setting.

Definition 4.1. Fix an integer $1 \leq k \leq n$. A *connected k -flag* of (G, q) is a (strictly increasing) sequence \mathcal{U} of subsets of $V(G)$

$$U_1 \subsetneq U_2 \subsetneq \cdots \subsetneq U_k = V(G)$$

such that $q \in U_1$ and, for all $1 \leq i \leq k - 1$, both $G[U_i]$ and $G[U_{i+1} \setminus U_i]$ are connected.

The set of all connected k -flags of (G, q) will be denoted by $\mathfrak{F}_k(G, q)$.

Remark 4.2. For a complete graph, $\mathfrak{F}_k(G, q)$ is simply the order complex of $\mathfrak{C}(G, q)$, but in general $\mathfrak{F}_k(G, q)$ is not a simplicial complex.

Notation. For convenience, whenever we use index 0 on a vertex set (e.g. U_0, V_0, W_0 , etc.) we mean the empty set.

Definition 4.3. Given $\mathcal{U} \in \mathfrak{F}_k(G, q)$ we define:


- (a) a “partial orientation” of G by orienting edges *from* U_i to $U_{i+1} \setminus U_i$ (for all $1 \leq i \leq k-1$) and leaving all other edges unoriented. We denote the resulting partially oriented graph by $G(\mathcal{U})$.
- (b) an effective divisor $D(\mathcal{U}) \in \text{Div}(G)$ given by (see (2.1)) $D(\mathcal{U}) := \sum_{i=1}^{k-1} D(U_{i+1} \setminus U_i, U_i)$.

Note that the partial orientation in (a) is always acyclic.

Example 4.4. Let G be the following graph on the vertices v_1, v_2, \dots, v_5 . We let v_1 be the distinguished vertex. Then the partial orientation associated to

$$\mathcal{U} : \{v_1\} \subset \{v_1, \mathbf{v}_2\} \subset \{v_1, v_2, \mathbf{v}_3, \mathbf{v}_4\} \subset \{v_1, v_2, v_3, v_4, \mathbf{v}_5\}$$

is depicted in the following figure. Note that $D(\mathcal{U}) = (v_2) + (v_3) + (v_4) + 2(v_5)$.



Remark 4.5. It is easy to check that $D(\mathcal{U}) = \sum_{v \in V(G)} (\text{indeg}_{G(\mathcal{U})}(v))(v)$, where $\text{indeg}_{G(\mathcal{U})}(v)$ denotes the number of oriented edges directed *to* v in $G(\mathcal{U})$.

4.2. Total ordering on $\mathfrak{F}_k(G, q)$. We endow each $\mathfrak{F}_k(G, q)$ with a *total ordering* \prec_k for all $1 \leq k \leq n$.

Let \preceq denote the ordering on $\mathfrak{C}^{\text{op}}(G, q)$ (the opposite poset of $\mathfrak{C}(G, q)$) given by reverse inclusion:

$$U \preceq V \iff U \supseteq V .$$

Definition 4.6. We fix, once and for all, a *total ordering* extending \preceq . By a slight abuse of notation, \preceq will be used to denote this total ordering extension. In particular, \prec will denote the associated strict total order.

We consider one of the natural “lexicographic extensions” (more precisely, the reverse lexicographic extension) of \prec to the set of connected k -flags.

Definition 4.7. For $\mathcal{U} \neq \mathcal{V}$ in $\mathfrak{F}_k(G, q)$ written as

$$\mathcal{U} : U_1 \subsetneq U_2 \subsetneq \dots \subsetneq U_k = V(G)$$

$$\mathcal{V} : V_1 \subsetneq V_2 \subsetneq \dots \subsetneq V_k = V(G)$$

we say $\mathcal{U} \prec_k \mathcal{V}$ if for the maximum $1 \leq \ell \leq k-1$ with $U_\ell \neq V_\ell$ we have $U_\ell \prec V_\ell$.

As usual, we write $\mathcal{U} \preceq_k \mathcal{V}$ if and only if $\mathcal{U} \prec_k \mathcal{V}$ or $\mathcal{U} = \mathcal{V}$.

Lemma 4.8. $(\mathfrak{F}_k(G, q), \preceq_k)$ is a totally ordered set.

Proof. Let

$$\begin{aligned}\mathcal{U} &: U_1 \subsetneq U_2 \subsetneq \cdots \subsetneq U_k = V(G) , \\ \mathcal{V} &: V_1 \subsetneq V_2 \subsetneq \cdots \subsetneq V_k = V(G) , \\ \mathcal{W} &: W_1 \subsetneq W_2 \subsetneq \cdots \subsetneq W_k = V(G) .\end{aligned}$$

If $\mathcal{U} \neq \mathcal{V}$ then there is an index $1 \leq \ell \leq k - 1$ with $U_\ell \neq V_\ell$. Since \prec is a strict total ordering we have either $U_\ell \prec V_\ell$ or $V_\ell \prec U_\ell$. It follows from the definition that if $\mathcal{U} \neq \mathcal{V}$ then either $\mathcal{U} \prec_k \mathcal{V}$ or $\mathcal{V} \prec_k \mathcal{U}$ (i.e., \prec_k is trichotomous). It remains to show that \prec_k is transitive. Assume that $\mathcal{U} \prec_k \mathcal{V}$ and $\mathcal{V} \prec_k \mathcal{W}$. Let $1 \leq \ell_1 \leq k - 1$ and $1 \leq \ell_2 \leq k - 1$ be such that

$$\begin{aligned}U_{\ell_1} \prec V_{\ell_1} \text{ and } U_i = V_i \text{ for } \ell_1 < i \leq k , \\ V_{\ell_2} \prec W_{\ell_2} \text{ and } V_i = W_i \text{ for } \ell_2 < i \leq k .\end{aligned}$$

There are three cases:

- If $\ell_1 = \ell_2$ we have $U_{\ell_1} \prec V_{\ell_1} \prec W_{\ell_1}$ and $U_i = W_i$ for $\ell_1 < i \leq k$,
- If $\ell_1 < \ell_2$ we have $U_{\ell_2} = V_{\ell_2} \prec W_{\ell_2}$ and $U_i = W_i$ for $\ell_2 < i \leq k$,
- If $\ell_1 > \ell_2$ we have $U_{\ell_1} \prec V_{\ell_1} = W_{\ell_1}$ and $U_i = W_i$ for $\ell_1 < i \leq k$.

Therefore in any case $\mathcal{U} \prec_k \mathcal{W}$. \square

4.3. Equivalence relation on $\mathfrak{F}_k(G, q)$. It is easy to find two different connected k -flags having identical associated partially oriented graphs.

Example 4.9. If the connected k -flag

$$\mathcal{U} : U_1 \subsetneq \cdots \subsetneq U_{i-1} \subsetneq (U_{i-1} \cup A_i) \subsetneq (U_{i-1} \cup A_i \cup A_{i+1}) \subsetneq \cdots \subsetneq U_k$$

is such that A_{i+1} is disjoint from A_i , and $d(A_{i+1}, A_i) = 0$ (i.e., A_{i+1} is not connected to A_i) then


$$\mathcal{V} : U_1 \subsetneq \cdots \subsetneq U_{i-1} \subsetneq (U_{i-1} \cup A_{i+1}) \subsetneq (U_{i-1} \cup A_{i+1} \cup A_i) \subsetneq \cdots \subsetneq U_k$$

is a different connected k -flag, but $G(\mathcal{U})$ and $G(\mathcal{V})$ coincide.

Example 4.10. Let G be the following graph on the vertices v_1, v_2, \dots, v_5 . We fix v_1 as the distinguished vertex. Consider the following connected flags:

- $\mathcal{U} : \{v_1\} \subset \{v_1, \mathbf{v}_2\} \subset \{v_1, v_2, \mathbf{v}_3\} \subset \{v_1, v_2, v_3, \mathbf{v}_4, \mathbf{v}_5\}$
- $\mathcal{V} : \{v_1\} \subset \{v_1, \mathbf{v}_3\} \subset \{v_1, \mathbf{v}_2, v_3\} \subset \{v_1, v_2, v_3, \mathbf{v}_4, \mathbf{v}_5\}$.

Then, as we see, their associated graphs coincide.



This example motivates the following definition.

Definition 4.11. Two k -flags $\mathcal{U}, \mathcal{V} \in \mathfrak{F}_k(G, q)$ are called *equivalent* if the associated partially oriented graphs $G(\mathcal{U})$ and $G(\mathcal{V})$ coincide.

Remark 4.12.

- (i) This equivalence relation is easily seen to be the transitive closure of the equivalences described in Example 4.9.
- (ii) When $k = 2$ each equivalence class contains a unique element.

Notation. The set of all equivalence classes in $\mathfrak{F}_k(G, q)$ will be denoted by $\mathfrak{E}_k(G, q)$. The set $\mathfrak{S}_k(G, q)$ will denote the set of minimal representatives of the classes in $\mathfrak{E}_k(G, q)$ with respect to \prec_k .

Example 4.13. Here we list all acyclic (partial) orientations associated to equivalence classes of connected flags on the 4-cycle graph on the vertices 1, 2, 3, 4. The vertex 1 is chosen as the distinguished vertex (see Figures 1, 2, 3).




FIGURE 1. Acyclic orientations corresponding to 2-partitions of C_4




FIGURE 2. Acyclic orientations corresponding to 3-partitions of C_4

4.4. Main properties of $\mathfrak{S}_k(G, q)$. This section, while elementary, is the most technical part of the paper. The reader is invited to draw graphs and Venn diagrams to follow the proofs. The main ingredients needed for our main theorems are Definition 4.16, Proposition 4.18, Proposition 4.20, Corollary 4.21, Lemma 4.24, and Proposition 4.25. Other results are used to establish these main ingredients.




FIGURE 3. Acyclic orientations corresponding to 4-partitions of C_4

Lemma 4.14. *Let*

$$\mathcal{U} : U_1 \subsetneq U_2 \subsetneq \cdots \subsetneq V(G)$$

be an element of $\mathfrak{S}_k(G, q)$. Let

$$\mathcal{W} : W_1 \subsetneq W_2 \subsetneq \cdots \subsetneq V(G)$$

be an element of $\mathfrak{F}_{k'}(G, q)$.

- (a) *If $k' = k - 1$ and $W_i = U_{i+1}$ for $1 \leq i \leq k - 1$ then $\mathcal{W} \in \mathfrak{S}_{k-1}(G, q)$.*
- (b) *If $k' = k - 1$ and $W_1 = U_1 \cup (U_3 \setminus U_2)$ and $W_i = U_{i+1}$ for $2 \leq i \leq k - 1$ then $\mathcal{W} \in \mathfrak{S}_{k-1}(G, q)$.*
- (c) *If $k' = k$ and $W_1 \subseteq U_1$ and $W_i = U_i$ for $2 \leq i \leq k$ then $\mathcal{W} \in \mathfrak{S}_k(G, q)$.*

Proof. The strategy of the proof is similar for all three parts. Namely in each case, for the sake of contradiction, we assume that

$$\mathcal{W}' : W'_1 \subsetneq W'_2 \subsetneq \cdots \subsetneq V(G)$$

is another connected flag equivalent to \mathcal{W} which precedes \mathcal{W} in the total ordering. We will then find a connected flag \mathcal{U}' equivalent to \mathcal{U} such that $\mathcal{U}' \prec_k \mathcal{U}$.

We first note that in all cases, since $G(\mathcal{W})$ and $G(\mathcal{W}')$ coincide and $q \in W_1, W'_1$, we must have $W_1 = W'_1$. Let $\ell \geq 2$ be such that $W'_\ell \prec W_\ell$ and $W'_i = W_i$ for $\ell < i$.

(a) The ordered collection $(W'_j \setminus W'_{j-1})_{j=2}^{k-1}$ is a permutation of the ordered collection $(W_j \setminus W_{j-1})_{j=2}^{k-1} = (U_j \setminus U_{j-1})_{j=3}^k$. Also $W'_1 \setminus U_1 = W_1 \setminus U_1 = U_2 \setminus U_1$. It follows that

$$\mathcal{U}' : U_1 \subsetneq W'_1 \subseteq W'_2 \subseteq \cdots \subsetneq W'_{k-2} \subsetneq W'_{k-1} = V(G)$$

is a connected k -flag equivalent to \mathcal{U} . But then $W'_\ell \prec W_\ell$ and $W'_i = W_i$ for $\ell < i \leq k - 1$ implies $\mathcal{U}' \prec_k \mathcal{U}$, a contradiction.

(b) The ordered collection $(W'_j \setminus W'_{j-1})_{j=2}^{k-1}$ is a permutation of the ordered collection $(W_j \setminus W_{j-1})_{j=2}^{k-1} = (U_j \setminus U_{j-1})_{j=2,4,5,\dots,k}$; here we used $W_2 \setminus W_1 = U_3 \setminus (U_1 \cup (U_3 \setminus U_2)) = U_2 \setminus U_1$. Also $W'_1 \setminus U_1 = W_1 \setminus U_1 = (U_1 \cup (U_3 \setminus U_2)) \setminus U_1 = U_3 \setminus U_2$. It follows that

$$\mathcal{U}' : U_1 \subsetneq W'_1 \subseteq W'_2 \subseteq \cdots \subsetneq W'_{k-2} \subsetneq W'_{k-1} = V(G) .$$

is a connected k -flag equivalent to \mathcal{U} . But then $W'_\ell \prec W_\ell$ and $W'_i = W_i$ for $\ell < i \leq k - 1$ implies $\mathcal{U}' \prec_k \mathcal{U}$, a contradiction.

(c) The ordered collection $(W'_j \setminus W'_{j-1})_{j=2}^k$ is a permutation of the ordered collection $(W_j \setminus W_{j-1})_{j=2}^k$. Therefore $W_2 \setminus W_1 = W'_t \setminus W'_{t-1}$ for some $t \geq 2$. Note that if $t \geq 3$ then $W'_{t-1} \neq W_{t-1}$; for $t \geq 3$ we have $W_2 \subseteq W_{t-1}$ but $W_2 \not\subseteq W'_{t-1}$. In particular we deduce that $\max(2, t - 1) \leq \ell$.

Now let $A = U_1 \setminus W_1$.

• For $0 \leq i \leq t-1$ the sets A and W'_i are disjoint. This is because $A \subseteq W_2 \setminus W_1 = W'_t \setminus W'_{t-1}$, so $A \cap W'_{t-1} = \emptyset$.

• For $2 \leq i \leq t-1$ there is no edge between A and $W'_i \setminus W'_{i-1}$. This is because if there is an edge connecting $u \in W'_i \setminus W'_{i-1}$ and $v \in A$ then there is an oriented edge from $u \in W'_i \setminus W'_{i-1}$ to $v \in W'_t \setminus W'_{t-1}$ in $G(\mathcal{W}')$. But $G(\mathcal{W}') = G(\mathcal{W})$ and the oriented edge from u to v cannot appear in $G(\mathcal{W})$ because $v \in W_2 \setminus W_1$ and $u \notin W_1$.

We now define

$$\mathcal{U}' : U'_1 \subsetneq U'_2 \subsetneq \cdots \subsetneq V(G)$$

by letting

$$U'_i = \begin{cases} W'_i \cup A, & \text{if } 1 \leq i \leq t-1; \\ W'_i, & \text{if } t \leq i \leq k. \end{cases}$$

Note that

• $U'_1 = W'_1 \cup A = U_1$. This is because $A = U_1 \setminus W_1$ and $W_1 = W'_1$.

• All subgraphs $G[U'_i]$ are connected: for $1 \leq i \leq t-1$ since $W'_1 \cup A = U_1$ is connected, we know that A is connected to W'_i at least via $W'_1 \subset W'_i$.

Moreover

• For $2 \leq i \leq t-1$ we have $U'_i \setminus U'_{i-1} = (W'_i \cup A) \setminus (W'_{i-1} \cup A) = W'_i \setminus W'_{i-1}$. This follows from the fact that A is disjoint from W'_{i-1} and W'_i .

• $U'_t \setminus U'_{t-1} = W'_t \setminus (W'_{t-1} \cup A) = (W_2 \setminus W_1) \setminus A = (U_2 \setminus W_1) \setminus (U_1 \setminus W_1) = U_2 \setminus U_1$.

• For $t+1 \leq i \leq k$ we have $U'_i \setminus U'_{i-1} = W'_i \setminus W'_{i-1}$.

Recall that the ordered collection $(W'_j \setminus W'_{j-1})_j$ is a permutation of the ordered collection $(W_j \setminus W_{j-1})_j$ and that $W_j \setminus W_{j-1} = U_j \setminus U_{j-1}$ for $3 \leq j \leq k$.

It follows that $\mathcal{U}' \in \mathfrak{F}_k(G, q)$ and that $(U'_j \setminus U'_{j-1})_1^k$ is a permutation of $(U_j \setminus U_{j-1})_1^k$.

Moreover \mathcal{U}' is equivalent to \mathcal{U} . To see this first observe that the only difference between $G(\mathcal{U})$ and $G(\mathcal{W})$ is that we orient the edges from U_1 to $U_2 \setminus U_1$ in $G(\mathcal{U})$ and we orient the edges from W_1 to $U_2 \setminus W_1$ in $G(\mathcal{W})$ (other oriented edges are identical). Similarly, since there is no edge between A and $W'_i \setminus W'_{i-1}$ for $2 \leq i \leq t-1$, the only difference between $G(\mathcal{U}')$ and $G(\mathcal{W}')$ is that we orient the edges from $U'_1 = U_1$ to $U'_t \setminus U'_{t-1} = U_2 \setminus U_1$ in $G(\mathcal{U}')$ and we orient the edges from $W'_1 = W_1$ to $W'_t \setminus W'_{t-1} = U_2 \setminus W_1$ in $G(\mathcal{W}')$ (other oriented edges are identical). Since $G(\mathcal{W})$ and $G(\mathcal{W}')$ coincide it follows that $G(\mathcal{U})$ and $G(\mathcal{U}')$ coincide.

Finally we show that $\mathcal{U}' \prec_k \mathcal{U}$. Recall that $\ell \geq 2$ and $\ell \geq t-1$.

• If $\ell \geq t$, then $U'_\ell = W'_\ell \prec W_\ell = U_\ell$ and $U'_i = W'_i = W_i = U_i$ for $\ell < i \leq k$.

• If $\ell = t-1$, then $U'_\ell = W'_\ell \cup A \prec W'_\ell \prec W_\ell = U_\ell$ and $U'_i = W'_i = W_i = U_i$ for $\ell < i \leq k$. Therefore in any case $\mathcal{U}' \prec_k \mathcal{U}$, a contradiction. \square

Proposition 4.15. *Let*

$$U_1 \subsetneq U_2 \subsetneq \cdots \subsetneq V(G)$$

$$V_1 \subsetneq V_2 \subsetneq \cdots \subsetneq V(G)$$

be two elements of $\mathfrak{S}_k(G, q)$ such that $U_i = V_i$ for $2 \leq i \leq k$. Let $A = U_2 \setminus U_1$, $B = V_2 \setminus V_1$, $C = U_1 \cap V_1$, and assume at least one of the following holds:

- (i) $G[C]$ is not connected.
- (ii) $G[A \cup B]$ is connected.

Then

$$W_1 \subsetneq W_2 \subsetneq \cdots \subsetneq V(G)$$

is also an element of $\mathfrak{S}_k(G, q)$, where $W_i = U_i$ for $2 \leq i \leq k$ and W_1 is (the vertex set of) the connected component of $G[C]$ that contains q .

Proof. It is enough to check that $G[W_2 \setminus W_1]$ is connected. Then the assertion follows by Lemma 4.14 (c), since $W_1 \subset U_1$.

Write

$$(4.1) \quad W_2 \setminus W_1 = A \cup B \cup (C \setminus W_1) ,$$

and write U_1 and V_1 as the disjoint unions:

$$(4.2) \quad U_1 = (U_1 \setminus V_1) \cup (C \setminus W_1) \cup W_1 \quad \text{and} \quad V_1 = (V_1 \setminus U_1) \cup (C \setminus W_1) \cup W_1 .$$

(i) Assume that $G[C]$ is not connected, that is, $C \setminus W_1 \neq \emptyset$. Since $G[U_1]$ is connected and there are no edges between $C \setminus W_1$ and W_1 , it follows from (4.2) that there must be an edge between each connected component $C \setminus W_1$ and $U_1 \setminus V_1$. As $U_1 \setminus V_1 \subseteq B$ we conclude that every connected component of $C \setminus W_1$ has an edge to B . Similarly, every connected component of $C \setminus W_1$ has an edge to A . Since $G[A]$ and $G[B]$ are both connected it follows from (4.1) that $G[W_2 \setminus W_1]$ is connected.

(ii) Assume that $G[A \cup B]$ is connected. We may assume $C \setminus W_1 = \emptyset$, since otherwise the result follows from (i) above. But then (4.1) becomes $W_2 \setminus W_1 = A \cup B$ and therefore $G[W_2 \setminus W_1]$ is connected. \square

Given an element in $\mathfrak{S}_k(G, q)$ there is a canonical way to obtain two related elements in $\mathfrak{S}_{k-1}(G, q)$. To state this result we first need a definition.

Definition 4.16. Given $\mathcal{U} \in \mathfrak{F}_k(G, q)$, the elements $\mathcal{U}^{(1)}, \mathcal{U}^{(2)} \in \mathfrak{F}_{k-1}(G, q)$ are obtained from \mathcal{U} by removing the first and second elements in the following appropriate sense. Let

$$\mathcal{U} : U_1 \subsetneq U_2 \subsetneq \cdots \subsetneq V(G) .$$

(a) $\mathcal{U}^{(1)}$ will denote

$$U_2 \subsetneq U_3 \subsetneq U_4 \subsetneq \cdots \subsetneq V(G) .$$

(b) $\mathcal{U}^{(2)}$ will denote

$$\begin{cases} U_1 \subsetneq U_3 \subsetneq U_4 \subsetneq \cdots \subsetneq V(G), & \text{if } G[U_3 \setminus U_1] \text{ is connected;} \\ \text{or} \\ (U_1 \cup (U_3 \setminus U_2)) \subsetneq U_3 \subsetneq U_4 \subsetneq \cdots \subsetneq V(G), & \text{if } G[U_3 \setminus U_1] \text{ is not connected.} \end{cases}$$

We remark that $\mathcal{U}^{(1)}$ and $\mathcal{U}^{(2)}$ are essential in the expression of our minimal free resolutions (see (5.3)).

Remark 4.17. For part (b) in Definition 4.16 we note that $U_3 \setminus U_1 = (U_3 \setminus U_2) \cup (U_2 \setminus U_1)$ and by assumption $G[U_3 \setminus U_2]$ and $G[U_2 \setminus U_1]$ are both connected and nonempty. So $G[U_3 \setminus U_1]$ is connected if and only if there are some edges connecting $U_3 \setminus U_2$ to $U_2 \setminus U_1$. Moreover, if there are no edges connecting $U_3 \setminus U_2$ to $U_2 \setminus U_1$ then there must be some edges connecting $U_3 \setminus U_2$ to U_1 . Since $G[U_3]$ is connected and $U_3 = (U_3 \setminus U_2) \cup (U_2 \setminus U_1) \cup U_1$ we get $G[U_1 \cup (U_3 \setminus U_2)]$ is connected.

Proposition 4.18. *If $\mathcal{U} \in \mathfrak{S}_k(G, q)$ then*

- (a) $\mathcal{U}^{(1)} \in \mathfrak{S}_{k-1}(G, q)$.
- (b) $\mathcal{U}^{(2)} \in \mathfrak{S}_{k-1}(G, q)$.
- (c) $\mathcal{U}^{(1)} \prec_{k-1} \mathcal{U}^{(2)}$.

Proof. Let

$$\mathcal{U} : U_1 \subsetneq U_2 \subsetneq \cdots \subsetneq V(G) .$$

Part (a) is exactly Lemma 4.14 (a). For part (b), if $G[U_3 \setminus U_1]$ is not connected then the result follows from Lemma 4.14 (b). If $G[U_3 \setminus U_1]$ is connected the result follows from part (a) and Lemma 4.14 (c).

For part (c) we first note that if $G[U_3 \setminus U_1]$ is connected $\mathcal{U}^{(1)} \prec_{k-1} \mathcal{U}^{(2)}$ follows directly from definitions. If $G[U_3 \setminus U_1]$ is not connected assume that $\mathcal{U}^{(2)} \prec_{k-1} \mathcal{U}^{(1)}$. Then one can easily see that

$$\mathcal{U}' : U_1 \subsetneq U_1 \cup (U_3 \setminus U_2) \subsetneq U_3 \subsetneq U_4 \subsetneq \cdots \subsetneq V(G)$$

is a connected k -flag equivalent to \mathcal{U} with $\mathcal{U}' \prec_k \mathcal{U}$ which is a contradiction. \square

There is a nice converse to Proposition 4.18 which is our next result.

Proposition 4.19. *Assume that $\mathcal{U} \in \mathfrak{F}_k(G, q)$ and the following three conditions hold:*

- (i) $\mathcal{U}^{(1)} \in \mathfrak{S}_{k-1}(G, q)$.
- (ii) $\mathcal{U}^{(2)} \in \mathfrak{S}_{k-1}(G, q)$.
- (iii) $\mathcal{U}^{(1)} \prec_{k-1} \mathcal{U}^{(2)}$.

Then $\mathcal{U} \in \mathfrak{S}_k(G, q)$.

Proof. Let

$$\mathcal{U} : U_1 \subsetneq U_2 \subsetneq \cdots \subsetneq V(G) .$$

To simplify the notation we let $A = U_2 \setminus U_1$ and $B = U_3 \setminus U_2$. For the sake of contradiction, we assume that

$$\mathcal{U}' : U'_1 \subsetneq U'_2 \subsetneq \cdots \subsetneq V(G)$$

is another connected flag equivalent to \mathcal{U} which precedes \mathcal{U} in the total ordering \prec_k . We will then find a connected flag equivalent to $\mathcal{U}^{(j)}$ (for $j = 1$ or 2) preceding it in the total ordering \prec_{k-1} .

We note that since $G(\mathcal{U})$ and $G(\mathcal{U}')$ coincide and $q \in U_1$ and $q \in U'_1$, we must have $U_1 = U'_1$. Let $\ell \geq 2$ be such that $U'_\ell \prec U_\ell$ and $U'_i = U_i$ for $i > \ell$.

We first show that $\ell > 2$. If $\ell = 2$, then $U'_1 = U_1$, $U'_2 = U_1 \cup (U_3 \setminus U_2)$ and $U'_i = U_i$ for all $i > 2$. If there is at least one edge between A and B , then it is oriented from B to A in $G(\mathcal{U}')$. But that edge must be oriented from A to B in $G(\mathcal{U})$ which contradicts $G(\mathcal{U}) = G(\mathcal{U}')$. If there exists no edge between A and B , then

$$\mathcal{U}^{(2)} : (U_1 \cup (U_3 \setminus U_2)) \subsetneq U_3 \subsetneq U_4 \subsetneq \cdots \subsetneq V(G) .$$

Now our assumption that $\mathcal{U}^{(1)} \prec_{k-1} \mathcal{U}^{(2)}$ implies that $U_2 \prec U_1 \cup (U_3 \setminus U_2)$. Thus $\mathcal{U} \prec_k \mathcal{U}'$ which is a contradiction. Thus we must have $\ell > 2$.

The ordered collection $(U'_j \setminus U'_{j-1})_{j=2}^k$ is a permutation of the ordered collection $(U_j \setminus U_{j-1})_{j=2}^k$. Then $A = U'_t \setminus U'_{t-1}$ and $B = U'_s \setminus U'_{s-1}$ for some $t, s \geq 2$. Note that if $t \geq 3$ then $U'_{t-1} \neq U_{t-1}$; for $t \geq 3$ we have $U_2 \subseteq U_{t-1}$ but $U_2 \not\subseteq U'_{t-1}$ (since $A \not\subseteq U'_{t-1}$). On the other hand, if $s \geq 4$ then $U'_{s-1} \neq U_{s-1}$; for $s \geq 4$ we have $U_3 \subseteq U_{s-1}$ but $U_3 \not\subseteq U'_{s-1}$ (since $B \not\subseteq U'_{s-1}$). In particular we deduce that $\ell \geq \max(s-1, t-1)$.

We first show that $\mathcal{U}^{(1)} \in \mathfrak{S}_{k-1}(G, q)$ implies $t > s$. Since all edges connecting A and $V(G) \setminus U_2$ are oriented from A to $V(G) \setminus U_2$ in $G(\mathcal{U})$, and $G(\mathcal{U}) = G(\mathcal{U}')$ we conclude that there is no edge between A and $U'_{t-1} \setminus U'_1$ if $t > 2$. Now we define

$$\mathcal{W} : W_1 \subsetneq W_2 \subsetneq \cdots \subsetneq W_{k-1} = V(G)$$

by letting

$$W_i = \begin{cases} U'_i \cup A, & \text{if } 1 \leq i \leq t-1; \\ U'_{i+1}, & \text{if } t \leq i \leq k-1. \end{cases}$$

Note that

- $W_1 = U'_1 \cup A = U_2$. This is because $A = U_2 \setminus U_1$ and $U_1 = U'_1$.
- The subgraphs $G[W_i]$ are all connected; for $1 \leq i \leq t-1$ since $U'_1 \cup A = U_2$ is connected, we know that A is connected to U'_i at least via $U'_1 \subset U'_i$.

Moreover

- For $2 \leq i \leq t-1$ we have $W_i \setminus W_{i-1} = (U'_i \cup A) \setminus (U'_{i-1} \cup A) = U'_i \setminus U'_{i-1}$. This follows from the fact that A is disjoint from U'_{i-1} and U'_i .
- $W_t \setminus W_{t-1} = U'_{t+1} \setminus (U'_{t-1} \cup A) = U'_{t+1} \setminus U'_t$.
- For $t+1 \leq i \leq k-1$ we have $W_i \setminus W_{i-1} = U'_{i+1} \setminus U'_i$.

This implies that the ordered collection $(W_i \setminus W_{i-1})_1^{k-1}$ is a permutation of the ordered collection $U_2 \cup (U'_j \setminus U'_{j-1})_{j \in \{2, \dots, k\} \setminus \{t\}}$ which is a permutation of ordered collection $U_2 \cup (U_j \setminus U_{j-1})_{j=3}^k$ of $\mathcal{U}^{(1)}$. We show that \mathcal{W} is equivalent to $\mathcal{U}^{(1)}$: first note that the only difference between $G(\mathcal{U})$ and $G(\mathcal{U}^{(1)})$ is that we orient the edges from U_1 to $U_2 \setminus U_1$ in $G(\mathcal{U})$ but we keep these edges unoriented in $G(\mathcal{U}^{(1)})$ (other oriented edges are identical). If $t = 2$ then $U'_2 = U_2$ and all oriented edges from A to $V(G) \setminus U_2$ in $G(\mathcal{U}^{(1)})$ are also in $G(\mathcal{W})$. Other edges are identical, since $G(\mathcal{U})$ and $G(\mathcal{U}')$ coincide. If $t > 2$ then there is no edge between A and $U'_c \setminus U'_1$ for $2 \leq c \leq t-1$. Therefore the only difference between $G(\mathcal{U}')$ and $G(\mathcal{W})$ is that we orient the edges from $U'_1 = U_1$ to $U'_t \setminus U'_{t-1} = U_2 \setminus U_1$ in $G(\mathcal{U}')$ and we keep these edges unoriented in $G(\mathcal{W})$. Other edges are identical, since $G(\mathcal{U})$ and $G(\mathcal{U}')$ coincide. Thus it follows that $G(\mathcal{U}^{(1)})$ and $G(\mathcal{W})$ coincide.

Note that the i th element in $\mathcal{U}^{(1)}$ is U_{i+1} for all i . If $\ell > t-1$, then $W_{\ell-1} = U'_\ell \prec U_\ell$ and $W_i = U_{i+1}$ for $i > \ell$. Thus $\mathcal{W} \prec_{k-1} \mathcal{U}^{(1)}$ which is a contradiction by our assumption that $\mathcal{U}^{(1)}$ belongs to $\mathfrak{S}_{k-1}(G, q)$. Therefore we have $\ell = t-1$. This also implies that $t > s$.

Now we consider two cases:

- If $G[U_3 \setminus U_1]$ is connected we have

$$\mathcal{U}^{(2)} : U_1 \subsetneq U_3 \subsetneq U_4 \subsetneq \cdots \subsetneq V(G) .$$

In this case $\mathcal{U}^{(2)} \in \mathfrak{S}_{k-1}(G, q)$ implies that $U_3 \setminus U_1 = (U_3 \setminus U_2) \cup (U_2 \setminus U_1) = A \cup B$ is connected. So there is at least one edge between A and B . This edge must have opposite orientations in $G(\mathcal{U})$ and $G(\mathcal{U}')$, which is a contradiction because $G(\mathcal{U}) = G(\mathcal{U}')$.

- If $G[U_3 \setminus U_1]$ is not connected we have

$$\mathcal{U}^{(2)} : (U_1 \cup (U_3 \setminus U_2)) \subsetneq U_3 \subsetneq U_4 \subsetneq \cdots \subsetneq V(G) .$$

In this case we need to do more work. We define

$$\mathcal{W}' : W'_1 \subsetneq W'_2 \subsetneq \cdots \subsetneq W'_{k-1} = V(G)$$

by letting

$$W'_i = \begin{cases} U'_i \cup B, & \text{if } 1 \leq i \leq s-1; \\ U'_{i+1}, & \text{if } s \leq i \leq k-1. \end{cases}$$

Note that

- $W'_1 = U'_1 \cup B = U_1 \cup (U_3 \setminus U_2)$. This is because $B = U_3 \setminus U_2$ and $U_1 = U'_1$.
- The subgraphs $G[W'_i]$ are all connected; for $1 \leq i \leq s-1$ since $U'_1 \cup B$ is connected (by our assumption that $\mathcal{U}^{(2)}$ belongs to $\mathfrak{S}_{k-1}(G, q)$), we know that B is connected to U'_i at least via $U'_1 \subset U'_i$.

Moreover

- For $2 \leq i \leq s-1$ we have $W'_i \setminus W'_{i-1} = (U'_i \cup B) \setminus (U'_{i-1} \cup B) = U'_i \setminus U'_{i-1}$. This follows from the fact that B is disjoint from U'_{i-1} and U'_i .
- $W'_s \setminus W'_{s-1} = U'_{s+1} \setminus (U'_{s-1} \cup B) = U'_{s+1} \setminus U'_s$.
- For $s+1 \leq i \leq k-1$ we have $W'_i \setminus W'_{i-1} = U'_{i+1} \setminus U'_i$.

This implies that the ordered collection $(W'_i \setminus W'_{i-1})_1^{k-1}$ is a permutation of the ordered collection $(U_1 \cup B) \cup (U'_j \setminus U'_{j-1})_{j \in \{2, \dots, k\} \setminus \{s\}}$ which is a permutation of the ordered collection for $\mathcal{U}^{(2)}$. Now we check that \mathcal{W}' is equivalent to $\mathcal{U}^{(2)}$: the only difference between $G(\mathcal{U})$ and $G(\mathcal{U}^{(2)})$ is that we orient the edges from U_1 to $U_3 \setminus U_2$ in $G(\mathcal{U})$ but we keep these edges unoriented in $G(\mathcal{U}^{(2)})$ (other oriented edges are identical). Similarly, since there is no edge between B and $U'_c \setminus U'_2$ for $2 \leq c \leq s-1$, the only difference between $G(\mathcal{U}')$ and $G(\mathcal{W}')$ is that we orient edges from $U'_1 = U_1$ to $U'_s \setminus U'_{s-1} = U_3 \setminus U_2$ in $G(\mathcal{U}')$ and we keep these edges unoriented in $G(\mathcal{W}')$ (other oriented edges are identical). Since $G(\mathcal{U})$ and $G(\mathcal{U}')$ coincide it follows that $G(\mathcal{U}^{(2)})$ and $G(\mathcal{W}')$ coincide.

Now our assumption that $U'_\ell \prec U_\ell$ and $\ell > s-1$ implies that $U'_\ell = W'_{\ell-1} \prec U_\ell$ and $W'_i = U_{i+1}$ for $i > \ell-1$. Note that the i th element in $\mathcal{U}^{(2)}$ is U_{i+1} for $i > 1$. Thus $\mathcal{W}' \prec_{k-1} \mathcal{U}^{(2)}$ which is a contradiction. \square

Proposition 4.20. *Let X_1, X_2, Y_1, Y_2 be four nonempty subsets of $V(G)$ such that:*

- (1) $G[X_1], G[X_2], G[Y_1]$, and $G[Y_2]$ are connected.
- (2) $X_1 \cap X_2 = \emptyset$ and $Y_1 \cap Y_2 = \emptyset$,
- (3) $X_1 \cup X_2 = Y_1 \cup Y_2$,
- (4) $X_2 \cap Y_2 \neq \emptyset$.

Then $D(X_1, X_2) \leq D(Y_1, Y_2)$ implies $X_1 = Y_1$ and $X_2 = Y_2$.

Proof. First we show that we must have $Y_2 \subseteq X_2$. Assume that $Y_2 \not\subseteq X_2$. Then for any $v \in X_1 \setminus Y_1$ we have $D(X_1, X_2)(v) \leq D(Y_1, Y_2)(v) = 0$ because $D(Y_1, Y_2)$ is supported on Y_1 . Therefore there is no edge between $X_1 \cap Y_2$ (i.e. the subset $X_1 \setminus Y_1 \subseteq X_1$) and $X_2 \cap Y_2$ (a subset of X_2). Note that $Y_2 = (X_1 \cap Y_2) \cup (X_2 \cap Y_2)$. Since $X_2 \cap Y_2 \neq \emptyset$, the assumption $X_1 \setminus Y_1 \neq \emptyset$ results in $G[Y_2]$ being disconnected which is a contradiction.

So we may assume $Y_2 \subseteq X_2$. Then for any $v \in X_1 \subseteq Y_1$ the set of all edges connecting v to a vertex in X_2 contains the set of all edges connecting v to a vertex in Y_2 . Therefore we have $D(Y_1, Y_2)(v) \leq D(X_1, X_2)(v)$. Comparing this with the inequality in the assumption we get $D(Y_1, Y_2)(v) = D(X_1, X_2)(v)$ for all $v \in X_1$. This means that there cannot be any edge connecting X_1 to $X_2 \setminus Y_2$. Since $Y_1 = X_1 \cup (X_2 \setminus Y_2)$, if $X_2 \setminus Y_2 \neq \emptyset$ then $G[Y_1]$ would be disconnected. So we must have $Y_2 = X_2$ and so $Y_1 = X_1$. \square

Corollary 4.21. *Let*

$$\mathcal{U} : U_1 \subsetneq U_2 \subsetneq \cdots \subsetneq V(G)$$

$$\mathcal{V} : V_1 \subsetneq V_2 \subsetneq \cdots \subsetneq V(G)$$

be two elements of $\mathfrak{S}_k(G, q)$. If for some $2 \leq i \leq k$

$$U_i = V_i \quad \text{and} \quad D(U_i \setminus U_{i-1}, U_{i-1}) \leq D(V_i \setminus V_{i-1}, V_{i-1})$$

then $U_{i-1} = V_{i-1}$.

Proof. Let $X_1 = U_i \setminus U_{i-1}$, $X_2 = U_{i-1}$, $Y_1 = V_i \setminus V_{i-1}$, and $Y_2 = V_{i-1}$ in Proposition 4.20. Note that $X_2 \cap Y_2 \neq \emptyset$ because $q \in U_1 \subseteq U_{i-1}$ and similarly $q \in V_1 \subseteq V_{i-1}$. \square

Definition 4.22. Write $\mathcal{W}, \mathcal{V} \in \mathfrak{S}_k(G, q)$ as

$$\mathcal{W} : W_1 \subsetneq W_2 \subsetneq \cdots \subsetneq W_k = V(G) ,$$

$$\mathcal{V} : V_1 \subsetneq V_2 \subsetneq \cdots \subsetneq V_k = V(G) .$$

Assume that $W_i = V_i$ for $i \geq 2$. We define an effective divisor

$$\mathcal{K}(\mathcal{W}, \mathcal{V}) := \max(D(W_2 \setminus W_1, W_1), D(V_2 \setminus V_1, V_1)) ,$$

where \max denotes the entry-wise maximum.

We remark that the notion $\mathcal{K}(\mathcal{W}, \mathcal{V})$ is essential in the study of our ideals and modules using Gröbner theory (see proofs of Theorem 5.1 and Theorem 5.3).

The following lemma gives an alternate formula for computing $\mathcal{K}(\mathcal{W}, \mathcal{V})$ which is sometimes more convenient.

Lemma 4.23. *For $\mathcal{W}, \mathcal{V} \in \mathfrak{S}_k(G, q)$ as in Definition 4.22, we have the following alternate formula:*

$$\begin{aligned} \mathcal{K}(\mathcal{W}, \mathcal{V}) = & \max(D(W_2 \setminus (W_1 \cup V_1), W_1), D(W_2 \setminus (W_1 \cup V_1), V_1)) \\ & + D(V_1 \setminus W_1, W_1) + D(W_1 \setminus V_1, V_1) . \end{aligned}$$

Proof. Let

$$\mathcal{K} = \max(D(W_2 \setminus W_1, W_1), D(V_2 \setminus V_1, V_1)) ,$$

$$\mathcal{K}' = \max(D(W_2 \setminus (W_1 \cup V_1), W_1), D(W_2 \setminus (W_1 \cup V_1), V_1)) + D(V_1 \setminus W_1, W_1) + D(W_1 \setminus V_1, V_1) .$$

Note that W_2 is the disjoint union of sets $W_1 \cap V_1$, $W_1 \setminus V_1$, $V_1 \setminus W_1$, and $W_2 \setminus (W_1 \cup V_1)$.

- If $v \in W_1 \cap V_1$ then $\mathcal{K}(v) = \mathcal{K}'(v) = 0$.
- If $v \in W_1 \setminus V_1$ then $\mathcal{K}(v) = \mathcal{K}'(v) = D(W_1 \setminus V_1, V_1)(v)$.
- If $v \in V_1 \setminus W_1$ then $\mathcal{K}(v) = \mathcal{K}'(v) = D(V_1 \setminus W_1, W_1)(v)$.
- If $v \in W_2 \setminus (W_1 \cup V_1)$ then

$$\mathcal{K}(v) = \mathcal{K}'(v) = \max(D(W_2 \setminus (W_1 \cup V_1), W_1), D(W_2 \setminus (W_1 \cup V_1), V_1))(v) .$$

\square

Lemma 4.24. *For $\mathcal{U} \in \mathfrak{S}_k(G, q)$ of the form*

$$\mathcal{U} : U_1 \subsetneq U_2 \subsetneq \cdots \subsetneq U_k = V(G)$$

we have

$$\mathcal{K}(\mathcal{U}^{(1)}, \mathcal{U}^{(2)}) = D(U_2 \setminus U_1, U_1) + D(U_3 \setminus U_2, U_2) .$$

Proof. From Definition 4.16, Remark 4.17, and Definition 4.22 we need to compute $\max(\alpha, \beta)$ where

$$\alpha = D(U_3 \setminus U_2, U_2) \quad \text{and} \quad \beta = \begin{cases} D(U_3 \setminus U_1, U_1), & \text{if } G[U_3 \setminus U_1] \text{ is connected;} \\ \text{or} \\ D(U_2 \setminus U_1, U_1), & \text{if } G[U_3 \setminus U_1] \text{ is not connected.} \end{cases}$$

Since $D(U_3 \setminus U_1, U_1) = D(U_3 \setminus U_2, U_1) + D(U_2 \setminus U_1, U_1)$ and $D(U_2 \setminus U_1, U_1) \geq D(U_3 \setminus U_2, U_2)$, it follows that in either case

$$\max(\alpha, \beta) = D(U_2 \setminus U_1, U_1) + D(U_3 \setminus U_2, U_2) .$$

□

We end this section by the following result which uses (and generalizes) many results of this section. This result plays a crucial role in the proof of Theorem 5.3.

Proposition 4.25. *Fix $\mathcal{W} \in \mathfrak{S}_k(G, q)$ and define*

$$\mathfrak{N}_{\mathcal{W}} = \{\mathcal{V} \in \mathfrak{S}_k(G, q) : \mathcal{V}^{(1)} = \mathcal{W}^{(1)} \text{ and } \mathcal{W} \prec_k \mathcal{V}\} .$$

For any $\mathcal{V} \in \mathfrak{N}_{\mathcal{W}}$ there exists a $\mathcal{W}' \in \mathfrak{N}_{\mathcal{W}}$ such that

- (i) $\mathcal{K}(\mathcal{W}, \mathcal{W}') \leq \mathcal{K}(\mathcal{W}, \mathcal{V})$,
- (ii) $\mathcal{U}^{(1)} = \mathcal{W}$ and $\mathcal{U}^{(2)} = \mathcal{W}'$ for some $\mathcal{U} \in \mathfrak{S}_{k+1}(G, q)$.

Proof. Fix $\mathcal{V} \in \mathfrak{N}_{\mathcal{W}}$. Consider the following subset of $\text{Div}(G)$ containing $\mathcal{K}(\mathcal{W}, \mathcal{V})$:

$$Q = \{\mathcal{K}(\mathcal{W}, \mathcal{V}') : \mathcal{V}' \in \mathfrak{N}_{\mathcal{W}} \text{ and } \mathcal{K}(\mathcal{W}, \mathcal{V}') \leq \mathcal{K}(\mathcal{W}, \mathcal{V})\} .$$

This is a nonempty finite set of effective divisors, so it has some minimal elements with respect to the partial ordering \leq on $\text{Div}(G)$. Choose the largest (with respect to the total ordering \prec_k) element $\mathcal{W}' \in \mathfrak{N}_{\mathcal{W}}$ such that $\mathcal{K}(\mathcal{W}, \mathcal{W}')$ is a minimal element of Q . Write

$$\mathcal{W} : W_1 \subsetneq W_2 \subsetneq \cdots \subsetneq W_k = V(G)$$

$$\mathcal{W}' : W'_1 \subsetneq W_2 \subsetneq \cdots \subsetneq W_k = V(G) ,$$

and to simplify the notation let $A = W_2 \setminus W_1$, $B = W_2 \setminus W'_1$, and $C = W_1 \cap W'_1$. Let X be (the vertex set of) the connected component of $G[C]$ containing q , and define the (auxiliary) k -flag (i.e. increasing sequence of subsets with no connectivity assumption)

$$\mathcal{X} : X \subsetneq W_2 \subsetneq W_3 \subsetneq \cdots \subsetneq W_k = V(G) .$$

Claim. If $\mathcal{X} \in \mathfrak{S}_k(G, q)$ then $W'_1 \subsetneq W_1$.

Assume that $\mathcal{X} \in \mathfrak{S}_k(G, q)$. Since $\mathcal{W} \prec_k \mathcal{W}'$ we have $W_1 \neq W'_1$ and therefore $X \subsetneq W_1$. It follows that $\mathcal{W} \prec_k \mathcal{X}$ and $\mathcal{X} \in \mathfrak{N}_{\mathcal{W}}$. We also have $X \subseteq W'_1$ and $\mathcal{W}' \preceq_k \mathcal{X}$.

We will show that

$$\mathcal{K}(\mathcal{W}, \mathcal{X}) \leq \mathcal{K}(\mathcal{W}, \mathcal{W}') .$$

Once this is shown, the claim is proved; if $\mathcal{K}(\mathcal{W}, \mathcal{X}) < \mathcal{K}(\mathcal{W}, \mathcal{W}')$ then $\mathcal{K}(\mathcal{W}, \mathcal{W}')$ is not minimal which is a contradiction. If $\mathcal{K}(\mathcal{W}, \mathcal{X}) = \mathcal{K}(\mathcal{W}, \mathcal{W}')$ then \mathcal{X} also realizes a minimal divisor and $\mathcal{W}' \preceq_k \mathcal{X}$. This contradicts the definition of \mathcal{W}' unless $\mathcal{W}' = \mathcal{X}$, which means $X = W'_1$ and hence $W'_1 \subsetneq W_1$.

By the formula in Lemma 4.23 we obtain

$$\begin{aligned}
\mathcal{K}(\mathcal{W}, \mathcal{X}) &= \max(D(W_2 \setminus (W_1 \cup X), W_1), D(W_2 \setminus (W_1 \cup X), X)) \\
&\quad + D(X \setminus W_1, W_1) + D(W_1 \setminus X, X) \\
&= \max(D(W_2 \setminus W_1, W_1), D(W_2 \setminus W_1, X)) + D(W_1 \setminus X, X) \\
&= D(W_2 \setminus W_1, W_1) + D(W_1 \setminus X, X) \\
&= D(W_2 \setminus (W_1 \cup W'_1), W_1) + D(W'_1 \setminus W_1, W_1) + D(W_1 \setminus X, X) .
\end{aligned}$$

Compare this with

$$\begin{aligned}
\mathcal{K}(\mathcal{W}, \mathcal{W}') &= \max(D(W_2 \setminus (W_1 \cup W'_1), W_1), D(W_2 \setminus (W_1 \cup W'_1), W'_1)) \\
&\quad + D(W'_1 \setminus W_1, W_1) + D(W_1 \setminus W'_1, W'_1) .
\end{aligned}$$

Since there is no edge between $C \setminus X$ and X we have

$$D(W_1 \setminus X, X) = D(W_1 \setminus C, X) = D(W_1 \setminus W'_1, X) \leq D(W_1 \setminus W'_1, W'_1) .$$

We get $\mathcal{K}(\mathcal{W}, \mathcal{X}) \leq \mathcal{K}(\mathcal{W}, \mathcal{W}')$, and the claim is proved.

We now define

$$\mathcal{U} : C \subsetneq W_1 \subsetneq W_2 \subsetneq \cdots \subsetneq W_k = V(G) .$$

By definition $\mathcal{U}^{(1)} = \mathcal{W}$. We will show that $\mathcal{U} \in \mathfrak{S}_{k+1}(G, q)$ and $\mathcal{U}^{(2)} = \mathcal{W}'$. Since $\mathcal{W}' \in \mathfrak{N}_{\mathcal{W}}$ implies $\mathcal{W} \prec_k \mathcal{W}'$, by Proposition 4.19 it suffices to prove $\mathcal{U} \in \mathfrak{F}_{k+1}(G, q)$ and $\mathcal{U}^{(2)} = \mathcal{W}'$. Recall from Proposition 4.15 that if $G[C]$ is not connected then we have $\mathcal{X} \in \mathfrak{S}_k(G, q)$. By **Claim** above, we then must have $C = W_1 \cap W'_1 = W'_1$ which is connected. Therefore, we assume that $G[C]$ is connected. Therefore to show $\mathcal{U} \in \mathfrak{F}_{k+1}(G, q)$ we only need to check that $G[W_1 \setminus C]$ is connected; all other connectivities are guaranteed by the assumption that $\mathcal{W} \in \mathfrak{S}_k(G, q)$.

We need to consider two cases:

- If $G[A \cup B]$ is not connected: First we note that since $G[A]$ and $G[B]$ are both connected, if $A \cap B \neq \emptyset$ then A would be connected to B via $A \cap B$, contradicting the fact that $G[A \cup B]$ is not connected. So we must have $A \cap B = \emptyset$ or equivalently $W_2 = W_1 \cup W'_1$. Now

$$W_1 \setminus C = W_1 \setminus W'_1 = W_2 \setminus W'_1$$

which is connected since $\mathcal{W}' \in \mathfrak{S}_k(G, q)$.

Now $\mathcal{W}' = \mathcal{U}^{(2)}$ by noticing that

$$W'_1 = (W_1 \cap W'_1) \cup (W_2 \setminus W_1) = U_1 \cup (U_3 \setminus U_2) .$$

- If $G[A \cup B]$ is connected: Proposition 4.15 implies that $\mathcal{X} \in \mathfrak{S}_k(G, q)$. Now by the claim above, we then must have $C = W_1 \cap W'_1 = W'_1$ and so

$$\mathcal{U} : W'_1 \subsetneq W_1 \subsetneq W_2 \subsetneq \cdots \subsetneq W_k = V(G) .$$

Thus $\mathcal{U}^{(2)} = \mathcal{W}'$ by definition.

If $G[W_1 \setminus W'_1]$ is not connected, let $Y \neq \emptyset$ be the vertex set of one of the connected components of $G[W_1 \setminus W'_1]$, and consider the flag

$$\mathcal{Y} : W'_1 \cup Y \subsetneq W_2 \subsetneq W_3 \subsetneq \cdots \subsetneq W_k = V(G) .$$

Then $\mathcal{Y} \in \mathfrak{F}_k(G, q)$ because

- $G[W'_1 \cup Y]$ is connected: W_1 is connected and is the disjoint union $W'_1 \cup Y \cup W_1 \setminus (W'_1 \cup Y)$. Since there are no edges between Y and $W_1 \setminus (W'_1 \cup Y)$, there must be some edges connecting Y to W'_1 .
- $G[W_2 \setminus (W'_1 \cup Y)]$ is connected: $W_2 \setminus W'_1$ is connected and is the disjoint union $(W_2 \setminus W_1) \cup Y \cup (W_1 \setminus (W'_1 \cup Y))$. Since there are no edges between Y and $W_1 \setminus (W'_1 \cup Y)$, there must be some edges connecting $W_2 \setminus W_1$ to $W_1 \setminus (W'_1 \cup Y)$. So $(W_2 \setminus W_1) \cup (W_1 \setminus (W'_1 \cup Y)) = W_2 \setminus (W'_1 \cup Y)$ is connected.

Since $W'_1 \cup Y \subseteq W_1$ by comparing \mathcal{Y} and \mathcal{W} and using Lemma 4.14(c) we get $\mathcal{Y} \in \mathfrak{S}_k(G, q)$. We also note that $\mathcal{W} \prec_k \mathcal{Y}$. Consequently $\mathcal{Y} \in \mathfrak{R}_{\mathcal{W}}$.

However $\mathcal{K}(\mathcal{W}, \mathcal{Y}) < \mathcal{K}(\mathcal{W}, \mathcal{W}')$ which contradicts the definition of \mathcal{W}' . To see this compare

$$\begin{aligned} \mathcal{K}(\mathcal{W}, \mathcal{X}) &= D(W_2 \setminus W_1, W_1) + D(W_1 \setminus (W'_1 \cup Y), W'_1 \cup Y) \\ &= D(W_2 \setminus (W_1 \cup W'_1), W_1) \\ &\quad + D(W'_1 \setminus W_1, W_1) + D(W_1 \setminus (W'_1 \cup Y), W'_1 \cup Y) \end{aligned}$$

with

$$\begin{aligned} \mathcal{K}(\mathcal{W}, \mathcal{W}') &= \max(D(W_2 \setminus (W_1 \cup W'_1), W_1), D(W_2 \setminus (W_1 \cup W'_1), W'_1)) \\ &\quad + D(W'_1 \setminus W_1, W_1) + D(W_1 \setminus W'_1, W'_1) . \end{aligned}$$

Here we have

$$\begin{aligned} D(W_1 \setminus (W'_1 \cup Y), W'_1 \cup Y) &= D(W_1 \setminus (W'_1 \cup Y), W'_1) \\ &< D(W_1 \setminus (W'_1 \cup Y), W'_1) + D(Y, W'_1) \\ &= D(W_1 \setminus W'_1, W'_1) . \end{aligned}$$

The first equality is because there are no edges between $W_1 \setminus (W'_1 \cup Y)$ and Y . The strict inequality is because we have shown above that there are edges connecting Y to W'_1 .

Therefore $G[W_1 \setminus W'_1]$ is connected, and $\mathcal{U} \in \mathfrak{F}_{k+1}(G, q)$ which is what we wanted. \square

5. SYZYGIES AND FREE RESOLUTIONS FOR I_G AND $\text{in}(I_G)$

Let K be a field and let $R = K[\mathbf{x}]$ be the polynomial ring in n variables $\{x_v : v \in V(G)\}$. Recall from §2.2.1 that $K[\mathbf{x}]$ has a natural \mathbf{A} -grading, where \mathbf{A} can be replaced by \mathbb{Z} , $\text{Div}(G)$, or $\text{Pic}(G)$. Recall that for $\mathbf{A} = \mathbb{Z}$ and $\mathbf{A} = \text{Pic}(G)$ the ideal I_G is graded.

Let the monomial ordering $<$ on R be as in Definition 2.4. Recall that this ordering depends on the choice of the fixed vertex q . The following theorem is essentially in [7, Theorem 14]. Here we state and prove the theorem in a language that suggests a generalization.

Theorem 5.1. *Fix a pointed graph (G, q) and let $\mathbf{A} = \mathbb{Z}$ or $\mathbf{A} = \text{Pic}(G)$. A minimal \mathbf{A} -homogeneous Gröbner bases of $(I_G, <)$ is*

$$\mathbf{G}(G, q) = \{\mathbf{x}^{D(U_2 \setminus U_1, U_1)} - \mathbf{x}^{D(U_1, U_2 \setminus U_1)} : U_1 \subsetneq U_2 = V(G) \text{ is in } \mathfrak{S}_2(G, q)\} .$$

Moreover $\text{LM}(\mathbf{x}^{D(U_2 \setminus U_1, U_1)} - \mathbf{x}^{D(U_1, U_2 \setminus U_1)}) = \mathbf{x}^{D(U_2 \setminus U_1, U_1)}$.

Proof. To simplify the notation for a subset $A \subseteq V(G)$ we use $\bar{A} = V(G) \setminus A$. Since $q \in U_1$ it follows from the definition of $<$ that $\mathbf{x}^{D(U_1, \bar{U}_1)} < \mathbf{x}^{D(\bar{U}_1, U_1)}$.

We first prove

$$\mathbf{G}'(G, q) = \{\mathbf{x}^{D(U_2 \setminus U_1, U_1)} - \mathbf{x}^{D(U_1, U_2 \setminus U_1)} : U_1 \subsetneq U_2 = V(G), q \in U_1\}$$

forms a Gröbner bases of I_G . We will call a sequence of subsets $U_1 \subsetneq U_2 = V(G)$ with $q \in U_1$ a 2-flag of (G, q) . Note that for a 2-flag there is no connectivity assumption on $G[U_1]$ or on $G[U_2 \setminus U_1]$.

As usual, we use Buchberger's criterion. Let $f = \mathbf{x}^{D(\bar{U}, U)} - \mathbf{x}^{D(U, \bar{U})}$ and $g = \mathbf{x}^{D(\bar{V}, V)} - \mathbf{x}^{D(V, \bar{V})}$ be two elements of $\mathbf{G}'(G, q)$. Define the effective divisor $D' \in \text{Div}(G)$ by

$$D' = \max(D(\bar{U}, U), D(\bar{V}, V)) = \mathcal{K}(\mathcal{U}, \mathcal{V}) .$$

In the language of chip-firing games, D' is the minimal divisor that allows one to "fire" either the set \bar{U} or the set \bar{V} and still have an effective divisor as outcome, that is,

$$D' - \Delta(\chi_{\bar{U}}) \geq 0 \text{ and } D' - \Delta(\chi_{\bar{V}}) \geq 0 .$$

Buchberger's s -polynomial is

$$\text{spoly}(f, g) = \mathbf{x}^{D' - D(\bar{U}, U)} f - \mathbf{x}^{D' - D(\bar{V}, V)} g = \mathbf{x}^{D_1} - \mathbf{x}^{D_2} ,$$

where $D_1 = D' - D(\bar{V}, V) + D(V, \bar{V}) = D' - \Delta(\chi_{\bar{V}})$ is the effective divisor obtained from D' by firing the set \bar{V} . Similarly $D_2 = D' - D(\bar{U}, U) + D(U, \bar{U}) = D' - \Delta(\chi_{\bar{U}})$ is the effective divisor obtained from D' by firing the set \bar{U} . It follows from this interpretation that

$$(5.1) \quad D_1 - \Delta(\chi_{\bar{U} \setminus \bar{V}}) = D_2 - \Delta(\chi_{\bar{V} \setminus \bar{U}}) = D' - \Delta(\chi_{\bar{U} \cup \bar{V}}) \geq 0 .$$

The reason is the net effect of firing first the set \bar{V} and then the set $\bar{U} \setminus \bar{V}$ is the same as firing the set $\bar{U} \cup \bar{V}$; chips going along edges connecting \bar{V} and $\bar{U} \setminus \bar{V}$ cancel each other.

Without loss of generality we assume that $\text{LM}(\text{spoly}(f, g)) = \mathbf{x}^{D_1}$. It follows from (5.1) that we can reduce it by $h_1 = \mathbf{x}^{D(\bar{U} \setminus \bar{V}, U \cup \bar{V})} - \mathbf{x}^{D(U \cup \bar{V}, \bar{U} \setminus \bar{V})} \in \mathbf{G}'(G, q)$ associated to the 2-flag $(U \cup \bar{V}) \subsetneq V(G)$, and get:

$$\begin{aligned} \text{spoly}(f, g) - \mathbf{x}^{D_1 - D(\bar{U} \setminus \bar{V}, U \cup \bar{V})} h_1 &= \mathbf{x}^{D_1 - \Delta(\chi_{\bar{U} \setminus \bar{V}})} - \mathbf{x}^{D_2} \\ &= \mathbf{x}^{D' - \Delta(\chi_{\bar{U} \cup \bar{V}})} - \mathbf{x}^{D' - \Delta(\chi_{\bar{U}})} . \end{aligned}$$

The leading monomial is now $\mathbf{x}^{D' - \Delta(\chi_{\bar{U}})}$. Again, it follows from (5.1) that we can reduce this by $h_2 = \mathbf{x}^{D(\bar{V} \setminus \bar{U}, \bar{U} \cup V)} - \mathbf{x}^{D(\bar{U} \cup V, \bar{V} \setminus \bar{U})} \in \mathbf{G}'(G, q)$ associated to the 2-flag $(\bar{U} \cup V) \subsetneq V(G)$, and get:

$$\begin{aligned} \text{spoly}(f, g) - \mathbf{x}^{D_1 - D(\bar{U} \setminus \bar{V}, U \cup \bar{V})} h_1 - \mathbf{x}^{D_2 - D(\bar{V} \setminus \bar{U}, \bar{U} \cup V)} h_2 &= \mathbf{x}^{D' - \Delta(\chi_{\bar{U} \cup \bar{V}})} - \mathbf{x}^{D' - \Delta(\chi_{\bar{U} \cup \bar{V}})} \\ &= 0 \end{aligned}$$

completing the proof that $\mathbf{G}'(G, q)$ is a Gröbner basis.

Finally we show that if we only consider the flags in $\mathfrak{S}_2(G, q)$ then we get a minimal Gröbner bases. We show this by successively removing the binomials which are not coming from connected 2-flags. There are two steps:

- If $U_1 \subsetneq V(G)$ is a 2-flag which is not in $\mathfrak{S}_2(G, q)$ then there exists another 2-flag $V_1 \subsetneq V(G)$ such that $\mathbf{x}^{D(\bar{V}_1, V_1)} \mid \mathbf{x}^{D(\bar{U}_1, U_1)}$.

- If U_1 is not connected let V_1 be the connected component of U_1 containing q ; then \bar{V}_1 is the union of \bar{U}_1 and $U_1 \setminus V_1$ (i.e. other connected components of U_1). There is no edge between V_1 and other connected components of U_1 so for $v \in U_1 \setminus V_1$ we have $0 = D(\bar{V}_1, V_1)(v) \leq D(\bar{U}_1, U_1)(v)$. Since $\bar{U}_1 \subseteq \bar{V}_1$, for $v \in \bar{U}_1$ we have $D(\bar{V}_1, V_1)(v) \leq D(\bar{U}_1, U_1)(v)$.
- If \bar{U}_1 is not connected let V_1 be the complement of any connected component of \bar{U}_1 . In this case $D(\bar{V}_1, V_1)(v) = D(\bar{U}_1, U_1)(v)$ for all $v \in \bar{V}_1$.

• If $U_1 \subsetneq V(G)$ is in $\mathfrak{S}_2(G, q)$ then its binomial cannot be removed. Otherwise, there exists a different 2-flag $V_1 \subsetneq V(G)$ in $\mathfrak{S}_2(G, q)$ such that $\mathbf{x}^{D(\bar{V}_1, V_1)} \mid \mathbf{x}^{D(\bar{U}_1, U_1)}$ which is a contradiction by Corollary 4.21.

Homogeneity with respect to the \mathbb{Z} and $\text{Pic}(G)$ gradings is obvious. \square

Remark 5.2. It is easy to check with examples (e.g. a path) that $\mathbf{G}(G, q)$ is generally not the *reduced* Gröbner bases for $(I_G, <)$.

Theorem 5.1 can be rephrased as having a bijection between $\mathfrak{S}_2(G, q)$ and $\mathbf{G}(G, q)$. The following theorem gives a generalization of this fact.

Theorem 5.3. *Fix a pointed graph (G, q) and let $\mathbf{A} = \mathbb{Z}$ or $\mathbf{A} = \text{Pic}(G)$. For each $k \geq 0$ there exists a natural injection*

$$\psi_k : \mathfrak{S}_{k+2}(G, q) \hookrightarrow \text{syz}_k(\mathbf{G}(G, q))$$

such that

- (i) For some module ordering $<_k$, the set $\mathbf{G}_k(G, q) := \text{Image}(\psi_k)$ forms a minimal \mathbf{A} -homogeneous Gröbner bases of $(\text{syz}_k(\mathbf{G}(G, q)), <_k)$,
- (ii) For $\mathcal{U} \in \mathfrak{S}_{k+2}(G, q)$ of the form $U_1 \subsetneq U_2 \subsetneq \cdots \subsetneq V(G)$ we have

$$(5.2) \quad \text{LM}(\psi_k(\mathcal{U})) = \mathbf{x}^{D(U_2 \setminus U_1, U_1)} [\psi_{k-1}(\mathcal{U}^{(1)})] .$$

Proof. For consistency in the notation we define $\text{syz}_{-1}(\mathbf{G}(G, q)) = \{0\}$ and the map

$$\psi_{-1} : \mathfrak{S}_1(G, q) \hookrightarrow \{0\}$$

sends the canonical connected 1-flag $V(G)$ to 0.

The proof is by induction on $k \geq 0$.

Base case. For $k = 0$ the result is proved in Theorem 5.1. Here $\mathbf{G}_0(G, q) = \mathbf{G}(G, q)$ and $<_0$ is $<$, and

$$\begin{aligned} \psi_0 : \mathfrak{S}_2(G, q) &\hookrightarrow \text{syz}_0(\mathbf{G}(G, q)) = I_G \\ (U_1 \subsetneq U_2) &\mapsto (\mathbf{x}^{D(U_2 \setminus U_1, U_1)} - \mathbf{x}^{D(U_1, U_2 \setminus U_1)})[0] , \end{aligned}$$

and $\text{LM}(\psi_k(\mathcal{U})) = \mathbf{x}^{D(U_2 \setminus U_1, U_1)}[0]$.

Induction hypothesis. Now let $k > 0$ and assume that there exists a bijection

$$\psi_{k-1} : \mathfrak{S}_{k+1}(G, q) \rightarrow \mathbf{G}_{k-1}(G, q) \subseteq \text{syz}_{k-1}(\mathbf{G}(G, q))$$

such that $\mathbf{G}_{k-1}(G, q)$ forms a minimal homogeneous Gröbner bases of $\text{syz}_{k-1}(\mathbf{G}(G, q))$ with respect to $<_{k-1}$, and (5.2) holds for the leading monomials.

Via the bijection ψ_{k-1} , the set $\mathbf{G}_{k-1}(G, q)$ inherits a total ordering \prec'_{k-1} from the total ordering \prec_{k+1} on $\mathfrak{S}_{k+1}(G, q)$, that is

$$f \prec'_{k-1} h \text{ in } \mathbf{G}_{k-1}(G, q) \iff \psi_{k-1}^{-1}(f) \prec_{k+1} \psi_{k-1}^{-1}(h) \text{ in } \mathfrak{S}_{k+1}(G, q).$$

Inductive step. Given $\mathcal{U} \in \mathfrak{S}_{k+2}(G, q)$ let $\mathcal{U}^{(1)}$ and $\mathcal{U}^{(2)}$ be as defined in Definition 4.16. We *define*

$$(5.3) \quad \begin{aligned} \psi_k : \mathfrak{S}_{k+2}(G, q) &\rightarrow \text{syz}_k(\mathbf{G}(G, q)) \\ \mathcal{U} &\mapsto s(\psi_{k-1}(\mathcal{U}^{(1)}), \psi_{k-1}(\mathcal{U}^{(2)})) . \end{aligned}$$

In the following $\mathcal{U}, \mathcal{V} \in \mathfrak{S}_{k+2}(G, q)$ are of the form

$$\begin{aligned} U_1 \subsetneq U_2 \subsetneq \cdots \subsetneq V(G) \\ V_1 \subsetneq V_2 \subsetneq \cdots \subsetneq V(G) . \end{aligned}$$

The result follows from a series of claims.

Claim 1. ψ_k is a well-defined.

By Proposition 4.18

$$\mathcal{U}^{(1)} \in \mathfrak{S}_{k+1}(G, q), \quad \mathcal{U}^{(2)} \in \mathfrak{S}_{k+1}(G, q), \quad \mathcal{U}^{(1)} \prec_{k+1} \mathcal{U}^{(2)} .$$

So by the induction hypothesis

$$\psi_{k-1}(\mathcal{U}^{(1)}), \psi_{k-1}(\mathcal{U}^{(2)}) \in \mathbf{G}_{k-1}(G, q)$$

and by the definition of the total ordering on $\mathbf{G}_{k-1}(G, q)$ we have

$$\psi_{k-1}(\mathcal{U}^{(1)}) \prec'_{k-1} \psi_{k-1}(\mathcal{U}^{(2)}) .$$

Let $\mathcal{U}^{(1,1)} := (\mathcal{U}^{(1)})^{(1)}$ and $\mathcal{U}^{(2,1)} := (\mathcal{U}^{(2)})^{(1)}$. It is apparent from Definition 4.16 that

$$\mathcal{U}^{(1,1)} = \mathcal{U}^{(2,1)} .$$

By the induction hypothesis and (5.2), $\text{LM}(\psi_{k-1}(\mathcal{U}^{(1)}))$ and $\text{LM}(\psi_{k-1}(\mathcal{U}^{(2)}))$ are both multiples of the same free bases element $[\psi_{k-2}(\mathcal{U}^{(1,1)})] = [\psi_{k-2}(\mathcal{U}^{(2,1)})]$. It follows that

$$s(\psi_{k-1}(\mathcal{U}^{(1)}), \psi_{k-1}(\mathcal{U}^{(2)})) \in \mathcal{S}(\mathbf{G}_{k-1}(G, q)) \subset \text{syz}_k(\mathbf{G}(G, q))$$

is well-defined (see Theorem 3.2).

Claim 2. $\mathbf{G}_k(G, q) := \text{Image}(\psi_k)$ consists of homogeneous elements.

Since $\psi_{k-1}(\mathcal{U}^{(1)})$ and $\psi_{k-1}(\mathcal{U}^{(2)})$ are homogeneous by the induction hypothesis, it follows that $s(\psi_{k-1}(\mathcal{U}^{(1)}), \psi_{k-1}(\mathcal{U}^{(2)}))$ is also homogeneous.

Claim 3. $\text{LM}(\psi_k(\mathcal{U})) = \mathbf{x}^{D(U_2 \setminus U_1, U_1)}[\psi_{k-1}(\mathcal{U}^{(1)})]$.

From Lemma 3.3 it suffices to show that $D(U_2 \setminus U_1, U_1) = \max(\alpha, \beta) - \alpha$ where

$$\text{LM}(\psi_{k-1}(\mathcal{U}^{(1)})) = \mathbf{x}^\alpha[\psi_{k-2}(\mathcal{U}^{(1,1)})] \quad , \quad \text{LM}(\psi_{k-1}(\mathcal{U}^{(2)})) = \mathbf{x}^\beta[\psi_{k-2}(\mathcal{U}^{(2,1)})] .$$

But this is precisely Lemma 4.24.

Claim 4. ψ_k is injective.

If $\mathcal{U}, \mathcal{V} \in \mathfrak{S}_{k+2}(G, q)$ are such that $\psi_k(\mathcal{U}) = \psi_k(\mathcal{V})$ then their leading monomials should be equal:

$$\mathbf{x}^{D(U_2 \setminus U_1, U_1)}[\psi_{k-1}(\mathcal{U}^{(1)})] = \mathbf{x}^{D(V_2 \setminus V_1, V_1)}[\psi_{k-1}(\mathcal{V}^{(1)})] .$$

Therefore $\psi_{k-1}(\mathcal{U}^{(1)}) = \psi_{k-1}(\mathcal{V}^{(1)})$ and $D(U_2 \setminus U_1, U_1) = D(V_2 \setminus V_1, V_1)$. By the induction hypothesis ψ_{k-1} is injective which implies $\mathcal{U}^{(1)} = \mathcal{V}^{(1)}$ and $D(U_2 \setminus U_1, U_1) = D(V_2 \setminus V_1, V_1)$. It follows from Corollary 4.21 that $U_1 = V_1$ and $\mathcal{U} = \mathcal{V}$.

Our last claim below will finish the inductive step.

Claim 5. $\text{Image}(\psi_k)$ forms a minimal homogeneous Gröbner bases of $\text{syz}_k(\mathbf{G}(G, q))$ with respect to $<_k$ obtained from $<_{k-1}$ according to (3.1).

We have already shown in the proof of Claim 1 that $\text{Image}(\psi_k) \subseteq \mathcal{S}(\mathbf{G}_{k-1}(G, q))$. By Theorem 3.2 and Remark 3.5 it remains to show that

(I) $0 \notin \text{Image}(\psi_k)$.

- (II) For any element $s(f, h) \in \mathcal{S}(\mathbf{G}_{k-1}(G, q))$ there exists an element $g \in \text{Image}(\psi_k)$ such that $\text{LM}(g) \mid \text{LM}(s(f, h))$.
- (III) For any two elements $g, g' \in \text{Image}(\psi_k)$, if $\text{LM}(g) \mid \text{LM}(g')$ then $g = g'$.

(I) follows immediately from Claim 3 above.

Proof of (II). By the induction hypothesis $f = \psi_{k-1}(\mathcal{W})$ and $h = \psi_{k-1}(\mathcal{V})$ for two $\mathcal{W} \prec_{k+1} \mathcal{V}$ in $\mathfrak{S}_{k+1}(G, q)$ such that $\mathcal{W}^{(1)} = \mathcal{V}^{(1)}$. We need to find $\mathcal{U} \in \mathfrak{S}_{k+2}(G, q)$ such that

$$\text{LM}(s(\mathcal{U}^{(1)}, \mathcal{U}^{(2)})) \mid \text{LM}(s(\psi_{k-1}(\mathcal{W}), \psi_{k-1}(\mathcal{V}))) .$$

We use Proposition 4.25. From the previous paragraph it follows

$$\mathcal{V} \in \mathfrak{N}_{\mathcal{W}} = \{\mathcal{X} \in \mathfrak{S}_{k+1}(G, q) : \mathcal{W}^{(1)} = \mathcal{X}^{(1)} \text{ and } \mathcal{W} \prec_{k+1} \mathcal{X}\} .$$

Hence there exists a $\mathcal{W}' \in \mathfrak{N}_{\mathcal{W}}$ such that $\mathcal{K}(\mathcal{W}, \mathcal{W}') \leq \mathcal{K}(\mathcal{W}, \mathcal{V})$, and $\mathcal{U}^{(1)} = \mathcal{W}$ and $\mathcal{U}^{(2)} = \mathcal{W}'$ for some $\mathcal{U} \in \mathfrak{S}_{k+1}(G, q)$.

By (5.2) and Lemma 4.24 (or Claim 3 above) we have

$$\text{LM}(\psi_k(\mathcal{U})) = \mathbf{x}^{\mathcal{K}(\mathcal{W}, \mathcal{W}') - \alpha} [\psi_{k-1}(\mathcal{W})] ,$$

$$\text{LM}(s(\psi_{k-1}(\mathcal{W}), \psi_{k-1}(\mathcal{V}))) = \mathbf{x}^{\mathcal{K}(\mathcal{W}, \mathcal{V}) - \alpha} [\psi_{k-1}(\mathcal{W})] ,$$

where $\alpha = D(U_3 \setminus U_2, U_2)$. Therefore

$$\text{LM}(\psi_k(\mathcal{U})) \mid \text{LM}(s(\psi_{k-1}(\mathcal{W}), \psi_{k-1}(\mathcal{V}))) .$$

Proof of (III). We need to show that for any $\mathcal{U}, \mathcal{V} \in \mathfrak{S}_{k+2}(G, q)$ with $\mathcal{U}^{(1)} = \mathcal{V}^{(1)}$, if $\text{LM}(\psi_k(\mathcal{U}) \mid \text{LM}(\psi_k(\mathcal{V})))$ then $\mathcal{U} = \mathcal{V}$.

From (5.2) $\text{LM}(\psi_k(\mathcal{U}) \mid \text{LM}(\psi_k(\mathcal{V})))$ is equivalent to $D(U_2 \setminus U_1, U_1) \leq D(V_2 \setminus V_1, V_1)$. This together with $\mathcal{U}^{(1)} = \mathcal{V}^{(1)}$ implies $\mathcal{U} = \mathcal{V}$ by Corollary 4.21. \square

Remark 5.4. In Theorem 5.3 if we replace $\mathbf{G}(G, q)$ with

$$\{\mathbf{x}^{D(U_2 \setminus U_1, U_1)} : U_1 \subsetneq U_2 = V(G) \text{ is in } \mathfrak{S}_2(G, q)\} ,$$

(i.e. the initial terms of the Gröbner bases constructed in Theorem 5.1) and replace ψ_0 with

$$\begin{aligned} \mathfrak{S}_2(G, q) &\hookrightarrow \text{in}(I_G) \\ (U_1 \subsetneq U_2) &\mapsto \mathbf{x}^{D(U_2 \setminus U_1, U_1)}[0] , \end{aligned}$$

then the exact same statement and proof are correct for the case of $\text{in}(I_G)$. As a corollary the exact same recipe gives a free resolution for $\text{in}(I_G)$ as well.

6. MINIMALITY OF THE RESOLUTION OF I_G AND $\text{in}(I_G)$

In Theorem 5.3 and Remark 5.4 we constructed free resolutions for the ideals I_G and $\text{in}(I_G)$. In this section we take a close look at (5.3) to show that the constructed resolutions are indeed minimal. Note that the basis elements of the free module F_k correspond to elements of $\mathfrak{S}_{k+2}(G, q)$. We show that for any $\mathcal{U} \in \mathfrak{S}_{k+2}(G, q)$ the corresponding basis element $[\psi_k(\mathcal{U})]$ maps to a combination of basis elements $[\psi_{k-1}(\mathcal{V})]$, where each \mathcal{V} is obtained from \mathcal{U} by *merging* (§6.2) appropriate connected parts. Moreover, the coefficients appearing in this combination are all non-units and, therefore, the constructed resolution is minimal (Theorem 6.17).


6.1. **Contraction map.** To understand merging, we first need to study contraction maps.

Definition 6.1. Assume that $\mathcal{U} \in \mathfrak{S}_k(G, q)$. Let G/\mathcal{U} be the graph obtained from G by contracting the unoriented edges of $G(\mathcal{U})$ and let $\phi : G \rightarrow G/\mathcal{U}$ be the contraction map. More precisely, G/\mathcal{U} is the graph on the vertices u_1, \dots, u_k corresponding to the collection $(U_i \setminus U_{i-1})_{i=1}^k$, i.e. $u_i = \phi(U_i \setminus U_{i-1})$. For any edge between $U_i \setminus U_{i-1}$ and $U_j \setminus U_{j-1}$ there is an edge between u_i and u_j .

Example 6.2. Let G be the graph in Example 4.4. For

$$\mathcal{U} : \{v_1\} \subset \{v_1, v_2\} \subset \{v_1, v_2, v_3, v_4\} \subset \{v_1, v_2, v_3, v_4, v_5\}$$

the graph G/\mathcal{U} depicted in the following figure in which $u_1 = v_1$, $u_2 = v_2$, the vertex u_3 corresponds to $U_3 \setminus U_2 = \{v_3, v_4\}$, and u_4 corresponds to $U_4 \setminus U_3 = \{v_5\}$.



Remark 6.3. The contraction map $\phi : G \rightarrow G/\mathcal{U}$ induces the map

$$\phi_* : \text{Div}(G) \rightarrow \text{Div}(G/\mathcal{U}) \quad \text{with} \quad \phi_*\left(\sum_{v \in V(G)} a_v(v)\right) = \sum_{v \in V(G)} a_v(\phi(v)) .$$

If the indices i and j are given, we obtain two divisors

$$D'(u_i, u_j) \in \text{Div}(G/\mathcal{U}) \quad \text{and} \quad D(U_i \setminus U_{i-1}, U_j \setminus U_{j-1}) \in \text{Div}(G)$$

which are related by the map ϕ_* (see (2.1)). Here we use the notation $D'(\cdot, \cdot)$ for divisors on G/\mathcal{U} and $D(\cdot, \cdot)$ for divisors on G .

In particular, an ordering on the vertices of G/\mathcal{U} gives an ordering on the collection of subsets $(U_i \setminus U_{i-1})_{i=1}^k$ of $V(G)$. By Definition 4.3 we get a divisor D' on G/\mathcal{U} and a divisor D on G , and $\phi_*(D) = D'$.

Remark 6.4. We also have the map $\phi^* : \mathfrak{F}_s(G/\mathcal{U}, u_1) \rightarrow \mathfrak{F}_s(G, q)$ induced by sending each vertex of G/\mathcal{U} to its preimage under ϕ . The map ϕ and the total ordering \preceq on $\mathfrak{C}^{\text{op}}(G, q)$ (as in Definition 4.6) give a total ordering \preceq' on $\mathfrak{C}^{\text{op}}(G/\mathcal{U}, u_1)$. The ordering \preceq' induces a strict total ordering \prec'_ℓ on $\mathfrak{F}_\ell(G/\mathcal{U}, u_1)$ compatible with the total ordering on connected flags on (G, q) ; that is, $\mathcal{X} \prec'_\ell \mathcal{Y}$ if and only if $\phi^*(\mathcal{X}) \prec_\ell \phi^*(\mathcal{Y})$. Therefore, we get a map

$$(6.1) \quad \phi^* : \mathfrak{S}_s(G/\mathcal{U}, u_1) \rightarrow \mathfrak{S}_s(G, q) .$$

This gives a one-to-one correspondence between the elements $\mathcal{V}' \in \mathfrak{S}_s(G/\mathcal{U}, u_1)$ and the elements $\mathcal{V} \in \mathfrak{S}_s(G, q)$. Under this correspondence, for any $u_i \in V_j' \setminus V_{j-1}'$ we have $U_i \setminus U_{i-1} \subseteq V_j \setminus V_{j-1}$ and thus $V_j = \bigcup_{u_i \in V_j'} (U_i \setminus U_{i-1})$. For any element \mathcal{V} in the image of ϕ^* the preimage \mathcal{V}' is obtained by $V_j' = \{u_i : u_i = \phi(U_i \setminus U_{i-1}) \text{ and } U_i \setminus U_{i-1} \subseteq V_j\}$. In particular, \mathcal{U} itself is in the image of ϕ^* .

The following example explains the notation introduced in the above remarks.

Example 6.5. In Example 6.2 the ordering u_1, u_2, u_3, u_4 on $V(G/\mathcal{U})$ induces the ordering $U_1 \setminus U_0, U_2 \setminus U_1, U_3 \setminus U_2, U_4 \setminus U_3$ on the collection $(U_i \setminus U_{i-1})_{i=1}^4$ of $V(G)$ which corresponds to \mathcal{U} . Also corresponding to the ordering u_1, u_2, u_3, u_4 on $V(G/\mathcal{U})$ we get the divisor $D' = (u_2) + 2(u_3) + 2(u_4)$ on G/\mathcal{U} , where $\phi_*(D') = D(\mathcal{U})$.

We consider

$$\mathcal{V}' : \{u_1\} \subset \{u_1, \mathbf{u}_3\} \subset \{u_1, \mathbf{u}_2, u_3, \mathbf{u}_4\}$$

in $\mathfrak{S}_3(G/\mathcal{U}, u_1)$. Then $\phi^*(\mathcal{V}') = \mathcal{V}$, where

$$\mathcal{V} : \{v_1\} \subset \{v_1, \mathbf{v}_3, \mathbf{v}_4\} \subset \{v_1, \mathbf{v}_2, v_3, v_4, \mathbf{v}_5\}.$$

More precisely, $V_1 = U_1 \setminus U_0, V_2 = (U_1 \setminus U_0) \cup (U_3 \setminus U_2)$, and $V_3 = (U_2 \setminus U_1) \cup (U_4 \setminus U_3)$.

6.2. Mergeable parts. Given $\mathcal{U} \in \mathfrak{S}_k(G, q)$ of the form

$$U_1 \subsetneq U_2 \subsetneq \cdots \subsetneq U_k = V(G)$$

it is sometimes more convenient to work with the connected partition given by $A_\ell := U_\ell \setminus U_{\ell-1}$ (for $1 \leq \ell \leq k$).

Recall for any $\mathcal{U} \in \mathfrak{S}_k(G, q)$ we get a partial orientation of G which we denoted by $G(\mathcal{U})$ in Definition 4.3. This partial orientation is acyclic with unique source on the underlying partition graph G/\mathcal{U} (Definition 6.1). This means that the underlying partition graph does not contain any directed cycle and it has a unique source on the vertex corresponding to A_1 . More generally, we say a partial orientation is *acyclic* if the associated oriented partition graph (obtained by contracting all unoriented edges) is acyclic. Equivalently, a partial orientation is acyclic if replacing every undirected edge with two antiparallel edges yields an acyclic directed graph. Recall that associated to each partial orientation we get a divisor as in Remark 4.5.


Definition 6.6. Let $\mathcal{U} \in \mathfrak{S}_k(G, q)$ and $A_\ell := U_\ell \setminus U_{\ell-1}$ (for $1 \leq \ell \leq k$) as before. We set $o_0(\mathcal{U}) := G(\mathcal{U})$. For $j > 0$ the partial orientation $o_j(\mathcal{U})$ is defined inductively as follows: we obtain $o_j(\mathcal{U})$ from $o_{j-1}(\mathcal{U})$ by reversing the orientation of the edges between A_j and $V(G) \setminus A_j$ in $o_{j-1}(\mathcal{U})$.

Note that, when all oriented edges are directed away from A_j , reversing the orientation of the edges between A_j and $V(G) \setminus A_j$ in $o_{j-1}(\mathcal{U})$ is equivalent to performing a chip-firing move, in which all vertices in A_j borrow chips from their neighbors in $V(G) \setminus A_j$. Note that $o_j(\mathcal{U})$ is well-defined since all edges are directed away from A_j in $o_{j-1}(\mathcal{U})$.

Definition 6.7. Let $\mathfrak{c}(\mathcal{U})$ denote the set consisting of all partial orientations $o_j(\mathcal{U})$ of G .

Example 6.8. Let G be the 4-cycle on the vertices 1, 2, 3, 4 such that 1 is the distinguished vertex. Let \mathcal{U} be the connected flag $\mathcal{U} : \{1\} \subset \{1, 2\} \subset \{1, 2, 3\} \subset \{1, 2, 3, 4\}$. Then in the following we depict the graph corresponding to $o_j(\mathcal{U})$ for all j (see Definition 6.6). Note that $o_4(\mathcal{U}) = G(\mathcal{U})$.

For disjoint subsets $A, B \subset V(G)$ let $E(A, B)$ denote the set of edges between A and B .



Definition 6.9. Let $\mathcal{U} \in \mathfrak{S}_k(G, q)$ and $A_\ell := U_\ell \setminus U_{\ell-1}$ (for $1 \leq \ell \leq k$) as before, and assume that there are some edges connecting A_i and A_j , that is, $E(A_i, A_j) \neq \emptyset$.

- (i) We say A_i is *mergeable* with A_j in $G(\mathcal{U})$ if all edges in $E(A_i, A_j)$ are oriented from A_i to A_j and the partial orientation obtained from $G(\mathcal{U})$ by removing the orientations on $E(A_i, A_j)$ is acyclic.

Note that in this case $i < j$. We let $\text{Merge}(\mathcal{U}; A_i, A_j) \in \mathfrak{S}_{k-1}(G, q)$ denote the corresponding unique connected $(k-1)$ -flag whose connected parts are A_ℓ (for $\ell \neq i, j$) and $A_i \cup A_j$.

- (ii) We say A_i is *mergeable* with A_j in $o_j(\mathcal{U})$ where $i > j > 0$, if the partial orientation obtained by removing the orientations on $E(A_i, A_j)$ in $o_j(\mathcal{U})$ results in an acyclic partial orientation. Note that $E(A_i, A_j)$ are oriented from A_i to A_j in $o_j(\mathcal{U})$.

Let $\text{Merge}(o_j(\mathcal{U}); A_i, A_j) \in \mathfrak{S}_{k-1}(G, q)$ denote the connected partition of G whose connected parts are A_ℓ (for $\ell \neq i, j$) and $A_i \cup A_j$, together with the acyclic (partial) orientation obtained from $o_j(\mathcal{U})$ by removing the orientations on $E(A_i, A_j)$. As usual one obtains an associated divisor by reading the indegrees in this new partial orientation. This gives a *maximal reduced divisor* (see §7.1) on the associated graph of partitions via the map ϕ (see Remark 6.3). This maximal reduced divisor gives a total ordering on the vertices of the graph of partitions (e.g., by performing *Dhar's algorithm* – see §7.1 and [2]). Consider the induced partial orientation of G obtained in this way, and let $\text{Merge}(\mathbf{c}(\mathcal{U}); A_i, A_j) \in \mathfrak{S}_{k-1}(G, q)$ denote the associated connected flag.

Definition 6.10. For $\mathcal{U} \in \mathfrak{S}_k(G, q)$ we associate two subsets of $\mathfrak{S}_{k-1}(G, q)$ as follows:




- (i) $\mathfrak{J}(\mathcal{U}) := \{\mathcal{W} : \mathcal{W} = \text{Merge}(\mathcal{U}; A_i, A_j), \text{ for } A_i, A_j \text{ mergeable in } G(\mathcal{U})\}$.
(ii) $\mathfrak{B}(\mathcal{U}) := \{\mathcal{W} : \mathcal{W} = \text{Merge}(\mathbf{c}(\mathcal{U}); A_i, A_j), \text{ for } A_i, A_j \text{ mergeable in } o_j(\mathcal{U}) \text{ or } G(\mathcal{U})\}$.

It immediately follows from the definitions that $\mathfrak{J}(\mathcal{U}) \subseteq \mathfrak{B}(\mathcal{U})$. As we will see soon, $\mathfrak{B}(\mathcal{U})$ is related to the differential maps in our resolution of the binomial ideal I_G and $\mathfrak{J}(\mathcal{U})$ is related to the differential maps in our resolution of the monomial ideal $\text{in}(I_G)$.

Example 6.11. Let G be of the 4-cycle on the vertices 1, 2, 3, 4 in which we fix 1 be the distinguished vertex. Let \mathcal{U} be the connected flag $\mathcal{U} : \{1\} \subset \{1, 2\} \subset \{1, 2, 3\} \subset \{1, 2, 3, 4\}$. Here we list the elements of $\mathfrak{J}(\mathcal{U})$.

Example 6.12. Returning to Example 6.8, i.e., for $\mathcal{U} : \{1\} \subset \{1, 2\} \subset \{1, 2, 3\} \subset \{1, 2, 3, 4\}$, we list the acyclic orientations $\text{Merge}(\mathbf{c}(\mathcal{U}); A_i, A_j)$ corresponding to the elements of $\mathfrak{B}(\mathcal{U}) \setminus \mathfrak{J}(\mathcal{U})$ which have been obtained from connected partitions $\text{Merge}(o_j(\mathcal{U}); A_i, A_j)$.

Lemma 6.13. Let $\mathcal{U} \in \mathfrak{S}_k(G, q)$ and assume that $E(A_i, A_j) \neq \emptyset$.



(a) Assume that $\text{Merge}(\mathcal{U}; A_i, A_j) \in \mathfrak{I}(\mathcal{U})$. Then there exists $B_i \supseteq A_i$ such that

$$\text{Merge}(\mathcal{U}^{(1)}; B_i, A_j) \in \mathfrak{I}(\mathcal{U}^{(1)}) \quad \text{or} \quad \text{Merge}(\mathcal{U}^{(2)}; B_i, A_j) \in \mathfrak{I}(\mathcal{U}^{(2)}) .$$

(b) Assume that $\text{Merge}(o_j(\mathcal{U}); A_i, A_j) \in \mathfrak{B}(\mathcal{U}) \setminus \mathfrak{I}(\mathcal{U})$ for $j > 0$. Then

- $\text{Merge}(o_{j-1}(\mathcal{U}^{(1)}); A_i, A_j) \in \mathfrak{B}(\mathcal{U}^{(1)})$, or
- there exist $\mathcal{W} \in \mathfrak{I}(\mathcal{U})$, $B_i \supseteq A_i$ and $B_j \supseteq A_j$ such that

$$\text{Merge}(o_1(\mathcal{W}); B_i, B_j) \in \mathfrak{B}(\mathcal{W}) .$$

Proof. (a) Since A_1 is a source in the partial orientation and cannot appear in any directed cycle we have the following:

- (i) if $A_i \neq A_1, A_2$ then $\text{Merge}(\mathcal{U}^{(1)}; A_i, A_j) \in \mathfrak{I}(\mathcal{U}^{(1)})$.
- (ii) if $A_i = A_1$ then $\text{Merge}(\mathcal{U}^{(1)}; A_1 \cup A_2, A_j) \in \mathfrak{I}(\mathcal{U}^{(1)})$.
- (iii) if $A_i = A_2$
 - if $E(A_2, A_3) = \emptyset$ then $\text{Merge}(\mathcal{U}^{(2)}; A_2, A_j) \in \mathfrak{I}(\mathcal{U}^{(2)})$.
 - if $E(A_2, A_3) \neq \emptyset$ then $\text{Merge}(\mathcal{U}^{(2)}; A_2 \cup A_3, A_j) \in \mathfrak{I}(\mathcal{U}^{(2)})$.

In other words, in each case, we can find a $B_i \supseteq A_i$ such that $\text{Merge}(\mathcal{U}^{(1)}; B_i, A_j) \in \mathfrak{J}(\mathcal{U}^{(1)})$ or $\text{Merge}(\mathcal{U}^{(2)}; B_i, A_j) \in \mathfrak{J}(\mathcal{U}^{(2)})$.

(b) Assume that $\text{Merge}(o_j(\mathcal{U}); A_i, A_j) \in \mathfrak{B}(\mathcal{U}) \setminus \mathfrak{J}(\mathcal{U})$. First assume that G/\mathcal{U} is a star graph, i.e., for each pair $\ell_1, \ell_2 > 1$ of indices $E(A_{\ell_1}, A_{\ell_2}) = \emptyset$. Then $A_j = A_1$ and A_ℓ is mergeable with A_j in $G(\mathcal{U})$ for all $\ell > 1$. In particular, A_2 is mergeable with $A_1 \cup A_3$ in $\mathcal{U}^{(2)}$ and A_i is mergeable with $A_1 \cup A_2$ in $\mathcal{U}^{(1)}$ for $i > 2$ which is what we want.

Now we assume that $E(A_{\ell_1}, A_{\ell_2}) \neq \emptyset$ for some ℓ_1, ℓ_2 . Then we define

$$r := \min\{p : E(A_p, A_\ell) \neq \emptyset \text{ have the same orientations in } G(\mathcal{U}) \text{ and } o_j(\mathcal{U}) \text{ for some } \ell\}$$

and

$$s := \min\{\ell : E(A_r, A_\ell) \neq \emptyset \text{ have the same orientations in } G(\mathcal{U}) \text{ and } o_j(\mathcal{U})\}.$$

Note that our assumption on $E(A_{\ell_1}, A_{\ell_2})$ shows that these sets are nonempty. We also have $A_j \neq A_r$ since the outdegree of each vertex of A_j in $o_j(\mathcal{U})$ is zero, but there are some edges going out from A_r to A_s . We set $\mathcal{W} = \text{Merge}(\mathcal{U}; A_r, A_s)$. In the following we consider all possible cases to show that $\mathcal{W} \in \mathfrak{J}(\mathcal{U})$ with the desired properties:

- (i) $A_i \neq A_1$: then the edges between A_1 and A_2 are oriented from A_1 to A_2 in $o_j(\mathcal{U})$ and $G(\mathcal{U})$. We first reverse the orientation of the edges between $A_1 \cup A_2$ and $V(G) \setminus (A_1 \cup A_2)$. If $A_j = A_2$ then A_i is mergeable with $A_1 \cup A_2$ in $o_1(\mathcal{U}^{(1)})$ and so $\text{Merge}(o_1(\mathcal{U}^{(1)}); A_1 \cup A_2, A_i) \in \mathfrak{B}(\mathcal{U}^{(1)})$. Assume that $A_j \neq A_2$. Then we let the vertices of A_3 borrow from their neighbors in $V(G) \setminus A_3$ in order to get $o_2(\mathcal{U}^{(1)})$ which differs with $o_3(\mathcal{U})$ just by merging the parts A_1 and A_2 . We continue the same process of chip firing on the parts of A_4, \dots, A_j step-by-step in order to get $o_{j-1}(\mathcal{U}^{(1)})$ which can be obtained from $o_j(\mathcal{U})$ just by merging the parts A_1 and A_2 . This shows that A_i is mergeable with A_j in $o_{j-1}(\mathcal{U}^{(1)})$ as well, and so $\text{Merge}(o_{j-1}(\mathcal{U}^{(1)}); A_i, A_j) \in \mathfrak{B}(\mathcal{U}^{(1)})$.
- (ii) $A_i = A_1$: then the same argument as case (i) shows the following cases can happen:
 - if $A_j \neq A_r, A_s$ then $\text{Merge}(o_1(\mathcal{W}); A_1, A_j) \in \mathfrak{B}(\mathcal{W})$.
 - if $A_j = A_r$ or $A_j = A_s$ then $\text{Merge}(\mathfrak{c}(\mathcal{W}); A_1, A_r \cup A_s) \in \mathfrak{B}(\mathcal{W})$. □

There is a nice converse to Lemma 6.13(a). Our next result shows that the mergeable parts of \mathcal{U} can be obtained from mergeable parts of the canonical flags $\mathcal{U}^{(1)}$ and $\mathcal{U}^{(2)}$.

Lemma 6.14. *There exists a one-to-one correspondence between elements $\mathfrak{J}(\mathcal{U}^{(1)}) \cup \mathfrak{J}(\mathcal{U}^{(2)})$ and elements of $\mathfrak{J}(\mathcal{U})$.*

Proof. Let $\mathcal{U} \in \mathfrak{S}_k(G, q)$. Corresponding to each pair of mergeable parts in $G(\mathcal{U}^{(1)})$ or $G(\mathcal{U}^{(2)})$ we will find a unique pair of mergeable parts in $G(\mathcal{U})$. Assume that $\mathcal{W} \in \mathfrak{J}(\mathcal{U}^{(1)}) \cup \mathfrak{J}(\mathcal{U}^{(2)})$. Then we consider the following cases:

- $\mathcal{W} \in \mathfrak{J}(\mathcal{U}^{(1)})$: Since A_1 is a source in the partial orientation and cannot appear in any directed cycle we have the following:

- (i) if $\text{Merge}(\mathcal{U}^{(1)}; A_i, A_j) \in \mathfrak{J}(\mathcal{U}^{(1)})$ then A_i is mergeable with A_j in $G(\mathcal{U})$.
- (ii) if $\text{Merge}(\mathcal{U}^{(1)}; A_1 \cup A_2, A_i) \in \mathfrak{J}(\mathcal{U}^{(1)})$ then
 - $\text{Merge}(\mathcal{U}; A_2, A_i) \in \mathfrak{J}(\mathcal{U})$ if $E(A_2, A_i) \neq \emptyset$.
 - $\text{Merge}(\mathcal{U}; A_1, A_i) \in \mathfrak{J}(\mathcal{U})$ if $E(A_2, A_i) = \emptyset$.

- $\mathcal{W} \in \mathfrak{J}(\mathcal{U}^{(2)})$: First of all note that $\text{Merge}(\mathcal{U}^{(2)}; A_1, A_2) \in \mathfrak{J}(\mathcal{U})$. Then we have the following cases:

- (i) if $\text{Merge}(\mathcal{U}^{(2)}; A_i, A_j) \in \mathfrak{J}(\mathcal{U}^{(2)})$ then $\text{Merge}(\mathcal{U}; A_i, A_j) \in \mathfrak{J}(\mathcal{U})$.
- (ii) if $\text{Merge}(\mathcal{U}^{(2)}; A_2 \cup A_3, A_i) \in \mathfrak{J}(\mathcal{U}^{(2)})$ then
 - $\text{Merge}(\mathcal{U}; A_3, A_i) \in \mathfrak{J}(\mathcal{U})$ if $E(A_3, A_i) \neq \emptyset$.
 - $\text{Merge}(\mathcal{U}; A_2, A_i) \in \mathfrak{J}(\mathcal{U})$ if $E(A_3, A_i) = \emptyset$.
- (iii) if $\text{Merge}(\mathcal{U}^{(2)}; A_1 \cup A_3, A_i) \in \mathfrak{J}(\mathcal{U}^{(2)})$ then
 - $\text{Merge}(\mathcal{U}; A_3, A_i) \in \mathfrak{J}(\mathcal{U})$ if $E(A_3, A_i) \neq \emptyset$.
 - $\text{Merge}(\mathcal{U}; A_1, A_i) \in \mathfrak{J}(\mathcal{U})$ if $E(A_3, A_i) = \emptyset$.

In Lemma 6.13(a) we have already shown that each element $\text{Merge}(\mathcal{U}; A_i, A_j)$ of $\mathfrak{J}(\mathcal{U})$ corresponds to an element of $\mathfrak{J}(\mathcal{U}^{(1)}) \cup \mathfrak{J}(\mathcal{U}^{(2)})$. \square

6.3. Incidence function. Signs of the summands in the image of the basis elements under differential maps can be read from incidence functions as follows.

Assume $\mathcal{U} \in \mathfrak{S}_k(G, q)$ for $3 \leq k \leq n$. For $\mathcal{W} \in \mathfrak{B}(\mathcal{U})$ we want to define an *incidence value* $\epsilon(\mathcal{U}, \mathcal{W}) \in \{-1, +1\}$. For this we look at two set of natural permutations on parts of \mathcal{U} and on parts of \mathcal{W} . Let $\mathcal{W} = \text{Merge}(\mathfrak{c}(\mathcal{U}); A_i, A_j)$.

- Let $\delta(\mathcal{U}) = (A_1, A_2, \dots, A_k)$ and $\delta(\mathcal{W}) = (A_{\ell_1}, A_{\ell_2}, \dots, A_i \cup A_j, \dots, A_{\ell_{k-1}})$ denote the permutations corresponding to the fixed ordering of parts (as fixed by the choice of minimal representatives of the classes in $\mathfrak{E}_k(G, q)$ with respect to \prec_k).
- Let $\alpha(\mathcal{U}) = (A_i, A_j, A_{s_1}, \dots, A_{s_{k-2}})$ be an arbitrary permutation which fixes A_i and A_j at the beginning and is arbitrary otherwise. Correspondingly, we define $\alpha(\mathcal{W}) = (A_i \cup A_j, A_{s_1}, \dots, A_{s_{k-2}})$ compatible with $\alpha(\mathcal{U})$.

Definition 6.15. We define

- (i) $\epsilon(\mathcal{U}, \mathcal{W}) = \text{sgn}(\delta(\mathcal{U}), \alpha(\mathcal{U})) \text{sgn}(\delta(\mathcal{W}), \alpha(\mathcal{W}))$, where $\text{sgn}(\cdot, \cdot)$ denote the standard sign function for permutations.
- (ii) $\theta(\mathcal{U}, \mathcal{W}) = D(A_j, A_i)$.

The definition of $\epsilon(\mathcal{U}, \mathcal{W})$ is easily seen to be independent of the choice of $\alpha(\mathcal{U})$, because if $\alpha(\mathcal{U})$ is replaced with $\alpha'(\mathcal{U}) = (A_i, A_j, A_{t_1}, \dots, A_{t_{k-2}})$ then

$$\text{sgn}(\alpha'(\mathcal{U}), \alpha(\mathcal{U})) = \text{sgn}(\alpha'(\mathcal{W}), \alpha(\mathcal{W})) = \text{sgn}((A_{t_1}, \dots, A_{t_{k-1}}), (A_{s_1}, \dots, A_{s_{k-2}}))$$

and $\epsilon(\mathcal{U}, \mathcal{W})$ is multiplied by $\text{sgn}(\alpha'(\mathcal{U}), \alpha(\mathcal{U}))^2 = 1$. It is easy to see that $\theta(\mathcal{U}, \mathcal{W})$ is also well-defined, and is independent of the choice acyclic orientation on $\mathfrak{c}(\mathcal{U})$ where A_i and A_j are mergeable.

Proposition 6.16. Fix $3 \leq k \leq n$ and let $\mathcal{U} \in \mathfrak{S}_k(G, q)$. For any $\mathcal{W} \in \mathfrak{B}(\mathcal{U})$ and $\mathcal{X} \in \mathfrak{B}(\mathcal{W})$ there exists a unique $\mathcal{W}' \in \mathfrak{B}(\mathcal{U})$ such that $\mathcal{X} \in \mathfrak{B}(\mathcal{W}')$ and

$$\epsilon(\mathcal{U}, \mathcal{W})\epsilon(\mathcal{W}, \mathcal{X}) = -\epsilon(\mathcal{U}, \mathcal{W}')\epsilon(\mathcal{W}', \mathcal{X})$$

$$\theta(\mathcal{U}, \mathcal{W}) + \theta(\mathcal{W}, \mathcal{X}) = \theta(\mathcal{U}, \mathcal{W}') + \theta(\mathcal{W}', \mathcal{X}) .$$

Moreover, if $\mathcal{W} \in \mathfrak{J}(\mathcal{U})$ and $\mathcal{X} \in \mathfrak{J}(\mathcal{W})$ we have $\mathcal{W}' \in \mathfrak{J}(\mathcal{U})$ and $\mathcal{X} \in \mathfrak{J}(\mathcal{W}')$.

In particular we have

$$\sum_{\substack{\mathcal{W} \in \mathfrak{B}(\mathcal{U}) \\ \mathcal{X} \in \mathfrak{B}(\mathcal{W})}} \epsilon(\mathcal{U}, \mathcal{W})\epsilon(\mathcal{W}, \mathcal{X}) \mathbf{x}^{\theta(\mathcal{U}, \mathcal{W})} \mathbf{x}^{\theta(\mathcal{W}, \mathcal{X})} [\psi(\mathcal{X})] = 0$$

and

$$\sum_{\substack{\mathcal{W} \in \mathfrak{J}(\mathcal{U}) \\ \mathcal{X} \in \mathfrak{J}(\mathcal{W})}} \epsilon(\mathcal{U}, \mathcal{W})\epsilon(\mathcal{W}, \mathcal{X}) \mathbf{x}^{\theta(\mathcal{U}, \mathcal{W})} \mathbf{x}^{\theta(\mathcal{W}, \mathcal{X})} [\psi(\mathcal{X})] = 0 .$$

Proof. Let the connected parts of \mathcal{U} be A_ℓ for $1 \leq \ell \leq k$. Assume that $\mathcal{W} = \text{Merge}(\mathbf{c}(\mathcal{U}); A_i, A_j)$ and $\mathcal{X} = \text{Merge}(\mathbf{c}(\mathcal{W}); B_r, B_s)$. Since connected parts B_r and B_s of \mathcal{W} are among A_ℓ (for $\ell \neq i, j$) and $A_i \cup A_j$, we need to consider three cases.

- $B_r = A_r, B_s = A_s$: In this case A_r and A_s are mergeable in $\mathbf{c}(\mathcal{U})$ and we let $\mathcal{W}' = \text{Merge}(\mathbf{c}(\mathcal{U}); A_r, A_s)$. Then clearly $\mathcal{X} = \text{Merge}(\mathbf{c}(\mathcal{W}'); A_i, A_j)$. It follows that

$$\theta(\mathcal{U}, \mathcal{W}) + \theta(\mathcal{W}, \mathcal{X}) = \theta(\mathcal{U}, \mathcal{W}') + \theta(\mathcal{W}', \mathcal{X}) = D(A_j, A_i) + D(A_s, A_r) .$$

There is a unique $\mathcal{W}' \neq \mathcal{W}$ with this property because there are only two ways to merge A_i with A_j and A_r with A_s . Let

$$\begin{aligned} \alpha(\mathcal{U}) &= (A_i, A_j, A_r, A_s, \dots) & , & & \alpha'(\mathcal{U}) &= (A_r, A_s, A_i, A_j, \dots), \\ \alpha(\mathcal{W}) &= (A_i \cup A_j, A_r, A_s, \dots) & , & & \alpha'(\mathcal{W}') &= (A_r \cup A_s, A_i, A_j, \dots), \\ \beta(\mathcal{W}) &= (A_r, A_s, A_i \cup A_j, \dots) & , & & \beta'(\mathcal{W}') &= (A_i, A_j, A_r \cup A_s, \dots), \\ \beta(\mathcal{X}) &= (A_r \cup A_s, A_i \cup A_j, \dots) & , & & \beta'(\mathcal{X}) &= (A_i \cup A_j, A_r \cup A_s, \dots). \end{aligned}$$

From Definition 6.15 we know

$$\begin{aligned} \epsilon(\mathcal{U}, \mathcal{W})\epsilon(\mathcal{W}, \mathcal{X}) &= \text{sgn}(\delta(\mathcal{U}), \alpha(\mathcal{U})) \text{sgn}(\delta(\mathcal{W}), \alpha(\mathcal{W})) \text{sgn}(\delta(\mathcal{W}), \beta(\mathcal{W})) \text{sgn}(\delta(\mathcal{X}), \beta(\mathcal{X})) \\ \epsilon(\mathcal{U}, \mathcal{W}')\epsilon(\mathcal{W}', \mathcal{X}) &= \text{sgn}(\delta(\mathcal{U}), \alpha'(\mathcal{U})) \text{sgn}(\delta(\mathcal{W}'), \alpha'(\mathcal{W}')) \text{sgn}(\delta(\mathcal{W}'), \beta'(\mathcal{W}')) \text{sgn}(\delta(\mathcal{X}), \beta'(\mathcal{X})). \end{aligned}$$

The result follows from

$$\begin{aligned} \text{sgn}(\delta(\mathcal{W}), \alpha(\mathcal{W})) \text{sgn}(\delta(\mathcal{W}), \beta(\mathcal{W})) &= \text{sgn}(\alpha(\mathcal{W}), \beta(\mathcal{W})) = 1, \\ \text{sgn}(\delta(\mathcal{W}'), \alpha'(\mathcal{W}')) \text{sgn}(\delta(\mathcal{W}'), \beta'(\mathcal{W}')) &= \text{sgn}(\alpha'(\mathcal{W}'), \beta'(\mathcal{W}')) = 1, \\ \text{sgn}(\delta(\mathcal{U}), \alpha(\mathcal{U})) \text{sgn}(\delta(\mathcal{U}), \alpha'(\mathcal{U})) &= \text{sgn}(\alpha(\mathcal{U}), \alpha'(\mathcal{U})) = 1, \\ \text{sgn}(\delta(\mathcal{X}), \beta(\mathcal{X})) \text{sgn}(\delta(\mathcal{X}), \beta'(\mathcal{X})) &= \text{sgn}(\beta(\mathcal{X}), \beta'(\mathcal{X})) = -1. \end{aligned}$$

- $B_r = A_r, B_s = A_i \cup A_j$: There are two cases:

(1) $E(A_r, A_i) \neq \emptyset$ in which case we let $\mathcal{W}' = \text{Merge}(\mathbf{c}(\mathcal{U}); A_r, A_i)$, and we have $\mathcal{X} = \text{Merge}(\mathbf{c}(\mathcal{W}'); A_r \cup A_i, A_j)$,

(2) $E(A_r, A_i) = \emptyset$ in which case we must have $E(A_r, A_j) \neq \emptyset$ and we let $\mathcal{W}' = \text{Merge}(\mathbf{c}(\mathcal{U}); A_r, A_j)$. We then have $\mathcal{X} = \text{Merge}(\mathbf{c}(\mathcal{W}'); A_i, A_r \cup A_j)$.

In each case it follows that

$$\theta(\mathcal{U}, \mathcal{W}) + \theta(\mathcal{W}, \mathcal{X}) = \theta(\mathcal{U}, \mathcal{W}') + \theta(\mathcal{W}', \mathcal{X}) = D(A_j, A_i) + D(A_i, A_r) + D(A_j, A_r) .$$

There is a unique $\mathcal{W}' \neq \mathcal{W}$ with this property because there are only two ways to merge A_i, A_j and A_r .

We now verify the equality for the incidence function ϵ in case (1). Let

$$\begin{aligned} \alpha(\mathcal{U}) &= (A_i, A_j, A_r, \dots) & , & & \alpha'(\mathcal{U}) &= (A_r, A_i, A_j, \dots), \\ \alpha(\mathcal{W}) &= (A_i \cup A_j, A_r, \dots) & , & & \alpha'(\mathcal{W}') &= (A_r \cup A_i, A_j, \dots), \\ \beta(\mathcal{W}) &= (A_r, A_i \cup A_j, \dots) & , & & \beta'(\mathcal{W}') &= \alpha'(\mathcal{W}'), \\ \beta(\mathcal{X}) &= (A_r \cup A_i \cup A_j, \dots) & , & & \beta'(\mathcal{X}) &= \beta(\mathcal{X}). \end{aligned}$$

From Definition 6.15 we know

$$\begin{aligned} \epsilon(\mathcal{U}, \mathcal{W})\epsilon(\mathcal{W}, \mathcal{X}) &= \text{sgn}(\delta(\mathcal{U}), \alpha(\mathcal{U})) \text{sgn}(\delta(\mathcal{W}), \alpha(\mathcal{W})) \text{sgn}(\delta(\mathcal{W}), \beta(\mathcal{W})) \text{sgn}(\delta(\mathcal{X}), \beta(\mathcal{X})), \\ \epsilon(\mathcal{U}, \mathcal{W}')\epsilon(\mathcal{W}', \mathcal{X}) &= \text{sgn}(\delta(\mathcal{U}), \alpha'(\mathcal{U})) \text{sgn}(\delta(\mathcal{W}'), \alpha'(\mathcal{W}')) \text{sgn}(\delta(\mathcal{W}'), \beta'(\mathcal{W}')) \text{sgn}(\delta(\mathcal{X}), \beta'(\mathcal{X})). \end{aligned}$$

The result follows from

$$\text{sgn}(\delta(\mathcal{W}), \alpha(\mathcal{W})) \text{sgn}(\delta(\mathcal{W}), \beta(\mathcal{W})) = \text{sgn}(\alpha(\mathcal{W}), \beta(\mathcal{W})) = -1,$$

$$\begin{aligned}
\operatorname{sgn}(\delta(\mathcal{W}'), \alpha'(\mathcal{W}')) \operatorname{sgn}(\delta(\mathcal{W}'), \beta'(\mathcal{W}')) &= \operatorname{sgn}(\alpha'(\mathcal{W}'), \beta'(\mathcal{W}')) = 1, \\
\operatorname{sgn}(\delta(\mathcal{U}), \alpha(\mathcal{U})) \operatorname{sgn}(\delta(\mathcal{U}), \alpha'(\mathcal{U})) &= \operatorname{sgn}(\alpha(\mathcal{U}), \alpha'(\mathcal{U})) = 1, \\
\operatorname{sgn}(\delta(\mathcal{X}), \beta(\mathcal{X})) \operatorname{sgn}(\delta(\mathcal{X}), \beta'(\mathcal{X})) &= \operatorname{sgn}(\beta(\mathcal{X}), \beta'(\mathcal{X})) = 1.
\end{aligned}$$

For case (2) this verification is completely analogous. Let

$$\begin{aligned}
\alpha(\mathcal{U}) &= (A_i, A_j, A_r, \dots) \quad , \quad \alpha'(\mathcal{U}) = (A_r, A_j, A_i, \dots), \\
\alpha(\mathcal{W}) &= (A_i \cup A_j, A_r, \dots) \quad , \quad \alpha'(\mathcal{W}') = (A_r \cup A_j, A_i, \dots), \\
\beta(\mathcal{W}) &= (A_r, A_i \cup A_j, \dots) \quad , \quad \beta'(\mathcal{W}') = (A_i, A_r \cup A_j, \dots), \\
\beta(\mathcal{X}) &= (A_r \cup A_i \cup A_j, \dots) \quad , \quad \beta'(\mathcal{X}) = \beta(\mathcal{X}).
\end{aligned}$$

From Definition 6.15 we know

$$\begin{aligned}
\epsilon(\mathcal{U}, \mathcal{W})\epsilon(\mathcal{W}, \mathcal{X}) &= \operatorname{sgn}(\delta(\mathcal{U}), \alpha(\mathcal{U})) \operatorname{sgn}(\delta(\mathcal{W}), \alpha(\mathcal{W})) \operatorname{sgn}(\delta(\mathcal{W}), \beta(\mathcal{W})) \operatorname{sgn}(\delta(\mathcal{X}), \beta(\mathcal{X})), \\
\epsilon(\mathcal{U}, \mathcal{W}')\epsilon(\mathcal{W}', \mathcal{X}) &= \operatorname{sgn}(\delta(\mathcal{U}), \alpha'(\mathcal{U})) \operatorname{sgn}(\delta(\mathcal{W}'), \alpha'(\mathcal{W}')) \operatorname{sgn}(\delta(\mathcal{W}'), \beta'(\mathcal{W}')) \operatorname{sgn}(\delta(\mathcal{X}), \beta'(\mathcal{X})).
\end{aligned}$$

The result follows from

$$\begin{aligned}
\operatorname{sgn}(\delta(\mathcal{W}), \alpha(\mathcal{W})) \operatorname{sgn}(\delta(\mathcal{W}), \beta(\mathcal{W})) &= \operatorname{sgn}(\alpha(\mathcal{W}), \beta(\mathcal{W})) = -1, \\
\operatorname{sgn}(\delta(\mathcal{W}'), \alpha'(\mathcal{W}')) \operatorname{sgn}(\delta(\mathcal{W}'), \beta'(\mathcal{W}')) &= \operatorname{sgn}(\alpha'(\mathcal{W}'), \beta'(\mathcal{W}')) = -1, \\
\operatorname{sgn}(\delta(\mathcal{U}), \alpha(\mathcal{U})) \operatorname{sgn}(\delta(\mathcal{U}), \alpha'(\mathcal{U})) &= \operatorname{sgn}(\alpha(\mathcal{U}), \alpha'(\mathcal{U})) = -1, \\
\operatorname{sgn}(\delta(\mathcal{X}), \beta(\mathcal{X})) \operatorname{sgn}(\delta(\mathcal{X}), \beta'(\mathcal{X})) &= \operatorname{sgn}(\beta(\mathcal{X}), \beta'(\mathcal{X})) = 1.
\end{aligned}$$

• $B_s = A_r$, $B_r = A_i \cup A_j$: This case is proved precisely as the previous case by permuting the indices.

In all cases if $\mathcal{W} \in \mathfrak{J}(\mathcal{U})$ and $\mathcal{X} \in \mathfrak{J}(\mathcal{W})$ the constructed \mathcal{W}' is in $\mathfrak{J}(\mathcal{U})$. □

6.4. Differential maps and minimality of the free resolutions. We are now ready to use (5.3) and induction to give a precise description of the differential maps constructed in Theorem 5.3 (respectively, Remark 5.4) for I_G (respectively, $\operatorname{in}(I_G)$). The minimality of the constructed resolutions follows from this explicit description, as no units appear in the described differential maps. To simplify the notation we use ψ instead of ψ_k for all k (as defined in Theorem 5.3 and Remark 5.4).

Theorem 6.17. *Let $k \geq 0$ and $\mathcal{U} \in \mathfrak{S}_{k+2}(G, q)$.*

For I_G the differential maps given by (5.3) are of the form

$$\varphi_k([\psi(\mathcal{U})]) = \sum_{\mathcal{W} \in \mathfrak{B}(\mathcal{U})} \epsilon(\mathcal{U}, \mathcal{W}) \mathbf{x}^{\theta(\mathcal{U}, \mathcal{W})} [\psi(\mathcal{W})].$$

In particular, the set $\psi(\mathfrak{S}_{k+2}(G, q))$ minimally generates $\operatorname{syz}_k(\mathbf{G}(G, q))$.

Remark 6.18. For $\operatorname{in}(I_G)$ the differential maps given by (5.3) and the initial condition described in Remark 5.4 are of the form

$$\varphi_k([\psi(\mathcal{U})]) = \sum_{\mathcal{W} \in \mathfrak{J}(\mathcal{U})} \epsilon(\mathcal{U}, \mathcal{W}) \mathbf{x}^{\theta(\mathcal{U}, \mathcal{W})} [\psi(\mathcal{W})].$$

The proof is completely analogous to the binomial case, and is skipped here. We will not use this description when we discuss the Betti numbers because we instead appeal to Theorem 3.11.

Proof. For $\mathcal{U} \in \mathfrak{S}_{k+2}(G, q)$, from (5.3), we need to show that

$$(6.2) \quad s([\psi(\mathcal{U}^{(1)})], [\psi(\mathcal{U}^{(2)})]) = \sum_{\mathcal{W} \in \mathfrak{B}(\mathcal{U})} \epsilon(\mathcal{U}, \mathcal{W}) \mathbf{x}^{\theta(\mathcal{U}, \mathcal{W})} [\psi(\mathcal{W})] .$$

Note that this would prove the minimality of the resolution because \mathcal{W} is a connected flag and therefore $\theta(\mathcal{U}, \mathcal{W}) \neq 0$.

The proof is by induction on the number of vertices of the graph. The result is obvious for a graph with 2 vertices. Suppose the result holds for graphs with less than n vertices, and consider the graph G with n vertices. We need to show that all maps φ_k (for $0 \leq k \leq n-2$) are of the form (6.2). Fix a $\mathcal{U} \in \mathfrak{S}_{k+2}(G, q)$. We consider two cases:

- $0 \leq k < n-2$. We consider the graph G/\mathcal{U} on the vertex set $\{u_1, u_2, \dots, u_{k+2}\}$. This graph has fewer than n vertices because $k+2 < n$. Let $\mathcal{U}' \in \mathfrak{S}_{k+2}(G/\mathcal{U}, u_1)$ be the inverse image of \mathcal{U} under the map ϕ^* as described in Remark 6.4. By the induction hypothesis we have

$$(6.3) \quad \psi(\mathcal{U}') = \sum_{\mathcal{W}' \in \mathfrak{B}(\mathcal{U}')} \epsilon(\mathcal{U}', \mathcal{W}') \mathbf{x}^{\theta(\mathcal{U}', \mathcal{W}')} [\psi(\mathcal{W}')] .$$

For each \mathcal{W}' , let \mathcal{W} be the inverse image under the map ϕ^* . Then $\theta(\mathcal{U}, \mathcal{W})$ and $\theta(\mathcal{U}', \mathcal{W}')$ are related by the map ϕ_* of Remark 6.3, and there is a one-to-one correspondence between elements of $\mathfrak{B}(\mathcal{U}')$ and elements of $\mathfrak{B}(\mathcal{U})$. We claim that

$$(6.4) \quad \psi(\mathcal{U}) = \sum_{\mathcal{W} \in \mathfrak{B}(\mathcal{U})} \epsilon(\mathcal{U}, \mathcal{W}) \mathbf{x}^{\theta(\mathcal{U}, \mathcal{W})} [\psi(\mathcal{W})] ,$$

where $c(\mathcal{W}) = c(\mathcal{W}')$. To see this, we need to show that

$$(6.5) \quad \sum_{\mathcal{W} \in \mathfrak{B}(\mathcal{U})} \epsilon(\mathcal{U}, \mathcal{W}) \mathbf{x}^{\theta(\mathcal{U}, \mathcal{W})} \psi(\mathcal{W}) = 0 .$$

First note that (6.3) is equivalent to

$$(6.6) \quad \sum_{\mathcal{W}' \in \mathfrak{B}(\mathcal{U}')} \epsilon(\mathcal{U}', \mathcal{W}') \mathbf{x}^{\theta(\mathcal{U}', \mathcal{W}')} \psi(\mathcal{W}') = 0 .$$

We again use the induction hypothesis for $\psi(\mathcal{X}')$ to write

$$(6.7) \quad \sum_{\substack{\mathcal{W}' \in \mathfrak{B}(\mathcal{U}') \\ \mathcal{X}' \in \mathfrak{B}(\mathcal{W}')}} \epsilon(\mathcal{U}', \mathcal{W}') \epsilon(\mathcal{W}', \mathcal{X}') \mathbf{x}^{\theta(\mathcal{U}', \mathcal{W}')} \mathbf{x}^{\theta(\mathcal{W}', \mathcal{X}')} [\psi(\mathcal{X}')] = 0 .$$

Equation (6.7) implies that corresponding to each term $\mathbf{x}^{\theta(\mathcal{U}', \mathcal{W}'_1)} \mathbf{x}^{\theta(\mathcal{W}'_1, \mathcal{X}'_1)} [\psi(\mathcal{X}'_1)]$ there exists a term $\mathbf{x}^{\theta(\mathcal{U}', \mathcal{W}'_2)} \mathbf{x}^{\theta(\mathcal{W}'_2, \mathcal{X}'_2)} [\psi(\mathcal{X}'_2)]$ with opposite sign, with which it cancels. Therefore

$$(6.8) \quad E' := \theta(\mathcal{U}', \mathcal{W}'_1) + \theta(\mathcal{W}'_1, \mathcal{X}'_1) = \theta(\mathcal{U}', \mathcal{W}'_2) + \theta(\mathcal{W}'_2, \mathcal{X}'_2) \quad \text{and} \quad [\psi(\mathcal{X}'_1)] = [\psi(\mathcal{X}'_2)] .$$

Since ψ is injective (Theorem 5.3) we get $\mathcal{X}'_1 = \mathcal{X}'_2$. By the induction hypothesis we know

$$\theta(\mathcal{U}', \mathcal{W}'_1) = D'(u_i, u_j)$$

$$\theta(\mathcal{W}'_1, \mathcal{X}'_1) = \begin{cases} D'(u_r, u_s) & \text{or} \\ D'(\{u_i, u_j\}, u_r) & \text{or} \\ D'(u_r, \{u_i, u_j\}) & \end{cases}$$

for some distinct i, j, r, s (depending on what parts of \mathcal{W}' are merged to get $\mathcal{X}'_1 = \mathcal{X}'_2$).

Since E' can be written as a sum of $D'(u_a, u_b)$'s, once we recognize all a 's and b 's appearing in the sum, we can use Remark 6.3 and lift E' to some E as a sum of $D(U_a \setminus U_{a-1}, U_b \setminus U_{b-1})$'s. Then the same cancellations as in (6.6) occur in the left-hand side of (6.5) and we get zero.

- If $\mathcal{W}'_1 = \mathcal{W}'_2$, then $\theta(\mathcal{U}', \mathcal{W}'_1) = \theta(\mathcal{U}', \mathcal{W}'_2)$ and it follows from (6.8) that $\theta(\mathcal{W}'_1, \mathcal{X}'_1) = \theta(\mathcal{W}'_2, \mathcal{X}'_2)$. By looking at $D'(u_i, u_j)$ we can recognize u_i . By looking at the unique part in $\mathcal{W}'_1 = \mathcal{W}'_2$ that contains two elements, we recognize u_j . By looking at the vertices where $\theta(\mathcal{W}'_1, \mathcal{X}'_1) = \theta(\mathcal{W}'_2, \mathcal{X}'_2)$ is nonzero we can recognize u_r since $\mathcal{X}'_1 = \mathcal{X}'_2$.

- If $\mathcal{W}'_1 \neq \mathcal{W}'_2$, then we consider the following cases:

- (1) $\theta(\mathcal{W}'_1, \mathcal{X}'_1) = D'(u_r, u_s)$: we have $E' = D'(u_i, u_j) + D'(u_r, u_s)$. The places where E' is nonzero determine $\{u_i, u_r\}$. By looking at the two parts in $\mathcal{X}'_1 = \mathcal{X}'_2$ which contain precisely two vertices, we can distinguish $\{\{u_i, u_j\}, \{u_r, u_s\}\}$.

- (2) $\theta(\mathcal{W}'_1, \mathcal{X}'_1) = D'(\{u_i, u_j\}, u_r)$: we have $E' = D'(u_i, u_j) + D'(u_i, u_r) + D'(u_j, u_r)$. Since we know $\mathcal{X}'_1 = \mathcal{X}'_2$ we know $\{u_i, u_j, u_r\}$.

- if u_i and u_r are not adjacent: then u_r is the vertex where E' is zero. The vertex u such that $E'(u)$ is equal to the number of edges between u and $\{u_i, u_j, u_r\} \setminus u$ is u_i . The other vertex where E' is nonzero is u_j .

- if u_j and u_r are not adjacent: then u_i is the unique vertex where E' is nonzero. We do not need to distinguish between u_j and u_r because E' is of the form $E' = D'(u_i, u_j) + D'(u_i, u_r)$.

- (3) $\theta(\mathcal{W}'_1, \mathcal{X}'_1) = D'(u_r, \{u_i, u_j\})$: we have $E' = D'(u_i, u_j) + D'(u_r, u_i) + D'(u_r, u_j)$. Since we know $\mathcal{X}'_1 = \mathcal{X}'_2$ we know $\{u_i, u_j, u_r\}$. This case reduces to (2) by permuting the indices.

Therefore (6.5) holds. Note that $\mathcal{U}^{(1)}$ and $\mathcal{U}'^{(1)}$ (respectively, $\mathcal{U}^{(2)}$ and $\mathcal{U}'^{(2)}$) are related by the map ϕ^* (6.1). On the other hand by Remark 6.4 for each ℓ the total ordering \prec_ℓ corresponding to G and the total ordering \prec'_ℓ corresponding to G/\mathcal{U} (and so the term orderings $<_\ell$ and ℓ') are compatible. Hence it follows from the discussion above and Remark 6.4 that (6.5) is precisely coming from the s -polynomial computation.

- $k = n - 2$. Let $U_i \setminus U_{i-1} = \{v_i\}$ and for simplicity, $x_{v_i} = x_i$. By Theorem 5.3 $\psi(\mathcal{U}) = s(\psi(\mathcal{U}^{(1)}), \psi(\mathcal{U}^{(2)}))$. We directly apply the division algorithm to describe $s(\psi(\mathcal{U}^{(1)}), \psi(\mathcal{U}^{(2)}))$.

From the proof of Lemma 4.24 the coefficient of $[\psi(\mathcal{U}^{(1)})]$ is $\mathbf{x}^{\theta(\mathcal{U}, \mathcal{U}^{(1)})} = \mathbf{x}^{D(v_2, v_1)}$ and the coefficient of $[\psi(\mathcal{U}^{(2)})]$ is $\mathbf{x}^{\theta(\mathcal{U}, \mathcal{U}^{(2)})}$, where

$$\theta(\mathcal{U}, \mathcal{U}^{(2)}) = \begin{cases} D(v_3, v_2), & \text{if } v_2 \text{ and } v_3 \text{ are adjacent;} \\ D(v_3, v_1), & \text{if } v_2 \text{ and } v_3 \text{ are not adjacent.} \end{cases}$$

Now assume that

$$M := \text{LM}(\text{spoly}(\psi(\mathcal{U}^{(1)}), \psi(\mathcal{U}^{(2)}))) .$$

Since $\text{Image}(\psi)$ forms a minimal Gröbner bases of $(\text{syz}_{n-3}(\mathbf{G}(G, q)), <_{n-3})$ there exists an element $\mathcal{V} \in \mathfrak{S}_{n-1}(G, q)$ such that $\text{LM}(\psi(\mathcal{V}))$ divides M . We know from (5.2) that $\text{LM}(\psi(\mathcal{V})) = \mathbf{x}^{D(V_2 \setminus V_1, V_1)}[\psi(\mathcal{V}^{(1)})]$. Hence

$$M = \mathbf{x}^{\theta(\mathcal{U}, \mathcal{V}) + D(V_2 \setminus V_1, V_1)}[\psi(\mathcal{V}^{(1)})] \quad \text{for some } \theta(\mathcal{U}, \mathcal{V}) \geq 0 .$$

In the first step of the division algorithm we obtain

$$\mathbf{x}^{\theta(\mathcal{U}, \mathcal{U}^{(1)})} \psi(\mathcal{U}^{(1)}) - \mathbf{x}^{\theta(\mathcal{U}, \mathcal{U}^{(2)})} \psi(\mathcal{U}^{(2)}) + \epsilon(\mathcal{U}, \mathcal{V}) \mathbf{x}^{\theta(\mathcal{U}, \mathcal{V})} \psi(\mathcal{V}) .$$

The division algorithm proceeds by finding the leading monomial of the above expression and continuing similarly. To show (6.2) we show that in each step \mathcal{V} belongs to $\mathfrak{B}(\mathcal{U})$ and $\theta(\mathcal{U}, \mathcal{V})$ is of the form $D(v, v')$ for some vertices v and v' (and that all such terms appear). This is clearly correct for $\mathcal{V} = \mathcal{U}^{(1)}$ and $\mathcal{V} = \mathcal{U}^{(2)}$ as discussed above. Assume we are in i^{th} step. Recursively we may assume the leading monomial of the existing expression is a monomial of $\mathbf{x}^{D(v_i, v_j)} [\psi(\mathcal{W})]$.

Since $\mathcal{W} \in \mathfrak{S}_{n-1}(G, q)$ we can use the result of the previous case for $k = n - 3 < n - 2$ to write

$$\psi(\mathcal{W}) = \sum_{\mathcal{X} \in \mathfrak{B}(\mathcal{W})} \epsilon(\mathcal{W}, \mathcal{X}) \mathbf{x}^{\theta(\mathcal{W}, \mathcal{X})} [\psi(\mathcal{X})] ,$$

where $\theta(\mathcal{W}, \mathcal{X}) = D(W_r \setminus W_{r-1}, W_s \setminus W_{s-1})$ for some r, s .

Now let M be the term $\mathbf{x}^{D(v_i, v_j)} \mathbf{x}^{\theta(\mathcal{W}, \mathcal{X})} [\psi(\mathcal{X})]$ which is divisible by

$$\text{LM}(\psi(\mathcal{V})) = \mathbf{x}^{D(V_2 \setminus V_1, V_1)} [\psi(\mathcal{V}^{(1)})] .$$

Set $E' := D(v_i, v_j) + \theta(\mathcal{W}, \mathcal{X}) = D(v_i, v_j) + D(W_r \setminus W_{r-1}, W_s \setminus W_{s-1})$. Let $\{v, v'\}$ be the unique part of \mathcal{V} which contains two vertices. It is enough to show that $E' = D(V_2 \setminus V_1, V_1) + D(v, v')$ and $\mathcal{V} \in \mathfrak{B}(\mathcal{U})$. Depending on which parts of \mathcal{W} are merged to get \mathcal{X} we have the following cases:

Case 1. $\theta(\mathcal{W}, \mathcal{X}) = D(v_r, v_s)$: note that $\{v_r, v_s\}$ and $\{v_i, v_j\}$ are two disjoint parts of \mathcal{X} . Since V_2 has at least two elements we have $V_2 = \{v_i, v_j\}$ or $V_2 = \{v_r, v_s\}$. Now by noting that $v_1 \in V_2$ we can recognize V_2 . With no loss of generality assume that $V_2 = \{v_i, v_j\}$. Then we must have $v_j = v_1$; since $\mathbf{x}^{D(V_2 \setminus V_1, V_1)}$ divides $\mathbf{x}^{E'}$. Therefore $E' - D(V_2 \setminus V_1, V_1) = D(v_r, v_s)$. Note that $\mathcal{X} = \mathcal{V}^{(1)}$ implies that $\{v_r, v_s\}$ is the unique part of \mathcal{V} containing two vertices and so $D(v_r, v_s) = \theta(\mathcal{U}, \mathcal{V})$.

Case 2. $\theta(\mathcal{W}, \mathcal{X}) = D(v_r, \{v_i, v_j\})$: the ordered collection of $(X_i \setminus X_{i-1})_{i=1}^{n-2}$ is a permutation of the sets $\{v_t\}_{t \neq i, j, r}$ and $\{v_i, v_j, v_r\}$, since \mathcal{W} is obtained from \mathcal{U} by merging v_i, v_j and \mathcal{X} is obtained from \mathcal{W} by merging $v_r, \{v_i, v_j\}$ where v_r appears before $\{v_i, v_j\}$ in \mathcal{W} . So we have $V_2 = \{v_i, v_j, v_r\}$ and $v_1 \in V_2$. Therefore the following cases may occur:

- $V_1 = \{v_i, v_j\}$: then $D(V_2 \setminus V_1, V_1) = D(v_r, \{v_i, v_j\})$ and $E' - D(V_2 \setminus V_1, V_1) = D(v_i, v_j)$. Therefore $\theta(\mathcal{U}, \mathcal{V}) = D(v_i, v_j)$.
- $V_1 = \{v_j, v_r\}$: then $D(V_2 \setminus V_1, V_1) = D(v_i, \{v_j, v_r\})$ and v_j is adjacent to v_r . On the other hand, since $\text{LM}(\psi(\mathcal{V}))$ divides M we should have no edge between v_i and v_r . Therefore $E' - D(V_2 \setminus V_1, V_1) = D(v_r, v_j)$ which is equal to $\theta(\mathcal{U}, \mathcal{V})$.
- $V_1 = \{v_j\}$: then $D(V_2 \setminus V_1, V_1) = D(\{v_i, v_r\}, v_j)$ and v_i is adjacent to v_r . Therefore $\theta(\mathcal{U}, \mathcal{V}) = D(v_r, v_i)$.
- $v_j \in V_2 \setminus V_1$ and $v_i \in V_1$: the fact that v_i and v_j are adjacent implies that x_j divides M which is impossible since $\theta(\mathcal{U}, \mathcal{W})(v_j) = 0$.
- $V_1 = \{v_r\} = \{v_1\}$: then $D(V_2 \setminus V_1, V_1) = D(\{v_j, v_i\}, v_r)$ and two vertices v_i and v_r are adjacent. The number of edges between v_i and v_r is less than the number of edges between v_i and v_j . On the other hand, since $\text{LM}(\psi(\mathcal{V}))$ divides M we should have no edge between v_j and v_r .

Now we will show that this case cannot happen. First we note that M is equal to the term $\mathbf{x}^{D(v_1, v_i)} \mathbf{x}^{\theta(\mathcal{W}', \mathcal{X})} [\psi(\mathcal{X})]$ for some $\mathcal{W}' \in \mathfrak{B}(\mathcal{U})$ since M is a term corresponding to a summand of an element which was added in the previous steps of the division algorithm. This term is obtained by merging the parts $\{v_1\}$ and $\{v_i\}$, i.e., removing the orientations on the edges between v_1 and v_i . Thus $\{v_1, v_i\}$ is the unique part of \mathcal{W}' which contains two vertices. This implies that $\mathbf{x}^{\theta(\mathcal{U}, \mathcal{W}')} [\psi(\mathcal{W}')]$ has not been added in the previous steps of the division algorithm. Otherwise this term has been canceled with its corresponding dual term coming from Proposition 6.16. On the other hand $M' = \mathbf{x}^{\theta(\mathcal{U}, \mathcal{W}')} \text{LM}([\psi(\mathcal{W}')]])$ is not among the previous terms added in the division algorithm till this step (since otherwise M' could be the leading term). Note that $W'_1 = \{v_1, v_i\}$. Now assume that $W'_2 \setminus W'_1 = \{v_s\}$. If v_1 and v_s are adjacent then the dual element of M' coming from Proposition 6.16, is obtained by merging the parts $\{v_1\}$ and $\{v_s\}$ (in order to get $\mathcal{W}'' \in \mathfrak{A}(\mathcal{U})$) and then merging the parts $\{v_i\}$ and $\{v_1, v_s\}$ in \mathcal{W}'' . Note that by Lemma 6.13 we know that $\{v_1\}$ and $\{v_s\}$ are mergeable in $G(\mathcal{U}^{(1)})$ or $G(\mathcal{U}^{(2)})$ which implies that $\mathbf{x}^{\theta(\mathcal{U}, \mathcal{W}'')} [\psi(\mathcal{W}'')]$ has been already added in the previous steps of the division algorithm. However the term M' of this element has not been canceled with its dual term which is a summand of $\mathbf{x}^{\theta(\mathcal{U}, \mathcal{W}')} [\psi(\mathcal{W}')]$ which is a contradiction since $M' >_{\text{revlex}} M$.

Case 3. $\theta(\mathcal{W}, \mathcal{X}) = D(\{v_i, v_j\}, v_r)$: we have $E' = D(v_i, v_j) + D(v_i, v_r) + D(v_j, v_r)$. This case reduces to (2) by changing the indices.

As we see in all cases we add the term $\mathbf{x}^{\theta(\mathcal{U}, \mathcal{V})} \psi(\mathcal{V})$ to the s -polynomial which has the desired properties.

By Proposition 6.16 corresponding to each summand $\mathbf{x}^{\theta(\mathcal{U}, \mathcal{W})} \mathbf{x}^{\theta(\mathcal{W}, \mathcal{X})} [\psi(\mathcal{X})]$ where $\mathcal{W} \in \mathfrak{B}(\mathcal{U})$ and $\mathcal{X} \in \mathfrak{B}(\mathcal{W})$ there exists a unique $\mathcal{W}' \in \mathfrak{B}(\mathcal{U})$ such that $\mathcal{X} \in \mathfrak{B}(\mathcal{W}')$ and so there exists a unique element $\mathbf{x}^{\theta(\mathcal{U}, \mathcal{W}')} \mathbf{x}^{\theta(\mathcal{W}', \mathcal{X})} [\psi(\mathcal{X})]$ with different sign. On the other hand, by Lemma 6.14 all terms coming from elements of $\mathfrak{I}(\mathcal{U})$ are appearing in $s(\psi(\mathcal{U}^{(1)}), \psi(\mathcal{U}^{(2)}))$. By Lemma 6.13(b) corresponding to each element $\text{Merge}(o_j(\mathcal{U}); \{v_i\}, \{v_j\})$ of $\mathfrak{B}(\mathcal{U})$ there exists a $\mathcal{W} \in \mathfrak{I}(\mathcal{U})$ with a mergeable edge corresponding to $E(v_i, v_j)$. Our previous argument implies that the term associated to \mathcal{W} has been already added in the division algorithm. Therefore all terms corresponding to elements of $\mathfrak{B}(\mathcal{U})$ are added in the division algorithm as well, which completes the proof. \square

Corollary 6.19. *The Betti numbers of the ideals I_G and $\text{in}(I_G)$ are independent of the characteristic of the base field K .*

Remark 6.20. Note that in the proof of Theorem 5.3 we actually construct a free resolution for I_G according to Algorithm 1. Once we know that this resolution is minimal it follows that $\mathbf{G}_k(G, q) := \text{Image}(\psi_k)$ indeed forms a *minimal generating set* of $\text{syz}_k(\mathbf{G}(G, q))$ by Remark 3.9. Then Theorem 3.11 implies that the same statement is true for $\text{in}(I_G)$ as well.

Remark 6.21. As described in §1, the constructed minimal free resolutions are in fact supported on a cellular complex. In [23] we describe this geometric picture in detail.

The following example sums up all our notions by giving the explicit minimal free resolution for our running example, the 4-cycle graph.

Example 6.22. Returning to Example 6.8, (and Examples 6.11, 6.12) by Theorems 5.3 and 6.20 we have that

$$\begin{aligned} \varphi_2([\psi(\mathcal{U})]) &= x_2[\psi(\text{Merge}(\mathcal{U}; 1, 2))] - x_3[\psi(\text{Merge}(\mathcal{U}; 1, 3))] + x_4[\psi(\text{Merge}(\mathcal{U}; 3, 4))] \\ &\quad - x_4[\psi(\text{Merge}(\mathcal{U}; 2, 4))] + x_1[\psi(\text{Merge}(\mathfrak{c}(\mathcal{U}); 2, 1))] - x_1[\psi(\text{Merge}(\mathfrak{c}(\mathcal{U}); 3, 1))] \\ &\quad - x_2[\psi(\text{Merge}(\mathfrak{c}(\mathcal{U}); 4, 2))] + x_3[\psi(\text{Merge}(\mathfrak{c}(\mathcal{U}); 4, 3))] . \end{aligned}$$

Moreover for the ideal I_{C_4} the minimal free resolution R/I_{C_4} is

$$0 \rightarrow R(-4)^3 \xrightarrow{\varphi_2} R(-3)^8 \xrightarrow{\varphi_1} R(-2)^6 \xrightarrow{\varphi_0} R .$$

The matrix for the first differential map is

$$\varphi_0 : \left(\begin{array}{cccccccc} x_3x_4 - x_1x_2 & x_2x_4 - x_1x_3 & x_2x_3 - x_1^2 & x_4^2 - x_2x_3 & x_3^2 - x_1x_4 & x_2^2 - x_1x_4 & & \\ & & & & & & & \end{array} \right)$$

in which the columns correspond to the generators of I_{C_4} listed in the same order $x_3x_4 - x_1x_2, x_2x_4 - x_1x_3, \dots, x_2^2 - x_1x_4$.

The second differential map is presented by the matrix

$$\varphi_1 : \left(\begin{array}{cccccccc} -x_4 & 0 & x_2 & 0 & -x_3 & 0 & 0 & -x_1 \\ 0 & -x_4 & 0 & x_3 & 0 & -x_2 & -x_1 & 0 \\ 0 & 0 & -x_4 & -x_4 & 0 & 0 & -x_3 & -x_2 \\ x_3 & x_2 & 0 & 0 & -x_1 & -x_1 & 0 & 0 \\ -x_2 & 0 & 0 & -x_1 & x_4 & 0 & x_2 & 0 \\ 0 & -x_3 & -x_1 & 0 & 0 & x_4 & 0 & x_3 \end{array} \right)$$

where the columns of the above matrix correspond to the bases elements associated to connected flags

$$\text{Merge}(\mathcal{U}; 1, 2), \text{Merge}(\mathcal{U}; 1, 3), \text{Merge}(\mathcal{U}; 3, 4), \text{Merge}(\mathcal{U}; 2, 4)$$

listed in Example 6.11 and the connected flags

$$\text{Merge}(\mathfrak{c}(\mathcal{U}); 2, 1), \text{Merge}(\mathfrak{c}(\mathcal{U}); 3, 1), \text{Merge}(\mathfrak{c}(\mathcal{U}); 4, 2), \text{Merge}(\mathfrak{c}(\mathcal{U}); 4, 3)$$

listed in Example 6.12.

The last differential map is presented by the matrix

$$\varphi_2 : \left(\begin{array}{ccc} x_2 & 0 & -x_1 \\ -x_3 & -x_3 & 0 \\ x_4 & 0 & 0 \\ -x_4 & 0 & 0 \\ x_1 & x_2 & 0 \\ -x_1 & 0 & x_3 \\ -x_2 & -x_4 & x_2 \\ x_3 & x_3 & -x_4 \end{array} \right)$$

in which the first column corresponds to

$$\mathcal{U} : \{1\} \subset \{1, 2\} \subset \{1, 2, 3\} \subset \{1, 2, 3, 4\},$$

the second and third columns correspond to

$$\mathcal{U}_2 : \{1\} \subset \{1, 2\} \subset \{1, 2, 4\} \subset \{1, 2, 3, 4\} \text{ and } \mathcal{U}_3 : \{1\} \subset \{1, 3\} \subset \{1, 3, 4\} \subset \{1, 2, 3, 4\}.$$

Their corresponding acyclic orientations are listed in Figure 3.

7. BETTI NUMBERS

Let $\mathbf{A} = \mathbb{Z}$ or $\mathbf{A} = \text{Pic}(G)$. We let β_i and $\beta_{i,j}$ denote $\beta_i(R/I_G)$ and $\beta_{i,j}(R/I_G)$ respectively (for $i \geq 0$ and $j \in \mathbf{A}$). Note that by Remark 6.20 one might replace I_G with $\text{in}(I_G)$.

Proposition 7.1. *For all $i \geq 0$, $\beta_i = |\mathfrak{S}_{i+1}(G, q)| = |\mathfrak{E}_{i+1}(G, q)|$.*

Proof. The assertion follows by Theorem 5.3, Remark 5.4, and Theorem 6.17 and the fact that $\beta_i(R/\text{in}(I_G)) = \beta_{i-1}(\text{in}(I_G))$. \square

Remark 7.2. It follows from Proposition 7.1 that $|\mathfrak{S}_{i+1}(G, q)|$ is independent of q . It is a nice combinatorial exercise to show this directly.

Recall from §2.2.1 that for $D \in \text{Div}(G)$ we define

$$\deg_{\mathbf{A}}(D) = \deg_{\mathbf{A}}(\mathbf{x}^D) = \begin{cases} \deg(D), & \text{if } \mathbf{A} = \mathbb{Z}; \\ D, & \text{if } \mathbf{A} = \text{Div}(G); \\ [D], & \text{if } \mathbf{A} = \text{Pic}(G). \end{cases}$$

Definition 7.3. For $k \geq 1$ and $j \in \mathbf{A}$ define

$$\mathfrak{S}_{k,j}(G, q) = \{\mathcal{U} \in \mathfrak{S}_k(G, q) : \deg_{\mathbf{A}}(D(\mathcal{U})) = j\}$$

where $D(\mathcal{U})$ is defined in Definition 4.3.

We now strengthen Proposition 7.1 as follows.

Proposition 7.4. *For $\mathbf{A} = \mathbb{Z}$ or $\mathbf{A} = \text{Pic}(G)$*

$$\beta_{i,j} = |\mathfrak{S}_{i+1,j}(G, q)|$$

for all $i \geq 0$ and $j \in \mathbf{A}$.

Proof. By Theorem 5.3, Theorem 6.17, and Remark 3.9 the set $\psi_i(\mathfrak{S}_{i+2}(G, q))$ minimally generates the module $\text{syz}_i(\text{in}(I_G))$ for each $i \geq 0$, and we have

$$\beta_{i,j} = \beta_{i-1,j}(I_G) = |\{\psi_{i-1}(\mathcal{U}) : \deg(\psi_{i-1}(\mathcal{U})) = j \text{ for } \mathcal{U} \in \mathfrak{S}_{i+1,j}(G, q)\}|.$$

We first note that for $\mathcal{U} \in \mathfrak{S}_{i+1,j}(G, q)$ we have

$$D(\mathcal{U}) = \sum_{\ell=1}^{i+1} D(U_{\ell} \setminus U_{\ell-1}, U_{\ell-1}) \quad , \quad \mathbf{x}^{D(\mathcal{U})} = \prod_{\ell=1}^{i+1} \mathbf{x}^{D(U_{\ell} \setminus U_{\ell-1}, U_{\ell-1})}$$

and

$$(7.1) \quad \deg_{\mathbf{A}}(D(\mathcal{U})) = \deg_{\mathbf{A}}(\mathbf{x}^{D(\mathcal{U})}) = \sum_{i=1}^k \deg_{\mathbf{A}}(\mathbf{x}^{D(U_i \setminus U_{i-1}, U_{i-1})}).$$

We need to show that $\deg(\psi_{i-1}(\mathcal{U})) = \deg_{\mathbf{A}}(D(\mathcal{U}))$. The proof is by induction on $i \geq 0$. For $i = 0$ there is nothing to prove. Since $\psi_{i-1}(\mathcal{U})$ is homogeneous, by (5.2) and (7.1) we obtain

$$\begin{aligned} \deg_{\mathbf{A}}(\psi_{i-1}(\mathcal{U})) &= \deg_{\mathbf{A}}(\text{LM}(\psi_{i-1}(\mathcal{U}))) \\ &= \deg(\mathbf{x}^{D(U_2 \setminus U_1, U_1)}[\psi_{k-1}(\mathcal{U}^{(1)})]) \\ &= \deg_{\mathbf{A}}(\mathbf{x}^{D(U_2 \setminus U_1, U_1)}) + \deg_{\mathbf{A}}(\psi_{k-1}(\mathcal{U}^{(1)})) \\ &= \deg_{\mathbf{A}}(\mathbf{x}^{D(U_2 \setminus U_1, U_1)}) + \deg_{\mathbf{A}}(D(\mathcal{U}^{(1)})) \\ &= \deg_{\mathbf{A}}(D(\mathcal{U})). \end{aligned}$$

\square

Example 7.5. It follows from above descriptions that for the \mathbb{Z} -grading of R , $\beta_{i,j}$ can take nonzero values only if $0 \leq i \leq n-1$ and $0 \leq j \leq m$ where $n = |V(G)|$ and $m = |E(G)|$. Clearly $\beta_0 = 1$. Moreover $\beta_{n-1} = \beta_{n-1,m}$ which is equal to the number of acyclic orientations of G with unique source at q (see also Lemma 7.12). Since both R/I_G and $R/\text{in}(I_G)$ are Cohen-Macaulay (see, e.g., [23, Section 11.1]), it follows that the *Castelnuovo-Mumford regularity* of both R/I_G and $R/\text{in}(I_G)$ is equal to $g = m - n + 1$ (see, e.g., [11, page 69]).

Example 7.6. Let $G = K_n$ be the complete graph on n vertices. Let $\{A_1, A_2, \dots, A_k\}$ be any k -partition of $V(G)$ with $q \in A_1$. Then corresponding to each permutation $\delta = (i_1, i_2, \dots, i_{k-1})$ of $(2, 3, \dots, n)$ the strictly increasing k -flag

$$\mathcal{U}_\delta : U_1 \subsetneq U_2 \subsetneq \dots \subsetneq U_k = V(G)$$

is an element of $\mathfrak{S}_k(G, q)$ where $U_j = A_1 \cup A_{i_1} \cup \dots \cup A_{i_{j-1}}$ for each j . Therefore $|\mathfrak{S}_k(G, q)| = (k-1)! S(n, k)$ where $S(n, k)$ denotes the Stirling number of the second kind (i.e. the number of ways to partition a set of n elements into k nonempty subsets). In other words $\beta_i = i! S(n, i+1)$. See <http://oeis.org/A028246> for other interpretations of these numbers.

Example 7.7. Let G be a tree on n vertices. Let $\mathcal{U} \in \mathfrak{S}_k(G, q)$. For each $i > 1$ the part $U_i \setminus U_{i-1}$ is connected by exactly one edge to only one part $U_j \setminus U_{j-1}$ with $j < i$; otherwise we get a cycle in the graph. Therefore each element $\mathcal{U} \in \mathfrak{S}_k(G, q)$ is determined by the $k-1$ edges (of $n-1$ edges of G) between the partitions $(U_i \setminus U_{i-1})$'s of \mathcal{U} and $|\mathfrak{S}_k(G, q)| = \binom{n-1}{k-1}$. The fact that each edge contributes 1 to the degree of $\psi(\mathcal{U})$ means that $\beta_i = \beta_{i,i} = \binom{n-1}{i}$.

Example 7.8. Let $G = C_n$ be the cycle on n vertices. For simplicity of notation let $V(G) = [n]$. Then we will show, by induction on n , that for $k \geq 2$

$$|\mathfrak{S}_k(C_n, q)| = (k-1) \times \binom{n}{k}.$$

One can easily check the formula for $k=2$ and $k=3$. So we may assume that $k \geq 4$. Let $1 \leq i_1 < i_2 < \dots < i_k \leq n$. Then we consider the parts

$$A_1 := \{i_k, i_k+1, \dots, n\} \cup \{1, \dots, i_1-1\} \text{ and } A_{t+1} := \{i_t, i_t+1, \dots, i_{t+1}-1\} \text{ for } 1 \leq t \leq k-1$$

of the graph. Then there are three types of elements \mathcal{U} of $\mathfrak{S}_k(G, q)$ such that $(U_i \setminus U_{i-1})_{i=2}^k$ is a permutation of A_2, \dots, A_k :

- (1) $U_1 = A_1$ and $U_i = U_{i-1} \cup A_i$ for each $1 < i \leq k$.
- (2) $U_1 = A_1$ and $U_i = U_{i-1} \cup A_{k-i+2}$ for each $1 < i \leq k$.
- (3) $U_1 = A_1, U_2 \setminus U_1 = A_2, U_3 \setminus U_2 = A_k$: Then the number of k -connected flags of C_n with this partition set is equal to the number of $(k-3)$ -connected flags of C_{k-2} on the vertex set u_1, u_2, \dots, u_{k-2} where $q = u_{k-3}$ is associated to $A_1 \cup A_2 \cup A_k$ and u_i is associated to the part A_i for each $3 \leq i \leq k-1$. This number equals to $(k-3) \times \binom{k-3}{k-3}$ by the induction hypothesis.
- (4) $U_1 = A_1, U_2 \setminus U_1 = A_k, U_3 \setminus U_2 = A_2$: Similar to the previous case the number of k -connected flags of C_n with this partition set is equal to $(k-3) \times \binom{k-3}{k-3}$.

Now note that since A_2 and A_k are not adjacent just one of the two elements

$$\mathcal{U} : A_1 \subsetneq A_1 \cup A_2 \subsetneq A_1 \cup A_2 \cup A_k \subsetneq U_4 \subsetneq \dots \subsetneq U_k$$

and

$$\mathcal{U}' : A_1 \subsetneq A_1 \cup A_k \subsetneq A_1 \cup A_2 \cup A_k \subsetneq U_4 \subsetneq \dots \subsetneq U_k$$

will be in $\mathfrak{S}_k(G, q)$. This implies that $|\mathfrak{S}_k(G, q)| = (1 + 1 + (k - 3)) \times \binom{n}{k} = (k - 1) \times \binom{n}{k}$. We get $\beta_i = \beta_{i,i+1} = i \binom{n}{i+1}$ for $i \geq 1$.

For example for $G = C_5$ we have

$$\beta_0 = 1, \beta_1 = 10, \beta_2 = 20, \beta_3 = 15, \beta_4 = 4 .$$

Example 7.9. It follows from Proposition 7.4 that adding or removing parallel edges will not change β_i , since this process does not add/remove any element to/from the set $\mathfrak{S}_{i+1}(G, q)$. However, the graded Betti numbers $\beta_{i,j}$ do change by adding or removing parallel edges. For example, consider the theta graph G with two vertices u and v connected by m edges. Then $\mathfrak{S}_2(G, u)$ has the unique element $\{v\} \subsetneq \{u, v\}$ which implies that $\beta_1 = \beta_{1,m} = 1$.

7.1. Relation to maximal reduced divisors. Recall the definition of reduced divisors.

Definition 7.10. Let (Γ, v_0) be a pointed graph. A divisor $D \in \text{Div}(\Gamma)$ is called v_0 -reduced if it satisfies the following two conditions:

- (i) $D(v) \geq 0$ for all $v \in V(\Gamma) \setminus \{v_0\}$.
- (ii) For every nonempty subset $A \subseteq V(\Gamma) \setminus \{v_0\}$, there exists a vertex $v \in A$ such that $D(v) < \text{outdeg}_A(v)$.

These divisors arise precisely from the normal forms with respect to the Gröbner bases given in Theorem 5.1. There is a well-known algorithm due to Dhar for checking whether a given divisor is reduced (see, e.g., [2] and references therein).

Recall from Definition 6.1 that given $\mathcal{U} \in \mathfrak{S}_k(G, q)$ we obtain a graph G/\mathcal{U} from G by contracting all the unoriented edges of $G(\mathcal{U})$. The contraction map $\phi : G \rightarrow G/\mathcal{U}$ induces the map

$$\phi_* : \text{Div}(G) \rightarrow \text{Div}(G/\mathcal{U}) \quad \text{with} \quad \phi_*\left(\sum_{v \in V(G)} a_v(v)\right) = \sum_{v \in V(G)} a_v(\phi(v)).$$

Assume U_1 is the part of \mathcal{U} containing q and let $q' = \phi(U_1) \in V(G/\mathcal{U})$.

Lemma 7.11. $\phi_*(D(\mathcal{U})) = E + \mathbf{1}$, where E is a maximal q' -reduced divisor and $\mathbf{1}$ is the all-one divisor.

Proof. This follows from the well-known fact that Dhar's algorithm gives a one-to-one correspondence between acyclic orientations with unique source at v_0 and maximal v_0 -reduced divisors; given such an acyclic orientation the corresponding v_0 -reduced divisor is $\sum_{v \in V(\Gamma)} (\text{indeg}(v) - 1)(v)$ (see, e.g., [4]). The result now follows from Remark 4.5. \square

Since different acyclic orientations with unique source at q' give rise to inequivalent q' -reduced divisors we deduce that if $\mathcal{U}, \mathcal{V} \in \mathfrak{S}_k(G, q)$ and the graphs G/\mathcal{U} and G/\mathcal{V} coincide, then $\phi_*(D(\mathcal{U})) - \mathbf{1}$ and $\phi_*(D(\mathcal{V})) - \mathbf{1}$ are two inequivalent maximal reduced divisors. These observations lead to the following formula for Betti numbers which, in an equivalent form, was conjectured in [26] for I_G :

$$\begin{aligned} \beta_i &= \sum_{G/\mathcal{U}} |\{D : D \text{ is a maximal } v_0\text{-reduced divisor on } G/\mathcal{U}\}| \\ &= \sum_{G/\mathcal{U}} |\{\text{acyclic orientations of } G/\mathcal{U} \text{ with unique source at } v_0\}| \end{aligned}$$

where the sum is over all *distinct* contracted graphs G/\mathcal{U} as \mathcal{U} varies in $\mathfrak{S}_{i+1}(G, q)$, and v_0 is an arbitrary vertex of G/\mathcal{U} .

Here is another connection with reduced divisors. Hochster’s formula for computing the Betti numbers topologically (see, e.g., [21, Theorem 9.2]), when applied to I_G and the “nice” grading by $\text{Pic}(G)$, says that for each $\mathbf{j} \in \text{Pic}(G)$ the graded Betti number $\beta_{i,\mathbf{j}}(R/I_G)$ is the dimension of the i^{th} reduced homology of the simplicial complex

$$\Delta_{\mathbf{j}} = \{\text{supp}(E) : 0 \leq E \leq D' \in |\mathbf{j}|\}$$

where $|\mathbf{j}|$ denotes the linear system of $\mathbf{j} \in \text{Pic}(G)$. One can use this to give an alternate proof for the highest graded Betti numbers. The following is a simplification of the proof of [26, Theorem 7.7] (see also Example 7.5).

Lemma 7.12. *For $\mathbf{j} \in \text{Pic}(G)$, we have $\beta_{n-1,\mathbf{j}}(R/I_G) = 1$ if and only if*

$$\mathbf{j} \sim E + \mathbf{1}$$

where E is a maximal q -reduced divisor.

Proof. $\beta_{n-1,\mathbf{j}}(R/I_G) = 1$ if and only if $\Delta_{\mathbf{j}}$ is homotopy equivalent to an $(n - 1)$ -sphere. This is equivalent to the following two conditions.

- (1) $|\mathbf{j} - \mathbf{1}| = \emptyset$,
- (2) $|\mathbf{j} - \mathbf{1} + (v)| \neq \emptyset$ for any $v \in V(G)$.

Let E be the unique q -reduced divisor equivalent to $\mathbf{j} - \mathbf{1}$. Then (1) is equivalent to saying $E(q) \leq -1$. But (2) for $v = q$ would require $E(q) = -1$, and for $v \neq q$ would require that E be a maximal q -reduced divisor. This is because for all maximal reduced divisors the values of vertices $v \neq q$ add up to the same number $g = |E(G)| - |V(G)| + 1$. \square

Remark 7.13.

- (i) By Remark 6.20 one can use Proposition 7.4 to read all dimensions of the reduced homologies of $\Delta_{\mathbf{j}}$. Although we now know all the Betti numbers, giving an explicit bijection between connected flags and the bases of the reduced homologies of $\Delta_{\mathbf{j}}$ is an intriguing problem.
- (ii) In a recent work, Mania [17] studies the number of connected components of $\Delta_{\mathbf{j}}$. This gives an alternate proof that $\beta_1 = |\mathfrak{S}_2(G, q)|$.

REFERENCES

- [1] Matthew Baker and Serguei Norine. Riemann-Roch and Abel-Jacobi theory on a finite graph. *Adv. Math.*, 215(2):766–788, 2007.
- [2] Matthew Baker and Farbod Shokrieh. Chip-firing games, potential theory on graphs, and spanning trees. *J. Combin. Theory Ser. A*, 120(1):164–182, 2013.
- [3] Dave Bayer, Sorin Popescu, and Bernd Sturmfels. Syzygies of unimodular Lawrence ideals. *J. Reine Angew. Math.*, 534:169–186, 2001.
- [4] Brian Benson, Deeparnab Chakrabarty, and Prasad Tetali. G -parking functions, acyclic orientations and spanning trees. *Discrete Math.*, 310(8):1340–1353, 2010.
- [5] Norman Biggs. Algebraic potential theory on graphs. *Bull. London Math. Soc.*, 29(6):641–682, 1997.
- [6] Aldo Conca, Serkan Hoşten, and Rekha R. Thomas. Nice initial complexes of some classical ideals. In *Algebraic and geometric combinatorics*, volume 423 of *Contemp. Math.*, pages 11–42. Amer. Math. Soc., Providence, RI, 2006.
- [7] Robert Cori, Dominique Rossin, and Bruno Salvy. Polynomial ideals for sandpiles and their Gröbner bases. *Theoret. Comput. Sci.*, 276(1-2):1–15, 2002.
- [8] Deepak Dhar. Self-organized critical state of sandpile automaton models. *Phys. Rev. Lett.*, 64(14):1613–1616, 1990.

- [9] Anton Dochtermann and Raman Sanyal. Laplacian ideals, arrangements, and resolutions. Preprint available at [arXiv:1212.6244](https://arxiv.org/abs/1212.6244), 2012.
- [10] David Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. With a view toward algebraic geometry.
- [11] David Eisenbud. *The geometry of syzygies*, volume 229 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2005. A second course in commutative algebra and algebraic geometry.
- [12] Andrei Gabrielov. Abelian avalanches and Tutte polynomials. *Phys. A*, 195(1-2):253–274, 1993.
- [13] Andreas Gathmann and Michael Kerber. A Riemann-Roch theorem in tropical geometry. *Math. Z.*, 259(1):217–230, 2008.
- [14] Gert-Martin Greuel and Gerhard Pfister. *A Singular introduction to commutative algebra*. Springer, Berlin, extended edition, 2008. With contributions by Olaf Bachmann, Christoph Lossen and Hans Schönemann.
- [15] Yuji Kamoi. Noetherian rings graded by an abelian group. *Tokyo J. Math.*, 18(1):31–48, 1995.
- [16] Dino J. Lorenzini. Arithmetical graphs. *Math. Ann.*, 285(3):481–501, 1989.
- [17] Horia Mania. Wilmes’ conjecture and boundary divisors. Preprint available at [arXiv:1210.8109](https://arxiv.org/abs/1210.8109), 2012.
- [18] Madhusudan Manjunath, Frank-Olaf Schreyer, and John Wilmes. Minimal free resolutions of the G -parking function ideal and the toppling ideal. To appear in *Trans. Amer. Math. Soc.*, Preprint available at [arXiv:1210.7569](https://arxiv.org/abs/1210.7569), 2012.
- [19] Madhusudan Manjunath and Bernd Sturmfels. Monomials, binomials and Riemann–Roch. *J. Algebraic Combin.*, 37(4):737–756, 2013.
- [20] Grigory Mikhalkin and Ilia Zharkov. Tropical curves, their Jacobians and theta functions. In *Curves and abelian varieties*, volume 465 of *Contemp. Math.*, pages 203–230. Amer. Math. Soc., Providence, RI, 2008.
- [21] Ezra Miller and Bernd Sturmfels. *Combinatorial commutative algebra*, volume 227 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2005.
- [22] Fatemeh Mohammadi. Prime splittings of determinantal ideals. Preprint available at [arXiv:1208.2930](https://arxiv.org/abs/1208.2930), 2012.
- [23] Fatemeh Mohammadi and Farbod Shokrieh. Divisors on graphs, binomial and monomial ideals, and cellular resolutions. Preprint available at [arXiv:1306.5351](https://arxiv.org/abs/1306.5351), 2013.
- [24] Isabella Novik, Alexander Postnikov, and Bernd Sturmfels. Syzygies of oriented matroids. *Duke Math. J.*, 111(2):287–317, 2002.
- [25] Irena Peeva. *Graded syzygies*, volume 14 of *Algebra and Applications*. Springer-Verlag London Ltd., London, 2011.
- [26] David Perkinson, Jacob Perlman, and John Wilmes. Primer for the algebraic geometry of sandpiles. Preprint available at [arXiv:1112.6163](https://arxiv.org/abs/1112.6163), 2011.
- [27] Alexander Postnikov and Boris Shapiro. Trees, parking functions, syzygies, and deformations of monomial ideals. *Trans. Amer. Math. Soc.*, 356(8):3109–3142, 2004.
- [28] Michel Raynaud. Spécialisation du foncteur de Picard. *Inst. Hautes Études Sci. Publ. Math.*, (38):27–76, 1970.
- [29] Frank Schreyer. Die Berechnung von Syzygien mit dem verallgemeinerten Weierstrass’schen Divisionssatz. Diplom Thesis, University of Hamburg, Germany., 1980.
- [30] D. A. Spear. A constructive approach to commutative ring theory. In *Proceedings of the the 1977 MACSYMA Users’ Conference, NASA CP-2012*, pages 369–376, 1977.
- [31] John Wilmes. Algebraic invariants of sandpile graphs. Bachelors Thesis, Reed College, Portland, OR, 2010.

E-mail address: fatemeh.mohammadi716@gmail.com

FACHBEREICH MATHEMATIK UND INFORMATIK, PHILIPPS-UNIVERSITÄT, 35032 MARBURG, GERMANY

E-mail address: shokrieh@math.gatech.edu

GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GEORGIA 30332-0160, USA