



Communication

Where Have All the Spiders Gone? Observations of a Dramatic Population Density Decline in the Once Very Abundant Garden Spider, *Araneus diadematus* (Araneae: Araneidae), in the Swiss Midland

Martin Nyffeler ^{1,*} and Dries Bonte ²

¹ Department of Environmental Sciences, Section of Conservation Biology, University of Basel, CH-4056 Basel, Switzerland

² Department of Biology, Terrestrial Ecology Unit, Ghent University, 9000 Ghent, Belgium; Dries.Bonte@UGent.be

* Correspondence: martin.nyffeler@unibas.ch

Received: 6 March 2020; Accepted: 10 April 2020; Published: 15 April 2020



Abstract: Aerial web-spinning spiders (including large orb-weavers), as a group, depend almost entirely on flying insects as a food source. The recent widespread loss of flying insects across large parts of western Europe, in terms of both diversity and biomass, can therefore be anticipated to have a drastic negative impact on the survival and abundance of this type of spider. To test the putative importance of such a hitherto neglected trophic cascade, a survey of population densities of the European garden spider *Araneus diadematus*—a large orb-weaving species—was conducted in the late summer of 2019 at twenty sites in the Swiss midland. The data from this survey were compared with published population densities for this species from the previous century. The study verified the above-mentioned hypothesis that this spider’s present-day overall mean population density has declined alarmingly to densities much lower than can be expected from normal population fluctuations (0.7% of the historical values). Review of other available records suggested that this pattern is widespread and not restricted to this region. In conclusion, the decline of this once so abundant spider in the Swiss midland is evidently revealing a bottom-up trophic cascade in response to the widespread loss of flying insect prey in recent decades.

Keywords: bottom-up trophic cascade; low abundance; orb-weaving spiders; prey scarcity; western European landscape

1. Introduction

When the Krefeld Entomological Society, together with colleagues, published their famous long-term study (the “Krefeld study”) in 2017, it became known that the biomass of flying insects had declined by approx. 75% over the past three decades in over 60 nature protection areas of Germany [1]. Another long-term study (the “Munich study”) confirmed the Krefeld results, providing evidence that strong abundance declines of insect populations had occurred in farmland and forests across vast areas of Germany and Switzerland [2]. Similar abundance declines of insect populations have been documented in other long-term studies of other European and North American regions [3–9]. It is generally accepted that we now live in an era of global insect meltdown [9–13]. This has dramatic ecological implications: the fact that insects comprise the basis of many food chains and food webs [14,15] means that in a world without insects, countless insectivorous species will ultimately become extinct due to starvation [16].

In this context, aerial web-spinning spiders are a group of utmost interest. These spiders (including the orb-weaving species in the families Araneidae) trap flying insects with the aid of aerial webs [17–19]. In the temperate climatic zone, aerial web-spinning spiders feed for the most part on small dipterans, aphids, and hymenopterans (Figure 1), which are exactly the types of insects known to have most dramatically decreased in abundance and biomass in recent decades [1]. Aerial web-spinning spiders, because of their reliance upon flying prey, are therefore a highly vulnerable predator group in regions characterized by a high loss of flying insects, such as Germany and Switzerland.

While a larger number of studies on population declines of various groups of insects has been conducted in recent years (see references in [12]), potential cascading effects have only been documented so far for insectivorous birds [6]. No studies of this type have, to date, considered potential declines due to trophic cascades among arthropods, like aerial web-spinning spiders. Here we used the European garden spider *Araneus diadematus* as a model system to address whether evidence for population declines in aerial web-spinning spiders in the Swiss midland can be found (Figure 2). The female, 10–18 mm in length, reaches adulthood in late summer, at which time it spins (~30 cm diameter) orbs [20–22]. The webs are built 0.5–2 m off the ground [21]. This animal, with its conspicuous white, cross-shaped mark on the upper side of the abdomen, is one of the best known spider species in western Europe.

By comparing historical abundance data (20th century) with present-day data (2019), we examined whether the abundance of *Araneus diadematus* has changed over the past few decades. This work is based on extensive experience gathered during graduate research in the Swiss midland in the 20th century [20,21] and supplemented by a present-day population density survey in the same geographical area. To make the comparison more robust, the data base was expanded by including published population density values extracted from the scientific literature.




Figure 1. Prey composition (% by number) of the garden orb-weaver *Araneus diadematus* based on four field studies: (A) oat field near Zurich, Switzerland [21]; (B) fallow grassland near Zurich, Switzerland [21]; (C) fallow grassland near Jena, Germany [18]; (D) maize field margin near Munich [19]. Homoptera refers primarily to alate aphids.

2. Materials and Methods

2.1. Assessment of Population Densities of *Araneus diadematus* in 2019

Population density assessments were conducted at 20 locations in the Swiss midland, and the visual censusing technique described by Lubin (1978) [23] was applied at 18 locations. This technique was modified insofar as curved line transects were used in addition to straight line transects in some cases, depending on the paths used [24]. One of us (M.N.) slowly walked along the edge of a path and intensively searched by eye the vegetation within 1 m to one side of the path and from ground level to a height of 2 m (based on the fact that *Araneus diadematus* habitually constructs its webs at heights of 0.5–2 m off the ground [21]). Each adult *Araneus diadematus* web encountered within the transect was recorded. Transects 1 to 5 were 1000 m long, Transects 6 to 10 were 500 m long, and Transects 11 to 18 were 200 m long (see Table S1; transect lengths depended on the nature of the landscape). The search focused on webs of adult females in habitats typical for this species (i.e., gardens, parks, graveyards, hedgerows, forest edges, and forest trails; [25]). Only webs occupied by a spider were counted (regardless whether the spider was present in the web or hiding in a nearby retreat), since only those webs could be counted beyond any doubt as belonging to *Araneus diadematus*. All counts were conducted on rain-free afternoons in August and September 2019. It has been suggested in the literature that spider counts obtained using this method may result in underestimating the true population densities because of the spiders' cryptic lifestyles [26]. This might well be the case if densities of well-camouflaged, nocturnal, and/or small, immature spiders are included in the assessment [26]. However, in the case of *Araneus diadematus*, a bias of this type is highly unlikely because this is a species of which the adult females construct large, conspicuous webs which are unlikely to be overlooked by a keen observer. Lubin (1978) also stated that "... Visual censusing is an effective technique when applied to animals with conspicuous artifacts, such as most web spiders" [23]. The immature stages of *Araneus diadematus* are more likely to be overlooked with this method because their smaller-sized webs are less conspicuous; however, in the current study this objection was not relevant because immature spiders were not counted.

Additionally, web counts in two organic gardens, covering 330 m² and 800 m² ground area, respectively, were conducted. In these gardens, the vegetation and the exterior of the buildings were thoroughly searched for webs of adult female *Araneus diadematus* on several consecutive days in August and September 2019.

2.2. Literature Search for Historical Data on *Araneus diadematus* Population Densities

Araneus diadematus population density values assessed in the 1970s in the Zurich area were taken from two graduate research theses [20,21]. These were supplemented by data taken from the literature. Overall, a total of 12 reports (including 18 population density values) were gathered (see Table S1). In these studies, plots ranging from 17.5 to 3300 m² in ground area were used for the assessment of *Araneus diadematus* densities (see Table S1).

It is noticeable that the present habitat distribution was different from the habitats studied in the past, e.g., no fallow grassland was studied in 2019. The fallow grassland areas used in the Swiss studies 40 years ago [21] have fallen victim to urbanization and could not be surveyed. However, *Araneus diadematus* is a habitat generalist, widely distributed over a large number of different habitat types. All the study areas used in the present density assessment (Table 1) represent habitat types in which this species is typically found, such as forested areas [25]. The past century's spider density values for grasslands were not above that century's overall mean value (Table 1), and according to Hänggi et al. (1995) [25], tree- and shrub-dominated habitats appear to provide the most suitable sites for construction of the webs of this species. This implies that omission of grassland habitats in the 2019 density assessment should not have compromised our investigation.

2.3. Statistical Methods

The data in Table 1 were tested for normality using the Shapiro–Wilk Test Calculator (<http://www.statskingdom.com/320ShapiroWilk.html>), and this revealed that the data did not follow a normal distribution. The Mann–Whitney *U* test was applied to examine whether spider densities in 2019 ($N = 20$) differed statistically significantly from those in the 20th century ($N = 18$). The same test was used to examine whether numbers of prey per web in 2019 and in the 20th century differed statistically significantly. Medians are followed by interquartile ranges (IQR). In the past, multiple estimates of spider densities were probably also not normally distributed but were summarized as a mean. In order to make comparisons with the historical data, we also calculated a mean from our new data (while at the same time acknowledging that this is not the ideal measure).

3. Results

In Table 1, population densities of *Araneus diadematus* assessed in the Swiss midland in August/September 2019 are compared with 20th century population density data for this species from a variety of European locations. The table reveals that present-day population densities of adult female *Araneus diadematus* in the Swiss midland were generally extremely low (median = 0.0000 webs m^{-2} , IQR = 0.0000–0.0020 webs m^{-2} ; Table 1). In 2/3 of the 20 investigated transects, no webs of adult female *Araneus diadematus* could be found.

By contrast, historical population densities of *Araneus diadematus* (Figure 2A,B) in its typical habitats were considerably higher (median = 0.0670 webs m^{-2} , IQR = 0.0175–0.1835 webs m^{-2} ; Table 1). The difference between the 20th century European densities and the present-day Swiss densities (median = 0.0670 vs. 0.0000 webs m^{-2}) was statistically highly significant (Mann–Whitney *U* test, $n_1 = 18$, $n_2 = 20$, $Z = -4.0637$, $p < 0.0001$). The present-day Swiss overall mean density of *Araneus diadematus* was roughly 140 times lower than the 20th century European overall mean (Table 1). (If the seven Swiss 20th century values (median = 0.0300 webs m^{-2} ; Table 1) were exclusively compared with the twenty Swiss 21st century values (median = 0.0000 webs m^{-2} ; Table 1), the difference between the two groups was still statistically significant at $p < 0.05$.)

Furthermore, the webs contained significantly fewer prey compared to previous studies (Mann–Whitney *U* test, $n_1 = 20$, $n_2 = 18$, $Z = 4.5753$, $p < 0.0001$; Table S2). (These figures refer to mean numbers of prey per web counted in mid-afternoon as a proxy for the daily prey capture rate of large orb-weaving spiders (see Supplementary Materials; Table S2).)

Table 1. Population densities of adult female *Araneus diadematus* (18 historical vs. 20 recent values).
*Adults/subadults in September.

Habitat Type	Geographic Region	Period of Investigation	Spiders m^{-2}	Source
Historical data *				
Fallow grassland, plot 1	Switzerland	1979	0.000	[21]
Fallow grassland, plot 2	Switzerland	1979	0.030	[21]
Fallow grassland, plot 3	Switzerland	1979	0.070	[21]
Fallow grassland, plot 4	Switzerland	1979	0.000	[21]
Shrubs	Switzerland	1979	0.930	[21]
Garden	Switzerland	1974–75	0.064	[20]
Glasshouses botanical garden	Switzerland	1932	0.001	[27]
Fallow grassland	Germany	1990	0.138	[18]
Fallow grassland	Germany	1998	0.056	[28]
Fallow grassland	France	1984	0.023	[29]
Hedgerow/grassland	France	1977	0.135	[30]
Hedgerow	Italy	1990–91	0.0005	[31]
Pine stand	Poland	1979	0.076	[32]
Clearing-forest ecotone	Poland	1979	0.320	[32]
Scotch pine wood	Netherland	1946	0.130	[33]

Table 1. Cont.

Habitat Type	Geographic Region	Period of Investigation	Spiders m ⁻²	Source
Oak wood	Netherland	1946	0.400	[33]
Oak stand	UK	1954–56	0.410	[34]
Heathland	UK	1968	0.024 *	[35]
Overall mean			0.1560	
Median			0.0670	
IQR			0.0175–0.1835	
Recent data				
Forest road	Northeastern Switzerl.	2019	0.002	This paper
Suburb	Zurich region, Switzerl.	2019	0.000	This paper
Suburb	Zurich region, Switzerl.	2019	0.000	This paper
Suburb	Northwestern Switzerl.	2019	0.001	This paper
Hedgerow + shrubs along a river bank	Zurich region, Switzerl.	2019	0.002	This paper
Forest edge + suburb	Northwestern Switzerl.	2019	0.002	This paper
Forest edge	Northwestern Switzerl.	2019	0.000	This paper
Forest road	Northwestern Switzerl.	2019	0.000	This paper
Garden + shrubs	Northwestern Switzerl.	2019	0.000	This paper
Graveyard shrubs	Northwestern Switzerl.	2019	0.000	This paper
Forest road	Northwestern Switzerl.	2019	0.000	This paper
Forest edge	Northwestern Switzerl.	2019	0.000	This paper
Forest edge	Northwestern Switzerl.	2019	0.000	This paper
Graveyard shrubs	Northwestern Switzerl.	2019	0.005	This paper
Public park	Northwestern Switzerl.	2019	0.000	This paper
Hedgerow	Northwestern Switzerl.	2019	0.000	This paper
Hedgerow	Northwestern Switzerl.	2019	0.000	This paper
Hedgerow + reedbelt	Central Switzerl.	2019	0.000	This paper
Garden	Northeastern Switzerl.	2019	0.006	This paper
Garden	Northwestern Switzerl.	2019	0.003	This paper
Overall mean			0.0011	
Median			0.0000	
IQR			0.0000–0.0020	



Figure 2. (A,B) Adult female *Araneus diadematus* in web in a garden in Flawil, northeastern Switzerland (photo by Rätus Fischer).

4. Discussion

This study revealed that the large orb-weaving spider *Araneus diadematus* occurred in extraordinarily low population densities in the Swiss midland in 2019 (Table 1). *Araneus diadematus* spiders take down their webs during the night and rebuild them early in the morning. However, as laboratory experiments and field observations have revealed, not all individuals rebuild their webs the following day [30,36]. Some spiders, after consuming exceptionally large amounts of food at particularly favorable hunting sites, may cease feeding and rebuilding their webs for one or more days [36]. Thus, the percentage of spiders found with webs within an *A. diadematus* population is, in general, lower than 100% and usually in the range of 66% to 88%, depending on the availability of food at a particular time and location [30,36]. The question arises of whether the method of web counting applied in this study might have resulted in underestimation of the true population densities by overlooking a certain percentage of satiated spiders which had not rebuilt their webs at the time of the surveys. On this issue, it may be noted that densities assessed in the past century were for the most part also based on web counts [18,20,21,28–33,35], so the same methodological bias was involved at both time periods. Furthermore, since unsatiated spiders tend to rebuild their webs more frequently than fully satiated spiders [36,37], and since present-day spiders living in the era of insect loss are more likely to be unsatiated compared with half a century ago (Table S2), the frequency of web building nowadays might be expected to be rather higher. Thus, the possibility that the extraordinarily low present-day population densities were underestimations due to this particular type of methodological bias is highly unlikely.

In 2019, the webs appeared to be rather fragile compared with the stronger webs from previous decades, as is the case when malnourished spiders make webs with thinner threads (Nyffeler, pers. observations). This is in good agreement with laboratory experiments, in which the amount of thread produced is reduced if *Araneus diadematus* spiders are kept deprived of food [36] and it agrees with our observation that the webs contained significantly fewer prey compared to previous studies (Table S2). The overall impression gained during this survey was that there has been a paucity of prey throughout the entire study area in recent decades (Table S2), which has been confirmed by the observations of other researchers [2,38,39].

The abundance decline we show here of this once common spider in the Swiss midland is evidently revealing a bottom-up trophic cascade in response to prey scarcity recently documented in the same area, and more widely across western Europe [1,2,38,39]. The hypothesis that the availability of flying insect prey in the study area had drastically declined over the past decades is confirmed by the “windshield phenomenon” noticed throughout the Swiss midland (i.e., compared with previous decades, many times fewer flying insects are nowadays killed on the front windshields of cars [9,40]). This is in sharp contrast to the situation a few decades ago, when fairly frequent “wasteful killing” (and, coupled with it, “partial consumption”) of insects in *Araneus diadematus* webs at favorable sites could be observed (with capture rates of sometimes up to 1000 prey web⁻¹ day⁻¹; [41]). The reduced food intake of *Araneus diadematus* in recent years (Table S2) is likely to have negatively impacted the fecundity and survival of this spider [42,43], which in turn may have led to the abundance decline documented in this study (Table 1). Sublethal effects of chemical pollution may have additional negative impacts on this spider’s survival [44–49].

Aside from food scarcity due to insect population declines and chemical pollution, other negative environmental developments have additionally contributed to the decreasing population densities of large orb-weaving spider species in the western European landscape. These include adverse effects of modern forestry practices (e.g., removal of forest understorey and of shrubs at forest margins, large-scale aerial spraying for the purpose of caterpillar or bark beetle control), the transformation of traditional farmland into large-scale depauperate monocultures (accompanied by the loss of weedy field margins and hedges), the transformation of fallow grassland patches into construction sites, and the removal of shrubs and trees in urban and suburban gardens. That habitat conversion and degradation can have a strong detrimental effect on large orb-weaving spider abundance is well demonstrated by the

example of the fate of a plot of land located near Zurich-Oberengstringen (in between Highway A3 and the Limmat riverbank). In the 1970s, tall grassland interspersed with shrubs covered this plot, and at that time, large orb-weaving spiders occurred there in high numbers (up to 6 m^{-2} in small, local patches [50]). During a visit to this site, 40 years later, it became apparent that the land had been converted to a highway rest area with short-cut lawns and, as a result, the density of large orb-weaving spiders had declined to a very low value of 0.002 m^{-2} (Nyffeler, pers. observations). Similar habitat conversions resulting in the destruction of suitable habitats for large orb-weaving spiders have been noted elsewhere.

We of course realize that the recorded declines were not based on systematic long-term monitoring and that they could be attributed to normal population cycling. However, while we are not here able to provide a smoking gun, several elements support a systematic decline. First of all, the dramatic decline up to $< 1\%$ of the reference baseline from 40 years ago has never been observed at shorter temporal time frames, and reaches far beyond the typical natural fluctuations in population sizes (e.g., [51]), even when they are known to be linked to resource pulses (e.g., [52]). Second, records from other locations show that this decline is occurring at such large spatial scales that it can only be explained by large-scale, more global environmental drivers like land-use change, climate change, or the global use of pollutants. Indeed, the extraordinarily low population densities of *Araneus diadematus* in the Swiss midland observed in the 2019 survey (Table 1) are supported by statements of biology students at the University of Basel, according to whom generally very few large orb-weaving spiders have been spotted in recent years during walks through forests and fields in the Basel region. Furthermore, in mid-June 2017, a group of over 40 biology experts conducted an extensive faunistic survey in the grounds of the Merian Gardens, a park-like area covering $180,000\text{ m}^2$ located on the outskirts of Basel [53]. In the course of this survey, only three specimens of *A. diadematus* could be found over a time period of 24 h suggesting that nowadays this once “abundant garden spider” [54] must have become rare in that area. The only large orb-weaver species still found in high densities during the 2019 survey in the Swiss midland was the bridge spider *Larinioides sclopetarius* (Nyffeler, pers. observations). This nocturnal species, which lives near water and frequently builds its webs on street lights and illuminated bridge railings, still gets sufficient amounts of food in the form of flying adults of semiaquatic insects (chironomids, ephemeropterans, trichopterans, etc.) attracted in large numbers to the artificial light [55–57]. Because of its capability to exploit these artificially high prey densities, *L. sclopetarius* is—in contrast to other large orb-weavers—extremely successful in colonizing urban habitats in high density even in the 21st century, not only in Switzerland, but also in other parts of western Europe [56].

Similar extraordinarily low population densities of *Araneus diadematus* (0.0004 adult females m^{-2} at a landscape scale) were also recorded in an extensive survey in nine landscapes of the Ghent region, northern Belgium, in the summer of 2014 [49]. Here, likewise, densities were one order of magnitude or more lower than those recorded one decade earlier (e.g., during a survey in 2004–2006, densities at two locations were 10–20 times higher compared to those ten years later [58]; Bonte, pers. observations). Interestingly, while other orb-weaver species—especially the larger ones—showed dramatic declines in both abundance and species richness along an urbanization gradient [59,60], *A. diadematus* was found to reach similar low densities across this land-use gradient. In other words, local densities at spatial scales of approx. 100 km were alarmingly low in both the more natural and human-impacted regions, demonstrating regional declines beyond the scale of local land use. Thus, extraordinarily low present-day population densities of *Araneus diadematus* have been recorded in two different regions of western Europe, ≈ 500 km apart. These declines extend beyond local environmental changes at small scales and suggest common negative impacts of intensive urbanization, climate change, or any other large-scale stressor across the entire landscape over the past half century [61,62].

The present study and that from northern Belgium suggest that the notion that *Araneus diadematus* is an abundant spider, as pointed out in much of the literature [63,64], has these days turned out to be a myth—at least in highly urbanized western European landscapes such as the Swiss midland and

northern Belgium. So far, the apparent abundance decline of this species in western Europe has been ignored by the faunists in charge of compiling local Red Lists for spiders (e.g., [63,64]). Red Lists are based on the number of different locations within a landscape in which a species has been recorded, and not on absolute population densities. The fact that *Araneus diadematus* is still found in many places (although in highly reduced densities) may explain why it is still labeled an “abundant species” (e.g., [63,64]).

In contrast, present-day population densities of *Araneus diadematus* in some areas outside of continental western Europe still appear to be rather high. For instance, exterminators of the pest control firm Senske Services removed more than 75 *Araneus diadematus* spiders from the exterior of a “2500 square foot home” (232.5 m²) in Seattle (USA) in August 2016 [65]. Evidently, the spiders at this particular location were still capturing sufficient food in the form of flying insects to enable them to build up a population density roughly 290 times that of the present-day Swiss overall mean density. Future detailed assessments of the population densities of this spider species in different parts of its geographic range are urgently needed, and may provide important information on the extent of the loss of flying insects in various regions of western Europe and further afield. We would in this respect like to promote the use of new citizen-science tools such as SpiderSpotter [66] to achieve these highly needed insights, and especially to guide short-term action to mitigate alarming declines in these sentinel arthropod top predators.

5. Conclusions

The drastic decline in the abundance of the orb-weaving spider *Araneus diadematus* over the past half-century documented in this study (Table 1) apparently reveals a bottom-up trophic cascade in response to the widespread insect losses that have occurred across large parts of Europe in recent decades [1,2,4,6,9,39,67,68]. There is evidence that other groups of aerial web-spinning spiders, which likewise depend on flying insects as food [17–19,21,69–71], have also become much rarer over the recent past (Nyffeler, pers. observations). So, for instance, the mesh-web weaver *Dictyna uncinata* (together with other dictynid species), whose small, tangled webs were found in large numbers on the leaves of garden plants a few decades ago [71], has apparently become very rare these days (Nyffeler, pers. observations). The ongoing abundance decline of the spiders parallels the dramatic abundance declines in other insectivorous animals such as insectivorous birds, bats, frogs, and lizards documented in recent decades [72–74].

To sum up, the findings of this study support the notion by other researchers that we now live in the midst of an ecological crisis in which trophic webs are being eroded and degraded as a result of adverse, man-made environmental impacts [13,67,73,75,76]. If this disastrous trend cannot be halted or even reversed in the very near future, “entire ecosystems will collapse due to starvation” [16].

Supplementary Materials: The following are available online at <http://www.mdpi.com/2075-4450/11/4/248/s1>, Table S1: Area size (m²) of the study plots, Table S2: Mean number of prey per web counted in mid-afternoon as a proxy for the daily prey capture rate of large orb-weaving spiders in western European habitats: Historical values vs. present day values.

Author Contributions: Conceptualization, M.N. and D.B.; methodology, M.N.; software, M.N. and D.B.; validation: M.N. and D.B.; formal analysis: M.N. and D.B.; investigation, M.N. (in Switzerland) and D.B. (in Belgium); data curation, M.N.; writing—original draft preparation, M.N.; writing—review & editing, D.B. All authors have read and agreed to the published version of the manuscript.

Funding: This research was in part supported by the Swiss National Science Foundation (grant to G. Benz, Swiss Federal Institute of Technology Zurich) in as far as it refers to 20th century spider density assessments.

Acknowledgments: The authors wish to extend their gratitude to Rätus Fischer and Christlinde Ruchti for assistance in the field. We also thank two anonymous reviewers for their valuable comments. One of the authors (M.N.) also wishes to thank Georg Benz, Swiss Federal Institute of Technology Zurich, for his past support.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Hallmann, C.A.; Sorg, M.; Jongejans, E.; Siepel, H.; Hofland, N.; Schwan, H.; Stenmans, W.; Müller, A.; Sumser, H.; Hörrén, T.; et al. More than 75 percent decline over 27 years in total flying insect biomass in protected areas. *PLoS ONE* **2017**, *12*, e0185809. [[CrossRef](#)] [[PubMed](#)]
- Seibold, S.; Gossner, M.M.; Simons, N.K.; Blüthgen, N.; Müller, J.; Ambarli, D.; Ammer, C.; Bauhus, J.; Fischer, M.; Habel, J.C.; et al. Arthropod decline in grasslands and forests is associated with landscape-level drivers. *Nature* **2019**, *574*, 671–674. [[CrossRef](#)] [[PubMed](#)]
- Schuch, S.; Wesche, K.; Schaefer, M. Long-term decline in the abundance of leafhoppers and planthoppers (Auchenorrhyncha) in Central European protected dry grasslands. *Biol. Conserv.* **2012**, *149*, 75–83. [[CrossRef](#)]
- Hallmann, C.A.; Zeegers, T.; van Klink, R.; Vermeulen, R.; van Wielink, P.; Spijkers, H.; van Deijk, J.; van Steenis, W.; Jongejans, E. Declining abundance of beetles, moths and caddisflies in the Netherlands. *Insect. Conserv. Diver.* **2019**, *13*, 127–139. [[CrossRef](#)]
- Harris, J.E.; Rodenhouse, N.L.; Holmes, R.T. Decline in beetle abundance and diversity in an intact temperate forest linked to climate warming. *Biol. Conserv.* **2019**, *240*, 108219. [[CrossRef](#)]
- Møller, A.P. Parallel declines in abundance of insects and insectivorous birds in Denmark over 22 years. *Ecol. Evol.* **2019**, *9*, 6581–6587. [[CrossRef](#)]
- Wepprich, T.; Adrion, J.R.; Ries, L.; Wiedmann, J.; Haddad, N.M. 2019 Butterfly abundance declines over 20 years of systematic monitoring in Ohio, USA. *PLoS ONE* **2019**, *14*, e0216270. [[CrossRef](#)]
- Stepanian, P.M.; Entrekina, S.A.; Wainwright, C.E.; Mirkovic, D.; Tank, J.L.; Kelly, J.F. Declines in an abundant aquatic insect, the burrowing mayfly, across major North American waterways. *Proc. Natl. Acad. Sci. USA* **2020**, *117*, 2987–2992. [[CrossRef](#)]
- Tinsley-Marshall, P.; Skilbeck, A.; Riggs, A. *Monitoring of Ecosystem Function at Landscape-Scale Demonstrates Temporal Difference in Invertebrate Abundance in Kent and South-East England*; Report Kent Wildlife Trust: Maidstone, UK, 2020.
- Leather, S.R. “Ecological Armageddon”—More evidence for the drastic decline in insect numbers. *Ann. Appl. Biol.* **2017**, *172*, 1–3. [[CrossRef](#)]
- Samways, M.J. Addressing global insect meltdown. *Ecol. Citizen* **2019**, *3* (Suppl. A), 23–26.
- Sánchez-Bayo, F.; Wyckhuys, K.A. Worldwide decline of the entomofauna: A review of its drivers. *Biol. Conserv.* **2019**, *232*, 8–27. [[CrossRef](#)]
- Wagner, D.L. Insect declines in the Anthropocene. *Annu. Rev. Entomol.* **2020**, *65*, 457–480. [[CrossRef](#)] [[PubMed](#)]
- Nyffeler, M.; Birkhofer, K. An estimated 400–800 million tons of prey are annually killed by the global spider community. *Sci. Nat.* **2017**, *104*, 30. [[CrossRef](#)] [[PubMed](#)]
- Nyffeler, M.; Şekercioğlu, Ç.H.; Whelan, C.J. Insectivorous birds consume an estimated 400–500 million tons of prey annually. *Sci. Nat.* **2018**, *105*, 47. [[CrossRef](#)] [[PubMed](#)]
- Why Insect Populations are Plummeting –And Why It Matters. Available online: <https://www.nationalgeographic.com/animals/2019/02/why-insect-populations-are-plummeting-and-why-it-matters/> (accessed on 29 February 2020).
- Nyffeler, M.; Dean, D.A.; Sterling, W.L. Prey selection and predatory importance of orb-weaving spiders (Araneae: Araneidae, Uloboridae) in Texas cotton. *Environ. Entomol.* **1989**, *18*, 373–380. [[CrossRef](#)]
- Malt, S. Untersuchungen zur Rolle ausgewählter netzbauender Spinnen (Araneida) im trophischen Beziehungsgefüge von Halbtrockenrasen. Ph.D. Thesis, University of Jena, Jena, Germany, 1996.
- Ludy, C. Prey selection of orb-web spiders (Araneidae) on field margins. *Agric. Ecosyst. Environ.* **2007**, *119*, 368–372. [[CrossRef](#)]
- Nyffeler, M. Diel Activity Patterns, Prey Composition, and Prey Consumption Rate of the Orb-Weaving Spiders *Araneus diadematus* and *Nuctenea umbratica* in a Garden in Höngg (Zurich). Master’s Thesis, Swiss Federal Institute of Technology Zurich, Zurich, Switzerland, 1976.
- Nyffeler, M. Field Studies on the Ecological Role of the Spiders as Insect Predators in Agroecosystems (Abandoned Grassland, Meadows and Cereal Fields). Ph.D. Thesis, Swiss Federal Institute of Technology Zurich, Zurich, Switzerland, 1982.
- Bee, L.; Oxford, G.; Smith, H. *Britain’s Spiders: A Field Guide*; Princeton University Press: Princeton, NJ, USA, 2017.

23. Lubin, Y.D. Seasonal abundance and diversity of web-building spiders in relation to habitat structure on Barro Colorado Island, Panama. *J. Arachnol.* **1978**, *6*, 31–51.
24. Hiby, L.; Krishna, M.B. Line transect sampling from a curving path. *Biometrics* **2001**, *57*, 727–731. [[CrossRef](#)] [[PubMed](#)]
25. Hänggi, A.; Stöckli, E.; Nentwig, W. *Lebensräume Mitteleuropäischer Spinnen: Charakterisierung der Lebensräume der häufigsten Spinnenarten Mitteleuropas und der mit diesen Vergesellschafteten Arten*; Centre Suisse de Cartographie de la Faune: Neuchâtel, Switzerland, 1995.
26. Jocqué, R.; Samu, F.; Bird, T. Density of spiders (Araneae: Ctenidae) in Ivory Coast rainforests. *J. Zool.* **2005**, *266*, 105–110. [[CrossRef](#)]
27. Holzapfel, M. Die Gewächshausfauna des Berner Botanischen Gartens. *Rev. Suisse Zool.* **1932**, *39*, 325–374. [[CrossRef](#)]
28. Netzstandorte und Bestandesdichte von Radnetzspinnen auf einer Probefläche. Available online: <https://www.natur-erforschen.net/unterrichtsprojekte/spinnen-ruheue/projektbeschreibung-spinnen/netzstandorte.html> (accessed on 29 February 2020).
29. Pasquet, A. Distribution of two orb-weaving spiders *Araneus marmoreus* (Clerk) and *Araneus diadematus* (Clerk). *Biol. Behav./Biol. Comport.* **1984**, *9*, 321–331.
30. Le Berre, M.; Ramousse, R.; Le Guelte, L. Eco-éthologie des Argiopidae: 1. Evolution temporelle d'une population d'*Araneus diadematus* Clerck dans son milieu naturel. *Atti. Soc. Toscana Sci. Nat. P. V. Mem. Ser. B* **1981**, *88*, 72–83.
31. Fasola, M.; Mogavero, F. Structure and habitat use in a web-building spider community in northern Italy. *Boll. Zool.* **1995**, *62*, 159–166. [[CrossRef](#)]
32. Krzyzanowska, E. Field experiments on the filling of space with the community of *Araneus diadematus* (Clerck) (Aranei) in a pine stand. *Mem. Zool.* **1979**, *30*, 27–35.
33. Van der Hammen, L. The distribution of spiders and harvestmen (Chelicerata) in the Dutch National Park "De Hoge Veluwe". *Zool. Meded.* **1983**, *57*, 325–338.
34. Turnbull, A.L. The spider population of a stand of oak (*Quercus robur* L.) in Wytham Wood, Berks., England. *Can. Entomol.* **1960**, *92*, 110–124. [[CrossRef](#)]
35. Colebourn, P.H. The influence of habitat structure on the distribution of *Araneus diadematus* Clerck. *J. Anim. Ecol.* **1974**, *43*, 401–409. [[CrossRef](#)]
36. Witt, P.N.; Reed, C.F.; Peakall, D.B. *A Spider's Web: Problems in Regulatory Biology*; Springer: New York, NY, USA, 1968.
37. Vollrath, F.; Samu, F. The effect of starvation on web geometry in an orb-weaving spider. *Bull. Br. Arachnol. Soc.* **1997**, *10*, 295–297.
38. Insektenschwund in der Schweiz und mögliche Folgen für Gesellschaft und Wirtschaft. Available online: <http://www.akademien-schweiz.ch/dms/D/Aktuell/News/Faktenblatt-Insekten/Faktenblatt%20Insekten.pdf> (accessed on 29 February 2020).
39. Decline of Insectivorous Birds. Available online: <https://www.vogelwarte.ch/en/atlas/focus/decline-of-insectivorous-birdsn> (accessed on 29 February 2020).
40. Vogel, G. Where have all the insects gone? *Science* **2017**, *356*, 576–579. [[CrossRef](#)]
41. Kirchner, W. Bisher Bekanntes über die forstliche Bedeutung der Spinnen—Versuch einer Literaturanalyse. *Waldhygiene* **1964**, *5*, 161–198.
42. Robinson, M.H.; Robinson, B. Prey caught by a sample population of the spider *Argiope argentata* (Araneae: Araneidae) in Panama: A year's census data. *Zool. J. Linn. Soc.-Lond.* **1970**, *49*, 345–358. [[CrossRef](#)]
43. Leborgne, R.; Pasquet, A. Time of oviposition and reproductive success in *Argiope bruennichi* (Araneae: Araneidae). *Eur. J. Entomol.* **2005**, *102*, 169–174. [[CrossRef](#)]
44. Samu, F.; Vollrath, F. Spider orb web as bioassay for pesticide side effects. *Entomol. Exp. Appl.* **1992**, *62*, 117–124. [[CrossRef](#)]
45. Lengwiler, U.; Benz, G. Effects of selected pesticides on web building behaviour of *Larinioides sclopetarius* (Clerck)(Araneae, Araneidae). *J. Appl. Entomol.* **1994**, *117*, 99–108. [[CrossRef](#)]
46. Benamú, M.A.; Schneider, M.I.; Sánchez, N.E. Effects of the herbicide glyphosate on biological attributes of *Alpaida veniliae* (Araneae, Araneidae), in laboratory. *Chemosphere* **2010**, *78*, 871–876. [[CrossRef](#)]

47. Benamú, M.A.; Schneider, M.I.; González, A.; Sánchez, N.E. Short and long-term effects of three neurotoxic insecticides on biological and behavioural attributes of the orb-web spider *Alpaida veniliae* (Araneae, Araneidae): Implications for IPM programs. *Ecotoxicology* **2013**, *22*, 1155–1164. [CrossRef]
48. Pasquet, A.; Tupinier, N.; Mazzia, C.; Capowiez, Y. Exposure to spinosad affects orb-web spider (*Agalenatea redii*) survival, web construction and prey capture under laboratory conditions. *J. Pest Sci.* **2016**, *89*, 507–515. [CrossRef]
49. Dahirel, M.; De Cock, M.; Vantieghem, P.; Bonte, D. Urbanization-driven changes in web building and body size in an orb web spider. *J. Anim. Ecol.* **2019**, *88*, 79–91. [CrossRef]
50. Nyffeler, M.; Benz, G. Bemerkenswerte Spinnenfunde in Wiesen und Getreidefeldern bei Zürich. *Mitt. Schweiz. Ent. Ges.* **1982**, *55*, 97–100.
51. Green, D.M. The ecology of extinction: Population fluctuation and decline in amphibians. *Biol. Conserv.* **2003**, *111*, 331–343. [CrossRef]
52. Wolff, J.O. Population fluctuations of mast-eating rodents are correlated with production of acorns. *J. Mammal.* **1996**, *77*, 850–856. [CrossRef]
53. Eggenschwiler, L.; Hänggi, A. 24 Stunden Forschung live: Ergebnisse vom GEO-Tag der Natur in den Merian Gärten in Basel. *Mitt. Naturf. Ges. Basel* **2018**, *18*, 19–37.
54. Nyffeler, M. Eine Notiz zur ökologischen Bedeutung der Radnetzspinnen als Blattlausprädatoren in Gärten. *Mitt. Schweiz. Ent. Ges.* **1983**, *56*, 200.
55. Kleinteich, A. Life History of the Bridge Spider, *Larinioides sclopetarius* (Clerck, 1757). Ph.D. Thesis, University of Hamburg, Hamburg, Germany, 2010.
56. Kleinteich, A.; Schneider, J.M. Developmental strategies in an invasive spider: Constraints and plasticity. *Ecol. Entomol.* **2011**, *36*, 82–93. [CrossRef]
57. Richter, D.; Göbel, D.; Schmitt, M. Zum Beutespektrum der Brückenspinne (*Larinioides sclopetarius*; Araneae: Araneidae) im Ruhrgebiet. *Elektronische Aufsätze der Biologischen Station Westliches Ruhrgebiet* **2012**, *22*, 1–9.
58. Bonte, D.; Lanckacker, K.; Wiersma, E.; Lens, L. Web building flexibility of an orb-web spider in a heterogeneous agricultural landscape. *Ecography* **2008**, *31*, 646–653. [CrossRef]
59. Dahirel, M.; Dierick, J.; De Cock, M.; Bonte, D. Intraspecific variation shapes community-level behavioral responses to urbanization in spiders. *Ecology* **2017**, *98*, 2379–2390. [CrossRef]
60. Merckx, T.; Souffreau, C.; Kaiser, A.; Baardsen, L.F.; Backeljau, T.; Bonte, D.; Brans, K.I.; Cours, M.; Dahirel, M.; Debortoli, N.; et al. Body-size shifts in aquatic and terrestrial urban communities. *Nature* **2018**, *558*, 113–116. [CrossRef]
61. Antrop, M. Landscape change and the urbanization process in Europe. *Landscape Urban Plan.* **2004**, *67*, 9–26. [CrossRef]
62. Urbanisierungsgrad in der Schweiz nach Kantonen am 31. Dezember 2018. Available online: <https://de.statista.com/statistik/daten/studie/712490/umfrage/urbanisierung-in-der-schweiz-nach-kantonen/> (accessed on 29 February 2020).
63. Martin, D. *Rote Liste der Spinnen (Araneae) Mecklenburg-Vorpommerns*; Ministerium für Landwirtschaft, Umwelt und Verbraucherschutz Mecklenburg-Vorpommern: Schwerin, Germany, 2012.
64. Lemke, M.; Reinke, H.D.; Vahder, S.; Irmeler, U. *Die Spinnen Schleswig-Holsteins. Rote Liste*; Ministerium für Energiewende, Landwirtschaft, Umwelt und ländliche Räume des Landes Schleswig-Holstein: Kiel, Germany, 2013.
65. Pest Control in the Pacific Northwest. Available online: <https://www.pctonline.com/article/pest-control-pacific-northwest-allen/> (accessed on 29 February 2020).
66. SpiderSpotter. Available online: <https://www.spinnenspotter.be/en/> (accessed on 29 February 2020).
67. Plummeting insect numbers ‘threaten collapse of nature’. Available online: <https://www.theguardian.com/environment/2019/feb/10/plummeting-insect-numbers-threaten-collapse-of-nature> (accessed on 29 February 2020).
68. Møller, A.P. Quantifying rapidly declining abundance of insects in Europe using a paired experimental design. *Ecol. Evol.* **2020**, *10*, 2446–2451. [CrossRef]
69. Turnbull, A.L. The prey of the spider *Linyphia triangularis* (Clerck)(Araneae, Linyphiidae). *Can. J. Zool.* **1960**, *38*, 859–873. [CrossRef]
70. Kajak, A. An analysis of food relations between the spiders *Araneus cornutus* and *A. quadratus* and their prey in meadows. *Ekol. Pol. A* **1965**, *13*, 717–768.

71. Nyffeler, M. Über den Beutefang der Netzspinne *Dictyna uncinata* Thorell. Bachelor's Thesis, Department of Entomology, Swiss Federal Institute of Technology Zurich, Zurich, Switzerland, 1974.
72. Inger, R.; Gregory, R.; Duffy, J.P.; Stott, I.; Voříšek, P.; Gaston, K.J. Common European birds are declining rapidly while less abundant species' numbers are rising. *Ecol. Lett.* **2015**, *18*, 28–36. [[CrossRef](#)] [[PubMed](#)]
73. Lister, B.C.; Garcia, A. Climate-driven declines in arthropod abundance restructure a rainforest food web. *Proc. Natl. Acad. Sci. USA* **2018**, *115*, E10397–E10406. [[CrossRef](#)] [[PubMed](#)]
74. Rosenberg, K.V.; Dokter, A.M.; Blancher, P.J.; Sauer, J.R.; Smith, A.C.; Smith, P.A.; Stanton, J.C.; Panjabi, A.; Helft, L.; Parr, M.; et al. Decline of the North American avifauna. *Science* **2019**, *366*, 120–124. [[CrossRef](#)]
75. Dobson, A.; Lodge, D.; Alder, J.; Cumming, G.S.; Keymer, J.; McGlade, J.; Mooney, H.; Rusak, J.A.; Sala, O.; Wolters, V.; et al. Habitat loss, trophic collapse, and the decline of ecosystem services. *Ecology* **2006**, *87*, 1915–1924. [[CrossRef](#)]
76. Goulson, D. The insect apocalypse, and why it matters. *Curr. Biol.* **2019**, *29*, R967–R971. [[CrossRef](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).