

The impact of increased flooding occurrence on the mobility of potentially toxic elements in floodplain soil – a review

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Ponting, J., Kelly, T. J., Verhoef, A., Watts, M. J. and Sizmur, T. (2021) The impact of increased flooding occurrence on the mobility of potentially toxic elements in floodplain soil – a review. *Science of the Total Environment*, 754. 142040. ISSN 0048-9697 doi: <https://doi.org/10.1016/j.scitotenv.2020.142040> Available at <http://centaur.reading.ac.uk/92613/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.scitotenv.2020.142040>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in

the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1

2 The impact of increased flooding occurrence on the
3 mobility of potentially toxic elements in floodplain soil
4 – A review

5

6 Jessica Ponting ^{a,b}, Thomas J. Kelly ^{a,c}, Anne Verhoef ^a, Michael J. Watts ^b and
7 Tom Sizmur ^{a*}

8
9 ^a Department of Geography and Environmental Science, University of Reading, Reading, RG6
10 6DW, UK

11 ^b Inorganic Geochemistry, Centre for Environmental Geochemistry, British Geological
12 Survey, Keyworth, Nottingham, NG12 5GG, UK

13 ^c Department of Geography, Queen Mary University of London, Mile End Road, London, E1
14 4NS, UK

15
16 * Corresponding author

17 Department of Geography and Environmental Science

18 Russell Building

19 University of Reading

20 Reading

21 RG6 6DW, UK

22 t.sizmur@reading.ac.uk

23 +44(0)118 3788913

24

25

26

27

28 Abstract

29 The frequency and duration of flooding events are increasing due to land-use changes increasing run-
30 off of precipitation, and climate change causing more intense rainfall events. Floodplain soils situated
31 downstream of urban or industrial catchments, which were traditionally considered a sink of
32 potentially toxic elements (PTEs) arriving from the river reach, may now become a source of legacy
33 pollution to the surrounding environment if PTEs are mobilised by unprecedented flooding events.

34 When a soil floods, the mobility of PTEs can increase or decrease due to the net effect of five key
35 processes; (i) the soil redox potential decreases which can directly alter the speciation, and hence
36 mobility, of redox sensitive PTEs (e.g. Cr, As), (ii) pH increases which usually decreases the mobility of
37 metal cations (e.g. Cd^{2+} , Cu^{2+} , Ni^{2+} , Pb^{2+} , Zn^{2+}), (iii) dissolved organic matter (DOM) increases, which
38 chelates and mobilises PTEs, (iv) Fe and Mn hydroxides undergo reductive dissolution, releasing
39 adsorbed and co-precipitated PTEs, and (v) sulphate is reduced and PTEs are immobilised due to
40 precipitation of metal sulphides. These factors may be independent mechanisms, but they interact
41 with one another to affect the mobility of PTEs, meaning the effect of flooding on PTE mobility is not
42 easy to predict. Many of the processes involved in mobilising PTEs are microbially mediated,
43 temperature dependent and the kinetics are poorly understood.

44 Soil mineralogy and texture are properties that change spatially and will affect how the mobility of
45 PTEs in a specific soil may be impacted by flooding. As a result, knowledge based on one river
46 catchment may not be particularly useful for predicting the impacts of flooding at another site. This
47 review provides a critical discussion of the mechanisms controlling the mobility of PTEs in floodplain
48 soils. It summarises current understanding, identifies limitations to existing knowledge, and highlights
49 requirements for further research.

50

51 **Key words;** floodplain soil, flooding, climate change, potentially toxic elements, contamination,
52 mobility

53

54 1. Introduction

55 Flooding is a major event that currently affects an estimated 20 to 300 million people per year,
56 and accounts for around 40% of natural disasters occurring worldwide, threatening both social
57 security and sustainable development (Euripidou and Murray, 2004; Hirabayashi and Kanae, 2009).
58 Alterations to land use and land cover are having widespread implications for catchment
59 characteristics; with soil sealing and impermeable surfaces increasing surface run-off, as well as a
60 reduction of natural buffering environments such as forests and wetlands, meaning there is less
61 capacity to accommodate flood waters in the same river reach (Dadson et al., 2017; Kundzewicz et al.,
62 2014). There is growing evidence, from climate models, that short-term extreme weather events (e.g.
63 high-frequency rainstorms, heat waves and wind storms) are likely to become increasingly frequent
64 in many parts of the world, threatening the long-term functioning of the terrestrial system (Harvey et
65 al., 2019; Kharin et al., 2007; Madsen et al., 2014; Pendergrass, 2018; Stagl et al., 2014). It is likely that
66 populations will experience warmer and drier summers, and an increase in the intensity of heavy
67 rainfall, contributing to more frequent pluvial, fluvial, groundwater or coastal flooding, and resulting
68 in the occasional inundation of land that has rarely been flooded in the past (Barber et al., 2017;
69 Kundzewicz et al., 2014; Schaller et al., 2016). The likelihood of flooding is also determined by
70 antecedent soil moisture conditions. The proportion of soil pore space that is filled with water at any
71 given time is largely dependent on local hydrological processes and stores including; infiltration,
72 surface and sub-surface runoff (when rainfall intensity exceeds infiltration capacity), redistribution
73 and drainage to/from groundwater, evaporation, and transpiration (Stagl et al., 2014).

74 Soil contamination is among the most serious threats to soil resources globally (Nriagu et al., 2007;
75 Srivastava et al., 2017; Tóth et al., 2016b). Since many commercial, industrial, residential and
76 agricultural developments have historically been situated adjacent to rivers; they contribute to the
77 contamination of river sediments, and these sediments are often deposited onto the floodplain soils
78 downstream by overbanking river water during a flooding event (Arnell et al., 2015; Nshimiyimana et

79 al., 2014; Zhao et al., 1999). Here we use the term PTEs, also referred to in the literature as ‘trace
80 elements’ or ‘heavy metals’, to encompass all metals, metalloids, non-metals and other inorganic
81 elements in the soil–plant–animal system, of which their mobility and potential toxicity to that system
82 and/or humans is largely dependent upon their concentration, bioavailability and chemical form
83 (Hooda, 2010; Rodgers et al., 2015). The term "mobility" is a concept that has been frequently used
84 to estimate the risk of contamination from the soil to the surrounding environment by PTEs
85 (Domergue and Vedy, 1992). Here we define mobile PTEs as those elements that are dissolved in soil
86 porewater or associated with colloids and thus capable of leaching from the soil profile, or being taken
87 up into plants or soil organisms. The mobility and subsequent fate of PTEs in periodically (occasionally)
88 flooded soils (such as floodplain soils) are imperfectly understood. The legacy of historic
89 contamination and continuing increases in emissions from urban activities pose a serious
90 environmental threat globally (de Souza Machado et al., 2016; Srivastava et al., 2017). Human actions
91 to mitigate and adapt to the impacts of climate change may influence the fate of contaminants, with
92 climate change itself also potentially affecting the toxicity of the contaminants within the environment
93 (Stahl et al., 2013).

94 The aim of this literature review is to provide an understanding of the factors involved in the
95 mobility of PTEs in soil by pulling together interdisciplinary knowledge in this area. The review will first
96 consider in more detail the expected changes to global rainfall patterns, the implications of these
97 changes for flooding, and the role that floodplains play during inundation, as well as the changes they
98 undergo. The review will then showcase how PTEs have entered the floodplain soil and how flooding
99 influences soil biogeochemical processes which, in turn, influence PTEs mobility, using examples from
100 the literature. Finally, this knowledge is used to identify gaps that will help to make recommendations
101 for future research into the effects of flooding on the mobility and fate of PTEs.

102

103 1.1. Expected changes to global rainfall patterns and implications for flooding

104 Anthropogenic (human) activities including intensified land use; urbanisation, forestry,
105 cultivation, and fossil energy use have increased atmospheric greenhouse gas concentrations which
106 are driving changes in climate and leading to increases in rainfall intensity and surface run-off that are
107 associated with increased flood risk (Bronstert, 2003; Chang and Franczyk, 2008; Kharin et al., 2007;
108 Kundzewicz et al., 2014; Wheeler and Evans, 2009). Mean global temperatures have risen by 1.1 °C
109 since the end of the 19th century; the “Paris Climate Agreement” seeks to contain global mean
110 temperatures well below 2°C and, ambitiously, below 1.5°C (Alfieri et al., 2017; Bronstert, 2003;
111 Huddart et al., 2020; Mullan et al., 2019). The Intergovernmental Panel on Climate Change (IPCC) has
112 predicted that under the A1B (medium) emissions scenario, temperatures will increase between 1.1
113 and 6.4 °C by the year 2100, leading to an increase in atmospheric water holding capacity and therefore
114 variations to seasonal rainfall (Arnell et al., 2015; Bell et al., 2012; Chan et al., 2014; Clemente et al.,
115 2008; González-Alcaraz and van Gestel, 2015; Jenkins et al., 2009). It has been argued that we will
116 experience an intensification of short-duration heavy rainfall events rather than a uniform increase in
117 the daily average rainfall (Chan et al., 2014; Hirabayashi et al., 2008; Kharin et al., 2007; Kundzewicz
118 et al., 2014).

119 An IPCC Special Report (SREX) on climate extremes (IPCC, 2012) assessed it is *likely* there have
120 been statistically significant increases in the number of heavy precipitation events in more regions
121 than significant decreases, with strong regional and sub-regional variation. The observed changes to
122 precipitation extremes have been found to be far less spatially coherent or statistically significant
123 compared with changes found in temperature extremes (Kundzewicz et al., 2014). Projected scenarios
124 with 4°C warming showed more than 70% of the global population will face increased flood risk (Alfieri
125 et al., 2017). Increases in flood frequency are expected in; Europe, America, Southeast Asia, eastern
126 Africa, and Peninsular India. Populations in regions such as Bangladesh, Mumbai and Thailand are
127 potentially at higher risk from flooding due to predicted increases in rainfall, coupled with changes in
128 land use (e.g. irrigation schemes and construction of dams), and increasing population size requiring

129 rapid expansion of urban areas (Hirabayashi et al., 2013; Kundzewicz et al., 2014; Tockner et al., 2010).
130 However, climate projections tend to have relatively low levels of model confidence, particularly for
131 the prediction of fluvial floods because there is still relatively limited evidence and the causes of
132 regional changes to flood occurrence are complex (Hirabayashi et al., 2013).

133 Flooding tends to be heterogeneous as it is affected not only by variability of the climatological
134 and hydrological systems but also by land-use and the effect is has on the storage capacity of the
135 receiving catchment (storage and drainage basin conditions). Changes to the characteristics of
136 precipitation (the frequency, intensity and timing of rainfall) will have decisive implications for flood
137 risk (Bronstert, 2003; Hirabayashi and Kanae, 2009; Kundzewicz et al., 2014). However, pre-existing
138 high river levels and groundwater levels, as well as saturated soils are equally important to establish
139 the capacity of the receiving catchment to cope with further rainfall (Maggioni and Massari, 2018;
140 Wilby et al., 2008). The extent of flooding in a particular catchment will depend largely on the
141 topography (variation in elevation), along with vegetation type, proportion of land used for cultivation
142 and the extent of urbanised areas positioned upstream (Arnell et al., 2015; Bell et al., 2012; Bronstert,
143 2003; Chang and Franczyk, 2008; Kundzewicz et al., 2014; Qiao et al., 2019). Urbanisation is a global
144 issue; with more than half the world's population now living in cities, the process of urbanisation is
145 leading to greater human occupation of floodplains, often with inadequate drainage planning
146 (Kundzewicz et al., 2014; Pathirana et al., 2014).

147 The probability of flooding occurring in a particular region is often related to regional processes
148 like El Niño Southern Oscillation (ENSO) cycle and the North Atlantic Oscillation (NAO) that, in turn,
149 cause global impacts. The intensity (frequency and amplitude) of both ENSO and NAO are influenced
150 by other modes of variability, for example; Pacific Decadal Oscillation (PDO) and Interdecadal Pacific
151 Oscillation (IPO) which cause opposite atmospheric and sea surface temperatures and can therefore
152 determine the magnitude of floods (Grimm and Tedeschi, 2009; Johnson et al., 2020). ENSO is a rapid
153 warming of the sea surface temperature (by 1–5 °C) of the equatorial Pacific over the duration of a

154 few weeks, resulting in extreme rainfall and increased cyclone activity in some regions, and risk of
155 drought and forest fires in others (Berz et al., 2001; Grimm and Tedeschi, 2009; Karl and Trenberth,
156 2003; Kundzewicz et al., 2014; Tedeschi and Collins, 2016). Periods of extreme rainfall and subsequent
157 flooding have been found to correlate with ENSO events in North and South America as well as in
158 Africa (Berz et al., 2001; Brönnimann, 2007; Kundzewicz et al., 2014). NAO is an atmospheric pattern
159 that affects the severity of winter temperatures and precipitation over Europe and eastern North
160 America (Karl and Trenberth, 2003). Intense rainfall is a common cause of river basin flooding;
161 however, in high latitude regions it is changes in temperatures altering the timing of seasonal
162 snowmelt and causing glacier retreat that commonly causes flooding, for example in north-eastern
163 Europe, Central and South America, and in polar regions such as the Russian Arctic (Blöschl et al., 2017;
164 Hirabayashi et al., 2008; Kharin et al., 2007; Kundzewicz et al., 2014; Stagl et al., 2014). Rising global
165 sea-level (11-16cm in the 20th century and a further 0.5m predicted this century) will certainly increase
166 risk of flooding caused by tidal processes, with current estimates that 630 million people live on land
167 below projected annual flood levels for 2100 (Kulp and Strauss, 2019). While there is uncertainty
168 regarding the effect that future climate change will have on river levels (Prudhomme and Davies,
169 2009), changes made to land-use, and land cover, for example by urbanisation, will drive changes in
170 the local climate (at the kilometre scale) influencing the hydrometeorological regime and resulting in
171 more flooding (Foley et al., 2005; Hirabayashi and Kanae, 2009). Pathirana et al. (2014), using a 3D
172 atmospheric model coupled with a land surface model (WRF-ARW) in southern India, found that in
173 three out of four simulated cases there was a significant increase in local extreme rainfall when
174 urbanisation in the area increased. This work was conducted in southern India, however the model
175 could be applied and validated to other regions to establish whether this correlation is found globally.

176

177 1.2. The role of floodplains during floods

178 Floodplains are by definition dynamic environments subjected to fluctuations between flooding
179 and drying (Vijver et al., 2007). They are distinctive landscape features, often on low-lying ground, and

180 characterised by a high spatio-temporal heterogeneity (Schulz-Zunkel et al., 2015; Stuart and
181 Lapworth, 2011; Tockner et al., 2010; Tockner and Stanford, 2002). Periodic overbank inundation from
182 the adjacent watercourse, overland flow, subsurface flow, and changes to the groundwater levels
183 result in a constantly changing water balance and degree of floodplain saturation (Stuart and
184 Lapworth, 2011; Tockner and Stanford, 2002). Floodplain topography and variations in elevation are
185 usually slight but have an important effect on the degree of soil saturation across the floodplain,
186 depending on the overall water balance from surface and sub-surface run-off (Arnell et al., 2015;
187 Kundzewicz et al., 2014; Qiao et al., 2019).

188 There are various sources and pathways of water that can lead to the inundation of a floodplain,
189 including lateral overflow of rivers or lakes, rising groundwater, upland sources, and direct
190 precipitation. Several different factors and water sources normally contribute to a flooding event, thus
191 making flooding a complex phenomenon to study (Junk et al., 1989; Tockner and Stanford, 2002).
192 Fluvial flooding tends to occur when excessive rain falls over an extended period of time, leading to a
193 river exceeding its capacity, or because of heavy snow that subsequently melts and, via surface run-
194 off, rapidly fills the river channels when infiltration is low because of frozen soils below the snow layer
195 (Blöschl et al., 2017).

196 River flow regimes are affected by the increased rainfall and this also has the potential to affect
197 erosion and generate additional sediment loads and particulate organic matter (POM) for deposition
198 within river channels, lakes and estuaries (Arnell et al., 2015; Le Gall et al., 2018; Rinklebe and Du
199 Laing, 2011). Intense rainfall over a short timescale (usually less than six hours i.e. “flash floods”) can
200 also cause rivers to overbank leading to an intense, high velocity torrent of water that moves through
201 river beds, disturbing river sediments and potentially bringing more PTEs contamination with the
202 flood water, greatly influencing the contaminated status of the floodplain (Blöschl et al., 2017;
203 Maggioni and Massari, 2018). The water inundating the floodplain contains dissolved matter (i.e. free
204 ions, inorganic and organic complexes and uncharged molecules) as well as particulate matter (i.e.

205 large organic and inorganic polymers, oxides, clay minerals and organic matter) (Kirk, 2004). The
206 sediment loads travel at different rates due to their particle size, which reflects the texture of the river
207 bed and bank (Malmon et al., 2004). Approximately 90% of PTEs load has been associated with
208 sediment particles, with dissolved PTEs playing a comparatively minor role in pollutant transfer to
209 floodplains (Ciszewski and Grygar, 2016). There have been many fluvial geomorphology studies
210 showing how erosion and sedimentation have been influenced by climatic variability in the past (e.g.
211 Lewin and Macklin, 2010; Macklin and Rumsby, 2007; Mullan et al., 2019), indicating that rivers are
212 sensitive to climatic change (Arnell et al., 2015). Fluvial flooding is receiving increased scientific and
213 political interest because of the potential impact that climate change may have on this type of
214 flooding, with climate model projections showing an increased flood risk at a global scale
215 (Pappenberger et al., 2012; Wilby et al., 2008).

216 In floodplains that are underlain by permeable deposits, increased rainfall causes groundwater to
217 rise (leading to groundwater flooding), which can result from direct rainfall recharge, when the soil
218 water storage potential is exceeded, as well as flow into the floodplain sediments from rivers with
219 high water levels, and from areas inundated with fluvial flooding. However, good hydraulic connection
220 between river and aquifer means that the aquifer can drain quickly as fluvial flood waters recess.
221 Groundwater flooding in these settings is relatively short-lived compared with other groundwater
222 flood settings, for example in chalk catchments (MacDonald et al., 2012).

223 With increased frequency of rainfall events predicted, it has become widely recognised that
224 the storage of floodwater on floodplains can help to reduce the magnitude of a flood downstream.
225 Thus, floodplains are useful for flood risk management (Acreman et al., 2003; Vink and Meeussen,
226 2007). As a result, floodplains may be deliberately managed to allow flooding to occur through
227 engineered soakaways in order to protect an urban residential area (Lane, 2017; Wheater and Evans,
228 2009). It is important to understand the potential implications of these types of management practices

229 on mobilisation of PTEs that may be associated with the sediments deposited on the floodplain during
230 past flooding events.

231 1.3. Changes that floodplain soils undergo during and after inundation


232 Extreme rainfall events leading to flooding have generally been found to alter soil physical and
233 chemical properties and influence biological processes (Harvey et al., 2019). The fluctuations between
234 inundation and subsequent drying, associated with periodically flooded soils, are major drivers of
235 spatial and temporal differences in soil properties that affect the biogeochemical processes taking
236 place in floodplain soils (Schulz-Zunkel et al., 2015; Tockner et al., 2010). These changes include; a
237 decrease in redox potential (E_H), which leads to, for example, reduction of iron (Fe) and manganese
238 (Mn), which in turn can influence the soil pH (Rinklebe and Shaheen, 2017). Other processes affected
239 include sulphur-cycling, changes to the presence of chelating agents such as dissolved organic carbon,
240 mineralisation of POM and suppression of microbial activity (Poot et al., 2007; Puchalski, 2003; Schulz-
241 Zunkel et al., 2015; Schulz-Zunkel and Krueger, 2009). Ibragimow, Walna, and Siepak (2013) showed,
242 through analyses of fluvial samples before and after a flood, that the physicochemical properties (grain
243 size, E_H , pH, POM, and calcium carbonate contents) as well as the total and available concentration of
244 PTEs had changed. Harvey et al. (2019) found that after UK floods receded in the winter of 2013-14
245 there was a decrease in the soil bulk density, pH and available P. The flood was found to have had a
246 negative effect on the overlying vegetation and caused a shift in the microbial community structure.

247 Inundation during a flooding event can carry PTEs dissolved in rising groundwater and
248 potentially contaminated suspended sediment from upstream overbanking water, depositing this
249 onto the floodplain during a flood (Acreman et al., 2003; Bednářová et al., 2015; Du Laing et al., 2009;
250 Gröngröft et al., 2005; Junk et al., 1989; Rudiš et al., 2009; Tockner and Stanford, 2002; Weber et al.,
251 2009). Subsequently, this deposition of suspended riverine sediments/POM by flood water results in
252 the floodplain topsoil becoming a sink for PTEs (Du Laing et al., 2009; Frohne et al., 2011;
253 Nshimiyimana et al., 2014; Overesch et al., 2007; Rinklebe et al., 2007; Visser et al., 2012; Zhao and

254 Marriott, 2013). As a result, floodplain topsoil (uppermost 15cm) can often initially contain elevated
255 concentrations of PTEs such as the metalloid; arsenic (As), and metals; chromium (Cr), copper (Cu),
256 lead (Pb), and zinc (Zn), but later due to post-depositional reactions with organic matter/other organic
257 components the PTEs concentrations will vary (Adekanmbi et al., 2020; Ciszewski and Grygar, 2016;
258 Hurley et al., 2017; Izquierdo et al., 2013; Jiao et al., 2014; Kelly et al., 2020). When laboratory
259 experiments are undertaken on samples gathered from floodplain site, soils are collected as single or
260 composite samples, air or oven dried and then homogenised, resulting in a loss of soil stratigraphy
261 and therefore the potential differences in PTEs concentration with depth may be unaccounted for
262 (Ciszewski and Grygar, 2016). Zhao and Marriott, (2013) looked at PTEs concentrations along a vertical
263 profile and found that there were peak values at varying depths; affected by translocation and
264 duration of inundation. The process of breaking up of soil samples for laboratory experiments will
265 make interpretation of PTEs levels difficult. Kelly et al. (2020) took intact soil cores to overcome this
266 and more closely reflect natural samples, they too found the duration of inundation influenced the
267 fate of PTEs.

268 The biological health of floodplain soils is important as they act as an interface between terrestrial
269 and aquatic environments, therefore playing an important role in maintaining the environmental
270 quality of surface waters (Izquierdo et al., 2013; Stuart and Lapworth, 2011). Artificial or constructed
271 wetlands have been used for flood and pollution control; storing and filtering excess water to protect
272 rivers from various kinds of runoff e.g. high nutrient loads from farm land (Blackwell and Pilgrim, 2011;
273 Ellis et al., 2003; Rizzo et al., 2018). An example of this technology is demonstrated at the Rothamsted
274 Research North Wyke experimental farm (Pulley and Collins, 2019). Even if river and groundwater
275 water quality improves due to the implementation of more stringent environmental policy,
276 contaminated floodplains remain as a legacy of historic upstream pollution (Bradley and Cox, 1990;
277 Förstner, 2004; Kowalik et al., 2004). With increased frequency and duration of flooding, there is the
278 possibility that changes to soil properties and biogeochemical processes will ultimately lead to the
279 mobilisation of PTEs from floodplain soils (Ciszewski and Grygar, 2016; González-Alcaraz and van

280 Gestel, 2015). Therefore, historically contaminated floodplains may become a source of legacy
281 pollution to the surrounding environment (Kelly et al., 2020; Pulchalski, 2003; Schulz-Zunkel and
282 Krueger, 2009), as shown in Figure 1.



283 Figure 1: How floodplains may switch from being a sink of pollution to becoming a source of legacy
 284 pollution: A) PTEs contaminated river sediment (red) due to industry in the catchment upstream, B)
 285 heavy rainfall influences the receiving catchment (increased river flow and groundwater level), resulting
 286 in flooding and the deposition of contaminated sediment onto the adjacent floodplain; dissolved
 287 contaminants may also reach the floodplain surface via rising groundwater, C) Later, the river is
 288 uncontaminated (brown) due to rising environmental quality standards, with legacy of PTEs
 289 contamination (red) in the river bank and floodplain soil, D) heavy rainfall results in flooding of the
 290 contaminated floodplain, mobilisation of the legacy PTEs by desorption and resuspended particulate
 291 matter into the surrounding environment and thus making them potentially available for uptake by

292 *vegetation and soil organisms, as well as pollutant transfer leaching into the overlying flood water, the*
293 *groundwater, and ultimately the river. Created with BioRender.com.*
294

295 2.Impact of flooding on the mobility of potentially toxic elements in floodplain soil

296 2.1 PTEs in floodplain soil

297 Several PTEs are also essential nutrients that are required in low concentrations for healthy
298 functioning and reproduction of microorganisms, plants, and animals, although may become toxic in
299 high concentrations, these include; Cu, Cobalt (Co), Nickel (Ni), Vanadium (V), Zn, chlorine (Cl), Mn,
300 Fe, boron (B), and molybdenum (Mo) (Adamo et al., 2014; Hooda, 2010; Wyszowska et al., 2013).
301 Other PTEs are non-essential and can cause toxicity even when they are found at low concentrations,
302 these include; As, Pb,) and mercury (Hg); (Adamo et al., 2014; Nriagu et al., 2007; Wuana et al., 2011;
303 Wyszowska et al., 2013). Cadmium (Cd) is generally considered a non-essential element to soil
304 organisms, but it has been found to be beneficial to some microalgae (Xu et al., 2008) Chromium can
305 be considered a micronutrient but its toxicity depends on its valence state (i.e. Cr (VI) is the more
306 mobile and toxic form compared with Cr (III)). Redox potential therefore not only affects the mobility
307 of PTEs, but also their toxicity (Lee et al., 2005; Shahid et al., 2017). The consequences of PTEs
308 contamination of soils are rarely observed with immediate effect, rather they tend to cause delayed
309 adverse ecological changes, due to the fact that PTEs are persistent in the environment for long
310 periods, non-biodegradable and can only be bio-transformed through complex physico-chemical and
311 biological processes (Chrzan, 2016; Czech et al., 2014; Hooda, 2010). PTEs cause adverse ecological
312 effects on plants and organisms such as impacting their activity, growth rate/yield, metabolism and
313 reproduction, causing symptoms of physiological stress and potentially death. The extent of the
314 adverse effect depends on the exposure route (ingestion, dermal absorption or uptake of pore water)
315 and time, resistance (related to residence time of the PTEs in the environment) and detoxification
316 mechanisms of the plant or animal (Alloway, 2013; Eggleton and Thomas, 2004; Ehlers and Loibner,
317 2006; Hooda, 2010; Pan et al., 2018; Shahid et al., 2017; Winger et al., 1998). Leaching of PTEs from

318 the floodplain soil into the groundwater or river will also cause adverse effects to aquatic organism in
319 these environments (Zia et al., 2018).

320 PTEs are either present naturally in the floodplain soil from the underlying or upstream
321 geology and subsequent geogenic processes (e.g. weathering of parent material, emissions from
322 volcanoes, forest fires) or introduced by anthropogenic sources, including solid and dissolved inputs
323 from; aerial deposition, transport emissions, industrial, municipal and diffuse runoff from agricultural
324 practices landfills and sewage treatment facilities (Alloway, 1995; Álvarez-Ayuso et al., 2012). PTEs can
325 be adsorbed to colloidal suspended particulate material, transported in the river water and
326 accumulate in the floodplain soil during inundation (Du Laing et al., 2009; Frohne et al., 2011;
327 Peijnenburg et al., 2007; Rinklebe et al., 2007). PTEs have been found to be primarily associated with
328 fine-grained clay or silt minerals and can reside in the floodplain for longer when compared with river
329 sediments, as they are less likely to be susceptible to erosion (Lučić et al., 2019; Malmon et al., 2002).
330 Contamination of the floodplain soil may result from a point source such as a sewage treatment
331 facility, or from diffuse sources that have no specific point of discharge (e.g., agricultural applications).
332 Impacts of diffuse pollution are difficult to predict as they can be affected by weather systems,
333 meaning soils far from the source may be affected (Gregory et al., 2015; Neal et al., 1996). The
334 anticipated changes to intense rainfall may result in increased delivery of diffuse pollution to rivers
335 and groundwater (Arnell et al., 2015; Foulds et al., 2014), particularly as contaminated floodplain soils
336 may become a diffuse source of pollution themselves during a flooding event (Schulz-Zunkel and
337 Krueger, 2009).

338 2. 2. Influence of flooding on PTEs mobility

339 During a flooding event, biogeochemical processes occur in the floodplain soil at the oxic-
340 anoxic interface and in the anoxic layers. The kinetics of these processes are of great importance
341 because the location of the oxic-anoxic interface is subject to change due to fluctuating water table
342 levels (Du Laing et al., 2009; Puchalski, 2003). In their review of trace metal behaviour in floodplain

343 sediments, Du Laing et al. (2009) state that the spatial occurrence of processes affecting metal mobility
344 and availability is largely determined by the topography of the floodplain. Remobilisation of PTEs from
345 sediments into the overlying water column during a flooding event depends on the flood regime; the
346 frequency of these intense floods which flush or remobilise contaminated material as well as the
347 duration or alternation of flood with dry spells (Arnell et al., 2015; Foulds et al., 2014; González-Alcaraz
348 and van Gestel, 2015). Whilst research has suggested that the longer the flood duration, the greater
349 the metal mobility (Shaheen et al., 2014a, 2014b), Stafford et al. (2018) suggest that even short
350 periods of soil saturation can have an influence the solubility of PTEs.


351 There are conflicting results in the literature regarding the effect of flooding on the mobility
352 of PTEs, expressed by changes in PTEs concentration (increase or decrease), in floodplain soils (Table
353 S1). This may largely be the result of different site-specific conditions (e.g. soil pH, texture, mineralogy)
354 or different laboratory set-ups (e.g. submerging soils in deionised water, or the use of inert gas to
355 simulate the anoxic conditions of a flood), illustrating the complexity of the processes involved in
356 mediating PTEs mobility in floodplain soils (Abgottspon et al., 2015; Du Laing et al., 2007; Frohne et
357 al., 2011; Schulz-Zunkel et al., 2015). Many of the considerations in the literature are founded on
358 research of soils or sediments in microcosm experiments, which often involves homogenising the soil
359 samples, resulting in loss of natural soil structure, loss of roots and biota, short-exposure time to flood
360 conditions, and the control of variable factors such as temperature and soil water conditions (Frohne
361 et al., 2011; Rinklebe et al., 2010). Redox conditions are often simulated and controlled through
362 additions of O₂, to increase E_H, and N₂, to lower E_H (Frohne et al., 2014, 2011; Schulz-Zunkel et al.,
363 2015; Shaheen et al., 2016; Shaheen and Rinklebe, 2017). These differences make extrapolation of
364 these laboratory-based findings to field situations difficult (Hooda, 2010).

365 A key factor in determining the fate and transport of PTEs is their chemical form which, in
366 combination with environmental factors, can influence their mobility in the soil. The chemical form of
367 an element is often referred to as its “speciation”, “oxidation state”, or “valence” (Rodgers et al., 2015;

368 Wuana et al., 2011). There are important redox sensitive PTEs for which the oxidation state has a large
369 influence on solubility and mobility. For example, Cr(VI) is more mobile than Cr(III), but As(V) is less
370 mobile than As(III) (Frohne et al., 2015; Rinklebe et al., 2016; Schulz-Zunkel et al., 2015; Shaheen et
371 al., 2014b; Yang et al., 2015). Speciation of PTEs within the environment has a distinct influence upon
372 their behaviour; specifically, reactivity, toxicity, mobility and bioavailability within the floodplain (Du
373 Laing et al., 2009; Gambrell, 1994; Hooda, 2010; Rodgers et al., 2015). This understanding is important
374 for predicting the environmental impact of contaminated soils, although we are only beginning to
375 converge on consensus on how bioavailability or speciation soil tests can help with risk-assessments,
376 while this is slowly introduced into legislation (Cipullo et al., 2018; Naidu et al., 2015, 2008; Ng et al.,
377 2015).

378 2. 3. Changes to soil physical and chemical properties that influences PTEs mobility

379 Potentially toxic elements present in soils are often adsorbed to or protected within
380 aggregates that are stabilised by organic matter. During a flooding event, these particles may be
381 leached through the soil profile, or suspended in flood waters where they may be redistributed across
382 floodplain soils, or be carried downstream by the river, potentially contributing to river pollution of
383 the contamination of downstream floodplains. The solubility and therefore mobility of PTEs from the
384 soils to the surrounding environment depends largely on the intrinsic soil physical and chemical
385 properties (texture, availability of soil particulate surfaces and dissolved organic matter, salinity and
386 the presence of Fe/Mn oxides, carbonates, phosphates and sulphides) and a range of variables that
387 are directly affected by periodic inundation of the floodplain, including; soil pH, redox potential (E_H),
388 dissolved organic carbon (DOC) and the valance of individual PTEs (Adewuyi and Osobamiro, 2016;
389 Dawson et al., 2010; Du Laing et al., 2009; Frohne et al., 2015; González-Alcaraz and van Gestel, 2015;
390 Lee et al., 2005; Puchalski, 2003; Rinklebe and Du Laing, 2011; Schulz-Zunkel and Krueger, 2009;
391 Shaheen et al., 2016; Shaheen and Rinklebe, 2014; Steinnes, 2013). A conceptual model (Figure 2) has
392 been produced based on our literature review (Table S1) as a way of visualising the various factors
393 and processes influencing the solubility of PTEs in a floodplain soil.



401 **Figure 2: Conceptual model depicting the key processes influencing the solubility of Potentially Toxic**
 402 **Elements (PTEs) after a soil becomes flooded.**

403 *a) Oxygen is rapidly consumed by microbial and root respiration, decreasing the redox potential (E_H).*
 404 *b) Decreasing E_H can lead to redox sensitive elements (e.g. As and Cr) changing valence state, directly*
 405 *affecting solubility. c) Greater soil moisture brings dissolved organic matter (DOM) into solution. d)*
 406 *Reducing conditions (lower E_H) leads to the release of more DOM. e) Lower E_H results in the reduction*
 407 *of Fe and Mn, consuming protons (H^+) and increasing pH. f) an increase in pH often results in the release*
 408 *of more DOM. g) DOM acts as a chelating agent, forming soluble organo-metal complexes with PTEs*
 409 *desorbed from soil surfaces. h) as pH increases metal cations (e.g. Cu, Pb, Zn) are adsorbed on pH-*
 410 *dependent adsorption sites of particulate matter. i) as pH increases, anions and oxy-anions (e.g. As)*
 411 *are desorbed from pH-dependent adsorption sites. j) Dissolution of reducible Fe and Mn oxides is*
 412 *facilitated by increasing pH. k) Microbial reduction of Mn and Fe oxides increases their solubility and*
 413 *can cause reductive dissolution of co-precipitated PTEs. l) An increase in pH facilitates the precipitation*
 414 *of insoluble metal sulphides. m) Microbial reduction of sulphate results in the precipitation of metal*
 415 *sulphides. n) Release of adsorbed PTEs from soil surfaces increases PTEs solubility. o) Immobilisation*
 416 *of PTEs through adsorption processes reduces PTEs solubility. p) Reductive dissolution of PTEs*
 417 *associated with Fe and Mn oxides increases PTE solubility. q) Precipitation of PTEs as metal sulphides*
 418 *decreases PTEs solubility.*

419

420 Soil physical, chemical and biological processes determine the mobility and redistribution of

421 PTEs (Hooda, 2010). These processes include; sorption, desorption, dissolution and precipitation

422 (Puchalski, 2003; Wijngaard et al., 2017). Subsequently, PTEs are redistributed into different

423 geochemical fractions, associated with other soluble species, released from the soil matrix into the

424 soil solution or porewater, and transferred through the ecosystem and food web to other terrestrial

425 or riparian areas downstream from the floodplain; thus potentially becoming a risk to human and

426 environmental health (Adamo et al., 2014; Adewuyi and Osobamiro, 2016; Baran and Tarnawski, 2015;

427 Dang et al., 2002; Du Laing et al., 2009; Rinklebe et al., 2016; Schulz-Zunkel et al., 2015; Shaheen et
428 al., 2014a, 2014b; Sizmur et al., 2011). Sorption processes that control PTEs mobility and bioavailability
429 in soil are affected by the soil pH, redox and their interactions with other ions and substances present
430 in soil solution (Antoniadis et al., 2018; Frohne et al., 2011; Ostergren et al., 2000; Violante, 2013).

431 Sorption processes are influenced by the changing conditions that flooding brings, particularly
432 with regards to soil moisture content, temperature and redox potential. The mobility of PTEs in
433 flooded soils is closely related to changes in redox potential which, in turn, is altered by flooding. This
434 can have direct impacts on the mobility of redox sensitive PTEs (e.g. As and Cr). Inundation of soils
435 with floodwater may indirectly affect PTEs mobility and speciation because it also influences, the
436 population, community composition, and behaviour of invertebrates inhabiting the floodplain which,
437 in turn, influence the mobility of PTEs through their burrowing and bioturbation behaviour. For
438 example, earthworms are known to increase the mobility of PTEs due to passage through the
439 earthworm gut (Sizmur et al., 2011; Sizmur and Richardson, 2020) and their populations are suppressed
440 by flooding events (Plum and Filser, 2008; Kiss et al., in review). Bioturbation/bioirrigation behaviour
441 by chironomid larvae has been found to increase oxygen uptake at the soil/sediment-water interface,
442 promoting POM decomposition that results in the release of dissolved organic matter and subsequent
443 release of PTEs (He et al., 2019). Furthermore, the reduction of Mn and Fe can cause reductive
444 dissolution of co-precipitated PTEs, and an increase in pH facilitates the precipitation of PTEs as
445 insoluble sulphides. The mobility of PTEs can therefore increase or decrease due to the net effect of
446 these processes (Figure 2). Which process dominates will depend primarily on the mineralogy of the
447 soil.

448 The following sub-sections will explain how key soil physical and chemical properties are
449 affected by flooding and how this influences PTEs mobility, followed by a discussion on the role of soil
450 organisms and plants in mediating PTEs mobility in floodplain soils. Attention will be given to how each
451 of these factors influence each other to distinguish direct and indirect impacts on PTEs mobility.

452 *2.3.1 Soil texture and related properties*

453 Soil texture is a stable property that refers to the physical composition of mineral fragments;
454 sand, silt and clay and varies due to differences in underlying or upstream geology. The texture and
455 related clay mineralogy reflect the particle/pore size distribution and overall soil surface area
456 (Amacher et al., 1986) which, in turn, affects the soils' water holding capacity (WHC); the maximum
457 quantity of water a soil can potentially contain, also known as the field capacity (Stürck et al., 2014).
458 Therefore, soil physical properties play a role in flood duration because they determine the soils'
459 ability to receive (via infiltration) and drain water during a rainfall event (Rinklebe et al., 2007). Clayey
460 soils are likely to be saturated for longer than freely draining sandy soils (Sherene, 2010). Soil hydraulic
461 (water retention and hydraulic conductivity curve) as well as thermal properties (thermal conductivity
462 and heat capacity) affect the hydrothermal regime of the soil. Together these properties determine
463 the ease in which water, and dissolved PTEs, moves through the soil pore continuum, how much water
464 can be stored in the pore volume, and how soil temperature varies with depth. These properties are
465 strongly dependent on soil texture, pore size distribution and mineralogy (Hillel, 1998; Tack et al.,
466 2006; Thomas et al., 2016). Soil temperature affects the flow of water through the soil due to changes
467 in viscosity and hence affects infiltration calculations (Gao and Shao, 2015; Prunty and Bell, 2005), so
468 this is often corrected for when reporting hydraulic conductivity data (Thomas et al., 2016).

469 PTEs must be in the soluble phase or associated with colloids to be transported through the
470 soil. The soil properties will play a part in the movement of PTEs into and out of the soil solution. Clay
471 minerals and organic matter compounds have a large number of binding sites, so act as adsorption
472 surfaces for PTEs in soils. The type of clay mineral present (kaolinite, illite, montmorillonite etc.) will
473 also affect the specific surface area (Meegoda and Martin, 2019; Tack et al., 2006). As a result, soils
474 with high clay and silt (fine fractions) tend to retain higher amounts of PTEs, compared to course
475 textured sandy soils (Sherene, 2010; Zhao et al., 1999).

476 2.3.2 *Organic matter*

477 Soil POM, along with the surfaces of clay particles and Fe and Al oxides, acts as a binding phase
478 for PTEs due to the attraction of positively charged cations to negatively charged surfaces (Evans,
479 1989). Thus, dissolved organic matter raises the cation exchange capacity (CEC) of a soil, and is thus
480 considered to be an important factor controlling PTEs distribution and mobility in floodplain soils and
481 sediments (Baran and Tarnawski, 2015; Bufflap and Allen, 1995; Du Laing et al., 2009; Ehlers and
482 Loibner, 2006). The mechanisms that bind the PTEs with particulate and dissolved organic matter
483 include adsorption, complexation and chelation (Alvim Ferraz and Lourenço, 2000; He et al., 2019;
484 Selinus et al., 2005). Floodplains are subject to changing water table levels and occasional inundation
485 that brings about associated changes in redox conditions. This can result microbially mediated soil
486 POM degradation, either during prolonged periods of flooding or in the subsequent oxidising
487 conditions when the flood recedes, which releases organically bound PTEs, such as As, Cu, Co, Cr, Ni,
488 Pb, and Zn from the soil into the soil solution (Adewuyi and Osobamiro, 2016; Alvim Ferraz and
489 Lourenço, 2000; Dang et al., 2002; Kalbitz and Wennrich, 1998; Koretsky et al., 2007; Rinklebe and Du
490 Laing, 2011). Therefore, the extent to which flooding of soils results in the mobilisation of PTEs into
491 solution is mediated by the proportion of the PTEs that are associated with soil POM, and the
492 susceptibility of this organic matter to degradation (as a result of microbial activity (Fe(III) and Mn(IV)-
493 reducing micro-organisms) under reducing conditions. The free ions that are then in solution are highly
494 reactive with the solid phase and are thought to be a major determinant of bioavailability and causing
495 the most significant biological effects (Bufflap and Allen, 1995; Dang et al., 2002; Dawson et al., 2010;
496 Degryse et al., 2009; Lloyd, 2003).

497 2.3.3 *Salinity*

498 Salinity is proportional to the conductivity of a sample solution; which is a measure of its ability
499 to conduct or carry electric current and depends on the presence of charged ion species (anions and
500 cations) (Ander et al., 2016; de Souza Machado et al., 2018; De Vivo et al., 2008)). Increasing salinity
501 in flood water is associated with an increase in major cations that compete with PTEs for sorption

502 sites. This competition promotes PTEs desorption from the floodplain soil in the absence of sulphides
503 and hence increases total PTEs concentrations in the soil porewater (Rinklebe and Du Laing, 2011).
504 The presence of Ca-salts releases more PTEs into the soil solution compared with Na-salts that are less
505 competitive for sorption (Du Laing et al., 2009; Hahne and Kroontje, 1973).

506 Changes in salinity may affect the soil physical properties and result in a destabilisation of the
507 soil structure (Gregory et al., 2015). The salinity of the water causes a neutralisation of negatively
508 charged clay particles, followed by flocculation (particles attaching together) which increases the
509 deposition of sediments (along with the PTEs adsorbed to them) onto the floodplain. This process
510 results in the floodplain becoming a sink for PTEs (Rinklebe and Du Laing, 2011). An extended flood
511 duration, particularly when accompanied by low flow-rates (including stagnant water), results in
512 sedimentation of fine grain sediment and organic matter that may have PTEs bound (Ciszewski and
513 Grygar, 2016; Du Laing et al., 2009; Shaheen and Rinklebe, 2014).

514 2.3.4 Redox potential (E_H)

515 Waterlogging of soils generally results in a reduction in oxygen availability due to rapid
516 consumption of oxygen by soil microbial activity and root respiration (Du Laing et al., 2007; Rinklebe
517 and Du Laing, 2011) and because the dissolution of oxygen through water is many times slower than
518 through air (Alloway, 1995; Du Laing et al., 2009; Frohne et al., 2015; Schulz-Zunkel et al., 2015). The
519 soil microbial community (e.g. bacterial species such as *Thiobacillus ferrooxidans*, *Thiobacillus*
520 *thiooxidans* and *Leptospirillum ferrooxidans*) then uses alternative electron acceptors (such as nitrate,
521 sulphate and Fe/Mn oxides), in anaerobic respiration, which results in a decrease in redox potential
522 (E_H) (Maluckov, 2017) as the floodplain soils change from oxic ($[O_2] > 30 \mu\text{mol L}^{-1}$) to anoxic
523 ($[O_2] < 14 \mu\text{mol L}^{-1}$) conditions (Bellanger et al., 2004). Associated alkalinity generation drives increases
524 in soil pH, a change which can be observed after a few days (Du Laing et al., 2007; Johnston et al.,
525 2014; Karimian et al., 2017). Soil temperature has been found to dictate the rate and type of redox
526 reactions; with soils at low temperatures (1 - 4 °C) requiring greater durations of saturation (20 days)

527 before the onset of reducing conditions were seen, whereas soils at higher temperatures (above 9 °C)
528 only required 2 days of saturation (Vaughan et al., 2009).

529 Redox potential has important effects on the speciation of As, Cu and Cr, as well as N, S, Fe,
530 Mn, because these elements can exist in soils in more than one oxidation state (Selinus et al., 2005)
531 and solubility depends on oxidation state. Copper solubility decreases after reduction from Cu (II) to
532 Cu (I) under anaerobic conditions and the presence of electron donors (Fe (II)) and bacteria. However,
533 other PTEs such as Cd and Zn change valence state as a consequence of redox dependent pH changes,
534 complexation with organic matter or precipitation with Fe and Mn (hydr)oxides or sulphides (Du Laing
535 et al., 2009; Frohne et al., 2011).) Shaheen et al. (2014a) demonstrated that sufficient time is needed
536 for transformations between valence states to take place. For example, the oxidation of Cr from Cr
537 (III) to the highly mobile Cr (IV) form was found to be a slow process. This means that with shorter
538 flooding duration and quicker cycling between oxic and anoxic conditions, Cr mobility may be difficult
539 to predict.

540 The presence of variable charge minerals, such as Fe and Mn oxides, phosphates, carbonates
541 and sulphides provide a reaction surface for sorption processes, allowing PTEs to bind and become
542 immobilised (Antoniadis et al., 2018; De Jonge et al., 2012; Sipos et al., 2014; Violante, 2013). Reducing
543 conditions change the oxidation state of Fe and Mn, increase their solubility and may have indirect
544 effects (known as reductive dissolution) on the mobility of associated metal cations (e.g. As, Cd, Cu,
545 Ni, Pb, and Zn), releasing them from the solid phase to pore waters, depending on flood duration
546 (Abgottspon et al., 2015; Ciszewski and Grygar, 2016; Du Laing et al., 2009; Frohne et al., 2011;
547 Karimian et al., 2017; Rinklebe and Du Laing, 2011; Schulz-Zunkel et al., 2015; Shaheen et al., 2016,
548 2014b; Vaughan et al., 2009). Redox processes are a key factor for the reductive dissolution of Mn and
549 Fe (hydr)oxides, these processes are often catalysed by microorganisms and result in the release of
550 PTEs from the sediment (Du Laing et al., 2009; Frohne et al., 2011; Stafford et al., 2018; Yang et al.,
551 2015). Relatively insoluble Fe(III) and Mn(IV) prevail under aerobic soil conditions providing sorption

552 surfaces for many metals, whereas under anaerobic conditions Mn(IV) and Fe(III) are reduced to more
553 soluble forms (Mn(II) and Fe(II)) with consequential dissolution of Mn and Fe hydrous oxides, co-
554 sorbed PTEs ions (e.g. As, Cd, Cr, Ni and Pb), are released into soil solution (Simmler et al., 2017;
555 Stafford et al., 2018; Yang et al., 2015). After inundation, Fe and Mn may re-precipitate as oxides and
556 can bind (by desorption or co-precipitation) the trace metals back into the solid state (Ciszewski and
557 Grygar, 2016; Davranche et al., 2011; Du Laing et al., 2009).

558 Decreasing of E_H can initiate microbial sulphate reduction and this can reduce the mobility of
559 some PTEs (e.g. As, Cd, Cu, Cr, Ni and Pb) through coprecipitation of metal cations with sulphides
560 (Abgottspon et al., 2015; Borch et al., 2010; Weber et al., 2009), although many of these minerals are
561 metastable and so prone to change (Karimian et al., 2018). Yang et al., (2015) put mixed sediment
562 samples into a laboratory culture tanks and found that microbially induced release of sulphur with
563 subsequent As precipitation was more important for controlling As adsorption/desorption than
564 reductive dissolution of Fe/Mn oxides. As the flood recedes, the floodplain soils undergo drying and
565 aeration that change the conditions from anoxic back to oxic. The now oxic environment causes
566 sulphides to be oxidised, which then releases PTEs back into the pore waters (Abgottspon et al., 2015;
567 Du Laing et al., 2007; Frohne et al., 2011). In addition to this, when exposed to oxygen and water,
568 sulphides are oxidised to sulphates which leads to the formation of sulphuric acid thereby causing a
569 decrease in pH and release of the PTEs (Emerson et al., 2017; Forstner and Wattman, 1981). Frohne
570 et al. (2011) suggested that the mobility of Cd, Cu, Mn, Ni and Zn under oxidising conditions could be
571 attributed to dissolution of sulphides and the resulting release of those metals. The extent to which
572 the mineralogy of a floodplain soil is dominated by Fe/Mn oxides or sulphates may dictate whether
573 PTEs are mobilised or immobilised during inundation, and the extent to which this phenomenon is
574 reversed after floodwater recedes.

575 2.3.5 Soil pH

576 pH is a measure of the hydrogen ion concentration and can also be referred to as the degree
577 of acidity or alkalinity. The soil pH is affected by flooding because of a well-established correlation
578 between soil pH and changing redox conditions; as a soil becomes flooded, this creates reducing
579 conditions where (H^+ ions) are consumed (for example due to reduction of Fe and Mn oxides) and the
580 pH increases (Rinklebe and Shaheen, 2017; Weber et al., 2009). When the flood recedes, oxidation
581 processes produce protons and decrease the pH (Adewuyi and Osobamiro, 2016; Frohne et al., 2015,
582 2011; Rinklebe and Shaheen, 2017; Shaheen and Rinklebe, 2017). Furthermore, on exposure to the
583 atmosphere, when flooding recedes, dissolved organic carbon (DOC) is converted to CO_2 , which
584 dissolves into porewater as carbonic acid, subsequently further reducing the soil pH (Peacock et al.,
585 2015). However, this negative correlation between E_H and pH hasn't always been observed (Du Laing
586 et al., 2009; Frohne et al., 2015). This is because the degradation of POM such as plant residues, by
587 soil microbes, may increase the soil pH due to ammonification of the residue N (Xu et al., 2006).

588 As the pH changes, processes such as precipitation, co-precipitation and sorption/desorption
589 of PTEs from organic matter or clay minerals occur, altering the chemical composition as well as
590 reaction rates (Frohne et al., 2011). The soil pH plays an important role in mediating the mobility of
591 PTEs and their availability for plant uptake, as the protons compete with metal cations for exchange
592 sites on the surface of soils. Some of these exchange sites, particularly those associated with soil
593 organic matter, are pH-dependent and thus only become deprotonated at high pH. A decrease in pH
594 is generally accompanied by an increase in the mobility of most PTEs that are metal cations (e.g. Cd^{2+} ,
595 Cu^{2+} , Co^{2+} , Ni^{2+} , Pb^{2+} and Zn^{2+}) (Gröngröft et al., 2005; Sherene, 2010). Thus, as pH increases there is a
596 subsequent decrease in the mobility of these PTEs (Giacalone et al., 2005). The extent to which PTEs
597 mobility decreases in soils during flooding, due to a redox-induced increase in pH, is likely to depend
598 on the proportion of PTEs in the soil that are associated with pH-dependent exchange sites) which are
599 typically associated with soil organic matter) and the pH of the soil prior to the flooding event.

600 2.3.6 *Dissolved organic matter (DOM)*

601 The increase in pH of soil solutions with lower redox potential (reducing conditions) is often
602 accompanied by a release of dissolved organic matter (DOM) and the subsequent formation of soluble
603 organo-metal complexes (Abgottspon et al., 2015; Alvim Ferraz and Lourenço, 2000; Frohne et al.,
604 2011). The presence of DOM in floodplain soils acts as a chelating agent which has a strong binding
605 ability and increases the mobility of PTEs into pore waters and subsequently into river water or
606 groundwater (Dawson et al., 2010; Du Laing et al., 2009; Shaheen et al., 2014b). The greater the
607 concentration of DOM in porewater, the more PTEs that are held in solution, and (to maintain an
608 equilibrium) the more PTEs that desorb from the surfaces of the soil to replenish the free ion
609 concentrations in the porewater, thus increasing PTEs mobility. Greater concentrations of DOM have
610 been observed with decreasing E_H , which may be due to suppressed microbial carbon consumption
611 under anoxic conditions (Frohne et al., 2015). Shaheen et al. (2014a) highlighted that increases in DOM
612 associated with lower E_H may help to catalyse changes in the valence state of PTEs; for example, of
613 Chromium (III) to (VI).

614 2.3.7 *Temperature*

615 As temperatures are predicted to increase as a result of climate change, they may become a
616 factor that contributes to greater release of PTEs from the soil during a flood (Visser et al., 2012). Soils
617 are affected by variations in air temperature which, in turn, affects the rate of biogeochemical
618 processes during a flooding event, including decreasing redox potential and, ultimately, influences the
619 rate and extent to which PTEs are released/ desorbed from POM into surface water and groundwater
620 (Arnell et al., 2015; González-Alcaraz and van Gestel, 2015; Sánchez-Rodríguez et al., 2019; Shaheen
621 et al., 2016; Stahl et al., 2013). Increases in temperature raise the ion activity in soil solution, and also
622 make plants more active, which may lead to greater plant root uptake of soil water and
623 dissolved/labile PTEs within this water (Sherene, 2010). Arsenic release from flooded soils was found
624 to have temperature dependence, with As solubilisation increasing as temperature increased
625 (Simmler et al., 2017; Visser et al., 2012). Temperature increases are attributed to a decrease in the

626 water viscosity resulting in dissociation of molecules and a subsequent increase in the number of ions
627 in the solution. For every degree Celsius increase in temperature there is an observed increase in
628 electrical conductivity of 1.9% (Ander et al., 2016; Ma et al., 2011).

629 Many of the processes (e.g. redox reactions) described in the previous sections are microbially
630 mediated and temperature dependent, and so the extent to which they affect the mobility of PTEs
631 depends on their kinetics and the duration that floodplain soils are inundated. Changes in soil pH and
632 DOM have been shown to exert a greater influence than E_H on the mobility of PTEs when considering
633 shorter flood-dry cycles (Shaheen et al., 2014b, 2014a). However, Dang, Liu, and Haigh (2002) found
634 that with increasing flood duration, more trace elements were transformed from inert phase to
635 exchangeable fractions, increasing mobilisation. Soil redox processes are important for protecting
636 environmental health; however, the kinetics and mechanisms remain poorly characterised and
637 understood (Abgottspon et al., 2015; Borch et al., 2010; Pulchalski, 2003). Many of the studies
638 reviewed in this manuscript undertook experiments in the laboratory at temperatures that are higher
639 than the soil temperatures typically found *in-situ* at the location where the soil samples were
640 collected, and so the rates at which reactions occur and the subsequent mobilisation of PTEs may be
641 overestimated in these experiments. It is not possible to verify the extent of the overestimation, so
642 more *in-situ* experiments are needed to investigate and provide quantification of the differences
643 between laboratory and *in-situ* experiments.


644 Groundwater and river temperatures may have a strong effect on floodplain soil temperatures
645 (Andersen, 2018). In warmer seasons they will generally be cooler than soil temperatures, but the
646 opposite occurs in cooler seasons. Also, changes in soil moisture content, as a result of flood events,
647 will affect the soil thermal properties such as thermal conductivity and heat capacity (Lu et al., 2007),
648 thereby also affecting the spatial and temporal variation in the soils' temperature regime. A laboratory
649 microcosm experiment with mining-contaminated topsoil and subsoil samples saturated for up to 41
650 days at temperatures ranging between 10-25°C, found that soil temperature increased the

651 solubilisation of As, particularly in the topsoil when saturated for 1-2 weeks (Simmler et al., 2017).
652 This means, for some PTEs, flooding during warmer seasons may result in greater mobilisation than
653 flooding during cooler seasons. More seasonal field observations are required to understand the
654 subtle interactions and feedbacks between soil moisture, floodwater temperature, and mobility of
655 PTEs.

656 2. 4. Soil biological processes that influence PTEs mobility

657 *2.4.1 Soil organisms*

658 Floodplain soils contain a great diversity of organisms that are known to contribute to the
659 physical structure of the soil/sediment through bioturbation which influences the biogeochemical
660 cycling of PTEs through oxygen diffusion, redox gradient and decomposition of dissolved organic
661 matter (Classen et al., 2015; He et al., 2019; Hooda, 2010; Selinus et al., 2005). As the soil pore spaces
662 are filled with water, oxygen diffusion is low so microbial respiration relies on alternative electron
663 acceptors (e.g. NO_3^- , Mn, Fe and S), resulting in reducing conditions (decreasing E_H) that
664 simultaneously increase pH (Matern and Mansfeldt, 2016), and the changes to PTEs mobility (Figure
665 3) that are described in previous sections. Changes in the chemical speciation of PTEs can also occur
666 due to microbial processes in reducing conditions, for example, sulphate reducing bacteria can
667 methylate Hg in anoxic conditions (Ma et al., 2019).



668

669 *Figure 3: Soil microbial processes during inundation of floodplain soil influences mobility of PTEs; (left-*
 670 *hand side) generic metals with valence state (denoted by M^{n+}) are coprecipitated to Fe oxides and are*
 671 *released due to reductive dissolution and (right-hand side) sulphate reduction (sulphate to sulphide)*
 672 *results in some metal (denoted by MS) precipitation, some of the metal (M^{n+}) remains in the pore water.*
 673 *Created with BioRender.com.*

674

675 PTEs that are present in floodplain soils are often protected within the soils' aggregates, which
 676 are stabilised by POM. However, inundation can stimulate the soil microbial community, which is
 677 sensitive to disturbance, accelerating the refractory organic matter mineralisation and destabilisation
 678 of aggregates, exposing and increasing the mobility of PTEs in the soil (Du Laing et al., 2009; Gall et al.,
 679 2015; González-Macé et al., 2016; He et al., 2019; Rawlins et al., 2013). Tack et al. (2006) found that
 680 the drying of sandy soils caused an increase in soil solution metal concentrations, compared with the
 681 same soils maintained at field capacity. This observation was attributed to microbial effects, increasing
 682 the solubility of dissolved organic matter.

683 Flooding has been found to shift the soil biological community structure and function. These
 684 changes include a reduction of Gram-positive bacteria, mycorrhizal fungi and earthworms found under
 685 flooded conditions (Gregory et al., 2015; Harvey et al., 2019; Unger et al., 2009). Harvey et al., (2019)
 686 found that flooding induced short-term alterations to soil microbial biomass but these changes did

687 not persist in the long term; they concluded that temperate systems may be resilient to winter flood
688 stress. The seasonal timing of floods influences the effect that flooding has on the soil microbial
689 community, and so may result in different effects on, and recovery of, the soil microbial community.
690 Sánchez-Rodríguez et al., (2019) subjected a UK agricultural grassland soil in an intact laboratory
691 microcosm to flooding and found that summertime flooding (25°C), resulted in a loss of actinomycetes
692 and arbuscular mycorrhizal fungi, and that these changes persisted post-flood. They expected
693 microbial biomass to increase with flooding at higher temperatures, due to degradation of vegetation
694 releasing labile carbon. However, they found that maintaining live roots and an active rhizosphere
695 were more important for preserving the microbial community in grassland soils. Earthworms also play
696 a role in increasing the mobility and availability of PTEs in floodplain soil through their activity causing
697 changes to the soil microbial populations, pH, DOC or metal speciation (Sizmur et al., 2011; Sizmur
698 and Hodson, 2009) which in turn influences PTEs mobility as discussed in the above sections.

699 As the PTEs are released into the aqueous phase and mobilised in the environment, they
700 present a potential risk to soil organisms (Ehlers and Loibner, 2006; González-Alcaraz and van Gestel,
701 2015). Soil organisms uptake PTEs via ingestion of polluted soil, food or pore water and/or via dermal
702 uptake or absorption of soil water, with the soil water being the more important of the two pathways
703 (Chrzan, 2016; Hobbelen et al., 2006; Sivakumar and Subbhuraam, 2005). Vijver et al. (2007) found
704 that the frequency of flooding did not result in consistent changes in the internal PTEs concentrations
705 of earthworms. Earthworms accumulate PTEs in their chloragogenous tissue and have a mechanism
706 that allows them to regulate their internal PTEs concentrations, so when they are introduced to
707 contaminated soils the earthworms reach an equilibrium and when they are returned to
708 uncontaminated/"clean" soils they are able to detoxify and eliminate essential metals through
709 excretion (e.g. Cu and Zn), but not non-essential metals (e.g. Cd and Pb) as detoxification processes
710 involve sequestration within an inorganic matrix or organic ligand (Sizmur and Hodson, 2009;
711 Spurgeon and Hopkin, 1999). While microbes can tolerate larger quantities of essential PTEs, in excess
712 both essential and non-essential PTEs (e.g., Al, As, Cd, Hg, Pb, Zn) can adversely affect microbial

713 communities by altering community structure and taxonomic richness; reducing the microbial
714 biomass and lowering their enzyme activity which results in a decrease of soil diversity (Gadd, 2010;
715 Gall et al., 2015; Wuana et al., 2011).

716

717 2.4.2 Plants

718 In many cases, PTEs are concentrated in the upper part of the soil profile where roots reside,
719 meaning that increased mobility is likely to affect plants growing in floodplain soils. Wetland plants
720 growing on inundated floodplain soils can also affect the mobility of PTEs because they are specially
721 adapted to have air-filled tissues, or aerenchyma, which create patches of oxygenated soil around
722 their roots, resulting in an increase in the volume of the oxic/anoxic interface and remobilising PTEs
723 thus increasing their availability (Du Laing et al., 2009; Wright et al., 2017). However, in arable and
724 pasture fields that are generally drier, flooding can cause crops to become stressed, as they are not
725 adapted to wet soils. As oxygen levels decrease there is a build-up of carbon dioxide, methane and
726 nitrogen gases that leads to the roots suffocating and dying (Hippolyte et al., 2012).

727 It is well established that symbiotic fungi, associated with plant roots, regulate the supply of
728 micronutrients and reduce the uptake of non-essential PTEs by plants (Classen et al., 2015; Gadd,
729 2010; Tack, 2010). Plants, such as *Artemisia* and *Phalaris* species, on the floodplain excrete exudates
730 during inundation which stimulates the activity of microbial symbionts in the rhizosphere, allowing
731 PTEs to be taken up into the vegetation (Gall et al., 2015; Sullivan and Gadd, 2019; Violante et al.,
732 2010; Xu et al., 2020). PTEs are often accumulated in plant root tissues and can sometimes be
733 translocated into the plant shoots. However this is regulated in plants by the Casparian strip and
734 therefore limited (Hooda, 2010; Nouri et al., 2009; Shahid et al., 2017). The uptake and accumulation
735 of PTEs is element and plant-specific (Niu et al., 2007; Rinklebe et al., 2016; Tack, 2010; Violante et al.,
736 2010; Xu et al., 2020). The mobilisation and uptake of PTEs by plants may pose a potential
737 environmental risk (Shaheen and Rinklebe, 2014). European floodplains are most commonly used as
738 grassland for grazing cattle or hay production, whereas in other regions e.g. India, they are used for

739 crops like rice, which raises concerns for possible pollutant transfer from the floodplain soil into the
740 surrounding water bodies, then uptake and potential biomagnification of PTEs into the food chain
741 (Martin et al., 2014; Overesch et al., 2007; Tóth et al., 2016a). However, the hyperaccumulation of
742 PTEs by some plants (e.g. sunflower, mustard (Brassicaceae), alfalfa and Ricinus) has resulted in them
743 being considered for phytoremediation of contaminated floodplain soils (Gall et al., 2015; Niu et al.,
744 2007; Nouri et al., 2009; Shaheen et al., 2016; Violante et al., 2010).

745 Factors influencing plant uptake of PTEs include soil pH, electrical conductivity and the total
746 concentrations of PTEs in the soil (Nouri et al., 2009). PTEs uptake also depends on the concentrations
747 in the soil solution, governed by plant exudates and root-induced changes to pH and DOM (Gall et al.,
748 2015). Quantifying the total content of PTEs transferred into the food chain via plants growing on
749 contaminated soil is difficult (Gröngröft et al., 2005). The concentrations of PTEs found in floodplain
750 plants are not always directly reflected in the PTEs content found in the soil, due to both physiological
751 and biochemical differences between different plant species; for example differences in the age of the
752 plant biomass (seasonal trends in growth and therefore uptake of nutrients). Moreover, the rooting
753 depth influences metal mobilisation/immobilisation and element specific uptake into the roots which
754 also affects the transfer into the shoots (Chrzan, 2016; Overesch et al., 2007). Thapa et al. (2016) also
755 demonstrated a change in semi-arid Australian floodplain vegetation productivity in response to
756 flooding and drying cycles; flooding brings nutrients which increases net primary productivity. These
757 changes in vegetation productivity could also initiate structural changes in floodplain vegetation
758 communities in natural and semi-natural ecosystems (Overesch et al., 2007).

759 3. Summary and further research needs

760 3.1. Summary of current understanding

761 Floodplain soils downstream of urban catchments contain elevated concentrations of PTEs as
762 a legacy of human activity and these PTEs could potentially be remobilised by future flooding events.
763 A number of processes occur within the soil, ultimately determining PTEs fate. These processes

764 include: sorption, desorption, complexation, precipitation and dissolution, transport of water and
765 heat, and biological activity. The processes are influenced by the changing conditions that flooding
766 brings particularly with regards to soil moisture content, temperature and redox potential. The
767 mobility of PTEs in flooded soils is closely related to changes in redox potential which, in turn, is altered
768 by flooding. These changes can have direct impacts on the mobility of redox sensitive PTEs (e.g. As
769 and Cr). Furthermore, the reduction of Mn and Fe can cause reductive dissolution of co-precipitated
770 PTEs, but the reduction of sulphate can result in the precipitation of PTEs as insoluble metal sulphides.
771 Which of these processes dominates will depend on the mineralogy of the soil. PTEs precipitated as
772 metal sulphides may oxidise after floodwaters recede and mobilise, accelerated by the pH reduction
773 caused by production of sulfuric acid. There are important interactions between redox potential and
774 other soil properties, such as soil pH, moisture content, POM, DOM, temperature, and salinity which
775 also have a strong impact on PTEs mobility (Vaughan et al., 2009). Many of these reactions are
776 microbially mediated, temperature dependent and the kinetics in real-world scenarios are poorly
777 understood. However, it seems that changes associated with alterations to pH and dissolved organic
778 carbon are relatively fast, while changes to E_H are slower and only become apparent after extended
779 periods of flooding. In many cases, PTEs deposited due to legacy pollution events are concentrated in
780 the upper part of the soil profile, meaning that increased mobility is likely to affect plants growing in
781 floodplain soils and potentially lead to contamination of the surrounding environment, including
782 overlying surface waters.

783

784 3.2 Knowledge gaps and recommendations for future research

785 Floods are dynamic events that expose floodplain soils to water with rapidly changing flows,
786 chemical composition, and sediment load. They can be difficult to predict, due to their different types
787 (e.g. overbanking or groundwater flooding), and the high variation in their magnitude, duration, and
788 frequency of recurrence. Therefore, chemical, physical and biological data from floodplain soils
789 immediately before and immediately after a flooding event are often lacking. However these data


790 would provide the necessary insights into the factors and processes involved in altering the mobility
791 of PTEs during and after a real flooding event (Barber et al., 2017). The effect of flooding on PTEs
792 mobility can be difficult to predict due to there being several factors (e.g. speciation, release through
793 biological degradation and competitive action of other ions) or interactions between factors (e.g.
794 changes in E_H caused degradation of POM) influencing PTEs mobility (Tack and Verloo, 1995).

795 Contamination of soil with PTEs receives most attention in highly contaminated urban,
796 industrial, mining and waste disposal sites (Adamo et al., 2014; Resongles et al., 2015; Simmler et al.,
797 2017; Wuana et al., 2011) with relatively little attention given to more 'typical' floodplains
798 downstream of catchments with a history of urban and industrial development. Much of the work
799 conducted to date (see Table S1) has been undertaken in Europe, America, Canada, China, Indonesia,
800 Australia and New Zealand. Just over half of the studies cited in Table S1 were undertaken in Germany
801 and Belgium (52%), with a particular research effort around the River Elbe and Wupper River in
802 Germany (Du Laing et al., 2009; Förstner, 2004; Frohne et al., 2011; Overesch et al., 2007; Rennert et
803 al., 2017; Rinklebe et al., 2013; Shaheen et al., 2017). However, research examining the relationship
804 between PTEs mobility and flooding in other parts of the world that are expected to see an increase
805 in the frequency and magnitude of flooding events, for example in Asia, Africa and India, is limited.

806 A number of factors were identified that contribute to whether the mobility of PTEs will
807 increase or decrease during inundation of a floodplain, which may be interconnected or work in
808 combination to affect PTEs mobility. As a result, different soils with differing mineralogy and thus
809 different biogeochemical and physical properties, will likely respond differently to flooding. Individual
810 studies tend to focus on one floodplain site. However, knowledge based on one river catchment may
811 not be particularly useful for predicting the impacts of flooding at another site with different
812 mineralogy and physical and chemical characteristics. A more fundamental mechanistic understanding
813 is required to inform the development of predictive models. Therefore, more coordinated work
814 encompassing multiple contrasting sites is required to understand the relative importance of key soil

815 properties (e.g. mineralogy, POM, soil pH, texture; and how these affect derived soil properties such
816 as hydraulic and thermal soil properties) on influencing the impact of flooding on the mobility of PTEs.

817 Many of the findings in the literature are based on research of soils or sediments in laboratory-
818 based artificial flooding environments (Figure 4), which often involve; homogenisation of samples and
819 removal of plant roots, short-exposure time for soil microorganisms and incubation under controlled
820 conditions, such as temperature (often higher than *in-situ* temperatures) and soil water conditions
821 (often wetting the samples with deionised water which is slightly acidic) (Frohne et al., 2011; Izquierdo
822 et al., 2017; Rinklebe et al., 2010; Weber et al., 2009). This makes extrapolation of laboratory-based
823 findings to field situations difficult (Hooda, 2010). Attempts to model the concentration of PTEs in
824 floodplain pore waters have demonstrated the complexity of predicting how different variables such
825 as soil moisture content and temperature interact and alter mobility (Rennert et al., 2017), with site
826 or catchment-specific information being of great importance to establish and capture spatial
827 differences sufficiently (Schulz-Zunkel et al., 2015). While much research undertaken in controlled
828 conditions in laboratory microcosms is undoubtedly useful because independent replicates can be
829 assigned to treatments without confounding variables (Figure 4), there is a clear research need for on-
830 site experiments on the effect of flooding on PTEs mobility using real-time field-based observations
831 that capture the kinetics of processes before, during, and after a flooding event under ambient
832 temperatures and in geochemically contrasting soils.



833

834 *Figure 4: Strengths (+) and weaknesses (-) of laboratory-based studies for researching the impact of*
 835 *flooding on mobility of PTEs. Created with BioRender.com.*

836

837 Acknowledgements

838 Jessica Ponting is supported by a NERC SCENARIO PhD studentship, with CASE support from the British
 839 Geological Survey Universities Funding Initiative.

840

841 Supplementary Material

842 One supplementary table (Table S1) is provided

843

844 References

845 Abgottspon, F., Bigalke, M., Wilcke, W., 2015. Fast colloidal and dissolved release of trace elements
 846 in a carbonatic soil after experimental flooding. *Geoderma* 259–260, 156–163.
 847 <https://doi.org/10.1016/J.GEODERMA.2015.06.005>

- 848 Acreman, M.C., Riddington, R., Booker, D.J., 2003. Hydrological impacts of floodplain restoration: a
849 case study of the River Cherwell, UK, European Geosciences Union.
- 850 Adamo, P., Iavazzo, P., Albanese, S., Agrelli, D., De Vivo, B., Lima, A., 2014. Bioavailability and soil-to-
851 plant transfer factors as indicators of potentially toxic element contamination in agricultural
852 soils. *Sci. Total Environ.* 500–501, 11–22. <https://doi.org/10.1016/J.SCITOTENV.2014.08.085>
- 853 Adekanmbi, A.A., Shaw, L.J., Sizmur, T., 2020. Effect of Sieving on Ex Situ Soil Respiration of Soils
854 from Three Land Use Types. *J. Soil Sci. Plant Nutr.* 1–5. [https://doi.org/10.1007/s42729-020-](https://doi.org/10.1007/s42729-020-00177-2)
855 [00177-2](https://doi.org/10.1007/s42729-020-00177-2)
- 856 Adewuyi, G.O., Osobamiro, M.T., 2016. Chemical speciation and potential mobility of some toxic
857 metals in tropical agricultural soil. *Res. J. Environ. Toxicol.* 10, 159–165.
858 <https://doi.org/10.3923/rjet.2016.159.165>
- 859 Alfieri, L., Bisselink, B., Dottori, F., Naumann, G., de Roo, A., Salamon, P., Wyser, K., Feyen, L., 2017.
860 Global projections of river flood risk in a warmer world. *Earth's Futur.* 5, 171–182.
861 <https://doi.org/10.1002/2016EF000485>
- 862 Alloway, B.J., 2013. Heavy Metals in Soils, Heavy Metals in Soils. [https://doi.org/10.1007/978-94-](https://doi.org/10.1007/978-94-007-4470-7_10)
863 [007-4470-7_10](https://doi.org/10.1007/978-94-007-4470-7_10)
- 864 Alloway, B.J., 1995. Heavy Metals in Soils: Trace Metals and Metalloids in Soils and their
865 Bioavailability, 3rd ed. Springer, Heidelberg. <https://doi.org/10.1007/978-94-011-1344-1>
- 866 Álvarez-Ayuso, E., Otones, V., Murciego, A., García-Sánchez, A., Regina, I.S., 2012. Antimony, arsenic
867 and lead distribution in soils and plants of an agricultural area impacted by former mining
868 activities. *Sci. Total Environ.* 439, 35–43. <https://doi.org/10.1016/J.SCITOTENV.2012.09.023>
- 869 Alvim Ferraz, M.C.M., Lourenço, J.C.N., 2000. Influence of organic matter content of contaminated
870 soils on the leaching rate of heavy metals. *Environ. Prog.* 19, 53–58.
871 <https://doi.org/10.1002/ep.670190118>
- 872 Amacher, M.C., Kotuby-Amacher, J., Selim, H.M., Iskandar, I.K., 1986. Retention and release of
873 metals by soils — Evaluation of several models. *Geoderma* 38, 131–154.
874 [https://doi.org/10.1016/0016-7061\(86\)90011-X](https://doi.org/10.1016/0016-7061(86)90011-X)
- 875 Ander, E.L., Watts, M.J., Smedley, P.L., Hamilton, E.M., Close, R., Crabbe, H., Fletcher, T., Rimell, A.,
876 Studden, M., Leonardi, G., 2016. Variability in the chemistry of private drinking water supplies

877 and the impact of domestic treatment systems on water quality. *Environ. Geochem. Health* 38,
878 1313–1332. <https://doi.org/10.1007/s10653-016-9798-0>

879 Andersen, D.C., 2018. Flood effects on soil thermal regimes in contrasting cold-desert river
880 floodplains (Yampa and Green rivers, Colorado). *Ecohydrology* 11, e1939.
881 <https://doi.org/10.1002/eco.1939>

882 Antoniadis, V., Shaheen, S.M., Tsadilas, C.D., Selim, M.H., 2018. Zinc sorption by different soils as
883 affected by selective removal of carbonates and hydrous oxides. *Appl. Geochemistry* 88, 49–58.
884 <https://doi.org/10.1016/J.APGEOCHEM.2017.04.007>

885 Arnell, N.W., Halliday, S.J., Battarbee, R.W., Skeffington, R.A., Wade, A.J., 2015. The implications of
886 climate change for the water environment in England. *Prog. Phys. Geogr. Earth Environ.* 39, 93–
887 120. <https://doi.org/10.1177/0309133314560369>

888 Baran, A., Tarnawski, M., 2015. Assessment of heavy metals mobility and toxicity in contaminated
889 sediments by sequential extraction and a battery of bioassays. *Ecotoxicology* 24, 1279–1293.
890 <https://doi.org/10.1007/s10646-015-1499-4>

891 Barber, L.B., Paschke, S.S., Battaglin, W.A., Douville, C., Fitzgerald, K.C., Keefe, S.H., Roth, D.A., Vajda,
892 A.M., 2017. Effects of an Extreme Flood on Trace Elements in River Water - From Urban Stream
893 to Major River Basin. *Environ. Sci. Technol.* 51, 10344–10356.
894 <https://doi.org/10.1021/acs.est.7b01767>

895 Bednářová, Z., Komprdová, K., Kalábová, T., Sáňka, M., 2015. Impact of floods and their frequency on
896 content and distribution of risk elements in alluvial soils. *Water. Air. Soil Pollut.* 226.
897 <https://doi.org/10.1007/s11270-014-2253-x>

898 Bell, V.A., Kay, A.L., Cole, S.J., Jones, R.G., Moore, R.J., Reynard, N.S., 2012. How might climate
899 change affect river flows across the Thames Basin? An area-wide analysis using the UKCP09
900 Regional Climate Model ensemble. *J. Hydrol.* 442–443, 89–104.
901 <https://doi.org/10.1016/J.JHYDROL.2012.04.001>

902 Bellanger, B., Huon, S., Steinmann, P., Velasquez, F., Vall Es E, V., Arn, K., Clauer, N., Mariotti, A.E.,
903 2004. Oxic-anoxic conditions in the water column of a tropical freshwater reservoir (Peña-
904 Larga dam, NW Venezuela), Published in *Applied Geochemistry*.
905 <https://doi.org/10.1016/j.apgeochem.2003.11.007>

906 Berz, G., Kron, W., Loster, T., Rauch, E., Schimetschek, J., Schmieder, J., Siebert, A., Smolka, A., Wirtz,
907 A., 2001. World Map of Natural Hazards-A Global View of the Distribution and Intensity of
908 Significant Exposures, Natural Hazards.

909 Blackwell, M.S.A., Pilgrim, E.S., 2011. Ecosystem services delivered by small-scale wetlands. *Hydrol.*
910 *Sci. J.* 56, 1467–1484. <https://doi.org/10.1080/02626667.2011.630317>

911 Blöschl, G., Hall, J., Parajka, J., Perdigão, R.A.P., Merz, B., Arheimer, B., Aronica, G.T., Bilbashi, A.,
912 Bonacci, O., Borga, M., Čanjevac, I., Castellarin, A., Chirico, G.B., Claps, P., Fiala, K., Frolova, N.,
913 Gorbachova, L., Gül, A., Hannaford, J., Harrigan, S., Kireeva, M., Kiss, A., Kjeldsen, T.R.,
914 Kohnová, S., Koskela, J.J., Ledvinka, O., Macdonald, N., Mavrova-Guirguinova, M., Mediero, L.,
915 Merz, R., Molnar, P., Montanari, A., Murphy, C., Osuch, M., Ovcharuk, V., Radevski, I., Rogger,
916 M., Salinas, J.L., Sauquet, E., Šraj, M., Szolgay, J., Viglione, A., Volpi, E., Wilson, D., Zaimi, K.,
917 Živković, N., 2017. Changing climate shifts timing of European floods. *Science* (80-.). 357, 588–
918 590. <https://doi.org/10.1126/science.aan2506>

919 Borch, T., Kretzschmar, R., Skappler, A., Van Cappellen, P., Ginder-Vogel, M., Voegelin, A., Campbell,
920 K., 2010. Biogeochemical redox processes and their impact on contaminant dynamics. *Environ.*
921 *Sci. Technol.* <https://doi.org/10.1021/es9026248>

922 Bradley, S.B., Cox, J.J., 1990. The significance of the floodplain to the cycling of metals in the river
923 Derwent Catchment, U.K. *Sci. Total Environ.* 97–98, 441–454. [https://doi.org/10.1016/0048-](https://doi.org/10.1016/0048-9697(90)90255-S)
924 [9697\(90\)90255-S](https://doi.org/10.1016/0048-9697(90)90255-S)

925 Brönnimann, S., 2007. Impact of El Niño-Southern Oscillation on European climate. *Rev. Geophys.*
926 45, n/a-n/a. <https://doi.org/10.1029/2006RG000199>

927 Bronstert, A., 2003. *Floods and Climate Change: Interactions and Impacts, Risk Analysis.*

928 Bufflap, S.E., Allen, H.E., 1995. Sediment pore water collection methods for trace metal analysis: A
929 review. *Water Res.* 29, 165–177. [https://doi.org/10.1016/0043-1354\(94\)E0105-F](https://doi.org/10.1016/0043-1354(94)E0105-F)

930 Chan, S.C., Kendon, E.J., Fowler, H.J., Blenkinsop, S., Roberts, N.M., 2014. Projected increase in
931 summer and winter UK sub-daily precipitation extremes from high-resolution regional climate
932 models [WWW Document]. *Environ. Res. Lett.* URL
933 [https://www.researchgate.net/publication/265090772_Projected_increase_in_summer_and_](https://www.researchgate.net/publication/265090772_Projected_increase_in_summer_and_winter_UK_sub-daily_precipitation_extremes_from_high-resolution_regional_climate_models)
934 [winter_UK_sub-daily_precipitation_extremes_from_high-resolution_regional_climate_models](https://www.researchgate.net/publication/265090772_Projected_increase_in_summer_and_winter_UK_sub-daily_precipitation_extremes_from_high-resolution_regional_climate_models)
935 (accessed 3.2.20).

936 Chang, H., Franczyk, J., 2008. Climate Change, Land-Use Change, and Floods: Toward an Integrated
937 Assessment. *Geogr. Compass* 2, 1549–1579. <https://doi.org/10.1111/j.1749-8198.2008.00136.x>

938 Chrzan, A., 2016. Monitoring bioconcentration of potentially toxic trace elements in soils trophic
939 chains. *Environ. Earth Sci.* 75, 1–8. <https://doi.org/10.1007/s12665-016-5595-4>

940 Cipullo, S., Prpich, G., Campo, P., Coulon, F., 2018. Assessing bioavailability of complex chemical
941 mixtures in contaminated soils: Progress made and research needs. *Sci. Total Environ.* 615,
942 708–723. <https://doi.org/10.1016/j.scitotenv.2017.09.321>

943 Ciszewski, D., Grygar, T.M., 2016. A Review of Flood-Related Storage and Remobilization of Heavy
944 Metal Pollutants in River Systems. *Water. Air. Soil Pollut.* 227, 1–19.
945 <https://doi.org/10.1007/s11270-016-2934-8>

946 Classen, A.T., Sundqvist, M.K., Henning, J.A., Newman, G.S., Moore, J.A.M., Cregger, M.A.,
947 Moorhead, L.C., Patterson, C.M., 2015. Direct and indirect effects of climate change on soil
948 microbial and soil microbial-plant interactions: What lies ahead? *Ecosphere* 6, art130.
949 <https://doi.org/10.1890/ES15-00217.1>

950 Clemente, R., Dickinson, N.M., Lepp, N.W., 2008. Mobility of metals and metalloids in a multi-
951 element contaminated soil 20 years after cessation of the pollution source activity. *Environ.*
952 *Pollut.* 155, 254–261. <https://doi.org/10.1016/j.envpol.2007.11.024>

953 Czech, T., Baran, A., Wieczorek, J., 2014. Zawartość Metali Ciężkich W Glebach I Roślinach Z Terenu
954 Gminy Borzęcin (Województwo Małopolskie) Content of Heavy Metals in Soil and Plants From
955 an Area Borzęcin Municipality (Malopolska Province) 89–98.
956 <https://doi.org/10.12912/2081139X.20>

957 Dadson, S.J., Hall, J.W., Murgatroyd, A., Acreman, M., Bates, P., Beven, K., Heathwaite, L., Holden, J.,
958 Holman, I.P., Lane, S.N., O'connell, E., Penning-Rowell, E., Reynard, N., Sear, D., Thorne, C.,
959 Wilby, R., 2017. A restatement of the natural science evidence concerning catchment-based
960 “natural” flood management in the UK. *Proc. R. Soc. A* 473.
961 <https://doi.org/10.1098/rspa.2016.0706>

962 Dang, Z., Liu, C., Haigh, M.J., 2002. Mobility of heavy metals associated with the natural weathering
963 of coal mine spoils. *Environ. Pollut.* 118, 419–426. [https://doi.org/10.1016/S0269-7491\(01\)00285-8](https://doi.org/10.1016/S0269-7491(01)00285-8)

- 965 Davranche, M., Grybos, M., Gruau, G., Pédrot, M., Dia, A., Marsac, R., 2011. Rare earth element
966 patterns: A tool for identifying trace metal sources during wetland soil reduction. *Chem. Geol.*
967 284, 127–137. <https://doi.org/10.1016/j.chemgeo.2011.02.014>
- 968 Dawson, J.J.C., Tetzlaff, D., Carey, A.M., Raab, A., Soulsby, C., Killham, K., Meharg, A.A., 2010.
969 Characterizing Pb mobilization from upland soils to streams using ²⁰⁶Pb/²⁰⁷Pb isotopic ratios.
970 *Environ. Sci. Technol.* 44, 243–249. <https://doi.org/10.1021/es902664d>
- 971 De Jonge, M., Teuchies, J., Meire, P., Blust, R., Bervoets, L., 2012. The impact of increased oxygen
972 conditions on metal-contaminated sediments part I: Effects on redox status, sediment
973 geochemistry and metal bioavailability. *Water Res.* 46, 2205–2214.
974 <https://doi.org/10.1016/J.WATRES.2012.01.052>
- 975 de Souza Machado, A.A., Spencer, K., Kloas, W., Toffolon, M., Zarfl, C., 2016. Metal fate and effects
976 in estuaries: A review and conceptual model for better understanding of toxicity. *Sci. Total*
977 *Environ.* 541, 268–281. <https://doi.org/10.1016/J.SCITOTENV.2015.09.045>
- 978 de Souza Machado, A.A., Spencer, K.L., Zarfl, C., O’Shea, F.T., 2018. Unravelling metal mobility under
979 complex contaminant signatures. *Sci. Total Environ.* 622–623, 373–384.
980 <https://doi.org/10.1016/J.SCITOTENV.2017.11.239>
- 981 De Vivo, B., Belkin, H.E., Lima, A., 2008. *Environmental Geochemistry: Site Characterization, Data*
982 *Analysis And Case Histories*, *Environmental Geochemistry: Site Characterization, Data Analysis*
983 *and Case Histories*. Elsevier. <https://doi.org/10.1016/B978-0-444-53159-9.X0001-0>
- 984 Degryse, F., Smolders, E., Parker, D.R., 2009. Partitioning of metals (Cd, Co, Cu, Ni, Pb, Zn) in soils:
985 concepts, methodologies, prediction and applications - a review. *Eur. J. Soil Sci.* 60, 590–612.
986 <https://doi.org/10.1111/j.1365-2389.2009.01142.x>
- 987 Domergue, F.L., Vedy, J.C., 1992. Mobility of heavy metals in soil profiles. *Int. J. Environ. Anal. Chem.*
988 46, 13–23. <https://doi.org/10.1080/03067319208026993>
- 989 Du Laing, G., Rinklebe, J., Vandecasteele, B., Meers, E., Tack, F.M.G., 2009. Trace metal behaviour in
990 estuarine and riverine floodplain soils and sediments: A review. *Sci. Total Environ.* 407, 3972–
991 3985. <https://doi.org/10.1016/J.SCITOTENV.2008.07.025>
- 992 Du Laing, G., Vanthuyne, D.R.J., Vandecasteele, B., Tack, F.M.G., Verloo, M.G., 2007. Influence of
993 hydrological regime on pore water metal concentrations in a contaminated sediment-derived

- 994 soil. *Environ. Pollut.* 147, 615–625. <https://doi.org/10.1016/J.ENVPOL.2006.10.004>
- 995 Eggleton, J., Thomas, K. V., 2004. A review of factors affecting the release and bioavailability of
996 contaminants during sediment disturbance events. *Environ. Int.* 30, 973–980.
997 <https://doi.org/10.1016/J.ENVINT.2004.03.001>
- 998 Ehlers, G.A.C., Loibner, A.P., 2006. Linking organic pollutant (bio)availability with geosorbent
999 properties and biomimetic methodology: A review of geosorbent characterisation and
1000 (bio)availability prediction. *Environ. Pollut.* 141, 494–512.
1001 <https://doi.org/10.1016/J.ENVPOL.2005.08.063>
- 1002 Ellis, J.B., Shutes, R.B.E., Revitt, M.D., 2003. Constructed Wetlands and Links with Sustainable
1003 Drainage Systems.
- 1004 Emerson, Kodak U, Scholastica Bejor, E., Ekeng, E.E., Ogarekpe, Nkpa M, Emerson, K U, Bejor, E.S.,
1005 Ogarekpe, N M, Onuruka, A.U., 2017. TRANSPORT AND FATE OF SELECTED HEAVY METALS IN
1006 CIRCUMNEUTRAL RIVER ENVIRONMENT: A CASE STUDY OF THE RIVER NENT CUMBRIA,
1007 ENGLAND. *Int. J. Res.* 5, 159–169. <https://doi.org/10.5281/zenodo.818628>
- 1008 Euripidou, E., Murray, V., 2004. Public health impacts of floods and chemical contamination. *J. Public*
1009 *Health (Bangkok)*. 26, 376–383. <https://doi.org/10.1093/pubmed/fdh163>
- 1010 Evans, L.J., 1989. Chemistry of Metal Retention by Soils Chemistry of metal retention by soils Several
1011 processes are explained. *Artic. Environ. Sci. Technol.* <https://doi.org/10.1021/es00067a001>
- 1012 Foley, J.A., DeFries, R., Asner, G.P., Barford, C., Bonan, G., Carpenter, S.R., Chapin, F.S., Coe, M.T.,
1013 Daily, G.C., Gibbs, H.K., Helkowski, J.H., Holloway, T., Howard, E.A., Kucharik, C.J., Monfreda, C.,
1014 Patz, J.A., Prentice, I.C., Ramankutty, N., Snyder, P.K., 2005. Global consequences of land use.
1015 *Science (80-)*. <https://doi.org/10.1126/science.1111772>
- 1016 Förstner, U., 2004. Traceability of sediment analysis. *TrAC Trends Anal. Chem.* 23, 217–236.
1017 [https://doi.org/10.1016/S0165-9936\(04\)00312-7](https://doi.org/10.1016/S0165-9936(04)00312-7)
- 1018 Forstner, U., Wattman, G., 1981. *Metal Pollution in the Aquatic Environment.*, 2nd ed. Springer-
1019 Verlag, Berlin.
- 1020 Foulds, S.A., Griffiths, H.M., Macklin, M.G., Brewer, P.A., 2014. Geomorphological records of extreme
1021 floods and their relationship to decadal-scale climate change. *Geomorphology* 216, 193–207.
1022 <https://doi.org/10.1016/J.GEOMORPH.2014.04.003>

- 1023 Frohne, T., Diaz-Bone, R.A., Du Laing, G., Rinklebe, J., 2015. Impact of systematic change of redox
1024 potential on the leaching of Ba, Cr, Sr, and V from a riverine soil into water. *J. Soils Sediments*
1025 15, 623–633. <https://doi.org/10.1007/s11368-014-1036-8>
- 1026 Frohne, T., Rinklebe, J., Diaz-Bone, R.A., 2014. Contamination of Floodplain Soils along the Wupper
1027 River, Germany, with As, Co, Cu, Ni, Sb, and Zn and the Impact of Pre-definite Redox Variations
1028 on the Mobility of These Elements. *Soil Sediment Contam.* 23, 779–799.
1029 <https://doi.org/10.1080/15320383.2014.872597>
- 1030 Frohne, T., Rinklebe, J., Diaz-Bone, R.A., Du Laing, G., 2011. Controlled variation of redox conditions
1031 in a floodplain soil: Impact on metal mobilization and biomethylation of arsenic and antimony.
1032 *Geoderma* 160, 414–424. <https://doi.org/10.1016/J.GEODERMA.2010.10.012>
- 1033 Gadd, G.M., 2010. Metals, minerals and microbes: Geomicrobiology and bioremediation.
1034 *Microbiology.* <https://doi.org/10.1099/mic.0.037143-0>
- 1035 Gall, Jillian E, Boyd, Robert S, Rajakaruna, Nishanta, Gall, J E, Rajakaruna, N, Boyd, R S, 2015. Transfer
1036 of heavy metals through terrestrial food webs: a review, *Monitoring and Assessment.*
- 1037 Gambrell, R.P., 1994. Trace and Toxic Metals in Wetlands-A Review. *J. Environ. Qual.* 23, 883–891.
- 1038 Gao, H., Shao, M., 2015. Effects of temperature changes on soil hydraulic properties. *Soil Tillage Res.*
1039 153, 145–154. <https://doi.org/10.1016/j.still.2015.05.003>
- 1040 Giacalone, A., Gianguzza, A., Orecchio, S., Piazzese, D., Dongarrà, G., Sciarrino, S., Varrica, D., 2005.
1041 Metals distribution in the organic and inorganic fractions of soil: A case study on soils from
1042 Sicily. *Chem. Speciat. Bioavailab.* 17, 83–93. <https://doi.org/10.3184/095422905782774892>
- 1043 González-Alcaraz, M.N., van Gestel, C.A.M., 2015. Climate change effects on enchytraeid
1044 performance in metal-polluted soils explained from changes in metal bioavailability and
1045 bioaccumulation. *Environ. Res.* 142, 177–184. <https://doi.org/10.1016/j.envres.2015.06.027>
- 1046 González-Macé, O., Steinauer, K., Jousset, A., Eisenhauer, N., Scheu, S., 2016. Flood-induced changes
1047 in soil microbial functions as modified by plant diversity. *PLoS One* 11.
1048 <https://doi.org/10.1371/journal.pone.0166349>
- 1049 Gregory, A.S., Ritz, K., McGrath, S.P., Quinton, J.N., Goulding, K.W.T., Jones, R.J.A., Harris, J.A., Bol,
1050 R., Wallace, P., Pilgrim, E.S., Whitmore, A.P., 2015. A review of the impacts of degradation
1051 threats on soil properties in the UK. *Soil Use Manag.* 31, 1–15.

1052 <https://doi.org/10.1111/sum.12212>

1053 Grimm, A.M., Tedeschi, R.G., 2009. ENSO and extreme rainfall events in South America. *J. Clim.* 22,
1054 1589–1609. <https://doi.org/10.1175/2008JCLI2429.1>

1055 Gröngröft, A., Krüger, F., Grunewald, K., Meißner, R., Miehl, G., 2005. Plant and soil
1056 contamination with trace metals in the Elbe floodplains: A case study after the flood in August
1057 2002. *Acta Hydrochim. Hydrobiol.* 33, 466–474. <https://doi.org/10.1002/aheh.200400596>

1058 Hahne, H.C.H., Kroontje, W., 1973. Significance of pH and Chloride Concentration on Behavior of
1059 Heavy Metal Pollutants: Mercury(II), Cadmium(II), Zinc(II), and Lead(II). *J. Environ. Qual.* 2, 444–
1060 450. <https://doi.org/10.2134/jeq1973.00472425000200040007x>

1061 Harvey, R.J., Chadwick, D.R., Sánchez-Rodríguez, A.R., Jones, D.L., 2019. Agroecosystem resilience in
1062 response to extreme winter flooding. *Agric. Ecosyst. Environ.* 279, 1–13.
1063 <https://doi.org/10.1016/J.AGEE.2019.04.001>

1064 He, Y., Men, B., Yang, X., Li, Y., Xu, H., Wang, D., 2019. Relationship between heavy metals and
1065 dissolved organic matter released from sediment by bioturbation/bioirrigation. *J. Environ. Sci.*
1066 (China) 75, 216–223. <https://doi.org/10.1016/j.jes.2018.03.031>

1067 Hillel, D., 1998. *Introduction to Environmental Soil Physics*, 1st ed. Academic Press.

1068 Hippolyte, I., Jenny, C., Gardes L., Bakry F., Rivallan R., Pomies V., Cubry P., Tomekpe K., Risterucci A.
1069 M., Roux N., Rouard M., Arnaud E., Kolesnikova-Allen M., Perrier X., 2012. Foundation
1070 characteristics of edible *Musa* triploids revealed from allelic distribution of SSR markers. *Ann.*
1071 *Bot.* 109, 937–951. <https://doi.org/10.1093/aob>

1072 Hirabayashi, Y., Kanae, S., 2009. First estimate of the future global population at risk of flooding.
1073 *Hydrol. Res. Lett.* 3, 6–9. <https://doi.org/10.3178/hrl.3.6>

1074 Hirabayashi, Y., Kanae, S., Emori, S., Oki, T., Kimoto, M., 2008. Global projections of changing risks of
1075 floods and droughts in a changing climate. *Hydrol. Sci. J.* 53, 754–772.
1076 <https://doi.org/10.1623/hysj.53.4.754>

1077 Hirabayashi, Y., Mahendran, R., Koirala, S., Konoshima, L., Yamazaki, D., Watanabe, S., Kim, H.,
1078 Kanae, S., 2013. Global flood risk under climate change. *Nat. Clim. Chang.* 3, 816–821.
1079 <https://doi.org/10.1038/nclimate1911>

- 1080 Hobbelen, P.H.F., Koolhaas, J.E., van Gestel, C.A.M., 2006. Bioaccumulation of heavy metals in the
1081 earthworms *Lumbricus rubellus* and *Aporrectodea caliginosa* in relation to total and available
1082 metal concentrations in field soils. *Environ. Pollut.* 144, 639–646.
1083 <https://doi.org/10.1016/J.ENVPOL.2006.01.019>
- 1084 Hooda, P.S., 2010. Trace Elements in Soils, Trace Elements in Soils. John Wiley and Sons.
1085 <https://doi.org/10.1002/9781444319477>
- 1086 Huddart, D., Stott, T., Huddart, D., Stott, T., 2020. Climate Change and Adventure Tourism, in:
1087 Adventure Tourism. Springer International Publishing, pp. 437–469.
1088 https://doi.org/10.1007/978-3-030-18623-4_13
- 1089 Hurley, R.R., Rothwell, J.J., Woodward, J.C., 2017. Metal contamination of bed sediments in the
1090 Irwell and Upper Mersey catchments, northwest England: exploring the legacy of industry and
1091 urban growth. *J. Soils Sediments* 17, 2648–2665. <https://doi.org/10.1007/s11368-017-1668-6>
- 1092 Ibragimow, A., Walna, B., Siepak, M., 2013. Effects of Flooding on the Contamination of Floodplain
1093 Sediments with Available Fractions of Trace Metals (Western Poland), *J. Environ. Stud.*
- 1094 IPCC, 2012. Managing the Risks of Extreme Events and Disasters to Advance Climate Change
1095 Adaptation. A Special Report of Working Groups I and II of the Intergovernmental Panel on
1096 Climate Change. [Field, C.B., V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.
- 1097 Izquierdo, M., Tye, A.M., Chenery, S.R., 2017. Using isotope dilution assays to understand speciation
1098 changes in Cd, Zn, Pb and Fe in a soil model system under simulated flooding conditions.
1099 *Geoderma* 295, 41–52. <https://doi.org/10.1016/J.GEODERMA.2017.02.006>
- 1100 Izquierdo, M., Tye, A.M., Chenery, S.R., 2013. Lability, solubility and speciation of Cd, Pb and Zn in
1101 alluvial soils of the River Trent catchment UK. *Environ. Sci. Process. Impacts* 15, 1844–1858.
1102 <https://doi.org/10.1039/c3em00370a>
- 1103 Jenkins, G.J., Murphy, J.M., Sexton, D.M.H., Lowe, J.A., Jones, P., Kilsby, C.G., 2009. UK Climate
1104 Projection Briefing Report [WWW Document]. Met Off. Hadley Cent. URL
1105 [https://www.researchgate.net/publication/257343385_UK_Climate_Projection_Briefing_Repo](https://www.researchgate.net/publication/257343385_UK_Climate_Projection_Briefing_Report)
1106 [rt](https://www.researchgate.net/publication/257343385_UK_Climate_Projection_Briefing_Report) (accessed 3.2.20).
- 1107 Jiao, W., Ouyang, W., Hao, F., Huang, H., Shan, Y., Geng, X., 2014. Combine the soil water assessment
1108 tool (SWAT) with sediment geochemistry to evaluate diffuse heavy metal loadings at watershed

- 1109 scale. *J. Hazard. Mater.* 280, 252–259. <https://doi.org/10.1016/J.JHAZMAT.2014.07.081>
- 1110 Johnson, Z.F., Chikamoto, Y., Wang, S.Y.S., McPhaden, M.J., Mochizuki, T., 2020. Pacific decadal
1111 oscillation remotely forced by the equatorial Pacific and the Atlantic Oceans. *Clim. Dyn.* 55,
1112 789–811. <https://doi.org/10.1007/s00382-020-05295-2>
- 1113 Johnston, S.G., Burton, E.D., Aaso, T., Tuckerman, G., 2014. Sulfur, iron and carbon cycling following
1114 hydrological restoration of acidic freshwater wetlands. *Chem. Geol.* 371, 9–26.
1115 <https://doi.org/10.1016/j.chemgeo.2014.02.001>
- 1116 Junk, W.J., Bayley, P.B., Sparks, R.E., 1989. The Flood Pulse Concept in River-Floodplain Systems. *Can.*
1117 *Spec. Publ. Fish. Aquat. Sci.* 106, 110–127. https://doi.org/10.1007/978-3-662-03416-3_4
- 1118 Kalbitz, K., Wennrich, R., 1998. Mobilization of heavy metals and arsenic in polluted wetland soils
1119 and its dependence on dissolved organic matter. *Sci. Total Environ.* 209, 27–39.
1120 [https://doi.org/10.1016/S0048-9697\(97\)00302-1](https://doi.org/10.1016/S0048-9697(97)00302-1)
- 1121 Karimian, N., Johnston, S.G., Burton, E.D., 2018. Iron and sulfur cycling in acid sulfate soil wetlands
1122 under dynamic redox conditions: A review. *Chemosphere* 197, 803–816.
1123 <https://doi.org/10.1016/j.chemosphere.2018.01.096>
- 1124 Karimian, N., Johnston, S.G., Burton, E.D., 2017. Effect of cyclic redox oscillations on water quality in
1125 freshwater acid sulfate soil wetlands. *Sci. Total Environ.*
1126 <https://doi.org/10.1016/j.scitotenv.2016.12.131>
- 1127 Karl, T.R., Trenberth, K.E., 2003. Modern Global Climate Change. *Science* (80-.). 302, 1719–1723.
1128 <https://doi.org/10.1126/science.1090228>
- 1129 Kelly, T.J., Hamilton, E., Watts, M.J., Ponting, J., Sizmur, T., 2020. The effect of flooding and drainage
1130 duration on the release of trace elements from floodplain soils. *Environ. Toxicol. Chem.*
1131 etc.4830. <https://doi.org/10.1002/etc.4830>
- 1132 Kharin, V. V., Zwiers, F.W., Zhang, X., Hegerl, G.C., 2007. Changes in temperature and precipitation
1133 extremes in the IPCC ensemble of global coupled model simulations. *J. Clim.* 20, 1419–1444.
1134 <https://doi.org/10.1175/JCLI4066.1>
- 1135 Kirk, G., 2004. *The Biogeochemistry of Submerged Soils, The Biogeochemistry of Submerged Soils.*
1136 Wiley & Sons, Ltd: Chichester. <https://doi.org/10.1002/047086303x>

- 1137 Kiss, T.B.W., Chen, X., Ponting, J., Sizmur, T., Hodson, M.E., (In Review) Dual stresses of flooding and
1138 agricultural land use reduce earthworm populations more than the individual stressors. *Sci.*
1139 *Total Environ.* STOTEN-D-20-17138
- 1140 Koretsky, C.M., Haveman, M., Beuving, L., Cuellar, A., Shattuck, T., Wagner, M., 2007. Spatial
1141 variation of redox and trace metal geochemistry in a minerotrophic fen. *Biogeochemistry* 86,
1142 33–62. <https://doi.org/10.1007/s10533-007-9143-x>
- 1143 Kowalik, C., Kraft, J., Einax, J.W., 2004. The situation of the German Elbe tributaries - Development of
1144 the loads in the last 10 years. *Acta Hydrochim. Hydrobiol.* 31, 334–345.
1145 <https://doi.org/10.1002/aheh.200300507>
- 1146 Kulp, S.A., Strauss, B.H., 2019. New elevation data triple estimates of global vulnerability to sea-level
1147 rise and coastal flooding. *Nat. Commun.* 10, 1–12. <https://doi.org/10.1038/s41467-019-12808->
1148 [z](https://doi.org/10.1038/s41467-019-12808-z)
- 1149 Kundzewicz, Z.W., Kanae, S., Seneviratne, S.I., Handmer, J., Nicholls, N., Peduzzi, P., Mechler, R.,
1150 Bouwer, L.M., Arnell, N., Mach, K., Muir-Wood, R., Brakenridge, G.R., Kron, W., Benito, G.,
1151 Honda, Y., Takahashi, K., Sherstyukov, B., 2014. Flood risk and climate change: global and
1152 regional perspectives. *Hydrol. Sci. J.* 59, 1–28. <https://doi.org/10.1080/02626667.2013.857411>
- 1153 Lane, S.N., 2017. Natural flood management. *Wiley Interdiscip. Rev. Water* 4, e1211.
1154 <https://doi.org/10.1002/wat2.1211>
- 1155 Le Gall, M., Ayrault, S., Evrard, O., Laceby, J.P., Gateuille, D., Lefèvre, I., Mouchel, J.-M., Meybeck, M.,
1156 2018. Investigating the metal contamination of sediment transported by the 2016 Seine River
1157 flood (Paris, France). *Environ. Pollut.* 240, 125–139.
1158 <https://doi.org/10.1016/J.ENVPOL.2018.04.082>
- 1159 Lee, D.Y., Huang, J.C., Juang, K.W., Tsui, L., 2005. Assessment of phytotoxicity of chromium in
1160 flooded soils using embedded selective ion exchange resin method. *Plant Soil* 277, 97–105.
1161 <https://doi.org/10.1007/s11104-005-5997-7>
- 1162 Lewin, J., Macklin, M.G., 2010. Floodplain catastrophes in the UK Holocene: Messages for managing
1163 climate change. *Hydrol. Process.* 24, 2900–2911. <https://doi.org/10.1002/hyp.7704>
- 1164 Lloyd, J.R., 2003. Microbial reduction of metals and radionuclides. *FEMS Microbiol. Rev.* 27, 411–
1165 425. [https://doi.org/10.1016/S0168-6445\(03\)00044-5](https://doi.org/10.1016/S0168-6445(03)00044-5)

- 1166 Lu, S., Ren, T., Gong, Y., Horton, R., 2007. An Improved Model for Predicting Soil Thermal
1167 Conductivity from Water Content at Room Temperature. *Soil Sci. Soc. Am. J.* 71, 8–14.
1168 <https://doi.org/10.2136/sssaj2006.0041>
- 1169 Lučić, M., Jurina, I., Ščančar, J., Mikac, N., Vdović, N., 2019. Sedimentological and geochemical
1170 characterization of river suspended particulate matter (SPM) sampled by time-integrated mass
1171 flux sampler (TIMS) in the Sava River (Croatia). *J. Soils Sediments* 19, 989–1004.
1172 <https://doi.org/10.1007/s11368-018-2104-2>
- 1173 Ma, M., Du, H., Wang, D., 2019. Mercury methylation by anaerobic microorganisms: A review. *Crit.*
1174 *Rev. Environ. Sci. Technol.* 49, 1893–1936. <https://doi.org/10.1080/10643389.2019.1594517>
- 1175 Ma, R., McBratney, A., Whelan, B., Minasny, B., Short, M., 2011. Comparing temperature correction
1176 models for soil electrical conductivity measurement. *Precis. Agric.*
1177 <https://doi.org/10.1007/s11119-009-9156-7>
- 1178 MacDonald, D., Dixon, A., Newell, A., Hallaways, A., 2012. Groundwater flooding within an urbanised
1179 flood plain. *J. Flood Risk Manag.* 5, 68–80. <https://doi.org/10.1111/j.1753-318X.2011.01127.x>
- 1180 Macklin, M.G., Rumsby, B.T., 2007. *Changing Climate and Extreme Floods in the British Uplands, New*
1181 *Series.*
- 1182 Madsen, H., Lawrence, D., Lang, M., Martinkova, M., Kjeldsen, T.R., 2014. Review of trend analysis
1183 and climate change projections of extreme precipitation and floods in Europe. *J. Hydrol.* 519,
1184 3634–3650. <https://doi.org/10.1016/J.JHYDROL.2014.11.003>
- 1185 Maggioni, V., Massari, C., 2018. On the performance of satellite precipitation products in riverine
1186 flood modeling: A review. *J. Hydrol.* 558, 214–224.
1187 <https://doi.org/10.1016/J.JHYDROL.2018.01.039>
- 1188 Malmon, D. V., Dunne, T., Reneau, S.L., 2002. Predicting the Fate of Sediment and Pollutants in River
1189 Floodplains. *Environ. Sci. Technol* 36, 2026–2032. <https://doi.org/10.1021/ES010509+>
- 1190 Malmon, D. V., Reneau, S.L., Dunne, T., 2004. Sediment sorting and transport by flash floods. *J.*
1191 *Geophys. Res. Earth Surf.* 109, n/a-n/a. <https://doi.org/10.1029/2003jf000067>
- 1192 Maluckov, B.S., 2017. The Catalytic Role of *Acidithiobacillus ferrooxidans* for Metals Extraction from
1193 Mining - Metallurgical Resource. *Biodivers. Int. J.* 1.
1194 <https://doi.org/10.15406/bij.2017.01.00017>

- 1195 Martin, M., Bonifacio, E., Hossain, K.M.J., Huq, S.M.I., Barberis, E., 2014. Arsenic fixation and
1196 mobilization in the soils of the Ganges and Meghna floodplains. Impact of pedoenvironmental
1197 properties. *Geoderma* 228–229, 132–141. <https://doi.org/10.1016/j.geoderma.2013.09.020>
- 1198 Meegoda, J.N., Martin, L., 2019. In-Situ Determination of Specific Surface Area of Clays. *Geotech*
1199 *Geol Eng* 37, 465–474. <https://doi.org/https://doi.org/10.1007/s10706-018-0623-7>
- 1200 Mullan, D., Matthews, T., Vandaele, K., Barr, I.D., Swindles, G.T., Meneely, J., Boardman, J., Murphy,
1201 C., 2019. Climate impacts on soil erosion and muddy flooding at 1.5 versus 2°C warming. *L.*
1202 *Degrad. Dev.* 30, 94–108. <https://doi.org/10.1002/ldr.3214>
- 1203 Naidu, R., Channey, R., McConnell, S., Johnston, N., Semple, K.T., McGrath, S., Dries, V., Nathanail, P.,
1204 Harmsen, J., Pruszinski, A., MacMillan, J., Palanisami, T., 2015. Towards bioavailability-based
1205 soil criteria: past, present and future perspectives. *Environ. Sci. Pollut. Res.* 22, 8779–8785.
1206 <https://doi.org/10.1007/s11356-013-1617-x>
- 1207 Naidu, R., Semple, K.T., Megharaj, M., Juhasz, A.L., Bolan, N.S., Gupta, S.K., Clothier, B.E., Schulin, R.,
1208 2008. Bioavailability: Definition, assessment and implications for risk assessment. *Dev. Soil Sci.*
1209 32, 39–51. [https://doi.org/10.1016/S0166-2481\(07\)32003-5](https://doi.org/10.1016/S0166-2481(07)32003-5)
- 1210 Neal, C., Smith, C.J., Jeffery, H.A., Jarvie, H.P., Robson, A.J., 1996. Trace element concentrations in
1211 the major rivers entering the Humber estuary, NE England. *J. Hydrol.* 182, 37–64.
1212 [https://doi.org/10.1016/0022-1694\(95\)02940-0](https://doi.org/10.1016/0022-1694(95)02940-0)
- 1213 Ng, J.C., Juhasz, A., Smith, E., Naidu, R., 2015. Assessing the bioavailability and bioaccessibility of
1214 metals and metalloids. *Environ. Sci. Pollut. Res.* 22, 8802–8825.
1215 <https://doi.org/10.1007/s11356-013-1820-9>
- 1216 Niu, Z., Sun, L., Sun, T., Li, Y., Wang, H., 2007. Evaluation of phytoextracting cadmium and lead by
1217 sunflower, ricinus, alfalfa and mustard in hydroponic culture. *J. Environ. Sci.* 19, 961–967.
1218 [https://doi.org/10.1016/S1001-0742\(07\)60158-2](https://doi.org/10.1016/S1001-0742(07)60158-2)
- 1219 Nouri, J., Khorasani, N., Lorestani, B., Karami, M., Hassani, A.H., Yousefi, N., 2009. Accumulation of
1220 heavy metals in soil and uptake by plant species with phytoremediation potential. *Environ.*
1221 *Earth Sci.* 59, 315–323. <https://doi.org/10.1007/s12665-009-0028-2>
- 1222 Nriagu, J.O., Bhattacharya, P., Mukherjee, A.B., Bundschuh, J., Zevenhoven, R., Loeppert, R.H., 2007.
1223 Arsenic in soil and groundwater: an overview. *Trace Met. other Contam. Environ.* 9, 3–60.

- 1224 [https://doi.org/10.1016/S1875-1121\(06\)09001-8](https://doi.org/10.1016/S1875-1121(06)09001-8)
- 1225 Nshimiyimana, F.X., Faciu, M.-E., Abidi, A. El, Blidi, S. El, Fekhaoui, M., Ifrim, I.L., Soulaymani, A.,
1226 Lazar, G., 2014. ANALYSIS OF SEASONAL VARIATION ON DEGREE OF CONTAMINATION WITH
1227 HEAVY METALS IN AARJATE VILLAGE, MOROCCO. AN INDEX APPROACH. *Sci. Study Res. Chem.*
1228 *Chem. Eng. Biotechnol. Food Ind.* 15, 337–344.
- 1229 Ostergren, J.D., Brown, G.E., Parks, G.A., Persson, P., 2000. Inorganic ligand effects on Pb(II) sorption
1230 to goethite (α -FeOOH). II. Sulfate. *J. Colloid Interface Sci.* 225, 483–493.
1231 <https://doi.org/10.1006/jcis.1999.6702>
- 1232 Overesch, M., Rinklebe, J., Broll, G., Neue, H.U., 2007. Metals and arsenic in soils and corresponding
1233 vegetation at Central Elbe river floodplains (Germany). *Environ. Pollut.* 145, 800–812.
1234 <https://doi.org/10.1016/j.envpol.2006.05.016>
- 1235 Pan, L., Fang, G., Wang, Y., Wang, L., Su, B., Li, D., Xiang, B., 2018. Potentially toxic element pollution
1236 levels and risk assessment of soils and sediments in the upstream river, miyun reservoir, china.
1237 *Int. J. Environ. Res. Public Health* 15. <https://doi.org/10.3390/ijerph15112364>
- 1238 Pappenberger, F., Dutra, E., Wetterhall, F., Cloke, H.L., 2012. Deriving global flood hazard maps of
1239 fluvial floods through a physical model cascade. *Hydrol. Earth Syst. Sci.* 16, 4143–4156.
1240 <https://doi.org/10.5194/hess-16-4143-2012>
- 1241 Pathirana, A., Denekew, H.B., Veerbeek, W., Zevenbergen, C., Banda, A.T., 2014. Impact of urban
1242 growth-driven landuse change on microclimate and extreme precipitation - A sensitivity study.
1243 *Atmos. Res.* 138, 59–72. <https://doi.org/10.1016/j.atmosres.2013.10.005>
- 1244 Peacock, M., Freeman, C., Gauci, V., Lebron, I., Evans, C.D., 2015. Investigations of freezing and cold
1245 storage for the analysis of peatland dissolved organic carbon (DOC) and absorbance properties.
1246 *Environ. Sci. Process. Impacts* 17, 1290–1301. <https://doi.org/10.1039/c5em00126a>
- 1247 Peijnenburg, W.J.G.M., Zablotzkaja, M., Vijver, M.G., 2007. Monitoring metals in terrestrial
1248 environments within a bioavailability framework and a focus on soil extraction. *Ecotoxicol.*
1249 *Environ. Saf.* 67, 163–179. <https://doi.org/10.1016/J.ECOENV.2007.02.008>
- 1250 Pendergrass, A.G., 2018. What precipitation is extreme? *Science* (80-.). 360, 1072–1073.
1251 <https://doi.org/10.1126/science.aat1871>
- 1252 Plum, N.M., Filser, J., 2008. Worms and wetland water: the role of lumbricids and enchytraeids in

- 1253 nutrient mobilization from flooded soils, undefined.
- 1254 Poot, A., Gillissen, F., Koelmans, A.A., 2007. Effects of flow regime and flooding on heavy metal
1255 availability in sediment and soil of a dynamic river system. *Environ. Pollut.* 148, 779–787.
1256 <https://doi.org/10.1016/j.envpol.2007.01.045>
- 1257 Prudhomme, C., Davies, H., 2009. Assessing uncertainties in climate change impact analyses on the
1258 river flow regimes in the UK. Part 2: Future climate. *Clim. Change* 93, 197–222.
1259 <https://doi.org/10.1007/s10584-008-9461-6>
- 1260 Prunty, L., Bell, J., 2005. Soil Temperature Change over Time during Infiltration. *Soil Sci. Soc. Am. J.*
1261 69, 766–775. <https://doi.org/10.2136/sssaj2004.0219>
- 1262 Puchalski, W., 2003. THE IMPORTANCE OF ACCUMULATED ORGANIC MATTER IN NUTRIENT
1263 TRANSPORT AND STORAGE IN FLOODPLAINS, in: International Conference “EcoFlood: Towards
1264 Natural Flood Reduction Strategies”, Warsaw, EU Framework 5.
- 1265 Pulchalski, W., 2003. The Importance of Accumulated Organic Matter in Nutrient Transport and
1266 Storage in Floodplains [WWW Document]. URL
1267 https://www.researchgate.net/publication/237333548_THE_IMPORTANCE_OF_ACCUMULATE
1268 [D_ORGANIC_MATTER_IN_NUTRIENT_TRANSPORT_AND_STORAGE_IN_FLOODPLAINS](https://www.researchgate.net/publication/237333548_THE_IMPORTANCE_OF_ACCUMULATE) (accessed
1269 2.24.20).
- 1270 Pulley, S., Collins, A.L., 2019. Field-based determination of controls on runoff and fine sediment
1271 generation from lowland grazing livestock fields. *J. Environ. Manage.* 249, 109365.
1272 <https://doi.org/10.1016/j.jenvman.2019.109365>
- 1273 Qiao, P., Lei, M., Yang, S., Yang, J., Zhou, X., Dong, N., Guo, G., 2019. Development of a model to
1274 simulate soil heavy metals lateral migration quantity based on SWAT in Huanjiang watershed,
1275 China. *J. Environ. Sci.* 77, 115–129. <https://doi.org/10.1016/J.JES.2018.06.020>
- 1276 Rawlins, B.G., Wragg, J., Lark, R.M., 2013. Application of a novel method for soil aggregate stability
1277 measurement by laser granulometry with sonication. *Eur. J. Soil Sci.* 64, 92–103.
1278 <https://doi.org/10.1111/ejss.12017>
- 1279 Rennert, T., Rabus, W., Rinklebe, J., 2017. Modelling the concentrations of dissolved contaminants
1280 (Cd, Cu, Ni, Pb, Zn) in floodplain soils. *Environ. Geochem. Health* 39, 331–344.
1281 <https://doi.org/10.1007/s10653-016-9859-4>

- 1282 Resongles, E., Casiot, C., Freyrier, R., Le Gall, M., Elbaz-Poulichet, F., 2015. Variation of dissolved and
1283 particulate metal(loid) (As, Cd, Pb, Sb, Tl, Zn) concentrations under varying discharge during a
1284 Mediterranean flood in a former mining watershed, the Gardon River (France). *J. Geochemical*
1285 *Explor.* 158, 132–142. <https://doi.org/10.1016/J.GEXPLO.2015.07.010>
- 1286 Rinklebe, J., Du Laing, G., 2011. Factors Controlling the Dynamics of Trace Metals in Frequently
1287 Flooded Soils, in: *Dynamics and Bioavailability of Heavy Metals in the Rootzone*. CRC Press, pp.
1288 245–270. <https://doi.org/10.1201/b10796-10>
- 1289 Rinklebe, J., During, A., Overesch, M., Du Laing, G., Wennrich, R., Stärk, H.J.H.J., Mothes, S., 2010.
1290 Dynamics of mercury fluxes and their controlling factors in large Hg-polluted floodplain areas.
1291 *Environ. Pollut.* 158, 308–318. <https://doi.org/10.1016/j.envpol.2009.07.001>
- 1292 Rinklebe, J., Franke, C., Neue, H.-U., 2007. Aggregation of floodplain soils based on classification
1293 principles to predict concentrations of nutrients and pollutants. *Geoderma* 141, 210–223.
1294 <https://doi.org/10.1016/J.GEODERMA.2007.06.001>
- 1295 Rinklebe, J., Knox, A.S., Paller, M., Knox, A.S., Paller, M., 2016. Potential Mobility, Bioavailability, and
1296 Plant Uptake of Toxic Elements in Temporary Flooded Soils In book: *Trace elements in*
1297 *waterlogged soils and sediments*, 1st ed. CRC Press, Taylor & Francis Group, New York.
1298 <https://doi.org/10.1201/9781315372952-23>
- 1299 Rinklebe, J., Shaheen, S.M., 2017. Redox chemistry of nickel in soils and sediments: A review.
1300 *Chemosphere* 179, 265–278. <https://doi.org/10.1016/J.CHEMOSPHERE.2017.02.153>
- 1301 Rinklebe, J., Wennrich, R., Du Laing, G., Stark, H.J., Mothes, S., 2013. Mercury emissions from
1302 flooded soils and sediments in Germany are an underestimated problem: challenges for
1303 reliable risk assessments and management strategies. *E3S Web Conf.* 1.
- 1304 Rizzo, A., Bresciani, R., Masi, F., Boano, F., Revelli, R., Ridolfi, L., 2018. Flood reduction as an
1305 ecosystem service of constructed wetlands for combined sewer overflow. *J. Hydrol.* 560, 150–
1306 159. <https://doi.org/10.1016/j.jhydrol.2018.03.020>
- 1307 Rodgers, K., Hursthouse, A., Cuthbert, S., 2015. The Potential of Sequential Extraction in the
1308 Characterisation and Management of Wastes from Steel Processing: A Prospective Review. *Int.*
1309 *J. Environ. Res. Public Health* 12, 11724–11755. <https://doi.org/10.3390/ijerph120911724>
- 1310 Rudiš, M., Valenta, P., Valentová, J., Nol, O., 2009. Assessment of the deposition of polluted

- 1311 sediments transferred by a catastrophic flood and related changes in groundwater quality. *J.*
1312 *Hydrol.* 369, 326–335. <https://doi.org/10.1016/J.JHYDROL.2009.02.023>
- 1313 Sánchez-Rodríguez, A.R., Nie, C., Hill, P.W., Chadwick, D.R., Jones, D.L., 2019. Extreme flood events at
1314 higher temperatures exacerbate the loss of soil functionality and trace gas emissions in
1315 grassland. *Soil Biol. Biochem.* 130, 227–236. <https://doi.org/10.1016/J.SOILBIO.2018.12.021>
- 1316 Schaller, N., Kay, A.L., Lamb, R., Massey, N.R., Van Oldenborgh, G.J., Otto, F.E.L., Sparrow, S.N.,
1317 Vautard, R., Yiou, P., Ashpole, I., Bowery, A., Crooks, S.M., Haustein, K., Huntingford, C., Ingram,
1318 W.J., Jones, R.G., Legg, T., Miller, J., Skeggs, J., Wallom, D., Weisheimer, A., Wilson, S., Stott,
1319 P.A., Allen, M.R., 2016. Human influence on climate in the 2014 southern England winter floods
1320 and their impacts. *Nat. Clim. Chang.* 6, 627–634. <https://doi.org/10.1038/nclimate2927>
- 1321 Schulz-Zunkel, C., Krueger, F., 2009. Trace Metal Dynamics in Floodplain Soils of the River Elbe: A
1322 Review. *J. Environ. Qual.* 38, 1349–1362. <https://doi.org/10.2134/jeq2008.0299>
- 1323 Schulz-Zunkel, C., Rinklebe, J., Bork, H.-R., 2015. Trace element release patterns from three
1324 floodplain soils under simulated oxidized–reduced cycles. *Ecol. Eng.* 83, 485–495.
1325 <https://doi.org/10.1016/J.ECOLENG.2015.05.028>
- 1326 Selinus, O., Alloway, B., Centeno, J.A., Robert B. Finkelman, Fuge, R., Lindh, U., Smedley, P., 2005.
1327 *Essentials of Medical Geology: Impacts of the Natural Environment on Public Health.* Elsevier
1328 Academic Press.
- 1329 Shaheen, S.M., Frohne, T., White, J.R., DeLaune, R.D., 2017. Redox-induced mobilization of copper,
1330 selenium, and zinc in deltaic soils originating from Mississippi (U.S.A.) and Nile (Egypt) River
1331 Deltas: A better understanding of biogeochemical processes for safe environmental
1332 management. *J. Environ. Manage.* 186, 131–140.
1333 <https://doi.org/10.1016/J.JENVMAN.2016.05.032>
- 1334 Shaheen, S.M., Rinklebe, J., 2017. Sugar beet factory lime affects the mobilization of Cd, Co, Cr, Cu,
1335 Mo, Ni, Pb, and Zn under dynamic redox conditions in a contaminated floodplain soil. *J.*
1336 *Environ. Manage.* 186, 253–260. <https://doi.org/10.1016/j.jenvman.2016.07.060>
- 1337 Shaheen, S.M., Rinklebe, J., 2014. Geochemical fractions of chromium, copper, and zinc and their
1338 vertical distribution in floodplain soil profiles along the Central Elbe River, Germany. *Geoderma*
1339 228–229, 142–159. <https://doi.org/10.1016/J.GEODERMA.2013.10.012>

- 1340 Shaheen, S.M., Rinklebe, J., Frohne, T., White, J.R., DeLaune, R.D., 2016. Redox effects on release
1341 kinetics of arsenic, cadmium, cobalt, and vanadium in Wax Lake Deltaic freshwater marsh soils.
1342 *Chemosphere* 150, 740–748. <https://doi.org/10.1016/J.CHEMOSPHERE.2015.12.043>
- 1343 Shaheen, S.M., Rinklebe, J., Rupp, H., Meissner, R., 2014a. Temporal dynamics of pore water
1344 concentrations of Cd, Co, Cu, Ni, and Zn and their controlling factors in a contaminated
1345 floodplain soil assessed by undisturbed groundwater lysimeters. *Environ. Pollut.* 191, 223–231.
1346 <https://doi.org/10.1016/J.ENVPOL.2014.04.035>
- 1347 Shaheen, S.M., Rinklebe, J., Rupp, H., Meissner, R., 2014b. Lysimeter trials to assess the impact of
1348 different flood–dry-cycles on the dynamics of pore water concentrations of As, Cr, Mo and V in
1349 a contaminated floodplain soil. *Geoderma* 228–229, 5–13.
1350 <https://doi.org/10.1016/J.GEODERMA.2013.12.030>
- 1351 Shahid, M., Shamshad, S., Rafiq, M., Khalid, S., Bibi, I., Niazi, N.K., Dumat, C., Rashid, M.I., 2017.
1352 Chromium speciation, bioavailability, uptake, toxicity and detoxification in soil-plant system: A
1353 review. *Chemosphere* 178, 513–533. <https://doi.org/10.1016/J.CHEMOSPHERE.2017.03.074>
- 1354 Sherene, T., 2010. Mobility and Transport of Heavy Metals in Polluted Soil Environment. *Biol.*
1355 *Forum—An Int. J.* 2, 112–121.
- 1356 Simmler, M., Bommer, J., Frischknecht, S., Christl, I., Kotsev, T., Kretzschmar, R., 2017. Reductive
1357 solubilization of arsenic in a mining-impacted river floodplain: Influence of soil properties and
1358 temperature. *Environ. Pollut.* 231, 722–731. <https://doi.org/10.1016/J.ENVPOL.2017.08.054>
- 1359 Sipos, P., Choi, C., Németh, T., Szalai, Z., Póka, T., 2014. Relationship between iron and trace metal
1360 fractionation in soils. *Chem. Speciat. Bioavailab.* 26, 21–30.
1361 <https://doi.org/10.3184/095422914X13887685052506>
- 1362 Sivakumar, S., Subbhuraam, C.V., 2005. Toxicity of chromium(III) and chromium(VI) to the
1363 earthworm *Eisenia fetida*. *Ecotoxicol. Environ. Saf.* 62, 93–98.
1364 <https://doi.org/10.1016/J.ECOENV.2004.08.006>
- 1365 Sizmur, T., Hodson, M.E., 2009. Do earthworms impact metal mobility and availability in soil? – A
1366 review. *Environ. Pollut.* 157, 1981–1989. <https://doi.org/10.1016/J.ENVPOL.2009.02.029>
- 1367 Sizmur, T., Palumbo-Roe, B., Watts, M.J., Hodson, M.E., 2011. Impact of the earthworm *Lumbricus*
1368 *terrestris* (L.) on As, Cu, Pb and Zn mobility and speciation in contaminated soils. *Environ.*

- 1369 Pollut. 159, 742–748. <https://doi.org/10.1016/J.ENVPOL.2010.11.033>
- 1370 Sizmur, T., Richardson, J., 2020. Earthworms accelerate the biogeochemical cycling of potentially
1371 toxic elements: Results of a meta-analysis. *Soil Biol. Biochem.* 148, 107865.
1372 <https://doi.org/10.1016/j.soilbio.2020.107865>
- 1373 Spurgeon, D.J., Hopkin, S.P., 1999. Comparisons of metal accumulation and excretion kinetics in
1374 earthworms (*Eisenia fetida*) exposed to contaminated field and laboratory soils. *Appl. Soil Ecol.*
1375 11, 227–243.
- 1376 Srivastava, V., Sarkar, A., Singh, S.K., Singh, P.K., De Araujo, A.S.F., Singh, R.P., 2017. Agroecological
1377 Responses of Heavy Metal Pollution with Special Emphasis on Soil Health and Plant
1378 Performances. *Front. Environ. Sci* 5, 64. <https://doi.org/10.3389/fenvs.2017.00064>
- 1379 Stafford, A., Jeyakumar, P., Hedley, M., Anderson, C., 2018. Influence of Soil Moisture Status on Soil
1380 Cadmium Phytoavailability and Accumulation in Plantain (*Plantago lanceolata*). *Soil Syst.* 2, 9.
1381 <https://doi.org/10.3390/soils2010009>
- 1382 Stagl, J., Mayr, E., Koch, H., Hattermann, F.F., Huang, S., 2014. Effects of Climate Change on the
1383 Hydrological Cycle in Central and Eastern Europe, in: *Advances in Global Change Research*.
1384 Springer International Publishing, pp. 31–43. https://doi.org/10.1007/978-94-007-7960-0_3
- 1385 Stahl, R.G., Hooper, M.J., Balbus, J.M., Clements, W., Fritz, A., Gouin, T., Helm, R., Hickey, C., Landis,
1386 W., Moe, S.J., 2013. The influence of global climate change on the scientific foundations and
1387 applications of Environmental Toxicology and Chemistry: Introduction to a SETAC international
1388 workshop. *Environ. Toxicol. Chem.* 32, 13–19. <https://doi.org/10.1002/etc.2037>
- 1389 Steinnes, E., 2013. Heavy metal contamination of the terrestrial environment from long-range
1390 atmospheric transport: Evidence from 35 years of research in Norway. *E3S Web Conf.* 1, 35001.
1391 <https://doi.org/10.1051/E3SCONF/20130135001>
- 1392 Stuart, M., Lapworth, D., 2011. A review of processes important in the floodplain setting. *Br. Geol.*
1393 *Surv.* 32 (OR/11/030).
- 1394 Stürck, J., Poortinga, A., Verburg, P.H., 2014. Mapping ecosystem services: The supply and demand
1395 of flood regulation services in Europe. *Ecol. Indic.* 38, 198–211.
1396 <https://doi.org/10.1016/j.ecolind.2013.11.010>
- 1397 Sullivan, T.S., Gadd, G.M., 2019. Metal bioavailability and the soil microbiome, in: *Advances in*

- 1398 Agronomy. Academic Press Inc., pp. 79–120. <https://doi.org/10.1016/bs.agron.2019.01.004>
- 1399 Tack, F.M., Verloo, M.G., 1995. Chemical Speciation and Fractionation in Soil and Sediment Heavy
1400 Metal Analysis: A Review, Chemical. International Journal of Environmental Analytical
1401 Chemistry.
- 1402 Tack, F.M.G., 2010. Trace elements: General soil chemistry, principles and processes, in: Trace
1403 Elements in Soils. John Wiley and Sons, pp. 9–37. <https://doi.org/10.1002/9781444319477.ch2>
- 1404 Tack, F.M.G., Van Ranst, E., Lievens, C., Vandenberghe, R.E., 2006. Soil solution Cd, Cu and Zn
1405 concentrations as affected by short-time drying or wetting: The role of hydrous oxides of Fe
1406 and Mn. *Geoderma* 137, 83–89. <https://doi.org/10.1016/j.geoderma.2006.07.003>
- 1407 Tedeschi, R.G., Collins, M., 2016. The influence of ENSO on South American precipitation during
1408 austral summer and autumn in observations and models. *Int. J. Climatol.* 36, 618–635.
1409 <https://doi.org/10.1002/joc.4371>
- 1410 Thapa, R., Thoms, M.C., Parsons, M., Reid, M., 2016. Adaptive cycles of floodplain vegetation
1411 response to flooding and drying. *Earth Surf. Dyn.* 4, 175–191. [https://doi.org/10.5194/esurf-4-](https://doi.org/10.5194/esurf-4-175-2016)
1412 [175-2016](https://doi.org/10.5194/esurf-4-175-2016)
- 1413 Thomas, S.K., Severson, E.D., Galbraith, J.M., 2016. Measuring Saturated Hydraulic Conductivity in
1414 Soil. Virginia Coop. Extension, Blacksburg, VA. 1–13.
- 1415 Tockner, K., Lorang, M.S., Stanford, J.A., 2010. River flood plains are model ecosystems to test
1416 general hydrogeomorphic and ecological concepts. *River Res. Appl.* 26, 76–86.
1417 <https://doi.org/10.1002/rra.1328>
- 1418 Tockner, K., Stanford, J.A., 2002. Riverine flood plains: Present state and future trends. *Environ.*
1419 *Conserv.* 29, 308–330. <https://doi.org/10.1017/S037689290200022X>
- 1420 Tóth, G., Hermann, T., Da Silva, M.R., Montanarella, L., 2016a. Heavy metals in agricultural soils of
1421 the European Union with implications for food safety. *Environ. Int.* 88, 299–309.
1422 <https://doi.org/10.1016/J.ENVINT.2015.12.017>
- 1423 Tóth, G., Hermann, T., Szatmári, G., Pásztor, L., 2016b. Maps of heavy metals in the soils of the
1424 European Union and proposed priority areas for detailed assessment. *Sci. Total Environ.* 565,
1425 1054–1062. <https://doi.org/10.1016/J.SCITOTENV.2016.05.115>

- 1426 Unger, I.M., Kennedy, A.C., Muzika, R.-M., 2009. Flooding effects on soil microbial communities.
1427 *Appl. Soil Ecol.* 42, 1–8. <https://doi.org/10.1016/J.APSSOIL.2009.01.007>
- 1428 Vaughan, K.L., Rabenhorst, M.C., Needelman, B.A., 2009. Saturation and Temperature Effects on the
1429 Development of Reducing Conditions in Soils. *Soil Sci. Soc. Am. J.* 73, 663–667.
1430 <https://doi.org/10.2136/sssaj2007.0346>
- 1431 Vijver, M.G., Vink, J.P.M., Miermans, C.J.H., van Gestel, C.A.M., 2007. Metal accumulation in
1432 earthworms inhabiting floodplain soils. *Environ. Pollut.* 148, 132–140.
1433 <https://doi.org/10.1016/j.envpol.2006.10.046>
- 1434 Vink, J.P.M., Meeussen, J.C.L., 2007. BIOCHEM–ORCHESTRA: A tool for evaluating chemical
1435 speciation and ecotoxicological impacts of heavy metals on river flood plain systems. *Environ.*
1436 *Pollut.* 148, 833–841. <https://doi.org/10.1016/J.ENVPOL.2007.01.041>
- 1437 Violante, A., 2013. Elucidating Mechanisms of Competitive Sorption at the Mineral/Water Interface.
1438 *Adv. Agron.* 118, 111–176. <https://doi.org/10.1016/B978-0-12-405942-9.00003-7>
- 1439 Violante, A., Cozzolino, V., Perelomov, L., Caporale, A.G., Pigna, M., 2010. Mobility and bioavailability
1440 of heavy metals and metalloids in soil environments. *J. Soil Sci. Plant Nutr.* 10, 268–292.
1441 <https://doi.org/10.4067/S0718-95162010000100005>
- 1442 Visser, A., Kroes, J., Van Vliet, M.T.H., Blenkinsop, S., Fowler, H.J., Broers, H.P., 2012. Climate change
1443 impacts on the leaching of a heavy metal contamination in a small lowland catchment, in:
1444 *Journal of Contaminant Hydrology*. Elsevier, pp. 47–64.
1445 <https://doi.org/10.1016/j.jconhyd.2011.04.007>
- 1446 Weber, F.-A., Voegelin, A., Kretzschmar, R., 2009. Multi-metal contaminant dynamics in temporarily
1447 flooded soil under sulfate limitation. *Geochim. Cosmochim. Acta* 73, 5513–5527.
1448 <https://doi.org/10.1016/J.GCA.2009.06.011>
- 1449 Wheeler, H., Evans, E., 2009. Land use, water management and future flood risk. *Land use policy* 26,
1450 S251–S264. <https://doi.org/10.1016/J.LANDUSEPOL.2009.08.019>
- 1451 Wijngaard, R.R., van der Perk, M., van der Grift, B., de Nijs, T.C.M., Bierkens, M.F.P., 2017. The
1452 Impact of Climate Change on Metal Transport in a Lowland Catchment. *Water. Air. Soil Pollut.*
1453 228, 107. <https://doi.org/10.1007/s11270-017-3261-4>
- 1454 Wilby, R.L., Beven, K.J., Reynard, N.S., 2008. Climate change and fluvial flood risk in the UK: more of

- 1455 the same? *Hydrol. Process.* 22, 2511–2523. <https://doi.org/10.1002/hyp.6847>
- 1456 Winger, P.V. V, Lasier, P.J.J., Jackson, B.P.P., 1998. The Influence of Extraction Procedure on Ion
1457 Concentrations in Sediment Pore Water. *Arch. Environ. Contam. Toxicol.* 35, 8–13.
1458 <https://doi.org/10.1007/s002449900341>
- 1459 Wright, A.J., de Kroon, H., Visser, E.J.W., Buchmann, T., Ebeling, A., Eisenhauer, N., Fischer, C.,
1460 Hildebrandt, A., Ravenek, J., Roscher, C., Weigelt, A., Weisser, W., Voeselek, L.A.C.J., Mommer,
1461 L., 2017. Plants are less negatively affected by flooding when growing in species-rich plant
1462 communities. *New Phytol.* 213, 645–656. <https://doi.org/10.1111/nph.14185>
- 1463 Wuana, R.A., Okieimen, F.E., Montuelle, B., Steinman, A.D., 2011. Heavy Metals in Contaminated
1464 Soils: A Review of Sources, Chemistry, Risks and Best Available Strategies for Remediation. *Int.*
1465 *Sch. Res. Netw. ISRN Ecol.* 2011, 20. <https://doi.org/10.5402/2011/402647>
- 1466 Wyzkowska, J., Borowik, A., Kucharski, M., Kucharski, J., 2013. Effect of cadmium, copper and zinc
1467 on plants, soil microorganisms and soil enzymes. *J. Elem.*
1468 <https://doi.org/10.5601/jelem.2013.18.4.455>
- 1469 Xu, J., Zheng, L., Xu, L., Wang, X., 2020. Uptake and allocation of selected metals by dominant
1470 vegetation in Poyang Lake wetland: From rhizosphere to plant tissues. *Catena* 189, 104477.
1471 <https://doi.org/10.1016/j.catena.2020.104477>
- 1472 Xu, J.M., Tang, C., Chen, Z.L., 2006. The role of plant residues in pH change of acid soils differing in
1473 initial pH. *Soil Biol. Biochem.* 38, 709–719. <https://doi.org/10.1016/j.soilbio.2005.06.022>
- 1474 Xu, Y., Feng, L., Jeffrey, P.D., Shi, Y., Morel, F.M.M., 2008. Structure and metal exchange in the
1475 cadmium carbonic anhydrase of marine diatoms. *Nature* 452, 56–61.
1476 <https://doi.org/10.1038/nature06636>
- 1477 Yang, C., Li, S., Liu, R., Sun, P., Liu, K., 2015. Effect of reductive dissolution of iron (hydr)oxides on
1478 arsenic behavior in a water-sediment system: First release, then adsorption. *Ecol. Eng.* 83, 176–
1479 183. <https://doi.org/10.1016/j.ecoleng.2015.06.018>
- 1480 Zhao, Y., Marriott, S., Rogers, J., Iwugo, K., 1999. A preliminary study of heavy metal distribution on
1481 the floodplain of the River Severn, UK by a single flood event. *Sci. Total Environ.* 243–244, 219–
1482 231. [https://doi.org/10.1016/S0048-9697\(99\)00386-1](https://doi.org/10.1016/S0048-9697(99)00386-1)
- 1483 Zhao, Y., Marriott, S.B., 2013. Dispersion and Remobilisation of Heavy Metals in the River Severn

1484 System, UK. *Procedia Environ. Sci.* 18, 167–173.

1485 <https://doi.org/10.1016/J.PROENV.2013.04.022>

1486 Zia, A., van den Berg, L., Ahmad, M.N., Riaz, M., Zia, D., Ashmore, M., 2018. Controls on
1487 accumulation and soil solution partitioning of heavy metals across upland sites in United
1488 Kingdom (UK). *J. Environ. Manage.* 222, 260–267.

1489 <https://doi.org/10.1016/J.JENVMAN.2018.05.076>

1490

1491

1492

1493