Received: 29 October 2019

Revised: 10 March 2020

Accepted: 16 March 2020

DOI: 10.1002/mus.26864

## CLINICAL RESEARCH ARTICLE



## Fasciculations demonstrate daytime consistency in amyotrophic lateral sclerosis

James Bashford MRCP, PhD<sup>1</sup> | Urooba Masood MSc<sup>1</sup> | Aidan Wickham PhD<sup>2</sup> | Raquel Iniesta PhD<sup>3</sup> | Emmanuel Drakakis PhD<sup>2</sup> | Martyn Boutelle PhD<sup>2</sup> | Kerry Mills FRCP, PhD<sup>1</sup> | Chris Shaw FRCP, MD<sup>1</sup>

<sup>1</sup>UK Dementia Research Institute, Department of Basic and Clinical Neuroscience, Maurice Wohl Clinical Neuroscience Institute, Institute of Psychiatry, Psychology and Neuroscience, King's College, London, UK

<sup>2</sup>Department of Bioengineering, Imperial College London, London, UK

<sup>3</sup>Department of Biostatistics and Health Informatics, King's College, London, UK

## **Abstract**

**Introduction:** Fasciculations represent early neuronal hyperexcitability in amyotrophic lateral sclerosis (ALS). To aid calibration as a disease biomarker, we set out to characterize the daytime variability of fasciculation firing.

Methods: Fasciculation awareness scores were compiled from 19 ALS patients. In addition, 10 ALS patients prospectively underwent high-density surface electromyoanarabia (LIDCENAC) na agradina a fugus bisana and asatus angusina at thura tima a sainta