

Stable fractional matchings

IOANNIS CARAGIANNIS, University of Patras & CTI “Diophantus”, Greece

ARIS FILOS-RATSIKAS, École Polytechnique Fédérale de Lausanne, Switzerland

PANAGIOTIS KANELLOPOULOS, University of Patras & CTI “Diophantus”, Greece

ROHIT VAISH, Rensselaer Polytechnic Institute, USA

We study a generalization of the classical stable matching problem that allows for *cardinal* preferences (as opposed to ordinal) and *fractional* matchings (as opposed to integral). After observing that, in this cardinal setting, stable fractional matchings can have much higher social welfare than stable integral ones, our goal is to understand the computational complexity of finding an optimal (i.e., welfare-maximizing) or nearly-optimal stable fractional matching. We present simple approximation algorithms for this problem with weak welfare guarantees and, rather unexpectedly, we furthermore show that achieving better approximations is hard. This computational hardness persists even for approximate stability. To the best of our knowledge, these are the first computational complexity results for stable fractional matchings. En route to these results, we provide a number of structural observations.

1 INTRODUCTION

The stable matching problem [15] is one of the most extensively studied problems at the interface of economics and computer science, with notable practical applications such as matching students to schools [1], medical graduates to hospitals [29], and organ donors to patients [31]. The input to the problem consists of the preference lists of two sets of agents, commonly referred to as the *men* and the *women*. The goal is to find a *stable* matching, i.e., a matching in which no pair of man and woman prefer each other over their assigned partners.

The standard formulation of the stable matching problem involves two key assumptions: First, that the matching is *integral* (i.e., two agents are either completely matched or completely unmatched), and second, that agents have *ordinal* preferences (typically in the form of rank-ordered lists). Although these assumptions suffice in a number of applications (including those mentioned above), there are natural examples where they turn out to be inadequate. For instance, time-sharing applications [30] naturally give rise to fractional matchings: Imagine a scenario where a set of newly hired employees are matched with a set of supervisors. Assuming that each individual can spend one unit of time at work, an *integral* matching prescribes that every employee should work full-time with a single supervisor. On the other hand, *fractional* matchings allow the employees to divide their time in working with multiple supervisors, making them a more natural modeling choice in such situations.




Fig. 1. Instance with cardinal preferences.

Likewise, *ordinal* preferences, despite their simplicity and ease of elicitation, can often be quite restrictive. Indeed, in many real-world matching applications (e.g., labor markets), the outcomes experienced by the participants are inherently *cardinal* in nature (e.g., wages). In such settings, it is decidedly more natural to model the *intensity* of preferences, as has been noted in theory [3, 28] as well as lab experiments [12]. Encouraged by these insights, and the fact that cardinal utilities provide a clean and unambiguous way of comparing matching outcomes in terms of their social welfare, we consider a generalization of the stable matching model that allows for *fractional* matchings (as opposed to integral) and *cardinal* preferences (as opposed to ordinal).

More concretely, we consider a setting where the preferences are specified in terms of *valuations* (e.g., in the matching instance in Figure 1, m_1 values w_1 at 0 and w_1 values m_1 at 3). A fractional

matching is simply a convex combination of integral matchings, and an agent’s utility under a fractional matching is the appropriately weighted sum of its utilities under the constituent integral matchings. A fractional matching μ is *stable* if no pair of man and woman simultaneously derive greater utility in being integrally matched to each other than they do under μ .

The above generalization has a clear merit in terms of social welfare: Stable solutions under the generalized model can have greater welfare than those under the standard model as the following example illustrates.

Example 1. Consider the instance in Figure 1 with three men m_1, m_2, m_3 and three women w_1, w_2, w_3 . Among the six possible integral matchings, only $\mu_1 := \{(m_1, w_3), (m_2, w_2), (m_3, w_1)\}$ and $\mu_2 := \{(m_1, w_1), (m_2, w_3), (m_3, w_2)\}$ are stable. Indeed, μ_1 and μ_2 are the men-proposing and women-proposing Gale-Shapley matchings respectively [15]. The social welfare (i.e., sum of utilities of all agents) under these matchings is $\mathcal{W}(\mu_1) = \mathcal{W}(\mu_2) = 7$.

Define $\mu_3 := \{(m_1, w_1), (m_2, w_2), (m_3, w_3)\}$, and notice that $\mathcal{W}(\mu_3) = 8$. Now consider a *fractional* matching $\mu := \frac{1}{2}\mu_2 + \frac{1}{2}\mu_3$. The social welfare of μ is $\mathcal{W}(\mu) = \frac{1}{2}\mathcal{W}(\mu_2) + \frac{1}{2}\mathcal{W}(\mu_3) = 15/2 > 7 = \mathcal{W}(\mu_1) = \mathcal{W}(\mu_2)$, i.e., μ has a higher social welfare than any stable integral matching. Importantly, μ is a *stable fractional* matching. Indeed, under μ , the utilities of m_1, m_2, m_3, w_1, w_2 , and w_3 are 0, 1/2, 3/2, 3, 3/2, and 1 respectively. Thus, for every man-woman pair, at least one of the two agents meets the corresponding utility threshold for that pair, implying that μ is stable.

Overall, the instance in Figure 1 admits a stable fractional matching with strictly greater welfare than any stable integral matching. \square

Starting with the seminal work of Gale and Shapley [15], there is now an extensive literature on algorithms for computing stable solutions, including ones that optimize a variety of objectives pertaining to fairness and economic efficiency [16, 19, 23, 27, 34]. Most of these algorithms, however, are tailored to compute stable *integral* matchings. As Example 1 demonstrates, such algorithms could, in general, return highly suboptimal outcomes in our setting. Therefore, it becomes pertinent to understand the computational complexity of finding an “optimal” stable matching in the generalized model. Our work studies this question from the lens of the fundamental objective of social welfare, and asks the following natural question:

Can we efficiently compute an optimal or nearly optimal stable fractional matching?

1.1 Our results and roadmap

We formalize this question by defining the optimization problem OPTIMAL STABLE FRACTIONAL MATCHING. We demonstrate its importance by strengthening the observation in Example 1 and showing that the social welfare gap between the best stable fractional and best stable integral matchings can be arbitrarily large. We then mitigate this very positive message by identifying important characteristics of optimal and nearly-optimal stable fractional matchings. These include a non-convexity property and, more crucially, necessary conditions requiring that such matchings are formed by convex combinations of many unstable integral matchings. This already narrows down the range of tools we can use for the design of efficient (approximation) algorithms. Still, we present simple algorithms for OPTIMAL STABLE FRACTIONAL MATCHING with an approximation ratio of $1 + \sigma_{\max}/\sigma_{\min}$, where σ_{\max} and σ_{\min} represent the maximum and minimum positive valuation in the input instance. For the variant OPTIMAL ε -STABLE FRACTIONAL MATCHING, where the stability constraints are relaxed by an ε factor, an embarrassingly simple algorithm computes $1/\varepsilon$ -approximate solutions. Rather unexpectedly, we show that these approximation guarantees are almost best possible by polynomial-time algorithms (unless $P = NP$). To the best of our knowledge, these are the first computational complexity results for stable fractional matchings.

The rest of the paper is structured as follows. We present related work in the matching literature in Section 1.2. We continue in Section 2 with preliminary definitions and warm up with exponential-time algorithms that solve OPTIMAL STABLE FRACTIONAL MATCHING using linear programming. Structural properties of optimal and nearly-optimal solutions to OPTIMAL STABLE FRACTIONAL MATCHING are presented in Section 3. Our algorithms are presented in Section 4. Our inapproximability results are stated and partially proved in Section 5. Some proofs and additional material appear in appendix.

1.2 Further related work

The stable marriage problem has been extensively studied over the years [15, 17, 26, 32] with several interesting results in the original model of [15] as well as several of its variants [8, 18, 20, 22, 27]. While the Gale-Shapley algorithm can find a stable matching in polynomial time, several of these variants are computationally more challenging [27]; we refer the interested reader to the survey of Iwama and Miyazaki [21] for a more detailed exposition. Starting with the works of Vande Vate [35], Rothblum [33], and Roth et al. [30], there is by now a well-developed literature on the linear programming formulations of the stable matching problem [34, 36], and various combinatorial algorithms that optimize fairness-related objectives are also known [7, 14, 16, 19, 23]. However, the majority of this work studies integral stable matchings under ordinal preferences.

The theoretical and practical importance of modeling agents’ cardinal preferences in stable matching settings has been highlighted, among others, by Anshelevich et al. [4] and Pini et al. [28]. Anshelevich et al. [4] formulate the notions of exact/approximate stability in terms of cardinal preferences that are central to our work; see Definition 1. They study the “price of anarchy” for stable matchings (defined as the ratio of social welfare of the optimal matching and the worst stable integral matching) under various preference structures as well as its extensions to approximate stability. A similar approach has been adopted in related settings [2, 13]. Pini et al. [28] consider a notion of stability similar to [4], but focus on achieving economic efficiency (in particular, Pareto optimality and its variants) along with stability. They also study strategic aspects which are an exciting avenue for future research even in our model.

In terms of computing efficient (i.e., welfare-maximizing or cost-minimizing) stable matchings, Irving et al. [19], Manlove et al. [27], and Mai and Vazirani [24] provide efficient algorithms and/or inapproximability results, but crucially, these results apply to *integral* matchings only. To the best of our knowledge, the computational questions associated with computing stable *fractional* matchings have not been considered prior to this work.

Deligkas et al. [9] study matchings computed by the natural greedy algorithm in edge-weighted graphs. Among other results, they show that the problem of computing the maximum-weight greedy matching in a bipartite graph is NP-hard. The greedy matchings in their model are stable integral matchings in ours. Still, as we discuss in Appendix A.2, this result implies the NP-hardness of OPTIMAL STABLE FRACTIONAL MATCHING. We note that our inapproximability results are much stronger.

Our definition of fractional stability (in the presence of cardinal preferences) has appeared before in the economics literature [5, 10, 11, 25]. However, these papers focus primarily on the relationship among various notions of stability and economic efficiency, and do not consider computational questions. We remark here that the term “fractional stability” has been overloaded in the related literature, as it has also been used to refer to fractional matchings that only have integral stable matchings in their support [34]; note that stability in the latter context can be defined purely in terms of the ordinal preferences. This notion of “ex-post” stability is fundamentally different from ours, as it is precisely the existence of unstable integral matchings in the support of the stable fractional matching (see Proposition 3) that allows for the large gain in welfare in Example 1 and

Theorem 1. If one is interested in the ex-post notion of stability, then the computational problem clearly reduces to the case of integral matchings.

2 PRELIMINARIES

An instance of *Stable Matching problem with Cardinal preferences* (SMC) is given by the tuple $\langle M, W, U, V \rangle$, where $M := \{m_1, \dots, m_n\}$ and $W := \{w_1, \dots, w_n\}$ denote the set of n men and n women, respectively, and U and V are $n \times n$ matrices of non-negative rational numbers that specify the *valuations* of the agents. Specifically, $U(m, w)$ is the value derived by man m from his match with woman w , and $V(m, w)$ is the value derived by woman w from her match with man m . Many of our results will focus on two special classes of valuations, namely *binary* (where $U, V \in \{0, 1\}^{n \times n}$) and *ternary* valuations (where $U, V \in \{0, 1, \alpha\}^{n \times n}$ for some $\alpha > 1$).

We will often describe an SMC instance using its *graph representation*. An instance $\mathcal{I} = \langle M, W, U, V \rangle$ can be represented as a bipartite graph with vertex sets M and W , and an edge for every pair $(m, w) \in M \times W$ such that at least one of $U(m, w) > 0$ or $V(m, w) > 0$ holds. Each edge (m, w) in this graph has two valuations associated with it, namely $U(m, w)$ and $V(m, w)$.

A *fractional matching* $\mu : M \times W \rightarrow \mathbb{R}_{\geq 0}$ is an assignment of non-negative weights to all man-woman pairs such that $\sum_{w \in W} \mu(m, w) \leq 1$ for each $m \in M$ and $\sum_{m \in M} \mu(m, w) \leq 1$ for each $w \in W$. A fractional matching μ is said to be *complete* if $\sum_{w \in W} \mu(m, w) = 1$ for each man $m \in M$ and $\sum_{m \in M} \mu(m, w) = 1$ for each woman $w \in W$. An *integral matching* μ is a fractional matching with weights $\mu(m, w) \in \{0, 1\}$ for every pair (m, w) . With slight abuse of notation, we sometimes view an integral matching μ as a set of pairs and write $(m, w) \in \mu$ instead of $\mu(m, w) = 1$.

It is well-known, and follows from the Birkhoff-von Neumann (BvN) theorem, that a fractional matching μ can be decomposed into a convex combination of $k = O(n^2)$ integral matchings $\mu^{(1)}, \mu^{(2)}, \dots, \mu^{(k)}$ so that for every pair $(m, w) \in M \times W$, we have

$$\mu(m, w) = \sum_{j=1}^k \lambda_j \cdot \mu^{(j)}(m, w),$$

where $\lambda_j > 0$ for all $j \in \{1, \dots, k\}$ and $\sum_{j=1}^k \lambda_j = 1$. The set of integral matchings $\{\mu^{(1)}, \dots, \mu^{(k)}\}$ is called the *support* of the fractional matching μ . Note that the support need not be unique.

We proceed with the formal definitions of *stability* and *approximate stability*, which in turn use the definition of the *utility* derived by agents under a fractional matching. In particular, the utility derived by the man m under μ is given by $u_m(\mu) := \sum_{w \in W} U(m, w)\mu(m, w)$, and the utility derived by the woman w is given by $v_w(\mu) := \sum_{m \in M} V(m, w)\mu(m, w)$.

Definition 1 (Stability). Given a fractional matching μ , a man-woman pair (m, w) is said to be a *blocking pair* if $u_m(\mu) < U(m, w)$ and $v_w(\mu) < V(m, w)$. A fractional matching μ is *stable* if there are no blocking pairs, i.e., for each $(m, w) \in M \times W$, either $u_m(\mu) \geq U(m, w)$ or $v_w(\mu) \geq V(m, w)$.

Definition 2 (ε -Stability). Given any $\varepsilon \in [0, 1)$ and a fractional matching μ , a man-woman pair (m, w) is said to be *ε -blocking* if $u_m(\mu) < (1 - \varepsilon)U(m, w)$ and $v_w(\mu) < (1 - \varepsilon)V(m, w)$; otherwise, the pair is said to be *ε -stable*. A fractional matching μ is *ε -stable* if all pairs are ε -stable.

A stable fractional matching is also ε -stable for every $\varepsilon \geq 0$. The next statement follows from the seminal result of Gale and Shapley [15].

PROPOSITION 1. *Given any SMC instance \mathcal{I} , a stable fractional matching μ for \mathcal{I} always exists and can be computed in polynomial time.*

Proposition 1 was originally proven in [15] in the standard stable matching model with ordinal preferences and integral matchings. It is easy to see that given any SMC instance \mathcal{I} , if an integral matching μ is stable for an ordinal instance derived from \mathcal{I} (where the ordinal preferences of each

agent are consistent with its valuations, breaking ties arbitrarily), then it is also stable for the original instance \mathcal{I} .

Next, we define *social welfare*, which is a measure of the efficiency of a fractional matching.

Definition 3 (Social welfare). Given an SMC instance $\langle M, W, U, V \rangle$ and a fractional matching μ , the *social welfare* of μ is defined as

$$\mathcal{W}(\mu) := \sum_{m \in M} u_m(\mu) + \sum_{w \in W} v_w(\mu) = \sum_{m \in M} \sum_{w \in W} (U(m, w) + V(m, w))\mu(m, w).$$

An *optimal* matching is one with the highest social welfare among all fractional matchings. It follows from the BvN decomposition that there is always an integral optimal matching. Similarly, an *optimal stable* fractional matching (respectively, *optimal ε -stable* fractional matching) is one with the highest social welfare among all stable (respectively, all ε -stable) fractional matchings. We will use OPTIMAL STABLE FRACTIONAL MATCHING and OPTIMAL ε -STABLE FRACTIONAL MATCHING to refer to the corresponding optimization problems. For $\rho \in (0, 1]$, the term ρ -*efficient* refers to a stable (respectively, ε -stable) fractional matching with welfare at least ρ times the welfare of the optimal stable (respectively, ε -stable) fractional matching. Thus, an optimal stable (or ε -stable) fractional matching is 1-efficient.

2.1 Computing optimal stable fractional matchings

We will now discuss two exponential-time algorithms for OPTIMAL STABLE FRACTIONAL MATCHING. The first algorithm uses the following mixed integer linear program (OPT-Stab):

$$\begin{aligned} \text{(OPT-Stab)} \quad & \text{maximize} \quad \sum_{m \in M} u_m + \sum_{w \in W} v_w \\ & \text{subject to} \quad u_m \geq U(m, w)y(m, w) && \forall m \in M, w \in W && (1) \\ & \quad v_w \geq V(m, w)(1 - y(m, w)) && \forall m \in M, w \in W && (2) \\ & \quad u_m = \sum_{w \in W} U(m, w)\mu(m, w) && \forall m \in M && (3) \\ & \quad v_w = \sum_{m \in M} V(m, w)\mu(m, w) && \forall w \in W && (4) \\ & \quad \sum_{w \in W} \mu(m, w) \leq 1 && \forall m \in M && (5) \\ & \quad \sum_{m \in M} \mu(m, w) \leq 1 && \forall w \in W && (6) \\ & \quad \mu(m, w) \geq 0 && \forall m \in M, w \in W && (7) \\ & \quad y(m, w) \in \{0, 1\} && \forall m \in M, w \in W && (8) \end{aligned}$$

The non-negative weights $\mu(m, w)$ of man-woman pairs as well as the utilities $u_m := u_m(\mu)$ and $v_w := v_w(\mu)$ of the agents (set in equalities (3) and (4)) are the fractional variables of (OPT-Stab). The binary variables $y(m, w)$ encode the stability requirements for pair (m, w) in constraints (1) and (2). Indeed, by setting $y(m, w)$ to 1 or 0, we can require either $u_m(\mu) \geq U(m, w)$ or $v_w(\mu) \geq V(m, w)$. Constraints (5) and (6) ensure feasibility. By enumerating over all possible combinations of values for the binary variables $y(m, w)$ for $(m, w) \in M \times W$, we get 2^{n^2} different linear programs, and, clearly, at least one of them must have the optimal stable fractional matching as its optimal solution.

Our second algorithm is slightly faster and solves at most $O(n^n)$ linear programs. It exploits the following linear program (OPT-Thresh), which is defined using non-negative constants θ_m for $m \in M$ and θ_w for $w \in W$, which we call *utility thresholds*.

$$\begin{aligned} \text{(OPT-Thresh)} \quad & \text{maximize} \quad \sum_{m \in M} u_m + \sum_{w \in W} v_w \\ & \text{subject to} \quad u_m \geq \theta_m && \forall m \in M && (9) \\ & \quad v_w \geq \theta_w && \forall w \in W && (10) \end{aligned}$$

$$u_m = \sum_{w \in W} U(m, w) \mu(m, w) \quad \forall m \in M \quad (11)$$

$$v_w = \sum_{m \in M} V(m, w) \mu(m, w) \quad \forall w \in W \quad (12)$$

$$\sum_{w \in W} \mu(m, w) \leq 1 \quad \forall m \in M \quad (13)$$

$$\sum_{m \in M} \mu(m, w) \leq 1 \quad \forall w \in W \quad (14)$$

$$\mu(m, w) \geq 0 \quad \forall m \in M, w \in W \quad (15)$$

When all utility thresholds are set to zero, the solution of (OPT-Threshold) is an optimal (i.e., welfare-maximizing) fractional matching. Using (OPT-Threshold) to maximize social welfare under stability constraints is more challenging. We say that a set of utility thresholds is *stability-preserving* if for every pair of agents $m \in M$ and $w \in W$, either $\theta_m \geq U(m, w)$ or $\theta_w \geq V(m, w)$. Then, any fractional matching μ that is feasible for (OPT-Stab) is also feasible for (OPT-Threshold) for some stability-preserving set of utility thresholds. Conversely, any fractional matching μ that is feasible for (OPT-Threshold) with some set of stability-preserving utility thresholds is also feasible for (OPT-Stab). Therefore, in order to solve OPTIMAL STABLE FRACTIONAL MATCHING, it suffices to enumerate all $\mathcal{O}(n^n)$ sets of utility thresholds with $\theta_m \in \{U(m, w) : w \in W\}$ for every man $m \in M$, compute utility thresholds $\theta_w \in \{V(m, w) : m \in M\}$ for all $w \in W$ so that the utility thresholds are stability-preserving, and solve (OPT-Threshold). Among these solutions, the fractional matching with highest social welfare will be the solution of OPTIMAL STABLE FRACTIONAL MATCHING.

3 STRUCTURAL PROPERTIES

In this section, we present several observations about the structure of optimal and nearly-optimal stable fractional matchings. We begin by considerably strengthening our observation in Example 1 regarding the welfare gap between stable fractional and stable integral matchings.

THEOREM 1. *For every $\delta > 0$ and $\alpha \geq 2$, there exists an SMC instance with ternary valuations in $\{0, 1, \alpha\}$ and an optimal stable fractional matching μ^* such that any stable integral matching μ^s satisfies $\mathcal{W}(\mu^s) \leq (\alpha - \frac{1}{2} - \delta)^{-1} \mathcal{W}(\mu^*)$.*

We emphasize that Theorem 1 is a *positive* result as it establishes that stable fractional matchings can have much higher welfare than their integral counterparts, and highlights the importance of OPTIMAL STABLE FRACTIONAL MATCHING. The proof of Theorem 1 appears in Appendix A.1.

Our next observation (Proposition 2) shows that the set of stable fractional matchings can be *non-convex* even for binary valuations. Interestingly, this does not prevent us from efficiently solving OPTIMAL STABLE FRACTIONAL MATCHING in this setting (see Theorem 4 in Section 4).




Fig. 2. The SMC instances used in the proofs of Propositions 2 and 3.

PROPOSITION 2. *There exists an SMC instance with binary valuations for which the set of stable fractional matchings is non-convex.*

PROOF. Consider the instance $\mathcal{I} = \langle M, W, U, V \rangle$ with three men m_1, m_2, m_3 and three women w_1, w_2, w_3 , whose graph representation and agent valuations are shown in Figure 2a. Consider the integral matchings $\mu^{(1)} := \{(m_1, w_3), (m_2, w_1), (m_3, w_2)\}$ and $\mu^{(2)} := \{(m_1, w_2), (m_2, w_3), (m_3, w_1)\}$. It is easy to verify that both $\mu^{(1)}$ and $\mu^{(2)}$ are stable for \mathcal{I} . However, the fractional matching $\mu := 0.5\mu^{(1)} + 0.5\mu^{(2)}$ is not stable since (m_2, w_2) is a blocking pair; indeed, $0.5 = u_{m_2}(\mu) < U(m_2, w_2) = 1$ and $0.5 = v_{w_2}(\mu) < V(m_2, w_2) = 1$. \square

The structure of stable fractional matchings becomes much more interesting (and, as we will see in Section 5, also computationally unwieldy) when we move to *ternary* valuations. It turns out that the support of a stable fractional matching can comprise entirely of *unstable* integral matchings (Proposition 3), and its size can grow *linearly* with the input (Theorem 3). These observations pose major limitations on the set of algorithmic tools at our disposal.

PROPOSITION 3. *There exists an SMC instance with ternary valuations and a stable fractional matching μ such that every integral matching in any support of μ is unstable.*

PROOF. Consider the SMC instance $\mathcal{I} = \langle M, W, U, V \rangle$ with three men and three women shown in Figure 2b. The parameter $\alpha \geq 3$ is a constant. There are six different perfect integral matchings:

- Matching $\mu^{(1)}$, which consists of pairs (m_1, w_1) , (m_2, w_2) , and (m_3, w_3) and has a social welfare of 6. It is easy to verify that this is the unique stable integral matching. Also, any subset of $\mu^{(1)}$ is not stable as the pair that is missing from $\mu^{(1)}$ will be blocking.
- Matching $\mu^{(2)}$, which consists of pairs (m_1, w_2) , (m_2, w_3) , and (m_3, w_1) and has a social welfare of 2α . The matching is not stable since the pair (m_1, w_1) is blocking.
- Matching $\mu^{(3)}$, which consists of pairs (m_1, w_3) , (m_2, w_1) , and (m_3, w_2) and has a social welfare of 2α . It is not stable since (m_2, w_2) is blocking.
- Matching $\mu^{(4)}$, which consists of pairs (m_1, w_1) , (m_2, w_3) , and (m_3, w_2) and has a social welfare of $\alpha + 2$. It is not stable since (m_3, w_3) is blocking.
- Matching $\mu^{(5)}$, which consists of pairs (m_1, w_3) , (m_2, w_2) , and (m_3, w_1) and has a social welfare of $2\alpha + 2$. It is not stable since (m_1, w_1) is blocking.
- Matching $\mu^{(6)}$, which consists of pairs (m_1, w_2) , (m_2, w_1) , and (m_3, w_3) and has a social welfare of $\alpha + 2$. It is not stable since the pair (m_2, w_2) is blocking.

Consider the matching $\mu := \frac{1}{\alpha(\alpha-1)} \cdot \mu^{(2)} + \frac{1}{\alpha} \cdot \mu^{(3)} + \frac{\alpha-2}{\alpha-1} \cdot \mu^{(5)}$. It is easy to verify that μ is stable. Indeed, the utilities of the agents under μ are given by $u_{m_1}(\mu) = 0$, $u_{m_2}(\mu) = 1$, $u_{m_3}(\mu) = \alpha - 1$, $v_{w_1}(\mu) = 1$, $v_{w_2}(\mu) = \frac{\alpha-2}{\alpha-1}$ and $v_{w_3}(\mu) = \alpha - \frac{1}{\alpha-1}$. Notice that only the pairs (m_1, w_1) , (m_2, w_2) , and (m_3, w_3) need to be checked for stability, since any other pair has at least one agent with a valuation of zero (and, hence, the stability constraint for those pairs is trivially satisfied). For each of the pairs (m_1, w_1) , (m_2, w_2) , and (m_3, w_3) , there is some member of the pair that has a utility of at least 1 under μ , which meets the requisite stability threshold, implying that μ is stable. Finally, notice that $\mu(m_1, w_1) = 0$, which means that the unique stable integral matching $\mu^{(1)}$ cannot occur in a support of μ . \square

We remark that with some extra work, one can show that the matching μ in the proof of Proposition 3 is the unique optimal stable fractional matching.

As mentioned previously in Section 2, a (stable) fractional matching is the convex combination of at most n^2 integral ones. Theorem 2 provides a stronger bound on the support size of an *optimal* stable fractional matching.

THEOREM 2. *Given any SMC instance \mathcal{I} , there exists an optimal stable fractional matching for \mathcal{I} with at most $4n$ integral matchings in its support.*

PROOF. Let μ^* be an optimal stable fractional matching for \mathcal{I} . Recall from Section 2.1 that μ^* solves the program (OPT-Thresh) for some set of stability-preserving utility thresholds. Observe that (OPT-Thresh) has n^2 free variables (we ignore here the $2n$ variables u_m for $m \in M$ and v_w for $w \in W$, which depend on the remaining ones according to constraints (11) and (12)). Without loss of generality, μ^* is an optimal *extreme point* solution of (OPT-Thresh). That is, when (OPT-Thresh) is instantiated for μ^* , n^2 linearly independent inequality constraints become tight. Among them, at most $4n$ can correspond to the sets of constraints (9), (10), (13), and (14). The remaining ones must correspond to the set of constraints (15), implying that at least $n^2 - 4n$ free variables will be equal to zero. Thus, μ^* can assign positive weights to at most $4n$ man-woman pairs and, consequently, can have at most $4n$ integral matchings in its support. \square

Next we show that the bound in Theorem 2 is tight up to a constant factor.

THEOREM 3. *For every $\rho \in (0, 1]$, there exists a family of SMC instances with ternary valuations for which any support of a ρ -efficient stable fractional matching consists of $\Omega(\rho n)$ integral matchings.*

PROOF. Consider a family of SMC instances $\mathcal{I}_n = \langle M, W, U, V \rangle$ with $M = \{m_1, \dots, m_n\}$ and $W = \{w_1, \dots, w_n\}$, where n is odd. Let α be such that $\alpha > \max \left\{ n + 2, \frac{2n}{\rho(n-1)} \right\}$. The (ternary) valuations of the agents are defined as follows: For each $i \in \{1, 2, \dots, n\}$, $U(m_i, w_i) = V(m_i, w_i) = 1$. For each $i \in \{1, 2, \dots, \frac{n-1}{2}\}$, $U(m_{2i}, w_1) = U(m_{2i+1}, w_{2i}) = V(m_{2i}, w_{2i+1}) = \alpha$ and $V(m_{2i}, w_1) = V(m_{2i+1}, w_{2i}) = U(m_{2i}, w_{2i+1}) = 0$. Finally, $U(m_n, w_1) = 0$ and $V(m_n, w_1) = \alpha$. For all remaining pairs $(m, w) \in M \times W$, $U(m, w) = V(m, w) = 0$. Figure 3a illustrates the SMC instance \mathcal{I}_5 .




Fig. 3. Subfigure (a) illustrates the graph representation of the SMC instance \mathcal{I}_n described in the proof of Theorem 3 for $n = 5$. Subfigure (b) shows the matching μ^{opt} . Subfigures (c), (d), and (e) show the matchings $\mu^{(1)}$, $\mu^{(2)}$, and $\mu^{(3)}$, respectively. Dashed lines indicate zero-valuation pairs that do not appear in the graph representation.

Define $\mu^{\text{opt}} := \{(m_1, w_1)\} \cup \{(m_{2i}, w_{2i+1}), (m_{2i+1}, w_{2i}) : i \in \{1, 2, \dots, \frac{n-1}{2}\}\}$ (see Figure 3b). We also define a number of other integral matchings obtained by modifying μ^{opt} , as follows: For $i \in \{1, 2, \dots, \frac{n-1}{2}\}$, the matching $\mu^{(i)}$ (see Figures 3c and 3d) is the integral matching which is obtained from μ^{opt} by replacing $\{(m_1, w_1), (m_{2i}, w_{2i+1})\}$ with $\{(m_1, w_{2i+1}), (m_{2i}, w_1)\}$, i.e.,

$$\mu^{(i)} := (m_1, w_{2i+1}) \cup (m_{2i}, w_1) \cup (m_{2i+1}, w_{2i}) \cup \{(m_{2\ell}, w_{2\ell+1}) \cup (m_{2\ell+1}, w_{2\ell})\}_{\ell \in \{1, 2, \dots, \frac{n-1}{2}\} \setminus \{i\}}.$$

Also, the matching $\mu^{(\frac{n+1}{2})}$ (see Figure 3e) is the integral matching obtained from μ^{opt} by replacing $\{(m_1, w_1), (m_n, w_{n-1})\}$ with $\{(m_1, w_{n-1}), (m_n, w_1)\}$, i.e.,

$$\mu^{(\frac{n+1}{2})} := (m_1, w_{n-1}) \cup (m_{n-1}, w_n) \cup (m_n, w_1) \cup \{(m_{2\ell}, w_{2\ell+1}) \cup (m_{2\ell+1}, w_{2\ell})\}_{\ell \in \{1, 2, \dots, \frac{n-3}{2}\}}.$$

Now, consider the fractional matching $\mu := \sum_{i=1}^{\frac{n+1}{2}} \frac{1}{\alpha} \mu^{(i)} + (1 - \frac{n+1}{2\alpha}) \mu^{\text{opt}}$. Since $\alpha > n + 2 > \frac{n+1}{2}$, μ is well-defined and has the matchings μ^{opt} and $\mu^{(i)}$ for $i \in \{1, \dots, \frac{n+1}{2}\}$ in its support. Notice that $\mathcal{W}(\mu^{\text{opt}}) = (n-1)\alpha + 2$ and $\mathcal{W}(\mu^{(i)}) = (n-1)\alpha$ for all $i \in \{1, 2, \dots, \frac{n+1}{2}\}$. Thus, $\mathcal{W}(\mu) > (n-1)\alpha$.

It can be verified that μ is stable. Indeed, we only need to check the blocking condition for the pairs (m_i, w_i) with $i \in \{1, 2, \dots, n\}$. We have that $v_{w_1}(\mu) \geq 1$ (since $\mu^{(\frac{n+1}{2})}$ has weight $\frac{1}{\alpha}$ in μ and $V(m_n, w_1) = \alpha$), $u_{m_{2i}}(\mu) \geq 1$ for each $i \in \{1, 2, \dots, \frac{n-1}{2}\}$ (since $\mu^{(i)}$ has weight $\frac{1}{\alpha}$ in μ and $U(m_{2i}, w_1) = \alpha$), and $v_{w_{2i+1}}(\mu) \geq 1$ for each $i \in \{1, 2, \dots, \frac{n-1}{2}\}$ (since $\mu^{(i)}$ has weight $\frac{1}{\alpha}$ in μ and $V(m_{2i}, w_{2i+1}) = \alpha$). The welfare of the optimal stable fractional matching must therefore be at least $\mathcal{W}(\mu)$, and thus strictly greater than $(n-1)\alpha$.

We now claim that any ρ -efficient stable fractional matching μ' satisfies $\mu'(m_n, w_1) > 0$. Indeed, assuming otherwise that $\mu'(m_n, w_1) = 0$, the only pair that can give positive utility to man m_1 and woman w_1 is (m_1, w_1) . Hence, we must also have $\mu'(m_1, w_1) = 1$, and, as a result, $\mu'(m_{2i}, w_1) = 0$ for $i \in \{1, 2, \dots, \frac{n-1}{2}\}$. Then, the only pair that can give positive utility to man m_{2i} and woman w_{2i} is (m_{2i}, w_{2i}) , and hence, it must also be that $\mu'(m_{2i}, w_{2i}) = 1$. Consequently, the only pair that can give positive utility to man m_{2i+1} and woman w_{2i+1} is (m_{2i+1}, w_{2i+1}) and, hence, we must have $\mu'(m_{2i+1}, w_{2i+1}) = 1$. The welfare of matching μ' would then be $2n$, which is less than $\rho(n-1)\alpha$ by the assumed bound on α . In other words, the welfare of μ' would be less than ρ times the welfare of the stable fractional matching μ , contradicting the assumption that μ' is ρ -efficient.

The final step in the proof involves showing that for any stable fractional matching μ' with support of size at most $\frac{n-1}{2}\rho$, we must have $\mathcal{W}(\mu') < \rho(n-1)\alpha$; the desired bound on the support size would then follow from the contrapositive. Let $T := \{i \in \{1, 2, \dots, \frac{n-1}{2}\} : \mu'(m_{2i}, w_1) > 0\}$, and $\bar{T} := \{1, 2, \dots, \frac{n-1}{2}\} \setminus T$. Since μ' has support of size at most $\frac{n-1}{2}\rho$ and $\mu'(m_n, w_1) > 0$, it holds that $|T| \leq \frac{n-1}{2}\rho - 1$.

For every $i \in T$, the agents m_{2i} , w_{2i} , m_{2i+1} , and w_{2i+1} can together contribute at most 2α to the welfare. On the other hand, when $i \in \bar{T}$, we have $\mu'(m_{2i}, w_1) = 0$, and the only pair that can give positive utility to man m_{2i} and woman w_{2i} is (m_{2i}, w_{2i}) . Therefore, we must have that $\mu'(m_{2i}, w_{2i}) = 1$. Consequently, the only pair that can give positive utility to man m_{2i+1} and woman w_{2i+1} is (m_{2i+1}, w_{2i+1}) , and it follows that $\mu'(m_{2i+1}, w_{2i+1}) = 1$. Therefore, when $i \in \bar{T}$, the agents m_{2i} , w_{2i} , m_{2i+1} , and w_{2i+1} can together contribute at most 4 to the welfare. Taking the possible contribution of pair (m_1, w_1) into account, we have that

$$\mathcal{W}(\mu') \leq 2 + 2\alpha|T| + 4|\bar{T}| = 2n + (2\alpha - 4)|T| \leq 2n + \rho(n-1)\alpha - 2\alpha - 2(n-1)\rho + 4 < \rho(n-1)\alpha.$$

The equality follows from the definition of \bar{T} , the second inequality follows from the bound on $|T|$ above, and the third one from the definition of α . This completes the proof of Theorem 3. \square

Theorem 3 has an interesting algorithmic implication. As the support size can be large in any good approximation of the optimal stable fractional matching, it implies that *support enumeration* strategies—which have been proved useful in other contexts; see [6] and the references therein—will be ineffective in computing (even approximate) solutions of OPTIMAL STABLE FRACTIONAL MATCHING. A similar implication can be shown for optimal ε -stable fractional matchings. In contrast, as we will show in Section 4, ε -stable fractional matchings of *small* support can be easily computed, and provide nearly the best approximation ratios achievable by efficient algorithms (under standard complexity-theoretic assumptions).

4 ALGORITHMIC RESULTS

We begin the discussion of our algorithmic results with binary valuations. In this setting, OPTIMAL STABLE FRACTIONAL MATCHING reduces to computing a maximum weight matching on a specific weighted graph associated with the given instance.

THEOREM 4. *Given an SMC instance $\mathcal{I} = \langle M, W, U, V \rangle$ with binary valuations, an optimal stable fractional matching for \mathcal{I} can be computed in polynomial time.*

PROOF. Let G be the graph representation of \mathcal{I} . We assign to each edge (m, w) in G a weight $\gamma(m, w)$, as follows: $\gamma(m, w) = 2 + 1/n^2$ whenever $U(m, w) = V(m, w) = 1$, otherwise $\gamma(m, w) = 1$. Thus, for any matching μ in G , if n_μ denotes the number of agents (men and women) with utility 1 in the SMC instance \mathcal{I} , then $n_\mu \leq \sum_{(m,w) \in \mu} \gamma(m, w) < n_\mu + 1$.

Let μ be a maximum weight matching in G . Note that μ can be computed in polynomial time. Also, it follows from the above inequality that μ is an optimal matching for \mathcal{I} . We will now argue that μ is stable. Indeed, assuming otherwise, any blocking pair (m, w) must have $0 = u_m(\mu) < U(m, w) = 1$, $0 = v_w(\mu) < V(m, w) = 1$ and $(m, w) \notin \mu$. Thus, if μ contains one or both of the edges (m, w') and (m', w) for some $w' \neq w$ and $m' \neq m$, then we must have that $U(m, w') = 0$ and/or $V(m', w) = 0$. By our definition of weights, this would imply that $\gamma(m, w') = 1$ and/or $\gamma(m', w) = 1$. We can now replace one or both of these edges with the edge (m, w) , which has weight $\gamma(m, w) = 2 + 1/n^2$, and obtain a new matching with strictly larger weight—a contradiction. \square

Next, we consider general valuations and show how to exploit stable integral matchings to get an approximate solution for OPTIMAL STABLE FRACTIONAL MATCHING. Let σ_{\max} and σ_{\min} denote the largest and the smallest non-zero valuation among all agents in \mathcal{I} , respectively. We call a man-woman pair (m, w) *light* if either $U(m, w) = 0$ or $V(m, w) = 0$, and *heavy* otherwise. Given an SMC instance \mathcal{I} as input, our algorithm computes a stable integral matching for \mathcal{I} , say μ , in two steps: First, it computes a stable integral matching μ_1 using only the heavy pairs (and taking into account the stability constraints in heavy pairs only). Then, it *completes* the solution with a matching μ_2 of maximum welfare using the light pairs subject to feasibility constraints, i.e., using light pairs that do not share any agents with the pairs in μ_1 . The light pairs impose no additional constraints on stability, so the resulting matching is stable.

We will show that μ has approximation ratio $1 + \sigma_{\max}/\sigma_{\min}$. Let μ^{opt} be an optimal matching for \mathcal{I} . Also, let μ_1^{opt} be the set of pairs of μ^{opt} that share an agent with some pair of μ_1 , i.e., $\mu_1^{\text{opt}} := \{(m, w) \in \mu^{\text{opt}} : \text{at least one of } m \text{ or } w \text{ is matched under } \mu_1\}$. By definition of μ_2 , we have $\mathcal{W}(\mu_2) \geq \mathcal{W}(\mu^{\text{opt}} \setminus \mu_1^{\text{opt}})$. To complete the proof, we will need the following lemma.

LEMMA 1. $\mathcal{W}(\mu_1 \setminus \mu_1^{\text{opt}}) \geq (1 + \sigma_{\max}/\sigma_{\min})^{-1} \mathcal{W}(\mu_1^{\text{opt}} \setminus \mu_1)$.

PROOF. Our proof constructs a mapping in which every pair $(m, w) \in \mu_1^{\text{opt}} \setminus \mu_1$ is mapped to one of its agents, whom we will call the *witness* of the pair. The mapping is such that the utility of the witness in the matching $\mu_1 \setminus \mu_1^{\text{opt}}$ is at least $(1 + \sigma_{\max}/\sigma_{\min})^{-1} (U(m, w) + V(m, w))$. Note that once we establish the said mapping, the proof will follow, since each agent can be the witness of at most one pair of $\mu_1^{\text{opt}} \setminus \mu_1$ and $\mathcal{W}(\mu_1 \setminus \mu_1^{\text{opt}})$ is at least the total utility of the witnesses in $\mu_1 \setminus \mu_1^{\text{opt}}$.

Consider a light pair $(m, w) \in \mu_1^{\text{opt}} \setminus \mu_1$. The witness is an agent (m or w) who also belongs to a pair of $\mu_1 \setminus \mu_1^{\text{opt}}$; such an agent certainly exists by the definition of μ_1^{opt} . Since all pairs of $\mu_1 \setminus \mu_1^{\text{opt}}$ are heavy, the utility of the witness of (m, w) in $\mu_1 \setminus \mu_1^{\text{opt}}$ is at least $\sigma_{\min} = \frac{\sigma_{\min}}{\sigma_{\max}} (0 + \sigma_{\max}) \geq (1 + \sigma_{\max}/\sigma_{\min})^{-1} (U(m, w) + V(m, w))$, since (m, w) is light.

Now consider a heavy pair $(m, w) \in \mu_1^{\text{opt}} \setminus \mu_1$. If μ_1 contains a pair (m, w') with $U(m, w') \geq U(m, w)$, select agent m to be the witness, otherwise select agent w . Note that in the latter case,

stability of μ_1 implies the existence of $(m', w) \in \mu_1$ such that $V(m', w) \geq V(m, w)$. Hence, the utility of the witness of (m, w) in $\mu_1 \setminus \mu_1^{\text{opt}}$ is at least $\min\{U(m, w), V(m, w)\}$, which, in turn, is at least $(1 + \sigma_{\max}/\sigma_{\min})^{-1} (U(m, w) + V(m, w))$. \square

Now, Lemma 1 gives the desired approximation ratio, as follows:

$$\begin{aligned} \mathcal{W}(\mu) &= \mathcal{W}(\mu_1) + \mathcal{W}(\mu_2) = \mathcal{W}(\mu_1 \setminus \mu_1^{\text{opt}}) + \mathcal{W}(\mu_1 \cap \mu_1^{\text{opt}}) + \mathcal{W}(\mu_2) \\ &\geq (1 + \sigma_{\max}/\sigma_{\min})^{-1} \mathcal{W}(\mu_1^{\text{opt}} \setminus \mu_1) + \mathcal{W}(\mu_1^{\text{opt}} \cap \mu_1) + \mathcal{W}(\mu^{\text{opt}} \setminus \mu_1^{\text{opt}}) \\ &\geq (1 + \sigma_{\max}/\sigma_{\min})^{-1} \mathcal{W}(\mu^{\text{opt}}). \end{aligned}$$

For ternary valuations in $\{0, 1, \alpha\}$, the above algorithm gives a $(1 + \alpha)$ -approximation. An improved approximation for ternary valuations can be achieved using the following modification: When computing the stable integral matching, resolve ties in favour of the pairs (m, w) with the highest $U(m, w) + V(m, w)$. The next lemma establishes an improved approximation ratio of $\max\{2, \alpha\}$.

LEMMA 2. *The modified algorithm for SMC instances with ternary valuations in $\{0, 1, \alpha\}$ satisfies $\mathcal{W}(\mu_1 \setminus \mu_1^{\text{opt}}) \geq \min\{\frac{1}{2}, \frac{1}{\alpha}\} \mathcal{W}(\mu_1^{\text{opt}} \setminus \mu_1)$.*

PROOF. For a pair (m, w) of matching $\mu_1^{\text{opt}} \setminus \mu_1$, we use the term *neighborhood* to refer to the pairs of $\mu_1 \setminus \mu_1^{\text{opt}}$ that use agent m or w . We will show that the total utility from pairs in the neighborhood of (m, w) is at least $\min\{1, 2/\alpha\} (U(m, w) + V(m, w))$. Since each pair of $\mu_1 \setminus \mu_1^{\text{opt}}$ can be in the neighborhood of at most two pairs of $\mu_1^{\text{opt}} \setminus \mu_1$, this will give us the desired inequality.

Indeed, by the particular way we resolve ties in the ordinal preferences before computing the matching μ_1 , a heavy pair (m, w) in $\mu_1^{\text{opt}} \setminus \mu_1$ must have a pair of utility at least $U(m, w) + V(m, w)$ in its neighborhood. A light pair (m, w) has $U(m, w) + V(m, w) \leq \alpha$ and certainly has a heavy pair of utility at least 2 in its neighborhood. \square

The above discussion is summarized in the following statement.

THEOREM 5. *There is a polynomial-time algorithm which, given an SMC instance \mathcal{I} with an optimal matching μ^{opt} , computes a stable integral matching μ with $\mathcal{W}(\mu) \geq \min\{\frac{1}{2}, \frac{1}{\alpha}\} \mathcal{W}(\mu^{\text{opt}})$ if \mathcal{I} has ternary valuations in $\{0, 1, \alpha\}$, and $\mathcal{W}(\mu) \geq (1 + \sigma_{\max}/\sigma_{\min})^{-1} \mathcal{W}(\mu^{\text{opt}})$ in general, where σ_{\max} and σ_{\min} denote the highest and lowest non-zero valuation in \mathcal{I} , respectively.*

We conclude this section by considering approximate stability. For general valuations, we present a polynomial-time $1/\varepsilon$ -approximation algorithm for OPTIMAL ε -STABLE FRACTIONAL MATCHING, which constructs an ε -stable fractional matching with a small support by combining an optimal matching with a stable integral matching.

THEOREM 6. *There is a polynomial-time algorithm that given any SMC instance $\mathcal{I} = \langle M, W, U, V \rangle$ and any rational $\varepsilon \in [0, 1]$, computes a fractional matching μ that is ε -stable for \mathcal{I} such that $\mathcal{W}(\mu) \geq \varepsilon \mathcal{W}(\mu^{\text{opt}})$, where μ^{opt} is an optimal matching for \mathcal{I} .*

PROOF. Let μ^s be any stable integral matching and μ^{opt} be an optimal matching for \mathcal{I} . Note that both μ^s and μ^{opt} can be computed in polynomial time. We will show that $\mu := (1 - \varepsilon)\mu^s + \varepsilon\mu^{\text{opt}}$ satisfies the desired properties. Indeed, $\mathcal{W}(\mu) = (1 - \varepsilon)\mathcal{W}(\mu^s) + \varepsilon\mathcal{W}(\mu^{\text{opt}}) \geq \varepsilon\mathcal{W}(\mu^{\text{opt}})$. Furthermore, since μ^s is stable, we have that for any man-woman pair $(m, w) \in M \times W$, either $u_m(\mu^s) \geq U(m, w)$ or $v_w(\mu^s) \geq V(m, w)$. The former condition implies that $u_m(\mu) \geq (1 - \varepsilon)u_m(\mu^s) \geq (1 - \varepsilon)U(m, w)$, while the latter condition gives $v_w(\mu) \geq (1 - \varepsilon)V(m, w)$. Either way, the pair (m, w) is ε -stable. \square

In particular, Theorem 6 shows that a $\frac{1}{2}$ -stable fractional matching with welfare at least half of that of an optimal fractional matching (and therefore, that of an optimal stable fractional matching) can be computed in polynomial time. In Section A.4, we provide a slightly stronger welfare guarantee: There is a polynomial-time algorithm that computes a $\frac{1}{2}$ -stable fractional matching with welfare at least that of an optimal (exactly) stable fractional matching.

5 HARDNESS OF APPROXIMATION

In this section, we present our inapproximability statements, which are by far the technically most involved results in the paper. We present polynomial-time reductions which, given a 3SAT formula ϕ of a particular structure, construct SMC instances that simulate the evaluation of ϕ for every variable assignment. The constructed SMC instances consist of several gadgets including an *accumulator*. The simulation of the evaluation of ϕ by the SMC instance is such that:

- (a) when ϕ has a satisfying assignment, there is a stable (or ε -stable) fractional matching where the contribution of the agents in the accumulator gadget to the welfare can be large and dominates the contribution from the remaining SMC instance and
- (b) when ϕ is not satisfiable, the contribution of the accumulator and, subsequently, the total welfare of any stable (or ε -stable) fractional matching is very small.

Hence, distinguishing between SMC instances with stable (or ε -stable) fractional matchings of very high and very low welfare would allow us to decide 3SAT. We have two inapproximability statements: Theorem 7 for OPTIMAL STABLE FRACTIONAL MATCHING and Theorem 8 for OPTIMAL ε -STABLE FRACTIONAL MATCHING.

THEOREM 7. *For every constant $\delta > 0$, it is NP-hard to approximate OPTIMAL STABLE FRACTIONAL MATCHING for SMC instances with ternary valuations in $\{0, 1, \alpha\}$ to within a factor of (i) $\alpha - 1/2 - \delta$ if $\alpha = O(n)$, and (ii) $\Omega(n^{1-\delta})$ otherwise.*

THEOREM 8. *For any constants $\varepsilon \in (0, 0.03]$ and $\delta > 0$, it is NP-hard to approximate OPTIMAL ε -STABLE FRACTIONAL MATCHING to within a factor of $1/\varepsilon - \delta$.*

We will prove Theorem 7 here; the proof of Theorem 8, which uses similar gadgets but is slightly more involved, appears in Appendix A.3. Since the proof is long, we have divided it into three parts: the description of the reduction (Section 5.1), technical claims with gadget properties (Section 5.2), and the proof of the inapproximability result (Section 5.3).

5.1 The reduction

In particular, we present a polynomial-time reduction from 2P2N-3SAT, the special case of 3SAT consisting of 3-CNF clauses in which every variable appears four times: twice as a positive literal and twice as a negative one. 2P2N-3SAT is known to be NP-hard [37]. Our reduction takes as input an instance of 2P2N-3SAT consisting of N (boolean) variables x_1, x_2, \dots, x_N , and a 3-CNF formula ϕ with $L = 4N/3$ clauses c_1, c_2, \dots, c_L . Without loss of generality, we assume that each clause in ϕ consists of distinct literals.

Given the instance of 2P2N-3SAT, our reduction generates an instance $\mathcal{I} = \langle M, W, U, V \rangle$ of OPTIMAL STABLE FRACTIONAL MATCHING. As usual, we denote by n the number of men (or women) in \mathcal{I} . We will use a positive integer parameter k which will determine the size of n ; in particular, $n = O(N + k)$. We define \mathcal{I} by referring to its graph representation, which consists of *variable gadgets*, *clause gadgets*, *variable-clause connectors*, an *accumulator*, and *clause-accumulator connectors*. For each gadget, we classify the edges (i.e., man-woman pairs and their valuations) into the following three types:

- *man-heavy edges* (m, w) with $U(m, w) = \alpha$ and $V(m, w) = 0$,

- *woman-heavy* edges (m, w) with $U(m, w) = 0$ and $V(m, w) = \alpha$, and
- *balanced* edges (m, w) with $U(m, w) = V(m, w) = 1$.

Recall that any pair (m, w) that does not appear as an edge in the graph representation has $U(m, w) = V(m, w) = 0$.

The instance \mathcal{I} has a variable gadget for every variable x , which consists of five men $m_1^x, m_2^x, e_1^x, e_2^x, e_3^x$, four women $w_1^{\bar{x}}, w_2^{\bar{x}}, f_1^x, f_2^x$ and the ten balanced edges $(e_1^x, f_1^x), (m_1^x, f_1^x), (m_1^x, w_1^{\bar{x}}), (e_3^x, w_1^{\bar{x}}), (e_3^x, w_2^{\bar{x}}), (m_2^x, w_2^{\bar{x}}), (m_2^x, f_2^x), (e_2^x, f_2^x), (m_1^x, w_2^{\bar{x}})$, and $(m_2^x, w_1^{\bar{x}})$, as shown in Figure 4a.




Fig. 4. (a) The variable gadget corresponding to the variable x . (b) The clause gadget corresponding to the clause c and its CA-connector (m^c, w^c) . As a convention, we use circles to represent men and diamonds to represent women.

For every clause c , instance \mathcal{I} has a clause gadget with three men m^c, e_1^c, e_2^c , three women w_1^c, w_2^c, w_3^c , and the nine balanced edges between them, as shown in Figure 4b.

For every appearance of a literal in a clause, there is a variable-clause connector (or *VC-connector*). VC-connectors have different structure depending (1) on whether they correspond to positive or negative literals, and (2) on the value of α . In each case, we identify one edge of the VC-connector as the *input*, and either one or two edges as the *output*.

Specifically, for every positive literal x whose i -th appearance ($i \in \{1, 2\}$) is as the j -th literal ($j \in \{1, 2, 3\}$) of clause c , \mathcal{I} has a VC-connector defined as follows:

- When $\alpha \geq 2$, the VC-connector consists of a single woman-heavy edge between m_i^x (from the variable gadget corresponding to variable x) and w_j^c (from the clause gadget corresponding to clause c), as shown in Figure 5a. This edge is simultaneously the input and the output edge of the VC-connector.
- When $\alpha \in (3/2, 2)$, the VC-connector consists of woman $w^{x,c}$, man $m^{x,c}$, the woman-heavy edges $(m_i^x, w^{x,c})$ and $(m^{x,c}, w_j^c)$, and the balanced edge $(m^{x,c}, w^{x,c})$, as shown in Figure 5b. Here, $(m_i^x, w^{x,c})$ is the input and $(m^{x,c}, w_j^c)$ is the output edge.

For every negative literal \bar{x} whose i -th appearance ($i \in \{1, 2\}$) is as the j -th literal ($j \in \{1, 2, 3\}$) of clause c , \mathcal{I} has a VC-connector defined as follows:

- When $\alpha \geq 2$, the VC-connector consists of man $m^{\bar{x},c}$, woman $w^{\bar{x},c}$, the man-heavy edge $(m^{\bar{x},c}, w_i^{\bar{x}})$, the balanced edge $(m^{\bar{x},c}, w^{\bar{x},c})$, and the woman-heavy edge $(m^{\bar{x},c}, w_j^c)$, as shown in Figure 5c. Here, $(m^{\bar{x},c}, w_i^{\bar{x}})$ is the input and $(m^{\bar{x},c}, w_j^c)$ is the output edge.

- When $\alpha \in (3/2, 2)$, the VC-connector consists of three men $m_1^{\bar{x},c}$, $m_2^{\bar{x},c}$, $m_3^{\bar{x},c}$, three women $w_1^{\bar{x},c}$, $w_2^{\bar{x},c}$, $w_3^{\bar{x},c}$, the man-heavy edges $(m_1^{\bar{x},c}, w_i^{\bar{x}})$, $(m_3^{\bar{x},c}, w_1^{\bar{x},c})$, $(m_3^{\bar{x},c}, w_2^{\bar{x},c})$, the woman-heavy edges $(m_1^{\bar{x},c}, w_2^{\bar{x},c})$, $(m_2^{\bar{x},c}, w_j^c)$, $(m_3^{\bar{x},c}, w_j^c)$, and the balanced edges $(m_1^{\bar{x},c}, w_1^{\bar{x},c})$, $(m_2^{\bar{x},c}, w_2^{\bar{x},c})$, $(m_3^{\bar{x},c}, w_3^{\bar{x},c})$, as shown in Figure 5d. In this case, the VC-connector has one input edge $(m_1^{\bar{x},c}, w_i^{\bar{x}})$ and two output edges $(m_2^{\bar{x},c}, w_j^c)$ and $(m_3^{\bar{x},c}, w_j^c)$.




Fig. 5. VC-connectors corresponding to clause c and positive literal x for (a) $\alpha \geq 2$ and (b) $\alpha \in (3/2, 2)$, and to clause c and negative literal \bar{x} for (c) $\alpha \geq 2$ and (d) $\alpha \in (3/2, 2)$.

The accumulator (Figure 6) of instance \mathcal{I} has different structure depending on the value of α . Its size depends on the positive integer parameter k .

- When $\alpha \geq 2$ (see Figure 6a), the accumulator has man m_i and woman w_i for all $i \in \{1, \dots, k\}$, men e_i^1 and e_i^2 and woman f_i^1 for all $i \in \{1, \dots, k-1\}$, man e_i^3 and women f_i^2 and f_i^3 for all $i \in \{2, \dots, k\}$, and woman w^c for every clause c of ϕ . In addition, there are man-heavy edges (m_i, w_{i-1}) and (e_i^3, f_i^2) for all $i \in \{2, \dots, k\}$ and (e_i^2, w_i) for all $i \in \{1, \dots, k-1\}$, the balanced edges (m_1, w^c) for every clause c , which we call *tine* edges, (e_i^1, w_i) for all $i \in \{1, \dots, k-1\}$ and (m_i, f_i^2) for all $i \in \{2, \dots, k\}$, and the woman-heavy edges (m_i, w_i) for all $i \in \{1, \dots, k\}$, (e_i^1, f_i^1) for all $i \in \{1, \dots, k-1\}$, and (m_i, f_i^3) for all $i \in \{2, \dots, k\}$.
- When $\alpha \in (3/2, 2)$ (see Figure 6b), the accumulator has man m_i , woman w_i for $i = 1, \dots, k$, man e_i^1 and woman f_i^1 for $i = 1, \dots, k-1$, man e_i^2 and woman f_i^2 for $i = 2, \dots, k$, and woman w^c for every clause c of ϕ . In addition, it contains the man-heavy edges (m_i, w_{i-1}) and (e_i^2, f_i^2) for $i = 2, \dots, k$ and (m_i, f_{i-1}^2) for $i = 3, \dots, k$, the balanced edges (m_1, w^c) for every clause c (tine edges), (e_i^1, w_i) for $i = 1, \dots, k-1$ and (m_i, f_i^2) for $i = 2, \dots, k$, and the woman-heavy edges (m_i, w_i) for $i = 1, \dots, k$, and (e_i^1, f_i^1) and (e_i^1, w_{i+1}) for $i = 1, \dots, k-1$.




Fig. 6. The accumulator for the cases (a) $\alpha \geq 2$ and (b) $\alpha \in (3/2, 2)$.

Finally, instance \mathcal{I} has a clause-accumulator connector (or *CA-connector*) for every clause c of ϕ consisting of the woman-heavy edge (m^c, w^c) between the man m^c (from the clause gadget corresponding to clause c) and woman w^c (from the accumulator); see Figure 4b. Notice that the above construction has more women than men. To restore balance, we pad the instance with extra (isolated) men that neither value nor are valued by any other agent. This completes the construction of the reduced instance.

5.2 Gadget properties

We will now prove several important properties (Claims 9-12) of our construction.

CLAIM 9. For every variable x , a stable fractional matching μ satisfies at least one of the following:

- (1) $\mu(m_1^x, w_1^{\bar{x}}) + \mu(m_1^x, w_2^{\bar{x}}) + \mu(m_1^x, f_1^x) = 1$ and $\mu(m_2^x, w_1^{\bar{x}}) + \mu(m_2^x, w_2^{\bar{x}}) + \mu(m_2^x, f_2^x) = 1$.
- (2) $\mu(m_1^x, w_1^{\bar{x}}) + \mu(m_2^x, w_1^{\bar{x}}) + \mu(e_3^x, w_1^{\bar{x}}) = 1$ and $\mu(m_1^x, w_2^{\bar{x}}) + \mu(m_2^x, w_2^{\bar{x}}) + \mu(e_3^x, w_2^{\bar{x}}) = 1$.

PROOF. Suppose, for contradiction, that for some $i, j \in \{1, 2\}$, we have $\mu(m_i^x, w_1^{\bar{x}}) + \mu(m_i^x, w_2^{\bar{x}}) + \mu(m_i^x, f_1^x) < 1$ and $\mu(m_1^x, w_j^{\bar{x}}) + \mu(m_2^x, w_j^{\bar{x}}) + \mu(e_3^x, w_j^{\bar{x}}) < 1$. Then, both m_i^x and $w_j^{\bar{x}}$ will have utility strictly less than 1 under μ , and thus the pair $(m_i^x, w_j^{\bar{x}})$ would be blocking. \square

We remark that the two conditions in the statement of Claim 9 affect the weight of the input edges of the VC-connectors that are attached to the variable gadget in any stable fractional matching. In particular, condition (1) implies that the weight assigned to the input edges of the VC-connectors that correspond to the two appearances of the positive literal x in clauses must be 0. To see why, observe that these input edges are incident to nodes m_1^x and m_2^x , and the total weight of all edges incident to each of these nodes cannot exceed 1. Condition (2) has a similar implication for the edges associated with the negative literal \bar{x} .

CLAIM 10. *Any stable fractional matching that assigns a weight of 0 to the input edge of a VC-connector must assign a weight of 0 to its output edge(s) as well.*

PROOF. For $\alpha \geq 2$, the claim holds trivially for VC-connectors corresponding to positive literals (Figure 5a). Consider a VC-connector corresponding to a negative literal \bar{x} and a clause c containing it (Figure 5c). Observe that, besides the input edge $(m_1^{\bar{x},c}, w_1^{\bar{x}})$, the edge $(m_1^{\bar{x},c}, w_2^{\bar{x},c})$ is the only balanced or man-heavy edge that is incident to man $m_1^{\bar{x},c}$ and the only balanced (or woman-heavy) edge incident to woman $w_2^{\bar{x},c}$. Hence, stability of the edge $(m_1^{\bar{x},c}, w_2^{\bar{x},c})$ requires a weight of 1 assigned to it when the weight assigned to input edge $(m_1^{\bar{x},c}, w_1^{\bar{x}})$ is 0. Then, the output edge $(m_1^{\bar{x},c}, w_j^c)$, which is also incident to the node $m_1^{\bar{x},c}$, must have a weight of 0 as well.

We now consider the case $\alpha \in (3/2, 2)$. First consider a VC-connector corresponding to a positive literal x and a clause c containing it (Figure 5b). The edge $(m_i^{x,c}, w^{x,c})$ is the only balanced or man-heavy edge that is incident to man $m_i^{x,c}$ and, besides the input edge $(m_i^x, w^{x,c})$, the only balanced (or woman-heavy) edge incident to woman $w^{x,c}$. Hence, stability of edge $(m_i^{x,c}, w^{x,c})$ requires a weight of 1 assigned to it when the weight assigned to edge $(m_i^x, w^{x,c})$ is 0. Then, the output edge $(m_i^{x,c}, w_j^c)$, which is also incident to node $m_i^{x,c}$, must have a weight of 0 as well.

Finally, consider a VC-connector corresponding to a negative literal \bar{x} and a clause c containing it (Figure 5d). Observe that $(m_1^{\bar{x},c}, w_1^{\bar{x},c})$ is the only balanced or woman-heavy edge that is incident to woman $w_1^{\bar{x},c}$ and, besides the input edge $(m_1^{\bar{x},c}, w_1^{\bar{x}})$, the only balanced or man-heavy edge incident to man $m_1^{\bar{x},c}$. Also, $(m_2^{\bar{x},c}, w_2^{\bar{x},c})$ is the only balanced or man-heavy edge that is incident to man $m_2^{\bar{x},c}$ and, besides edge $(m_1^{\bar{x},c}, w_2^{\bar{x},c})$, the only balanced or woman-heavy edge to woman $w_2^{\bar{x},c}$. Furthermore, $(m_3^{\bar{x},c}, w_3^{\bar{x},c})$ is the only balanced or man-heavy edge that is incident to woman $w_3^{\bar{x},c}$ and, besides edges $(m_3^{\bar{x},c}, w_1^{\bar{x},c})$ and $(m_3^{\bar{x},c}, w_2^{\bar{x},c})$, the only balanced or man-heavy edge incident to man $m_3^{\bar{x},c}$.

Hence, stability of edge $(m_1^{\bar{x},c}, w_1^{\bar{x},c})$ requires a weight of 1 assigned to it when the weight assigned to edge $(m_1^{\bar{x},c}, w_1^{\bar{x}})$ is 0. Then, edges $(m_1^{\bar{x},c}, w_2^{\bar{x},c})$ and $(m_3^{\bar{x},c}, w_1^{\bar{x},c})$ must have a weight of 0. Then, stability of edge $(m_2^{\bar{x},c}, w_2^{\bar{x},c})$ requires a weight of 1 assigned to it and the edge $(m_3^{\bar{x},c}, w_2^{\bar{x},c})$ and the output edge $(m_2^{\bar{x},c}, w_j^c)$ must have a weight of 0. Then, stability of edge $(m_3^{\bar{x},c}, w_3^{\bar{x},c})$ requires a weight of 1 assigned to it. Hence, the output edge $(m_3^{\bar{x},c}, w_j^c)$ must have a weight of 0 as well. \square

CLAIM 11. *Any stable fractional matching that assigns a weight of 0 to all output edges of the VC-connectors of clause c must assign a weight of 0 to the CA-connector of clause c as well.*

PROOF. Let ℓ_1, ℓ_2 , and ℓ_3 be the literals of clause c . Consider, for the sake of contradiction, a stable fractional matching that assigns (1) a weight of 0 to all output edges of the VC-connectors

corresponding to literals ℓ_i and clause c and (2) strictly positive weight to edge (m^c, w^c) of the CA-connector for clause c . Note that condition (2) implies that the total weight on the edges (m^c, w_1^c) , (m^c, w_2^c) and (m^c, w_3^c) is strictly smaller than 1. Since these are the only balanced or man-heavy edges incident to man m^c , the stability of these edges is guaranteed by a utility of (at least) 1 for each of the agents w_1^c , w_2^c , and w_3^c . By condition (1) and since, besides the output edges of the VC-connectors, the edges (e_1^c, w_1^c) , (e_2^c, w_1^c) , and (m^c, w_1^c) are the only balanced or woman-heavy edges incident to agent w_1^c for $i \in \{1, 2, 3\}$, the weight assigned to these three edges is at least 1. Hence, the total weight on the nine edges of the clause gadget is at least 3, i.e., strictly more than 2 for the six edges incident to men e_1^c and e_2^c , violating the definition of a fractional matching. \square

CLAIM 12. *Any stable fractional matching that assigns a weight of 0 to some CA-connector must assign a total weight of 1 to the tine edges and a weight of 1 to every balanced edge of the accumulator.*

PROOF. Assume that a weight of 0 has been assigned to the edge $(m^{c'}, w^{c'})$ of the CA-connector corresponding to some clause c' . Since this is the only woman-heavy edge that is incident to agent $w^{c'}$ and there is no man-heavy edge incident to agent m_1 , stability on the edge $(m_1, w^{c'})$ requires that the total weight of the tine edges (m_1, w^c) (for every clause c) is (at least) 1. Hence, the weight of the edge (m_1, w_1) is 0. We will complete the proof by distinguishing between the two different accumulator structures, depending on whether $\alpha \geq 2$ or $\alpha \in (3/2, 2)$.

When $\alpha \geq 2$, it suffices to show that for $i = 1, \dots, k-1$, if the weight of edge (m_i, w_i) is 0, then the weight of the balanced edges (e_i^1, w_i) and (m_{i+1}, f_{i+1}^2) is 1 and the weight of edge (m_{i+1}, w_{i+1}) is 0. Indeed, observe that, edge (e_i^1, w_i) is the only balanced or man-heavy edge incident to man e_i^1 and, besides edge (m_i, w_i) , the only balanced or woman-heavy edge incident to woman w_i . Hence, the balanced edge (e_i^1, w_i) must have a weight of 1 and the edge (m_{i+1}, w_i) a weight of 0.

Then, edge (m_{i+1}, f_{i+1}^2) is the only balanced or woman-heavy edge incident to woman f_{i+1}^2 and, besides edge (m_{i+1}, w_i) , the only balanced or man-heavy edge incident to man m_{i+1} . Hence, the balanced edge (m_{i+1}, f_{i+1}^2) must have a weight of 1 and the edge (m_{i+1}, w_{i+1}) a weight of 0.

When $\alpha \in (3/2, 2)$, it suffices to show that for $i = 1, \dots, k-1$, if the weight of edge (m_i, w_i) and (if they exist) edges (e_{i-1}^1, w_i) and (m_{i+1}, f_i^2) is 0, then the weight of the balanced edges (e_i^1, w_i) and (m_{i+1}, f_{i+1}^2) is 1 and the weight of edges (m_{i+1}, w_{i+1}) , (e_i^1, w_{i+1}) , and (if it exists) (m_{i+2}, f_{i+1}^2) is 0. Indeed, observe that, edge (e_i^1, w_i) is the only balanced or man-heavy edge incident to man e_i^1 and, besides edge (m_i, w_i) and (if it exists) (e_{i-1}^1, w_i) , the only balanced or woman-heavy edge incident to woman w_i . Hence, the balanced edge (e_i^1, w_i) must have a weight of 1 and the edges (m_{i+1}, w_i) and (if it exists) (e_i^1, w_{i+1}) a weight of 0. Then, the edge (m_{i+1}, f_{i+1}^2) is, besides (m_{i+1}, f_i^2) and (m_{i+1}, w_i) , the only balanced or man-heavy edge incident to man m_{i+1} and the only balanced or woman-heavy edge incident to woman f_{i+1}^2 . Hence, the balanced edge (m_{i+1}, f_{i+1}^2) must have a weight of 1 and the edges (m_{i+1}, w_{i+1}) and (if it exists) (m_{i+2}, f_{i+1}^2) a weight of 0. \square

5.3 Proof of inapproximability

LEMMA 3. *If formula ϕ is not satisfiable, then any stable fractional matching of \mathcal{I} has welfare at most $80\alpha N + 4(k-1)$.*

PROOF. We will first show that if ϕ is not satisfiable, then any stable fractional matching of \mathcal{I} assigns weight 0 to some CA-connector. For the sake of contradiction, consider a stable fractional matching that assigns a strictly positive weight to all CA-connectors. We will construct a truth assignment for the formula ϕ (contradicting the assumption of the lemma) by repeating the following process for every clause c of ϕ : Let ℓ be a literal that appears in c such that the output edge(s) of the VC-connector, that corresponds to the appearance of ℓ in c , have strictly positive

total weight. By Claim 11, such a literal must exist. We set ℓ to 1 (true). For every variable that has not received a value in this way, we arbitrarily set it to 1.

The above assignment satisfies all the clauses. To show that it is also valid, we need to argue that there is no variable x such that both literals x and \bar{x} have been set to 1. Assume, to the contrary, that literal x is set to 1 due to its appearance in a clause c_1 , and literal \bar{x} is set to 1 due to its appearance in a different clause c_2 . Thus, in the above assignment, the output edge(s) of the VC-connector between the literal x and the clause c_1 , as well as the VC-connector between the literal \bar{x} and the clause c_2 have strictly positive (total) weight. By Claim 10, the input edges of both VC-connectors also have strictly positive weight. Let $i_1, i_2 \in \{1, 2\}$ be such that the i_1 -th appearance of x is in the clause c_1 and the i_2 -th appearance of \bar{x} is in the clause c_2 . Therefore, the said input edges are incident to the nodes $m_{i_1}^x$ and $w_{i_2}^{\bar{x}}$. Using Claim 9, we get that the total weight on the edges incident to one of $m_{i_1}^x$ or $w_{i_2}^{\bar{x}}$ exceeds 1, contradicting feasibility. Thus, the above assignment must be valid, which, in turn, implies that any stable fractional matching assigns weight 0 to some CA-connector.

By Claim 12, the contribution of the accumulator to the welfare is exactly $4k - 2$ (2 from the fine edges plus 2 from each balanced edge). Let us now consider the contribution of the edges that do not belong to the accumulator. This comprises of

- a total value of 20 for the ten balanced edges of each of the N variable gadgets,
- a total value of α (respectively, $2 + 2\alpha$) for the edges of each of the $2N$ VC-connectors corresponding to a positive literal when $\alpha \geq 2$ (respectively, $\alpha \in (3/2, 2)$),
- a total value of $2 + 2\alpha$ (respectively, $6 + 6\alpha$) for the edges of each of the $2N$ VC-connectors corresponding to a negative literal when $\alpha \geq 2$ (respectively, $\alpha \in (3/2, 2)$),
- a total value of $18 + \alpha$ for the nine balanced edges of each of the $4N/3$ clause gadgets and their corresponding CA-connectors.

It can be easily seen that $80\alpha N - 2$ is a (loose) upper bound on the total value from these edges. \square

LEMMA 4. *If ϕ is satisfiable, then there exists a stable fractional matching of \mathcal{I} with welfare at least $4(k - 1)(\alpha - 1/2)$.*

PROOF. Starting from a satisfying assignment for ϕ , we will construct a stable fractional matching μ in which the welfare of the accumulator gadget is at least $4(k - 1)(\alpha - 1/2)$.

Variable gadgets. For the edges of the variable gadget of the variable x , μ is defined as:

- If x is true, then $\mu(m_1^x, w_1^{\bar{x}}) = \mu(e_3^x, w_1^{\bar{x}}) = \mu(e_3^x, w_2^{\bar{x}}) = \mu(m_2^x, w_2^{\bar{x}}) = 1/2$, $\mu(e_1^x, f_1^x) = \mu(e_2^x, f_2^x) = 1$, and the remaining edges have weight 0.
- If x is false, then $\mu(e_3^x, w_1^{\bar{x}}) = \mu(e_3^x, w_2^{\bar{x}}) = 1/2$, $\mu(m_1^x, f_1^x) = \mu(m_2^x, f_2^x) = 1$, and the remaining edges have weight 0.

Clause gadgets and CA-connectors. For each clause, select one of the true literals (tie-break arbitrarily) and call it *active*. Note that each clause has an active literal in a satisfying assignment. Consider the clause c , and let ℓ_i be its active literal for some $i \in \{1, 2, 3\}$. Also, let $i_1, i_2 \in \{1, 2, 3\} \setminus \{i\}$ denote the other two indices. Set $\mu(e_1^c, w_{i_1}^c) = \mu(e_2^c, w_{i_2}^c) = 1$, and set the weight of the remaining balanced edges to 0. Assign a weight of 1 to the CA-connector, i.e., $\mu(m^c, w^c) = 1$.

VC-connectors. For every non-active VC-connector, set the weight of its balanced edges (if any) to 1 and the weight of the remaining edges to 0. For every active VC-connector corresponding to the i -th appearance of the positive literal x as the j -th literal of clause c ($i \in \{1, 2\}$, $j \in \{1, 2, 3\}$), the weights of its edges are as follows:

- When $\alpha \geq 2$, we set $\mu(m_i^x, w_j^c) = 1/2$.
- When $\alpha \in (3/2, 2)$, we set $\mu(m_i^x, w^{x,c}) = 1/2$, $\mu(m^{x,c}, w^{x,c}) = 1 - \alpha/2$, and $\mu(m^{x,c}, w_j^c) = 1/\alpha$.

For every active VC-connector corresponding to the i -th appearance of the negative literal x as the j -th literal of clause c ($i \in \{1, 2\}$, $j \in \{1, 2, 3\}$), the weights of its edges are as follows:

- When $\alpha \geq 2$, we set $\mu(m^{\bar{x},c}, w_i^{\bar{x}}) = \mu(m^{\bar{x},c}, w_j^c) = 1/2$ and $\mu(m^{\bar{x},c}, w^{\bar{x},c}) = 0$.
- When $\alpha \in (3/2, 2)$, we set $\mu(m_1^{\bar{x},c}, w_i^{\bar{x}}) = 1/2$, $\mu(m_1^{\bar{x},c}, w_1^{\bar{x},c}) = 1 - \alpha/2$, $\mu(m_1^{\bar{x},c}, w_2^{\bar{x},c}) = (\alpha - 1)/2$, $\mu(m_2^{\bar{x},c}, w_2^{\bar{x},c}) = 1 - (\alpha^2 - \alpha)/2$, $\mu(m_2^{\bar{x},c}, w_j^c) = 2/\alpha - 1$, $\mu(m_3^{\bar{x},c}, w_1^{\bar{x},c}) = 1/\alpha$, $\mu(m_3^{\bar{x},c}, w_2^{\bar{x},c}) = \mu(m_3^{\bar{x},c}, w_3^{\bar{x},c}) = 0$, $\mu(m_3^{\bar{x},c}, w_j^c) = 1 - 1/\alpha$.

Accumulator. We set $\mu(m_1, w^c) = 0$ for every tine edge (m_1, w^c) of the accumulator. Furthermore:

- When $\alpha \geq 2$, we set $\mu(m_i, w_i) = 1/\alpha$ for all $i \in \{1, \dots, k\}$, $\mu(e_i^2, w_i) = 1 - 2/\alpha$, $\mu(m_{i+1}, w_i) = 1/\alpha$, $\mu(e_i^1, f_i^1) = 1$, $\mu(e_i^1, w_i) = 0$ for all $i \in \{1, \dots, k-1\}$, $\mu(m_i, f_i^2) = 0$, $\mu(m_i, f_i^3) = 1 - 2/\alpha$, and $\mu(e_i^3, f_i^2) = 1$ for all $i \in \{2, \dots, k\}$. Among these, any edge with a positive weight is either man- or woman-heavy, and hence, its contribution to the social welfare is α times its weight. It can be verified that the total contribution is $4(k-1)(\alpha-1/2) + 1$.
- When $\alpha \in (3/2, 2)$, we set $\mu(m_1, w_1) = 1/\alpha$, $\mu(m_2, w_2) = \alpha + 1/\alpha - 2$, $\mu(m_i, w_i) = 1 - 1/\alpha$ for all $i \in \{3, \dots, k\}$, $\mu(m_{i+1}, w_i) = 1 - 1/\alpha$ for all $i \in \{1, \dots, k-1\}$, $\mu(e_i^1, w_i) = 0$ for all $i \in \{1, \dots, k-1\}$, $\mu(m_2, f_2^2) = 2 - \alpha$, $\mu(m_i, f_i^2) = 0$ for all $i \in \{3, \dots, k\}$, $\mu(e_i^1, f_i^1) = \alpha - 1$, $\mu(e_i^2, f_2^2) = \alpha - 2/\alpha$, $\mu(e_k^2, f_k^2) = 1$, $\mu(e_i^1, f_i^1) = 2 - 2/\alpha$ for all $i \in \{2, \dots, k-1\}$, $\mu(e_i^2, f_i^2) = 2 - 2/\alpha$ for all $i \in \{3, \dots, k-1\}$, $\mu(e_1^1, w_2) = 2 - \alpha$, $\mu(e_1^1, w_{i+1}) = 2/\alpha - 1$ for all $i \in \{2, \dots, k-1\}$, and $\mu(m_{i+1}, f_i^2) = 2/\alpha - 1$ for all $i \in \{2, \dots, k-1\}$. Except for the balanced edge (m_2, f_2^2) , every edge with a positive weight among the ones listed above is either man- or woman-heavy, and hence, its contribution to the social welfare is α times its weight. It can be verified that the total contribution in this case is $4(k-1)(\alpha-1/2) + 2\alpha^2 - 7\alpha + 7$.

In each case, the accumulator contributes at least $4(k-1)(\alpha-1/2)$ to the social welfare, as desired.

The feasibility of μ can be verified by inspection. To see why μ is stable, note that we only need to check for the balanced edges, as the man- or woman-heavy edges and the remaining pairs do not impose any constraints on stability. For the balanced edges, stability is established by the following series of observations (we will use the term ‘stabilized by’ to denote that an agent’s utility is at least 1): The variable gadget for the variable x (Figure 4a) is stabilized by the agents f_1^x, f_2^x, e_3^x along with m_1^x, m_2^x (if x is true) or $w_1^{\bar{x}}, w_2^{\bar{x}}$ (if x is false). The clause gadget for clause c (Figure 4b) with active index i (and non-active indices i_1 and i_2) is stabilized by the agents $e_i^c, e_{i_1}^c, w_i^c, w_{i_1}^c, w_{i_2}^c$; in particular, the edge (m^c, w_i^c) is stabilized by w_i^c because an active literal triggers the woman-heavy edge in the VC-connector. A VC connector is stabilized by $w^{x,c}$ (Figure 5b), $m^{\bar{x},c}$ (Figure 5c), or $m_1^{\bar{x},c}, w_2^{\bar{x},c}$, and $m_3^{\bar{x},c}$ (Figure 5d). Finally, the tine edges in the accumulator (Figure 6) are stabilized by w^{c_1}, \dots, w^{c_L} (because we trigger the CA-connector), and the remaining balanced edges are stabilized by w_i ’s and m_i ’s except for m_1 . Overall, μ is a feasible stable fractional matching. \square

We are ready to prove Theorem 7. If $\alpha < N^{1+1/\delta}$, we use our construction with any k satisfying $k-1 \geq \frac{20\alpha N(\alpha-1/2-\delta)}{\delta}$. It is easy to verify that the reduction is polynomial-time. Furthermore, from Lemma 3, we know that the welfare of μ when ϕ is not satisfiable is at most

$$80\alpha N + 4(k-1) \leq \frac{4(k-1)\delta}{\alpha-1/2-\delta} + 4(k-1) = \frac{4(k-1)(\alpha-1/2)}{\alpha-1/2-\delta}.$$

This number is at least $\alpha-1/2-\delta$ times smaller than the welfare of μ when ϕ is satisfiable (Lemma 4). This establishes the inapproximability bound in part (i) of Theorem 7.

If $\alpha \geq N^{1+1/\delta}$, we use our construction with $k = N^{1+1/\delta}$. Once again, the reduction is polynomial-time, and the instance \mathcal{I} has $n = \Theta(N^{1+1/\delta})$ men and women. Observe that $\alpha = \Omega(n)$, $k = \Theta(n)$,

and $N = O(n^\delta)$. Hence, the welfare of μ when ϕ is not satisfiable is at most

$$80\alpha N + 4(k-1) \leq 80\alpha N + 4N^{1+1/\delta} \leq 84\alpha N = O(\alpha n^\delta).$$

On the other hand, the welfare of μ when ϕ is satisfiable is at least $4(k-1)(\alpha-1/2)$, i.e., $\Omega(\alpha n)$. This establishes the bound in part (ii), and with it, completes the proof of Theorem 7.

ACKNOWLEDGMENTS

We are grateful to Elliot Anshelevich for bringing the work of Deligkas et al. [9] to our attention; to Argyrios Deligkas for sharing with us the full version of their paper [9], and to Haris Aziz for pointing us to the work of Manjunath [25].

REFERENCES

- [1] Atila Abdulkadiroğlu, Parag A Pathak, and Alvin E Roth. 2005. The New York City High School Match. *American Economic Review* (2005), 364–367.
- [2] Elliot Anshelevich, Onkar Bhardwaj, and Martin Hoefer. 2013. Friendship and Stable Matching. In *European Symposium on Algorithms*. Springer, 49–60.
- [3] Elliot Anshelevich and Sanmay Das. 2010. Matching, Cardinal Utility, and Social Welfare. *ACM SIGecom Exchanges* 9, 1 (2010), 4.
- [4] Elliot Anshelevich, Sanmay Das, and Yonatan Naamad. 2013. Anarchy, Stability, and Utopia: Creating Better Matchings. *Autonomous Agents and Multi-Agent Systems* 26, 1 (2013), 120–140.
- [5] Haris Aziz and Bettina Klaus. 2017. Random Matching under Priorities: Stability and No Envy Concepts. *arXiv preprint arXiv:1707.01231* (2017).
- [6] Siddharth Barman. 2018. Approximating Nash Equilibria and Dense Subgraphs via an Approximate Version of Carathéodory’s Theorem. *SIAM J. Comput.* 47, 3 (2018), 960–981.
- [7] Xujin Chen, Guoli Ding, Xiaodong Hu, and Wenan Zang. 2012. The Maximum-Weight Stable Matching Problem: Duality and Efficiency. *SIAM Journal on Discrete Mathematics* 26, 3 (2012), 1346–1360.
- [8] Ágnes Cseh, Robert W Irving, and David F Manlove. 2016. The Stable Roommates Problem with Short Lists. In *International Symposium on Algorithmic Game Theory*. Springer, 207–219.
- [9] Argyrios Deligkas, George B Mertzios, and Paul G Spirakis. 2017. The Computational Complexity of Weighted Greedy Matching. In *Proceedings of the 31st AAAI Conference on Artificial Intelligence*. AAAI Press, 317–325.
- [10] Battal Doğan and Kemal Yıldız. 2016. Efficiency and Stability of Probabilistic Assignments in Marriage Problems. *Games and Economic Behavior* 95 (2016), 47–58.
- [11] Federico Echenique and Alfred Galichon. 2017. Ordinal and Cardinal Solution Concepts for Two-Sided Matching. *Games and Economic Behavior* 101 (2017), 63–77.
- [12] Federico Echenique, Alistair J Wilson, and Leeat Yariv. 2016. Clearinghouses for Two-Sided Matching: An Experimental Study. *Quantitative Economics* 7, 2 (2016), 449–482.
- [13] Yuval Emek, Tobias Langner, and Roger Wattenhofer. 2015. The Price of Matching with Metric Preferences. In *Proceedings of the 23rd Annual European Symposium on Algorithms*, Vol. 9294. Springer, 459–470.
- [14] Tomás Feder. 1992. A New Fixed Point Approach for Stable Networks and Stable Marriages. *J. Comput. System Sci.* 45, 2 (1992), 233–284.
- [15] David Gale and Lloyd S Shapley. 1962. College Admissions and the Stability of Marriage. *The American Mathematical Monthly* 69, 1 (1962), 9–15.
- [16] Dan Gusfield. 1987. Three Fast Algorithms for Four Problems in Stable Marriage. *SIAM J. Comput.* 16, 1 (1987), 111–128.
- [17] Dan Gusfield and Robert W Irving. 1989. *The Stable Marriage Problem: Structure and Algorithms*. MIT Press.
- [18] Robert W Irving. 1994. Stable Marriage and Indifference. *Discrete Applied Mathematics* 48, 3 (1994), 261–272.
- [19] Robert W Irving, Paul Leather, and Dan Gusfield. 1987. An Efficient Algorithm for the “Optimal” Stable Marriage. *J. ACM* 34, 3 (1987), 532–543.
- [20] Robert W Irving, David F Manlove, and Gregg O’Malley. 2009. Stable Marriage with Ties and Bounded Length Preference Lists. *Journal of Discrete Algorithms* 7, 2 (2009), 213–219.
- [21] Kazuo Iwama and Shuichi Miyazaki. 2008. A Survey of the Stable Marriage Problem and its Variants. In *International Conference on Informatics Education and Research for Knowledge-Circulating Society (ICKS)*. IEEE, 131–136.
- [22] Kazuo Iwama, Shuichi Miyazaki, Yasufumi Morita, and David Manlove. 1999. Stable Marriage with Incomplete Lists and Ties. In *International Colloquium on Automata, Languages, and Programming*. Springer, 443–452.

- [23] Donald Ervin Knuth. 1997. *Stable Marriage and its Relation to Other Combinatorial Problems: An Introduction to the Mathematical Analysis of Algorithms*. Vol. 10. American Mathematical Soc.
- [24] Tung Mai and Vijay V Vazirani. 2018. A Natural Generalization of Stable Matching Solved via New Insights into Ideal Cuts. *arXiv preprint arXiv:1802.06621* (2018).
- [25] Vikram Manjunath. 2013. Stability and the Core of Probabilistic Marriage Problems. (2013).
- [26] David F Manlove. 2013. *Algorithmics of Matching under Preferences*. Vol. 2. World Scientific.
- [27] David F Manlove, Robert W Irving, Kazuo Iwama, Shuichi Miyazaki, and Yasufumi Morita. 2002. Hard Variants of Stable Marriage. *Theoretical Computer Science* 276, 1-2 (2002), 261–279.
- [28] Maria Silvia Pini, Francesca Rossi, K Brent Venable, and Toby Walsh. 2013. Stability, Optimality and Manipulation in Matching Problems with Weighted Preferences. *Algorithms* 6, 4 (2013), 782–804.
- [29] Alvin E Roth and Elliott Peranson. 1999. The Redesign of the Matching Market for American Physicians: Some Engineering Aspects of Economic Design. *American Economic Review* 89, 4 (1999), 748–780.
- [30] Alvin E Roth, Uriel G Rothblum, and John H Vande Vate. 1993. Stable Matchings, Optimal Assignments, and Linear Programming. *Mathematics of Operations Research* 18, 4 (1993), 803–828.
- [31] Alvin E Roth, Tayfun Sönmez, and M Utku Ünver. 2004. Kidney Exchange. *The Quarterly Journal of Economics* 119, 2 (2004), 457–488.
- [32] Alvin E Roth and Marilda A Oliveira Sotomayor. 1990. *Two-Sided Matching: A Study in Game-Theoretic Modeling and Analysis*. Cambridge University Press.
- [33] Uriel G Rothblum. 1992. Characterization of Stable Matchings as Extreme Points of a Polytope. *Mathematical Programming* 54, 1-3 (1992), 57–67.
- [34] Chung-Piaw Teo and Jay Sethuraman. 1998. The Geometry of Fractional Stable Matchings and its Applications. *Mathematics of Operations Research* 23, 4 (1998), 874–891.
- [35] John H Vande Vate. 1989. Linear Programming Brings Marital Bliss. *Operations Research Letters* 8, 3 (1989), 147–153.
- [36] Rakesh V Vohra. 2012. Stable Matchings and Linear Programming. *Current Science* 105, 9 (2012), 1051–1055.
- [37] Ryo Yoshinaka. 2005. Higher-order Matching in the Linear Lambda Calculus in the Absence of Constants is NP-complete. In *Proceedings of the 16th International Conference on Term Rewriting and Applications*. Springer-Verlag, 235–249.


A APPENDIX

A.1 Proof of Theorem 1

PROOF. We will use the SMC instance shown in Figure 7, where, for some k (to be determined later), we have man m_i and woman w_i for all $i \in \{1, \dots, k\}$, men e_i^1 and e_i^2 for all $i \in \{1, \dots, k-1\}$, woman f_i^1 for all $i \in \{2, \dots, k-1\}$, man e_i^3 and women f_i^2 and f_i^3 for all $i \in \{2, \dots, k\}$. The valuations of the agents are as shown in Figure 7. We note that the instance shown in Figure 7 is a slight modification of the *accumulator* gadget used in the proof of Theorem 7 (see Figure 6a).

Consider a stable integral matching μ^s . By the stability requirement for the pair (e_1^1, w_1) , μ^s should contain either the pair (m_1, w_1) or the pair (e_1^1, w_1) . Either of these pairs contribute at most α to the social welfare (recall that $\alpha \geq 2$). Once again, the stability requirement for the pairs (m_i, f_i^2) for all $i \in \{2, \dots, k\}$ and (e_i^1, w_i) for all $i \in \{2, \dots, k-1\}$ will force these pairs to be contained in μ as well. Each of these pairs contributes 2 to the social welfare, and thus, $\mathcal{W}(\mu^s) \leq 4k - 6 + \alpha$.

Define a stable fractional matching μ as follows: We set $\mu(m_i, w_i) = 1/\alpha$ for all $i \in \{1, \dots, k\}$, $\mu(e_i^2, w_i) = 1 - 2/\alpha$, $\mu(m_{i+1}, w_i) = 1/\alpha$, $\mu(e_i^1, w_i) = 0$ for all $i \in \{1, \dots, k-1\}$, $\mu(m_i, f_i^2) = 0$, $\mu(m_i, f_i^3) = 1 - 2/\alpha$, $\mu(e_i^3, f_i^2) = 1$ for all $i \in \{2, \dots, k\}$, and $\mu(e_i^1, f_i^1) = 1$, for all $i \in \{2, \dots, k-1\}$ (note that we use a similar matching in the proof of Lemma 4). It can be easily seen that the social welfare of μ is $\mathcal{W}(\mu) = 4k(\alpha - 1/2) - 5\alpha + 3$. It is also not hard to check that μ is stable. The theorem follows by setting k to be sufficiently large. \square



PROOF. Consider an optimal stable fractional matching μ^* that is non-integral (i.e., has support of size at least two), and let μ be an integral matching in a support of μ^* . We will show that μ is a stable integral matching. In this way, we will have that all matchings in the support of μ^* are optimal stable integral matchings.

Assume otherwise that μ is not stable, and let (m, w) be a blocking pair in μ with $U(m, w) = V(m, w) = 1$. This means that either μ contains no pair involving agents m and w , or any such pair (m', w') with either $m' = m$ or $w' = w$ satisfies $U(m', w') = V(m', w') = 0$. Then, by replacing such pairs with (m, w) in μ (and, subsequently, in the support of μ^*), we get a stable fractional matching of even higher welfare than μ^* contradicting its optimality.

Now assume that (m, w) is a blocking pair in μ with $U(m, w) = V(m, w) = \alpha$. This means that either μ has no pair that contains agents m and w , or any such pair (m', w') with $m' = m$ and $w' = w$ satisfies $U(m', w') = V(m', w') = 1$. But then, since μ participates in the support of μ^* (i.e., with strictly positive weight) and no valuation is higher than α in \mathcal{I} , the utility of both agents m and w in μ^* will be strictly smaller than α , contradicting its stability. \square

By Lemma 5 and the results in [9], we immediately have the following.

COROLLARY 1. OPTIMAL STABLE FRACTIONAL MATCHING is NP-hard.

A.3 Hardness for ε -stability

The proof of Theorem 8 follows along similar lines to the proof of Theorem 7. Again, we present a reduction from 2P2N-3SAT but now we begin by augmenting the instance in the following way. For each variable of the original instance, we create a clone-variable and for each clause of the original instance we create a clone-clause that contains the clone-variables corresponding to the variables of the initial clause. Each variable and its clone are *coupled variables* and, similarly, each clause and its clone are *coupled clauses*.

Let the modified input consist of N (boolean) variables x_1, x_2, \dots, x_N , and a 3-CNF formula ϕ with $4N/3$ clauses $c_1, c_2, \dots, c_{4N/3}$. Note that if ϕ is not satisfiable, then there exist at least two clauses that are not satisfied.

A.3.1 *The reduction.* Instance \mathcal{I} again consists of variable gadgets, clause gadgets, VC-connectors, an accumulator, and CA-connectors.

In particular, instance \mathcal{I} contains a variable gadget (Figure 4a) for every variable x , and a clause gadget for every clause c as in the proof of Theorem 7 (see Figure 4b). The VC-connectors are as those in the proof of Theorem 7 for the case $\alpha \geq 2$ (see Figures 5a and 5c), while the accumulator, apart from the balanced *tine* edges (m_1, w^c) for every clause c , now contains just a single edge (m_1, w_1) such that $U(m_1, w_1) = 0$ and $V(m_1, w_1) = \beta$, where the value of β will be set later. Finally, each CA-connector (m^c, w^c) corresponding to clause c , is modified so that $U(m^c, w^c) = 0$ and $V(m^c, w^c) = 1$. All other woman-heavy, man-heavy, and balanced edges not explicitly mentioned above are as in the proof of Theorem 7. We set $\alpha = 2(1 - \varepsilon)$ for the value used in man-heavy and woman-heavy edges, and observe that, when $\varepsilon < 0.03$, it holds $(3\alpha^2 + 4)\varepsilon < 1/2$.

A.3.2 *Gadget properties.* We have completed the description of the reduction. We are ready to prove important properties of the several gadgets; these follow as Claims 13-16.

CLAIM 13. For every variable x , an ε -stable fractional matching μ satisfies at least one of the following conditions:

- (1) $\mu(m_1^x, w_1^{\bar{x}}) + \mu(m_1^x, w_2^{\bar{x}}) + \mu(m_1^x, f_1^x) \geq 1 - \varepsilon$ and $\mu(m_2^x, w_1^{\bar{x}}) + \mu(m_2^x, w_2^{\bar{x}}) + \mu(m_2^x, f_2^x) \geq 1 - \varepsilon$.
- (2) $\mu(m_1^x, w_1^{\bar{x}}) + \mu(m_2^x, w_1^{\bar{x}}) + \mu(e_3^x, w_1^{\bar{x}}) \geq 1 - \varepsilon$ and $\mu(m_1^x, w_2^{\bar{x}}) + \mu(m_2^x, w_2^{\bar{x}}) + \mu(e_3^x, w_2^{\bar{x}}) \geq 1 - \varepsilon$.

PROOF. Assume otherwise that $\mu(m_i^x, w_1^{\bar{x}}) + \mu(m_i^x, w_2^{\bar{x}}) + \mu(m_i^x, f_1^x) < 1 - \varepsilon$ and $\mu(m_1^x, w_j^{\bar{x}}) + \mu(m_2^x, w_j^{\bar{x}}) + \mu(e_3^x, w_j^{\bar{x}}) < 1 - \varepsilon$ for $i, j \in \{1, 2\}$. Then, since instance \mathcal{I} contains no man-heavy edge (m_i^x, w) and no woman-heavy edge $(m, w_j^{\bar{x}})$, the pair $(m_i^x, w_j^{\bar{x}})$ is ε -blocking; a contradiction. \square

As before, the two conditions in the statement of Claim 13 affect the weight of the input edges of the VC-connectors that are attached to the variable gadget in any ε -stable fractional matching. So, condition (1) implies that the weight assigned to each input edge of the VC-connectors that correspond to the two appearances of the positive literal x in clauses must be at most ε . To see why, observe that these input edges are incident to nodes m_1^x and m_2^x , and the total weight of all edges incident to each of these nodes cannot exceed 1. Similarly, condition (2) implies that the weight assigned to each input edge of the VC-connectors that correspond to the two appearances of the negative literal \bar{x} in clauses must be at most ε .

CLAIM 14. *Any ε -stable fractional matching, that assigns a weight of at most ε to the input edge of a VC-connector, must assign a weight of at most $\alpha\varepsilon$ to its output edge as well.*

PROOF. The claim holds trivially for VC-connectors corresponding to positive literals. Consider a VC-connector corresponding to a negative literal \bar{x} and a clause c containing it. Observe that, besides the input edge $(m^{\bar{x},c}, w_i^{\bar{x}})$, the edge $(m^{\bar{x},c}, w^{\bar{x},c})$ is the only balanced or man-heavy edge that is incident to man $m^{\bar{x},c}$ and the only balanced (or woman-heavy) edge incident to woman $w^{\bar{x},c}$. Hence, ε -stability of edge $(m^{\bar{x},c}, w^{\bar{x},c})$ requires a weight of at least $1 - \varepsilon - \alpha\zeta$ assigned to it when the weight assigned to edge $(m^{\bar{x},c}, w_i^{\bar{x}})$ is $\zeta \leq \varepsilon$. Then, the output edge, which is also incident to node $m^{\bar{x},c}$, must have a weight of at most $\varepsilon + (\alpha - 1)\zeta \leq \alpha\varepsilon$. \square

CLAIM 15. *Any ε -stable fractional matching, that assigns a weight of at most $\alpha\varepsilon$ to each output edge of the VC-connectors corresponding to clause c , must assign a weight of at most $3(\alpha^2 + 1)\varepsilon$ to the CA-connector of clause c as well.*

PROOF. Let ℓ_1, ℓ_2 , and ℓ_3 be the literals of clause c . Consider (for the sake of contradiction) an ε -stable fractional matching μ that assigns (1) a weight of at most $\alpha\varepsilon$ to each output edge of the VC-connectors corresponding to literals ℓ_i and clause c and (2) a weight of more than $3(\alpha^2 + 1)\varepsilon$ to edge (m^c, w^c) of the CA-connector for clause c . Note that condition (2) implies that the total weight on the edges (m^c, w_1^c) , (m^c, w_2^c) and (m^c, w_3^c) is strictly smaller than $1 - 3(\alpha^2 + 1)\varepsilon$. Since these are the only balanced or man-heavy edges incident to man m^c , the ε -stability of these edges is guaranteed by a utility of (at least) $1 - \varepsilon$ for each of the agents w_1^c, w_2^c , and w_3^c .

By condition (1) and since, besides the output edges of the VC-connectors, the edges (e_1^c, w_i^c) , (e_2^c, w_i^c) , and (m^c, w_i^c) are the only balanced or woman-heavy edges incident to agent w_i^c for $i \in \{1, 2, 3\}$, the weight assigned to these three edges is at least $1 - (\alpha^2 + 1)\varepsilon$. Hence, the total weight on the nine edges of the clause gadget is at least $3 - 3(\alpha^2 + 1)\varepsilon$, i.e., strictly more than 2 for the six edges incident to men e_1^c and e_2^c , violating the definition of a fractional matching. \square

CLAIM 16. *Any ε -stable fractional matching, that assigns a weight of at most $3(\alpha^2 + 1)\varepsilon$ to at least two CA-connectors, must assign a total weight of $1 - \varepsilon$ to the tine edges and a weight of at most ε to the single edge of the accumulator.*

PROOF. Assume that a weight of at most $3(\alpha^2 + 1)\varepsilon$ has been assigned to the edges (m^{c_1}, w^{c_1}) and (m^{c_2}, w^{c_2}) of the CA-connectors corresponding to some clauses c_1 and c_2 . Since these are the only edges for which agents w^{c_1} and w^{c_2} have positive value, and there is no man-heavy edge incident to agent m_1 , stability on the edges (m_1, w^{c_1}) and (m_1, w^{c_2}) requires that the total weight of the tine edges (m_1, w^c) (for every clause c) is (at least) $1 - \varepsilon$. Indeed, since $(3\alpha^2 + 4)\varepsilon < 1/2$ it is not possible to guarantee stability of the edges (m_1, w^{c_1}) and (m_1, w^{c_2}) by assigning weight at least $1 - (3\alpha^2 + 4)\varepsilon$

to each of them so that both w^{c_1} and w^{c_2} have utility at least $1 - \varepsilon$. Hence, the weight of the edge (m_1, w_1) of the accumulator is at most ε . \square

A.3.3 Proof of inapproximability.

LEMMA 6. *If formula ϕ is not satisfiable, then any ε -stable fractional matching of \mathcal{I} has welfare at most $56\alpha N + \beta\varepsilon$.*

PROOF. We first show that, if ϕ is not satisfiable, then any ε -stable fractional matching of \mathcal{I} assigns weight at most $3(\alpha^2 + 1)\varepsilon$ to at least two CA-connectors. For the sake of contradiction, consider an ε -stable fractional matching that assigns weight at most $3(\alpha^2 + 1)\varepsilon$ to at most one CA-connector; let c be the relevant clause, if such a clause exists. We will construct a truth assignment for formula ϕ (contradicting the assumption of the lemma) by repeating the following process for every clause $c' \neq c$ of ϕ . Let ℓ be a literal that appears in c' such that the output edge of the VC-connector, that corresponds to the appearance of ℓ in c , has weight greater than $\alpha\varepsilon$. Recall that such a literal certainly exists by Claim 15. We set ℓ to 1 (true). For every variable that has not received a value in this way, we arbitrarily set it to 1.

The above assignment satisfies all clauses except possibly for clause c . Since clause c is coupled with another clause c' that is satisfied, it suffices to assign one of the variables appearing in c the same value as its corresponding coupled variable appearing in c' . To show that it is also valid, we need to argue that there is no variable x such that both literals x and \bar{x} have been set to 1. Assume otherwise that this is the case. Furthermore, assume that literal x was set to 1 due to its appearance in a clause c_1 , and that this is its i_1 -th appearance (with $i_1 \in \{1, 2\}$). Also, literal \bar{x} was set to 1 due to its appearance in a different clause c_2 , where \bar{x} makes its i_2 -th appearance (again, $i_2 \in \{1, 2\}$). Hence, the output edge of the VC-connector that corresponds to literal x and clause c_1 (respectively, the VC-connector that corresponds to literal \bar{x} and clause c_2) has weight greater than $\alpha\varepsilon$. Then, by Claim 14, the input edges of both VC-connectors have weight greater than ε . As these input edges are incident to nodes $m_{i_1}^x$ and $w_{i_2}^{\bar{x}}$, Claim 13 yields that the total weight in the edges incident to some of the nodes $m_{i_1}^x$ and $w_{i_2}^{\bar{x}}$ is strictly higher than 1, contradicting the definition of a fractional matching.

Since any ε -stable fractional matching assigns a weight of at most $3(\alpha^2 + 1)\varepsilon$ to at least two CA-connectors, by Claim 16, the contribution of the accumulator to the welfare is at most $2(1 - \varepsilon) + \beta\varepsilon$ ($2(1 - \varepsilon)$ from the tine edges plus $\beta\varepsilon$ from the accumulator). The upper bound follows by considering the sum of valuations of all agents for edges that do not belong to the accumulator. This sum consists of

- total value of 20 for the ten balanced edges of each of the N variable gadgets,
- total value of α for the edges of each of the $2N$ VC-connectors corresponding to a positive literal,
- total value of $2 + 2\alpha$ for the edges of each of the $2N$ VC-connectors corresponding to a negative literal,
- total value of 19 for the nine balanced edges of each of the $4N/3$ clause gadgets and their corresponding CA-connectors.

It can be easily seen that $56\alpha N - 2(1 - \varepsilon)$ is a (loose) upper bound on the total value from these edges. \square

LEMMA 7. *If ϕ is satisfiable, then there exists an ε -stable fractional matching on \mathcal{I} that has welfare at least β .*

PROOF. Consider an assignment of boolean values to the variables that satisfies ϕ . We construct an ε -stable fractional matching μ in \mathcal{I} so that the contribution of the accumulator gadget to the welfare is at least β .

Variable gadgets. The weights on the edges of the variable gadget corresponding to variable x are:

- $\mu(m_1^x, w_1^{\bar{x}}) = \mu(e_3^x, w_1^{\bar{x}}) = \mu(e_3^x, w_2^{\bar{x}}) = \mu(m_2^x, w_2^{\bar{x}}) = 1/2$, $\mu(e_1^x, f_1^x) = \mu(e_2^x, f_2^x) = 1$, and $\mu(m_1^x, f_1^x) = \mu(m_1^x, w_2^{\bar{x}}) = \mu(m_2^x, f_2^x) = \mu(m_2^x, w_1^{\bar{x}}) = 0$ if x is true, and
- $\mu(e_3^x, w_1^{\bar{x}}) = \mu(e_3^x, w_2^{\bar{x}}) = 1/2$, $\mu(m_1^x, f_1^x) = \mu(m_2^x, f_2^x) = 1$, $\mu(m_1^x, w_1^{\bar{x}}) = \mu(m_2^x, w_2^{\bar{x}}) = \mu(m_1^x, w_2^{\bar{x}}) = \mu(m_2^x, w_1^{\bar{x}}) = \mu(e_1^x, f_1^x) = \mu(e_2^x, f_2^x) = 0$ if x is false.

Clause gadgets and CA-connectors. For every clause we select arbitrarily one of the true literals of the clause and call it *active*; since the assignment satisfies ϕ , there is certainly such a literal. Consider clause c and let ℓ_i (with $i \in \{1, 2, 3\}$) be its active literal; let i_1 and i_2 be the indices from $\{1, 2, 3\}$ than are different than i . We set $\mu(e_1^c, w_{i_1}^c) = \mu(e_2^c, w_{i_2}^c) = 1$ and $\mu(e_1^c, w_i^c) = \mu(e_2^c, w_i^c) = \mu(e_2^c, w_{i_1}^c) = \mu(e_2^c, w_{i_2}^c) = \mu(m^c, w_i^c) = \mu(m^c, w_{i_1}^c) = \mu(m^c, w_{i_2}^c) = 0$. We also assign a weight of 1 to the CA-connector corresponding to c , i.e., $\mu(m^c, w^c) = 1$.

VC-connectors. For every non-active VC-connector, we set the weight of its balanced edge (if it exists) to 1 and the weight of the remaining edges to 0. For every active VC-connector corresponding to the i -th appearance of the positive literal x as the j -th literal of clause c ($i \in \{1, 2\}$, $j \in \{1, 2, 3\}$), we set $\mu(m_i^x, w_j^c) = 1/2$.

For every active VC-connector corresponding to the i -th appearance of the negative literal x as the j -th literal of clause c ($i \in \{1, 2\}$, $j \in \{1, 2, 3\}$), we set the weights of its edges as follows: $\mu(m_i^{\bar{x}, c}, w_j^{\bar{x}}) = \mu(m_i^{\bar{x}, c}, w_j^c) = 1/2$ and $\mu(m_i^{\bar{x}, c}, w_i^{\bar{x}, c}) = 0$.

Accumulator. We set $\mu(m_1, w^c) = 0$ for every tine edge (m_1, w^c) of the accumulator. Furthermore, we set $\mu(m_1, w_1) = 1$. So, the contribution of the accumulator to the social welfare is β , as desired.

It can be easily verified that the total weight of the edges that are incident to any node is at most 1. Hence, μ is a valid fractional matching. Regarding stability, it suffices to verify that either the man or the woman of a balanced pair has a utility of at least $1 - \varepsilon$. \square

We are ready to prove Theorem 8. We select a value of β such that $\beta \geq \frac{56\alpha N}{1/\varepsilon - \varepsilon - \delta}$. By Lemma 6, the welfare of μ if ϕ was not satisfiable would be at most

$$56\alpha N + \beta\varepsilon \leq \beta(1/\varepsilon - \varepsilon - \delta) + \beta\varepsilon = \beta(1/\varepsilon - \delta).$$

By Lemma 7, we have that the welfare of μ if ϕ was not satisfiable would be at least $1/\varepsilon - \delta$ times smaller than the welfare \mathcal{I} could have if ϕ was satisfiable.

A.4 Approximation Algorithm for $\frac{1}{2}$ -stability

We will now provide a polynomial-time algorithm that computes a $\frac{1}{2}$ -stable fractional matching with welfare at least that of an optimal (exactly) stable fractional matching. Notice that unlike Theorems 5 and 6, where the quality of the computed matching is compared to the optimal matching μ^{opt} , the guarantee in Theorem 17 is considerably weaker.

THEOREM 17. *Let \mathcal{I} be an SMC instance and μ^* be an optimal stable fractional matching for \mathcal{I} . Then, a $\frac{1}{2}$ -stable fractional matching μ that satisfies $\mathcal{W}(\mu) \geq \mathcal{W}(\mu^*)$ can be computed in polynomial time.*

PROOF. Consider the mixed integer linear program (OPT-Stab) from Section 2.1 for finding an optimal stable fractional matching for \mathcal{I} . Relaxing the integrality constraint (8) to $y(m, w) \in [0, 1]$

results in a linear program. Since a stable fractional matching always exists (see Proposition 1), this relaxation is feasible. Let μ be a solution of the relaxed program. Since $\max\{y(m, w), 1 - y(m, w)\} \geq \frac{1}{2}$, we have that for every man-woman pair $(m, w) \in M \times W$, either $u_m \geq \frac{1}{2}U(m, w)$ or $v_w \geq \frac{1}{2}V(m, w)$, implying that μ is $\frac{1}{2}$ -stable. It is also clear that $\mathcal{W}(\mu) \geq \mathcal{W}(\mu^*)$ since μ^* satisfies (OPT-Stab). \square