



## Research Article

### COMPARATIVE PHYSICOCHEMICAL, PHYTOCHEMICAL AND HIGH PERFORMANCE THIN LAYER CHROMATOGRAPHY EVALUATION OF HEART WOOD AND SMALL BRANCHES OF *AQUILARIA AGALLOCHA* ROXB.

Bhavana Srivastava<sup>1\*</sup>, V.C. Sharma<sup>2</sup>, H. Sharma<sup>2</sup>, P. Pant<sup>3</sup>, A. D. Jadhav<sup>4</sup>

<sup>1</sup>Research Officer, <sup>2</sup>Senior Research Fellow, <sup>4</sup>Research Officer (S-4), National Research Institute for Ayurveda Siddha Human Resource Development, Gwalior, (M.P), India.

<sup>3</sup>Assistant Director, Central Council for Research in Ayurvedic Sciences, Janakpuri, New Delhi, India.

#### ABSTRACT

*Aquilaria agallocha* Roxb. commonly called as Agarú is a medicinal plant. Heartwood of this tree is widely used in Ayurveda for various diseases. Removal of heart wood from trunk of this tree may make this plant weak and susceptible to damage due to which availability of this plant may be difficult in near future. Present study outlines the concept of plant part substitution. Heart wood and small branches of *A. agallocha* are compared on the basis of physicochemical analysis, phytochemical analysis, total phenolic contents, total flavonoid contents and high performance thin layer chromatography (HPTLC) to evaluate the possibilities of using small branches instead of heart wood. Physicochemical parameters of heartwood and small branches and phytochemical analysis of *n*-hexane, ethyl acetate and ethanol extract of both heart wood and small branches were carried out using standard methods. Total phenolics and total flavonoids were estimated spectrophotometrically using Folin-ciocalteu assay and aluminum chloride assay methods, respectively. CAMAG HPTLC system equipped with semi-automatic applicator was used for HPTLC of *n*-hexane, ethyl acetate and ethanol extracts of stem bark and small branches using suitable mobile phases. Results of phytochemical analysis and HPTLC of *n*-hexane, ethyl acetate and ethanol extracts showed many similarities which suggest that small branches may have nearly similar active potency like heart wood and may be used as a substitute of heart wood after comparison and confirmation of same for pharmacological activities.

**KEYWORDS:** *Aquilaria agallocha*, HPTLC profile, Physicochemical analysis, Phytochemical analysis.

#### INTRODUCTION

Medicinal plants are extensively utilized throughout the world in providing health care for both humans and animals not only in diseased condition but also as potential material for maintaining proper health. Hence the numbers of plant species are individually under attack to accomplish the global demand. *A. agallocha* (Family: Thymeleaceae) commonly called as *Agarú* is a precious medicinal plant widely used in Ayurveda. As per Ayurvedic literature, heart wood of this plant is used *Aksiroga*, *Svasa*, *Karna roga*, *Kustha* and *visa*<sup>[1]</sup>. It is also reported for analgesic, anti-inflammatory<sup>[2]</sup>, anxiolytic, anticonvulsant<sup>[3]</sup>, antioxidant<sup>[4]</sup> and antibacterial<sup>[5]</sup> activities. Several chemical constituents like aquillochin<sup>[6]</sup>, 7,8-dimethoxy-2-[2-(3'-acetoxyphenyl)ethyl]chromones, 6-methoxy-2-(2-phenylethyl)chromones, 6,7-dimethoxy-2-(2-phenylethyl)chromones, abietane ester<sup>[7]</sup>, gmelofuran, agarol, 5-iso-propyl-7-methyl-4,5,5a,6,7,8-hexahydro-3H-naphtho-[1,8-bc]-furan-8 $\alpha$ -hydroxy-3-one<sup>[8]</sup>,  $\alpha$ -agarofuran, 10-epi- $\gamma$ -eudesmol, oxoagarospirol sesquiterpenes<sup>[9]</sup>,  $\alpha$ -agarofuran, agarofuran, dihydroagarofuran<sup>[10]</sup>, norketoagarofuran, 4-hydroxydihydroagarofuran, 3, 4-dihydroxydihydroagarofuran<sup>[11]</sup>,  $\alpha$ -agarofuran,  $\beta$ -

agarofuran, norketoagarofuran, (-)-10-epi-  $\gamma$  eudesmol, agrosipiral, jinkohol, jinkoh-cremol, kusunol, dihydrokaranone, jinkohol II, oxo- agrosipiral<sup>[12]</sup>, apigenin-4', 7-dimethyl ether<sup>[13]</sup> etc. have identified and isolated from this plant.

Removal of heart wood from trunk of this tree may affect the survival of this plant due to which availability of this plant may be difficult in near future for use in Indian system of medicine. To safeguard the survival of this plant and to ensure the availability of heartwood as raw material to manufacturers and dealers of Ayurveda, Siddha and Unani drugs, there is strong need to explore the possibility of substitution of heartwood of the trunk with suitable alternate. An approach which would satisfy the necessities of sustainable harvesting, yet simultaneously provide for health care needs, would be the substitution of heart wood with aerial part of the same plant. Present study is an attempt to evaluate the possibilities of using small branches in place of heart wood. Standard physicochemical parameters of small branches of *A. agallocha* have not been prepared yet. So work is also carried out to establish preliminary physicochemical standards of small branches.

## MATERIAL AND METHODS

### Plant material

The heart wood and small branches of *A. Agallocha* were collected from Ayurvedic Regional Research Institute, Itanagar, Arunachal Pradesh, India, identified and authenticated by Mr. N.K. Pandey, Research officer (Botany), NRIASHRD, Gwalior.

### Instrumentation

CAMAG HPTLC system (Muttentz, Switzerland) equipped with semi automatic TLC applicator Linomat IV, twin trough plate development chamber, Win CATS software version 1.4.2 and Hamilton (Reno, Nevada, USA) Syringe (100  $\mu$ l).

### Material and reagents

All chemicals, reagents and solvents used during the experiments were of analytical grade and HPTLC plates were purchased from E. Merck Pvt. Ltd. (Mumbai, India).

### Physicochemical parameters

Heart wood and small branches were studied for various physicochemical standards like foreign matter, loss on drying at 105°C, total ash, acid-insoluble ash, alcohol soluble extractive, water soluble extractive and pH of 10% solution using standard methods<sup>[14, 15]</sup>.

### Preliminary phytochemical screening

*n*-Hexane, ethyl acetate and ethanol extract of both heart wood and small branches were screened for the presence of phenols, tannins, carbohydrates, saponins, alkaloids, proteins, flavonoids, steroids, furanoids, coumarin, quinone and terpenoids by the standard methods of Harbone<sup>[16]</sup> and Kokate<sup>[17]</sup>.

### Estimation of total phenolic and flavonoid content

Five grams of each of the shade-dried plant material was pulverized into coarse powder and subjected to ethanolic extraction using soxhlet apparatus. The extracts were concentrated to dryness. The dried residues were then dissolved in 100 ml of 95% ethanol. The extracts were used for total phenolic and flavonoid assay.

The total phenolics content was determined by using the Folin-Ciocalteu assay<sup>[18]</sup>. An aliquot (1 ml) of extracts or standard solution of tannic acid (20, 40, 60, 80 and 100  $\mu$ g/ml) was added to a 25 ml volumetric flask, containing 9 ml of distilled water. A reagent blank was prepared using distilled water. One millilitre of Folin-Ciocalteu phenol reagent was added to the mixture and shaken. After 5 min, 10 ml of 7% Na<sub>2</sub>CO<sub>3</sub> solution was added to the mixture. The volume was then made up to the mark. After incubation for 90 min at room temperature, the absorbance against the reagent blank was determined at 550 nm with an UV/Vis spectrophotometer. Total phenolics content was expressed as  $\mu$ g tannic acid equivalents (TAE).

Total flavonoid content was measured by the aluminum chloride colorimetric assay<sup>[19]</sup>. An aliquot (1 ml) of extracts or standard solutions of quercetin (20, 40, 60, 80 and 100  $\mu$ g/ml) was added to a 10 ml volumetric flask containing 4 ml of distilled water. To the flask, 0.30

ml of 5% NaNO<sub>2</sub> was added and after 5 min, 0.3 ml of 10% AlCl<sub>3</sub> was added. After 5 min, 2 ml of 1M NaOH was added and the volume was made up to 10 ml with distilled water. The solution was mixed and absorbance was measured against the blank at 510 nm. The total flavonoid content was expressed as  $\mu$ g quercetin equivalents (QE).

### HPTLC profiles

HPTLC studies were carried out following the method of Sethi<sup>[20]</sup>, Stahl<sup>[21]</sup> and Wagner et al<sup>[22]</sup>. The heart wood and small branches were powdered coarsely. Ten gram powdered samples of each of heart wood and small branches were accurately weighed and exhaustively extracted by *n*-hexane, ethyl acetate and ethanol (each 100 ml) separately using soxhlet apparatus. The extracts were filtered and concentrated under reduced pressure and made up to 10 ml in standard flasks separately.

The mobile phase used for developing the *n*-hexane and ethyl acetate extracts of heart wood and small branches was toluene: ethyl acetate: formic acid (7: 3: 0.1 v/v) and for ethanol extract of heart wood and small branches was toluene: ethyl acetate (7: 3 v/v).

The samples were spotted in the form of bands of width 10 mm with a 100  $\mu$ l Hamilton syringe on aluminum TLC plates pre-coated with Silica gel 60 F<sub>254</sub> of 0.2 mm thickness with the help of TLC semi-automatic applicator Linomat IV attached to CAMAG HPTLC system, which was programmed through Win CATS software version 1.4.2. 10  $\mu$ l of each extracts of heart wood and small branches were applied in two tracks as 10 mm bands at a spraying rate of 10 seconds/ $\mu$ l. Track 1 was heart wood and track 2 was small branches for each of the extracts applied.

Development of the plate up to a migration distance of 80 mm was performed at 27  $\pm$  2°C with mobile phase for each extracts in a CAMAG HPTLC chamber previously saturated for 30 min. After development the plate was dried at 60°C in an oven for 5 min and visualized under wavelength 254 nm and 366 nm for ultra violet detection. The developed plate was then dipped in anisaldehyde sulphuric acid reagent for derivatization and dried at 105°C in hot air oven till the colour of the band appears and visualized under white light. Images were captured by keeping the plates in photodocumentation chamber and R<sub>f</sub> values were recorded by Win CATS software.

## RESULTS AND DISCUSSION

Physicochemical parameters like foreign matter, loss on drying at 105°C, ash values, acid insoluble ash, extractive values and pH are given in Table 1. These data can be used for identification of the drug. Both the parts of *A. agallocha* were found to possess little moisture and hence can be stored at room temperature without fear of spoilage. Approximately same value of alcohol soluble extractives for both heart wood and small branches indicates the presence of approximately same amount of polar extractable compounds in heart wood and small branches.

**Table 1: Physicochemical Parameters of Heart Wood and Small Branches of *A. agallocha*.**

S. No.	Parameters	Results	
		Heart wood	Small branches
1.	Foreign matter (% w/w)	Nil	Nil
2.	Loss on drying (% w/w)	6.074	6.78
3.	Total ash (% w/w)	3.13	3.94
4.	Acid insoluble ash (% w/w)	0.62	0.53
5.	Alcohol soluble extractive value (% w/w)	1.70	1.88
6.	Water soluble extractive value (% w/w)	3.0	7.55
7.	pH of 10 % aqueous solution	6.53	5.75

Phytochemical analysis of different extracts of heart wood and small branches are shown in Table 2. Results reveal the presence of similar phytochemicals in heart wood and small branches except for quinone and furanoids which were found present only in ethanolic extract of heartwood.

**Table 2: Phytochemical Analysis of Extracts of Heart Wood and Small Branches of *A. agallocha*.**

Phytochemicals	Heart wood			Small branches		
	<i>n</i> -Hexane	Ethyl acetate	Ethanol	<i>n</i> -Hexane	Ethyl acetate	Ethanol
Phenols	-ve	-ve	+ve	-ve	-ve	+ve
Tannins	-ve	-ve	+ve	-ve	-ve	+ve
Alkaloids	-ve	-ve	+ve	-ve	-ve	+ve
Carbohydrates	-ve	+ve	+ve	-ve	+ve	+ve
Saponins	-ve	-ve	-ve	-ve	-ve	-ve
Proteins	+ve	+ve	+ve	+ve	+ve	+ve
Steroids	-ve	+ve	-ve	-ve	+ve	-ve
Flavonoids	-ve	-ve	+ve	-ve	-ve	+ve
Coumarin	-ve	-ve	+ve	-ve	-ve	+ve
Quinone	-ve	-ve	+ve	-ve	-ve	-ve
Furanoids	-ve	-ve	+ve	-ve	-ve	-ve
Terpenoids	-ve	-ve	+ve	-ve	-ve	+ve


Total amount of phenolics and flavonoids content of ethanolic extract of heart wood and small branches of *A. agallocha* are summarized in Table 3. Results indicate that in comparison to small branches, heart wood had the high total phenolic and flavonoid content.

**Table 3: Total Phenolic and Total Flavonoid Content of Ethanolic Extracts of Heart Wood and Small Branches of *A. agallocha*.**

S. No.	Plant parts	Total phenolics µg of TAE/10 g dry weight*	Total flavonoids µg of QE/10 g dry weight*
1.	Heart wood	233.0 ± 0.260	272.0 ± 0.133
2.	Small branches	19.7 ± 0.001	224.0 ± 1.105

\*Values are expressed as Mean ± SD

Comparative HPTLC profile of *n*-hexane, ethyl acetate and ethanol extracts of heart wood and small branches of *A. agallocha* were recorded to reveal the chemical pattern of each extract. The HPTLC profile of *n*-hexane extract of both heart wood and small branches (Figure 1 and Table 4) showed one and two bands, respectively out of which only one band at  $R_f$  0.68 (black) found similar when visualized under UV at 254 nm. At UV 366 nm, heart wood and small branches showed five and four bands, respectively out of which one band at  $R_f$  0.62 (florescent blue) found similar. Visualization under white light after derivatization with anisaldehyde sulphuric acid reagent both heart wood and small branches showed two bands with similar  $R_f$ . This indicates the presence of many similar compounds in hexane extract of heart wood and small branches.




**Figure 1: HPTLC Profile of *n*-Hexane Extracts of Heart Wood and Small Branches of *A. agallocha* (Track 1: Heart Wood, Track 2: Small Branches)**

**Table 4: R<sub>f</sub> Value of *n*-Hexane Extract of *A. agallocha*.**

S. No.	Wavelength	R <sub>f</sub> value	
		Heart wood	Heart wood
1.	254 nm	0.68	0.27, 0.68
2.	366 nm	0.26, 0.33, 0.47, 0.62, 0.77	0.33, 0.62, 0.68, 0.74
3.	Visible light after derivatization	0.26, 0.33	0.26, 0.33

HPTLC profile of ethyl acetate extract of heart wood and small branches (Figure 2 and Table 5) showed two and one bands, respectively at UV at 254 nm out of which one band at R<sub>f</sub> 0.25 (black) found similar. At UV 366 heart wood and small branches showed five and six bands, respectively out of which two bands at R<sub>f</sub> 0.25 and 0.62 found similar but color of these bands found different in heart wood and small branches. Visualization under white light after derivatization with anisaldehyde sulphuric acid reagent, both heart wood and small branches showed six bands with similar R<sub>f</sub>. This indicates the presence of many similar compounds also in ethyl acetate extract of heart wood and small branches.




**Figure 2: HPTLC Profile of Ethyl Acetate Extracts of Heart Wood and Small Branches of *A. agallocha* (Track 1: Heart Wood, Track 2: Small Branches)**



**Table 5: R<sub>f</sub> Value of Ethyl Acetate Extract of *A. agallocha*.**

S. No.	Wavelength	R <sub>f</sub> value	
		Heart wood	Small branches
1.	254 nm	0.25, 0.52	0.25
2.	366 nm	0.21, 0.25, 0.41, 0.47, 0.62	0.23, 0.25, 0.46, 0.62, 0.70, 0.77
3.	Visible light after derivatization	0.26, 0.40, 0.52, 0.64, 0.80, 0.88	0.26, 0.40, 0.52, 0.64, 0.80, 0.88

HPTLC profile of ethanol extract of heart wood and small branches (Figure 3 and Table 6) showed two bands and no band when visualized under UV at 254 nm. At UV 366 heart wood and small branches showed nine and six bands, respectively out of which four bands at R<sub>f</sub> 0.26 (florescent blue), 0.63 (florescent blue), 0.72 (red), 0.80 (red) found similar. Visualization under white light after derivatization with anisaldehyde sulphuric acid reagent, both heart wood and small branches showed two bands with similar R<sub>f</sub>. This indicates the presence of many similar compounds in ethanol extract of heart wood and small branches also.

**Figure 3: HPTLC Profile of Ethanol Extracts of Heart Wood and Small Branches of *A. agallocha* (Track 1: Heart Wood, Track 2: Small Branches)****Table 6: R<sub>f</sub> Value of Ethanol Extract of *A. agallocha*.**

S. No.	Wavelength	R <sub>f</sub> value	
		Heart wood	Small branches
1.	254 nm	0.26, 0.58	No band
2.	366 nm	0.04, 0.16, 0.26, 0.34, 0.42, 0.47, 0.63, 0.72, 0.80	0.26, 0.39, 0.42, 0.63, 0.72, 0.80
3.	Visible light after derivatization	0.26, 0.29	0.26, 0.29

**CONCLUSION**

The present study carried out in *A. agallocha* to evaluate the possibilities of using small branches in place of heartwood will help sustainable utilization. The study will also be useful in identification and quality control of drug and can provide standard HPTLC profiles with selected solvent system for proper identification/ authentication of drug. Similarity in HPTLC profiles of *n*-hexane, ethyl acetate and ethanol extracts of heart wood and small branches suggests that small branches may have nearly similar active potency like heart wood and

may provide the base for further study to use the small branches as a substitute of heart wood of *A. agallocha*.

**ACKNOWLEDGEMENT**

Authors are thankful to CCRAS, New Delhi for financial assistance and Director General, CCRAS, New Delhi for providing facilities.

**REFERENCES**

1. The Ayurvedic Pharmacopoeia of India, Ministry of Health and Family Welfare, Department of Indian

- System of Medicine and Homeopathy. Part I Vol-IV New Delhi: Government of India. 1st Edition 1999.
2. T. Chitre, P. Bhutada, K. Nandakumar, R. Somani, P. Miniyar, Y. Mundhada, S. Gore and K. Jain, Analgesic and anti-inflammatory activity of heartwood of *Aquilaria agallocha* in laboratory animals, *Pharmacologyonline*, 1, 2007, 288-298.
  3. T. Alla, M. Handral, K. Nandakumar, N. Venkatrao, Shalam, Satyanaryana and S. M. Shantakumar, Anxiolytic and anticonvulsant activity of alcoholic extract of heartwood of *Aquilaria agallocha* Roxb (Thymelaeaceae) in mice, *Pharmacologyonline*, 1, 2007, 564-572.
  4. P.B. Miniyar, T.S. Chitre, S.S. Karve, H.J. Deuskar and K.S. Jain, Anti-oxidant activity of ethyl acetate extract of *Aquilaria agallocha* on nitrite-induced methemoglobin formation, *International Journal of Green Pharmacy*, 2, 2008, 43-45.
  5. T.K. Ghosh, H. Rahman, D. Bardalai and F. Ali, In-vitro antibacterial study of *Aquilaria agallocha* heart wood oil and *Citrullus lanatus* seed oil, *Scholars Journal of Applied Medical Sciences*, 1(1), 2013, 13-15.
  6. P. Bhandari, P. Pant and R.P. Rastogi, Aquillochin, a coumarinolignan from *Aquilaria agallocha*, *Phytochemistry*, 21, 1982, 2147-2149.
  7. H.Z. Alkhatlan, H.M. AL-Hazimi, F.S. AL-Dhalaan and A.A. Mousa, Three 2(2phenylethyl) chromones and two terpenes from agarwood, *Natural Product Research*, 19(4), 2005, 367- 372.
  8. P. Pant and R.P. Rastogi, Agarol, a new sesquiterpene from *Aquilaria agallocha*, *Phytochemistry*, 19(8), 1980, 1869-1870.
  9. M. Ishihara, T. Tsuneya, M. Suga and K.Uneyama, Three sesquiterpenes from agar wood, *Phytochemistry*, 30, 1991, 563-566.
  10. M.L. Maheshwari, T.C. Jain, R.B. Bates and S.C. Bhattacharyya, Terpenoids-xli: structure and absolute configuration of  $\alpha$ -agarofuran, agarofuran and dihydroagarofuran, *Tetrahedron*, 19(6), 1963, 1079-1090.
  11. M.L. Maheshwari, K.R. Varma and S.C. Bhattacharyya, Terpenoids XLVII: structure and absolute configuration of norketoagarofuran, 4-hydroxydihydroagarofuran, 3, 4-dihydroxydihydroagarofuran and conversion of  $\beta$ -agarofuran to  $\alpha$ -agarofuran, *Tetrahedron*, 19(10), 1963, 1519-1525.
  12. K. Yoneda, E. Yamagata, T. Nakanishi, T. Nagashima, I. Kawasaki, T. Yoshida, I. Mori, I. Miura, Sesquiterpenoids in two different kinds of Agarwood, *Phytochemistry*, 23(9), 1984, 2068-2069.
  13. A. Saraswathy, B. Vidhya and K. Amala, Apigenin-4', 7-dimethyl ether, *Indian Drugs*, 49(10), 2012, 30-32.
  14. Mukharjee PK. Quality Control of Herbal Drugs. New Delhi: Business Horizons Pharmaceutical Publishers; 2008. p. 186.
  15. Indian Herbal Pharmacopoeia. Indian Drug Manufacturers' Association. Revised edition. Mumbai: Indian Herbal Pharmacopoeia; 2002. p. 493-4.
  16. J.B. Harborne, *Phytochemical methods. A guide to modern techniques of plant analysis*: Chapman and Hall Co., New York, 3rd Edition 1998.
  17. C.K. Kokate, A.P. Purohit, and S.B. Gokhale, *Pharmacognosy*. Nirali Prakashan, Pune, India, 39th Edition 2007.
  18. V.L. Singleton and Jr. J.A. Rossi, Colorimetry of total phenolics with phosphomolybdic phosphotungstic acid reagents, *American Journal of Enology Viticulture*, 16, 1965, 144-158.
  19. J. Zhishen, T. Mengcheng and W. Jianming, The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals, *Food Chemistry*, 64 (4), 1999, 555-559.
  20. P.D. Sethi, *High performance thin layer chromatography*: CBS Publishers and Distributors, Vol. X. New Delhi, Student 1st Edition 1996.
  21. I. Stahl, *Thin layer chromatography. A laboratory hand book*: Springer- Verlag Berlin. 1969.
  22. H. Wagner and S. Bladt, *Plant drug analysis a thin layer chromatography*: Atlas, Springer-Verlag. Germany. 2nd Edition 1984.

**Cite this article as:**

Bhavana Srivastava, V.C. Sharma, H. Sharma, P. Pant, A. D. Jadhav. Comparative Physicochemical, Phytochemical and High Performance Thin Layer Chromatography Evaluation of Heart Wood and Small Branches of *Aquilaria Agallocha* Roxb. *International Journal of Ayurveda and Pharma Research*. 2016;4(1):1-6.

**Source of support: Nil. Conflict of interest: None Declared**

**\*Address for correspondence**

**Dr. Bhavana Srivastava**

Research Officer

National Research Institute for Ayurveda Siddha Human Resource Development, Aamkho, Gwalior-474009, M.P, India.

Email: [bhavanakan@gmail.com](mailto:bhavanakan@gmail.com)