

**Review Article****ETHNOMEDICINAL USES IN SCORPION POISONING (*GONUSU VISHA*) IN SRI LANKAN  
TRADITIONAL MEDICINE****E.R.H.S.S. Ediriweera<sup>1\*</sup>, W. D. Ratnasooriya<sup>2</sup>, K. W. M. D. Kamathewatta<sup>3</sup>**<sup>1</sup>Professor, Department of Nidana Chikithsa, Institute of Indigenous Medicine, University of Colombo, Rajagiriya, Sri Lanka,<sup>2</sup>Senior Professor, Faculty of Allied Health Sciences, General Sir John Kotelawala Defence University, Ratmalana, Sri Lanka,<sup>3</sup>Temporary Lecturer, Department of Dravyaguna Vignana, Institute of Indigenous Medicine, University of Colombo, Rajagiriya, Sri Lanka.**ABSTRACT**

Scorpions use their sting as a defensive weapon or as a mechanism to incapacitate prey. The scorpion sting possesses venom and the character of the poison depends on the scorpion species. Some may lead to death. Ayurveda and Sri Lankan traditional physicians treat scorpion sting by administering medicines internally and externally, in various forms. Data was gathered from Ayurveda and Sri Lankan traditional medical books, through interviewing physicians, research journals and internet. Herbs, minerals and animal sources are used to treat scorpion sting. Analgesic, anti-inflammatory, antihistaminic, antipyretic, antispasmodic, antioxidant, cardioprotective, respiratory protective, neuroprotective properties and anti-scorpion venom activity of these ingredients are scientifically proven. Due to these properties, these ingredients are capable of overcoming the symptoms such as pain, swelling, redness, itching, spasms, neurotoxicity and also cardiac and respiratory failure occurring through scorpion sting. It is concluded that Ayurveda and Sri Lankan traditional physicians possess multi-faceted, effective treatment methods for scorpion stings which can be scientifically proven.

**KEYWORDS:** Scorpion poisoning, *Gonusu Visha*, Ethnomedicinal uses, Sri Lanka.**INTRODUCTION**

Scorpions are eight-legged carnivorous arthropods that are found worldwide. They are mainly found in the southern hemisphere in deserts and jungle habitats alike. [1] Scorpions are generally nocturnal, meaning that they spend the day under rocks and in crevices and then come out to hunt in the safety of darkness.

There are many species of scorpions' worldwide. Fet *et al.*, (2000) list 16 families, 154 genera, and 1252 species of extant scorpions. While the number of families is now revised to 17 families, and it is acknowledged that this number will likely increase over the next few years as the research at the suprageneric level continues. Seventeen families of scorpion are Buthidae, Microcharmidae, Pseudochactidae, Chaerilidae, Chactidae, Euscorpiidae, Scorpipidae, Superstitioniidae, Troglotayosicidae, Iuridae, Vaejovidae, Bothriuridae, Ischnuridae, Heteroscorpionidae, Urodacidae, Diplocentridae and Scorpionida. [2]

Sri Lanka is home for 18 scorpions under 3 families. Whilst all scorpions are venomous, only one

species native to Sri Lanka is fatal to humans. [3] *Buthoscorpio sarasinorum*, *Charmus laneaus*, *Isometrus basilicus*, *Isometrus besucheti*, *Isometrus garyi*, *Isometrus loebli*, *Isometrus thwaitesi*, *Lychas ceylonensis* and *Lychas srilankanensis* belonging to the family Buthidae, are endemic to Sri Lanka. [3] *Chaerilus ceylonensis* belongs to the family Chaerilidae and *Heterometrus gravimanus* belongs to the family Scorpionidae are also endemic to Sri Lanka. [3] *Hottentotta tamulu*, *Isometrus acanthuru*, *Isometrus maculates* and *Isometrus thurstoni* belonging to the family Buthidae, and *Heterometrus indus*, *Heterometrus spinifer* and *Heterometrus swammerdami* belongs to the family Scorpionidae are nonendemic to Sri Lanka. [3] *Hottentotta tamulus* is the only dangerously venomous scorpion in Sri Lanka. However, globally around 50 scorpion species are thought to carry enough venom that could cause more serious harm to humans and around half of those 50 scorpion species are believed to be capable of causing fatal results when having stung a human being. [1]

Scorpion poisoning is known as *Vrishchika Visha* in Ayurveda medicine and *Gonusu Visha* in Sri Lankan Traditional medicine. Sushruta describes three types of scorpions based on poison; that is *Manda Visha* (mildly poisonous), *Madhya Visha* (moderately poisonous) and *Maha Visha* (virulently poisonous).<sup>[4]</sup> Mild, moderate and virulent scorpion species are twelve, three and fifteen in number respectively and total number is thirty.<sup>[4]</sup> Mildly poisonous scorpions are originated from putrefied cow dung. Those who originated from wood and bricks have moderate poison and those originated from sloughed snake or other poisonous substances are of virulent nature.<sup>[4]</sup>

According to Sushruta, sting of mildly poisoning scorpion produces pain, trembling, stiffness in body and out flow of black blood. If stung in extremities, pain rushes upwards along with burning sensation, sweating, local inflammatory swelling and fever.<sup>[4]</sup> Moderately poisoning scorpion produces disorders similar to that of snakes, which they are born of. Further, inflammation of tongue, obstruction of swallowing food and severe fainting may present due to sting from moderately poisoning scorpion.<sup>[4]</sup> With virulently poisoning scorpion sting, the symptoms of snake poisoning emerges, eruptive boils appear with giddiness, burning sensation, fever and virulent discharge of black blood from the passages may present which leads to death of the patient.<sup>[4]</sup>

Scorpion sting cases can be categorized into two general types: those involving localized, transitory symptoms usually lasting from few minutes to several hours, and those involving systemic reactions.<sup>[2]</sup>

Localized responses are characterized by immediate pain followed by moderate swelling at the sting site. In some cases, the sting may result in a raised, reddened, indurated lesion, even in the case of relatively harmless scorpions (e.g. *Vaejovis carolinianus*).

In cases involving cytolytic toxins (e.g. scorpionids and ischnurids) swelling may persist up to 72hr, followed by development of haemorrhages and blood-filled blisters near the sting site. Sloughing of skin may occur, but this varies greatly in severity. Other localized effects include gooseflesh, sweating, and muscle spasm near the sting site.

In case of stings, pain usually radiates from the site of the sting up the affected limb. The pain tends to concentrate in the joint, especially the armpits and groin, and often crosses from one armpit to another.

In the case of systemic reactions, clinical signs and symptoms are highly variable, ranging from mild

to life threatening. Systemic reactions commonly are mild and are not necessarily indicators of a serious problem. Often there is no appreciable swelling or discoloration of the sting site. An intense aching and burning sensation may spread to adjacent tissues, which in turn often throb, sometimes becoming numb. The acute pain at the sting site turns into a chronic dull pain accompanied by a feeling of numbness around the edge of the sting site, which may persist for one to several days. Numbness in the face, mouth, and throat is fairly common. Muscles may become spasmodic, resulting in muscular twitching, slurred speech, difficulty swallowing, tightness or cramps in the chest and back, rapid heartbeat, and nausea. Often these systemic responses persist less than an hour after the sting and are not considered as serious.<sup>[2]</sup>

In more severe systemic reactions, neurological effects can lead to profuse sweating and salivation, restlessness, extreme nervousness, respiratory and cardiovascular problems, mental confusion, and convulsions. As the clinical symptom indicates, the principle components of the venom of dangerous scorpions are neurotoxins.

#### **Neurotoxins mechanism**

These toxins act on autonomic, sympathetic, and neuro-muscular systems, causing the wide range of systemic reactions reported in sting victims.

They act by disrupting voltage-sensitive sodium and potassium channels of nerves, which in turn causes neural depolarization, prolonged action potentials, repetitive firing, and uncontrolled release of vasodilators and neurotransmitters, which affect virtually every major organ system.

The effects on neurotransmitters result in a depletive release of catecholamines (e.g., adrenaline, noradrenaline) that severely damages the heart and other organs. The most commonly reported cause of death in scorpion sting cases is cardiac failure. In other cases, respiratory failure maybe the cause, especially in patients with upper respiratory infections or related problems. Death usually occurs several days after envenomation. If symptoms subside during the first 2-12 hr following a sting, the prognosis for recovery is generally good. Mortality rates are quite variable, depending on the species and amount of venom injected.

Scorpion venom is a very complex mixture of substances which differs significantly among various taxa, within families, and among genera. Differences also occur in different geographic populations of the same species and even within the same populations. The toxins are low-molecular weight proteins which are among the most powerful toxin known. They are

comparable in some species to the neurotoxins of certain deadly snakes.

Two recognized types of neurotoxins are;

- $\alpha$ - scorpion toxin, characteristic of the genera *Androctonus*, *Leiurus*, and *Buthus*.
- $\beta$ - scorpion toxin, characteristic of *Centruroides*.<sup>[2]</sup>

The effect of envenomation by any given scorpion species can differ significantly among individual cases, owing to wide range of contributing factors.<sup>[5]</sup>

Sri Lankan traditional medicine and Ayurveda describe various medicines to neutralize these toxins. Aim of the study is to collect Sri Lankan traditional medicines which are used in treatment of scorpion sting.

## MATERIAL AND METHODS

Sri Lankan traditional physicians treat scorpion bites successfully. Treatments used in scorpion bites were gathered from Ayurveda and Sri Lankan traditional medical books, interviewing physicians, research journals and internet.

## RESULTS

Ayurveda and Sri Lankan traditional medicine prescribe various medicines internally and externally.

### a) Oral administration

1. Cow's milk mixed with ghee or sugar cane jaggery should be given orally to reduce scorpion poisoning.<sup>[4,6]</sup>
2. Jaggery dissolved in cold water and mixed with *Chaturjata* (stem bark of *Cinnamomum zeylanicum*, leaves of *Cinnamomum tamala*, seeds of *Elettaria cardamomum* and stamens of *Mesua ferrea* flowers) should be drunk for scorpion bites.<sup>[4]</sup>
3. Juice extracted from fresh leaves of *Jasminum officinale* (Family: Oleaceae; Sinhala name: *Da Saman*) is mixed with ghee and given to drink in scorpion sting.<sup>[6,7]</sup>
4. Juice extracted from fresh leaves of *Fluggea leucopyrus* (Family: Celastraceae; Sinhala name: *Katupila*) is mixed with cane sugar and given to drink in scorpion sting.<sup>[7]</sup>
5. Juice extracted from entire plant of *Amaranthus viridis* (Family: Amaranthaceae; Sinhala name: *Koora Thampala*) is mixed with cane sugar and given to drink in scorpion sting.<sup>[7]</sup>
6. Rock salt is mixed with ghee and given to drink in scorpion sting.<sup>[7]</sup>
7. Thirty grams (6 Kalan) roots of *Toddalia asiatica* (Family: Rutaceae; Sinhala name: *Kudumirissa*) and *Gmelina arborea* (Family: Verbenaceae; Sinhala name: *Eth Demata*) are mixed with 1920 ml of water and boiled down to 240ml. 120 ml of

this decoction is mixed with sugar and given twice a day with sugar.<sup>[8]</sup>

8. Thirty grams (6 Kalan) roots of *Toddalia asiatica* (Family: Rutaceae; Sinhala name: *Kudumirissa*) and *Gmelina arborea* (Family: Verbenaceae; Sinhala name: *Eth Demata*) are mixed with 1920 ml of water and boiled down to 240ml. 120 ml of this decoction is mixed with sugar and given twice a day with sugar.<sup>[8]</sup>
9. *Panchangaya* (root, stem, leaves, flowers, pods) of *Cassia auriculata* (Family: Fabaceae; Sinhala name: *Ranawara*) are mixed with 1920 ml of water and boiled down to 240ml. 120 ml of this decoction is mixed with sugar and given twice a day.<sup>[8]</sup>

### b) *Nasyakarma* (Errhine therapy)

1. Tubers of *Cyperus rotundus* (Family: Cyperaceae; Sinhala name: *Kalanduru*), fresh leaves of *Calotropis gigantea* (Family: Asclepiadaceae; Sinhala name: *Wara*) and *Cleome gynandra* (Family: Cleomaceae; Sinhala name: *Wela*) are ground with water and juice is extracted. *Nasya Karma* is performed by instilling extracted juice into nostrils of the persons with scorpion sting.<sup>[7]</sup>
2. Fresh leaves of *Calotropis gigantea* (Family: Asclepiadaceae; Sinhala name: *Wara*) and *Cleome gynandra* (Family: Cleomaceae; Sinhala name: *Wela*) are ground with water and juice is extracted. *Nasya Karma* is performed using extracted juice.<sup>[9]</sup>
3. Seeds of *Butea monosperma* (Family: Fabaceae; Sinhala name: *Kala*) and *Pongamia pinnata* (Family: Fabaceae; Sinhala name: *Magul Karanda*) are rubbed on a stone with water. The resultant is used in *Nasya Karma* in the persons with scorpion sting.<sup>[7,9]</sup>
4. Roots of *Pongamia pinnata* (Family: Fabaceae; Sinhala name: *Magul Karanda*) are rubbed on a stone with water. The resultant is used in *Nasya Karma* in the persons who are having heaviness in the head and headache due to scorpion sting.<sup>[8]</sup>
5. Fresh rhizome of *Zingiber officinale* (Family: Zingiberaceae; Sinhala name: *Inguru*) is pounded with water and juice is extracted. This is used in *Nasya Karma* in the persons with scorpion sting.<sup>[7]</sup>
6. Roots bark of *Vitex negundo* (Family: Verbenaceae; Sinhala name: *Nika*) is grounded and juice is extracted. *Nasya Karma* is performed using this extracted juice in treatment of scorpion sting.<sup>[10]</sup>
7. Seeds of *Erythrina variegata* (Family: Fabaceae; Sinhala name: *Erabadu*) and *Pongamia pinnata* (Family: Fabaceae; Sinhala name: *Magul Karanda*) are ground with water, juice is



extracted but squeezing and use in *Nasya Karma* in the persons with scorpion sting.<sup>[10]</sup>

### c) Local application

#### On the site of scorpion sting

1. Resin of *Ferula foetida* (Family: Umbelliferae; Sinhala name: *Perunkayam*) is to be dissolved in water and applied on the site of the scorpion sting.<sup>[7]</sup>
2. Leaves of *Pongamia pinnata* (Family: Fabaceae; Sinhala name: *Magul Karanda*) are crushed and applied on the site of the scorpion sting.<sup>[9]</sup>
3. Apply *Cassia occidentalis* (Family: Fabaceae; Sinhala name: *Peni Tora*) leaves dipped in salt water to the site of the scorpion sting.<sup>[9]</sup>
4. Resin of *Ferula foetida* (Family: Umbelliferae; Sinhala name: *Perunkayam*) is ground with juice of betel leaves and applied on the site of the scorpion sting.<sup>[11]</sup>
5. Leaves of *Acalypha indica* (Family: Euphorbiaceae; Sinhala name: *Kuppameniya*) are ground with lime (Calcium oxide) and applied to site of the scorpion sting.<sup>[7,9,12]</sup>
6. Entire plant of *Acalypha indica* (Family: Euphorbiaceae; Sinhala name: *Kuppameniya*) are pounded and juice extracted. This juice is mixed with lime(Calcium oxide) and applied to site of the scorpion sting.<sup>[11]</sup>
7. Juice extracted from fresh leaves of *Datura metel* (Family: Solanaceae; Sinhala name: *Attana*) is applied on the site of the scorpion sting.<sup>[7,9,12]</sup>
8. Dried resin-like gums of the tree *Shorea oblongifolia* (Family: Dipterocarpaceae; Sinhala name: *Dummala*) is mixed with ghee and applied on the site of scorpion sting.<sup>[7]</sup>
9. *Calamus thwaitesii* (Family: Arecaceae; Sinhala name: *Ma-wewel*) are ground with coconut vinegar and applied on the site of scorpion sting.<sup>[7]</sup>
10. Roots of *Achyranthes aspera* (Family: Amaranthaceae; Sinhala name: *Ratkaralheba*) are crushed and applied on the site of the scorpion sting.<sup>[13]</sup>
11. Seeds of *Achyranthes aspera* (Family: Amaranthaceae; Sinhala name: *Ratkaralheba*) are ground together with water and applied on the site of the scorpion sting.<sup>[11]</sup>
12. Ash obtained by burning coir is mixed with Margosa oil and applied on the site of the scorpion sting.<sup>[13]</sup>
13. Ash obtained by burning coir is mixed with coconut oil and applied on the site of the scorpion sting.<sup>[11]</sup>
14. Apply coconut oil on the site of the scorpion sting.<sup>[14]</sup>
15. Jute rope is burnt and ash is mixed with coconut oil. This mixture is applied on the site of the scorpion sting.<sup>[15]</sup>
16. Roots of *Calotropis gigantea* (Family: Asclepiadaceae; Sinhala name: *Wara*) is ground with lime juice and applied on the site of the scorpion sting.<sup>[15]</sup>
17. Leaves of *Calotropis gigantea* (Family: Asclepiadaceae; Sinhala name: *Wara*) are ground with urine and applied on the site of the scorpion sting.<sup>[12, 16]</sup>
18. Tender leaves of *Calotropis gigantea* (Family: Asclepiadaceae; Sinhala name: *Wara*), fresh rhizome of *Curcuma longa* (Family: Zingiberaceae; Sinhala name: *Kaha* ) and common salt are ground with lime juice and applied on the site of the scorpion sting.<sup>[17]</sup>
19. Seeds of *Carum bulbocastanum* (Family: Umbelliferae; Sinhala name: *Kaluduru*) and *Cuminum cyminum* (Family: Apiaceae; Sinhala name: *Suduru*) and rock salt are ground together with ghee. This paste is applied on the site of the scorpion sting.<sup>[15]</sup>
20. Curd is applied on the site of the scorpion sting and made to lick it by a cat.<sup>[7,9, 18]</sup>
21. Latex of *Calotropis gigantea* (Family: Asclepiadaceae; Sinhala name: *Wara*) is ground with ghee and applied on the site of the scorpion sting.<sup>[7]</sup>
22. Fresh rhizome of *Curcuma longa* (Family: Zingiberaceae; Sinhala name: *Kaha*) is ground with ghee and applied on the site of the scorpion sting.<sup>[7]</sup>
23. Pericarp of *Terminalia chebula* (Family: Combretaceae; Sinhala name: *Aralu*), Mace and cane sugar jaggery are ground together and applied on the site of the scorpion sting.<sup>[7]</sup>
24. Seeds of *Butea monosperma* (Family: Fabaceae; Sinhala name: *Kela*) are ground with latex of *Calotropis gigantea* and applied on the site of the scorpion sting.<sup>[7]</sup>
25. Droppings of crows, old rat droppings and centipedes are ground together and applied on the site of the scorpion sting.<sup>[7]</sup>
26. Leaves of *Benincasa hispida* (Family: Cucurbitaceae.; Sinhala name: *Puhul*) are ground and juice is applied to site of the scorpion sting.<sup>[19]</sup>
27. Leaves of *Cassia tora* (Family: Fabaceae.; Sinhala name: *Pethi Thora*) are pounded with common salt, heated on a pan and applied to site of the scorpion sting. <sup>[9,19]</sup> Mudannayaka, 1962 mentions that dried rhizome of *Curcuma longa* (Family: Zingiberaceae; Sinhala name: *Kaha*) can be added to above preparation.<sup>[12]</sup>

28. Leaves of *Emilia sonchifoli* (Family: Celastraceae; Sinhala name: *Kadupahara*) are ground with *Thandulodaka* (water obtained by washing raw rice) and applied to site of the scorpion sting.<sup>[11]</sup>
29. Tender part of the crown of *Carica papaya* (Family: Caricaceae; Sinhala name: *Gaslabu*) is ground with lime (Calcium oxide) and applied to site of the scorpion sting.<sup>[20]</sup>
30. Stem bark of *Moringa oleifera* (Family: Moringaceae; Sinhala name: *Murunga*) is ground with asafetida and applied to site of the scorpion sting.<sup>[20]</sup>
31. Stem bark of *Moringa oleifera* (Family: Moringaceae; Sinhala name: *Murunga*) is ground and applied to site of the scorpion sting.<sup>[12]</sup>
32. Fresh leaves of *Cassia tora* (Family: Caesalpiniaceae; Sinhala name: *Pethi Thora*) are pounded, juice is extracted by squeezing mixed with salt water and boiled. This is applied to site of the scorpion sting especially to reduce swelling.<sup>[7]</sup>
33. Latex of *Calotropis gigantea* (Family: Asclepiadaceae; Sinhala name: *Wara*), sesame seeds and chimney shoot are ground with salt water and boiled with juice of leaves of *C. tora* and applied on the site of the scorpion sting especially to reduce swelling.<sup>[7]</sup>
34. *Hiriyal* (Chemical name: *Orpiment/Arsenic Sulphide*; Sinhala name; *Hiriyal*) is ground with Latex of *Calotropis gigantea* (Family: Asclepiadaceae; Sinhala name: *Wara*) and applied on the site of the scorpion sting especially to reduce swelling.<sup>[7]</sup>
35. Rhizomes of *Acorus calamus* (Family: Acoraceae; Sinhala name: *Wadakaha*), roots of *Vitex negundo* (Family: Verbenaceae; Sinhala name: *Nika*) and *Ficus religiosa* (Family: Moraceae; Sinhala name: *Bo*), seeds of *Adenanthera pavonina* (Family: Fabaceae; Sinhala name: *Madatiya*) and *Pushkara* (Chemical name: Borax; Sinhala name: *Pushkara*) are ground with cow's urine or human's urine and applied on the site of the scorpion sting.<sup>[10]</sup>
36. Rhizomes of *Acorus calamus* (Family: Acoraceae; Sinhala name: *Wadakaha*), roots of *Trianthema portulacastrum* (Family: Nyctaginaceae; Sinhala name: *Sarana*) and chimney soot are ground with bee's honey and applied on the site of the scorpion sting. Equal parts of bee's honey and ghee are taken, mixed together and the patient is made to consume the mixture.<sup>[10]</sup>
37. Roots bark of *Vitex negundo* (Family: Verbenaceae; Sinhala name: *Nika*) is ground with rock salt and applied on the site of the scorpion sting.<sup>[10]</sup>
38. Equal quantities of *Manosila* (Chemical name: Arsenic sulfide; Sinhala name: *Manosila*), *Ferula asafetida* (Family: Apiaceae; Sinhala name; *Perunkaayam*) and rock salt (Chemical Name: Halite/Sodium Chloride; Sinhala name; *Saindava Lavana*) are taken, ground with cow's urine and applied on the site of the scorpion sting.<sup>[9,10]</sup>
39. Equal quantities of *Yavakshara* (*Yavakshara* is an alkali preparation of prepared with entire plant of *Hordeum vulgare*) dried rhizome of *Curcuma longa* (Family: Zingiberaceae; Sinhala name: *Kaha*), fruits of *Piper longum* (Family: Piperaceae; Sinhala name: *Thippili*), stem of *Coscinium fenestratum* (Family: Menispermaceae; Sinhala name: *Venivelgata*) and kernel of *Abrus precatorius* (Family: Fabaceae; Sinhala name: *Olinda*) are ground with cow's urine and applied on the site of the scorpion sting.<sup>[10]</sup>
40. Seeds of *Calotropis gigantea* (Family: Asclepiadaceae; Sinhala name: *Wara*) and *Piper nigrum* (Family: Piperaceae; Sinhala name: *Gammiris*) are taken in equal weights and powdered together and applied on the site of the scorpion sting.<sup>[21]</sup>
41. Seeds of *Butea monosperma* (Family: Fabaceae; Sinhala name: *Kala*) and *Calotropis gigantea* (Family: Asclepiadaceae; Sinhala name: *Wara*) are pounded and applied on the site of the scorpion sting.<sup>[21]</sup>
42. Fresh leaves of *Jasminum officinale* (Family: Oleaceae; Sinhala name: *Da Saman*) are ground with lime juice and applied on the scorpion sting.<sup>[6]</sup>
43. Leaves of *Nymphaea nouchal* (Family: Nymphaeaceae; Sinhala name: *Upul*), roots of *Clitoria ternatea* (Family: Fabaceae; Sinhala name: *Katarodu*), fruits of *Piper retrofractum* (Family: Piperaceae; Sinhala name: *Siviya*) and chimney soot are ground with cow's urine and applied on the scorpion sting.<sup>[6]</sup>
44. Roots of *Trianthema decandra* (Family: Aizoaceae; Sinhala name: *Sarana*), rhizome of *Acorus calamus* (Family: Acoraceae; Sinhala name: *Wadakaha*), and chimney soot are ground with cow's urine and applied on the scorpion sting.<sup>[18]</sup>
45. Leaves of *Ocimum sanctum* (Family: Lamiaceae; Sinhala name: *Madurutala*), are ground with juice of *Citrus medica* (Family: Rutaceae; Sinhala name: *Lapnaran*) and cow's urine and warm the mixture and applied as a paste on the scorpion sting.<sup>[4]</sup>
46. Leaves of *Andrographis paniculata* (Family: Acanthaceae; Sinhala name: Heen Binkohomba) applied as a paste on the scorpion sting.<sup>[21]</sup>

## Other sites of the body

1. *Thalap* (a paste-like Sri Lankan preparation) is prepared using stem bark of *Schleichera oleosa* (Family: Sapindaceae; Sinhala name: *Kon Gas*) and applied on the flanks of person bitten by the scorpion.<sup>[14]</sup>

### d) Blow on to the wound

1. Roots of *Achyranthes aspera* (Family: Amaranthaceae; Sinhala name: *Ratkaralheba*) is to be chewed and the breath blown on to the site of the scorpion sting.<sup>[9,22]</sup>

### e) Blow into the ears

1. Chew some leaves of *Cajanus cajan* (Family: Fabaceae; Sinhala name: *Rata Thora*) and blow the breath into the ear of person bitten by the scorpion.<sup>[7]</sup>

2. Roots of *Achyranthes aspera* (Family: Amaranthaceae; Sinhala name: *Ratkaralheba*) or *Gossipium arboreum* (Family: Malvaceae; Sinhala name: *Kapu*) are to be chewed and the breath blown into the ear of person bitten by the scorpion.<sup>[7]</sup>

### f) Fumigation (*Dum ellema*)

1. Peacock feather should be dipped in ghee and is set afire. It is then extinguished and the smoke arising is used to fumigate the site of scorpion bite.<sup>[7, 9,12,21, 22,]</sup>

2. Peacock and rooster feathers are mixed with rock salt and sesame oil, set afire and then snuffed out. The site of scorpion sting is fumigated with the smoke arising from this.<sup>[7,9]</sup> Further, Sushruta describes to add ghee to above mixture.<sup>[4]</sup>

3. The casualty is covered with a loose cloth. Then turmeric powder is burnt in embers of coconut shells. The smoke arising is directed under the cloth so that the whole body and the site of the sting gets fumigated.<sup>[7]</sup>

4. Droppings from peacocks, pigeons, roosters and roots of *Saminea saman* (Family: Fabaceae; Sinhala name: *Mara*) are powdered together and burnt over embers. The body and the site of the sting is fumigated with the smoke arising.<sup>[7]</sup>

5. Seeds of *Albizia odoratissima* (Family: Fabaceae; Sinhala name: *Mahari/Huri Mara*), roots of *Punica granatum* (Family: Lythraceae; Sinhala name: *Delum*), latex of *Calotropis gigantea* (Family: Asclepiadaceae; Sinhala name: *Wara*) and *Gomeda* gem stone (Hessonite) are heated together and the site of the sting is fumigated.<sup>[7]</sup>

6. Equal quantities of leaves of *Azadirachta indica* (Family: Meliaceae; Sinhala name: *Kohomba*), stem bark of *Cocos nucifera* (Family: Arecaceae; Sinhala name: *Thambili*) and old latex of *Artocarpus heterophyllus* (Family: Moraceae;

Sinhala name: *Kos*) are taken, pounded, mixed with sesame oil and burnt in embers of coconut (Family: Aracaceae; Botanical name; *Cocos nucifera*; Sinhala name; *pol*) shells. The scorpion sting is fumigated with the smoke arising from this. After that tender leaves of *Pagiantha dichotoma* (Family: Apocynaceae; Sinhala name: *Divi Kaduru*) are ground with salt water and applied on the site of the sting fumigated.<sup>[10]</sup>

7. Fruits of *Artocarpus heterophyllus* (Family: Moraceae; Sinhala name: *Kos*), leaves or stem bark of *Azadirachta indica* (Family: Meliaceae; Sinhala name: *Kohomba*) and *Sesamum indicum* seeds (Family: Pedaliaceae; Sinhala name; *Thalata*) are wrapped in a cloth and soaked in ghee and burnt. Flames are extinguish and the site of the scorpion sting is fumigated.<sup>[22]</sup>

### g) Fomentation (*Ushma Sweda / Vedu Paama*)

1. Entire plants of *Tragia involucrate* (Family: Euphorbiaceae; Sinhala name: *WelKahambiliya*), *Mimosa pudica* (Family: Fabaceae; Sinhala name: *Nidikumba*) and *Pogostemon heyneanus* (Family: Labiatae; Sinhala name: *Kollan Gas*) are boiled in water. The vapour is to be directed to the scorpion sting. Then the boiled residue should be ground and applied to the site of the scorpion bite.<sup>[10]</sup>

2. Rhizomes of *Acorus calamus* (Family: Acoraceae; Sinhala name: *Wadakaha*), roots of *Vitex negundo* (Family: Verbenaceae; Sinhala name: *Nika*) and *Pongamia pinnata* (Family: Fabaceae; Sinhala name: *Magul Karanda*) are boiled in water. The vapour is to be directed to the scorpion sting. Then the boiled residue should be ground and applied to the site of the scorpion bite.<sup>[10]</sup>

3. Entire plant of *Achyranthes aspera* (Family: Amaranthaceae; Sinhala name: *Gaskaralheba*) are boiled in cow's urine. The vapour is to be directed to the scorpion sting. Then the boiled residue should be ground and applied to the site of the scorpion bite.<sup>[10]</sup>

4. Leaves of *Erythrina variegata* (Family: Fabaceae; Sinhala name: *Erabadu*) are boiled in salt water. The vapour is to be directed to the scorpion sting. Then the boiled residue should be ground and applied to the site of the scorpion bite.<sup>[10]</sup>

5. Entire plant of *Calotropis gigantea* (Family: Asclepiadaceae; Sinhala name: *Wara*) is boiled in human urine. The vapour is to be directed to the scorpion sting. Then boluses are prepared with warm residue and the site of sting is fomented. After the residue is ground with water and applied to the scorpion bite.<sup>[10]</sup>

6. Entire plant of *Calotropis gigantea* (Family: Asclepiadaceae; Sinhala name: *Wara*) is boiled in



water. The vapour is to be directed to the scorpion sting.<sup>[8]</sup>

7. The site of the scorpion bite should be fomented with heated leaves of *Calotropis gigantea* (Family: Asclepiadaceae; Sinhala name: *Wara*).<sup>[9,22]</sup>
8. Leaves of *Calotropis gigantea* (Family: Asclepiadaceae; Sinhala name: *Wara*) are pounded and boluses are prepared using piece of cloth. These boluses are heated using steamer and the scorpion bite should be fomented with the boluses.<sup>[12]</sup>
9. Rhizome of *Alpinia galanga* (Family: Zingiberaceae; Sinhala name: *Heen Aratta*) is warmed by keeping under the hot ash in a hearth. The site of the scorpion bite is fomented using heated rhizome of *A. galanga*.<sup>[10]</sup>

#### h) Chanting (*Mantra*)

1. The site of the scorpion sting is chanted while blowing with stem of *Amorphophallus paeoniifolius* (Family: Araceae; Sinhala name: Kidaran) with the following Chant 'Om Kdham Hum Yam Thram Dham Vam Vam Lam Ksham Pham Phem Om Aum Ham Hah Ha Mam Cham Om' <sup>[7]</sup>
2. The physician's hand should be chanted with 'Om Namo Jah Jah' Manta. Then the site of the scorpion sting should be wiped downwards with physician's hand to remove the poison. <sup>[7]</sup>
3. The physician's hand should be chanted with "Namo Nalowa Guruluwa Nugayen Gaseem Daveem Thata Visa Natiya" Mantra. Then wipe the site of the scorpion sting with physician's hand).<sup>[17]</sup>
4. Gently rub the site of scorpion sting with coconut oil while repeatedly chanting the following Mantra, until the pain stops.  
"Om Theri Theri Yesvah" <sup>[12]</sup>

#### i) Beneficial food

1. Stem bark of *Schleichera oleosa* (Family: Sapindaceae; Sinhala name: *Kon*) is mixed with water and cooked until it become thick. It is called *Kon Thalapa*. This is given as a food to person bitten by the scorpion.<sup>[14]</sup>

#### DISCUSSION

According to the present study, 56 herbs, three minerals and two ingredients of animal origin are used in treatment of scorpion bites by Ayurveda and traditional physicians in Sri Lanka. Some of the medicines are administered orally and most of the medicines are applied externally on the site of scorpion bite or other places of the body. Some medicines are blown on to the wound or blown into the ears. Further, fomentation (*Ushma Sweda / Vedu Paama*), fumigation (*Dum Ellema*) and *Nasya Karma*

(Errhine therapy) are also performed to treat scorpion bites. In addition to these therapies, occult chanting (*Mantras*) are used to remove poison from the body and this is known as Visha Bassima. Various foods beneficial in scorpion bites are also described in texts.

Many plants used in Sri Lanka for treatment of scorpion bites are also used in other countries. *Amaranthus viridis*, *Achyranthes aspera*, *Calotropis gigantea*, and *Ferula foetida* are used to treat scorpion bites in Pakistan.<sup>[23]</sup> *Amaranthus viridis*, *Carum carvi*, *Achyranthes aspera*, *Cyperus rotundus*, *Abrus precatorius* and *Ricinus communis* are used to treat scorpion bites in Saudi Arabia.<sup>[24]</sup>

Entire plant of *Acalypha indica* is used to treat scorpion bites. Analgesic and anti-inflammatory activities.<sup>[25]</sup> of entire plant, antihistaminic, antipyretic and cardioprotective effects of leaves<sup>[26-28]</sup> antioxidant effect of aerial part<sup>[29]</sup> and Neuroprotective property of roots<sup>[30]</sup> of *A. indica* are scientifically proven.

Entire plant, roots and seeds of *Achyranthes aspera* are used to treat scorpion bites. Analgesic and anti-inflammatory activities<sup>[31]</sup> of roots and leaves, antihistaminic and antispasmodic<sup>[32]</sup> and cardio protective properties of entire plant<sup>[33]</sup>, antipyretic property of leaves<sup>[34]</sup> and antioxidant property of entire plant<sup>[35]</sup> of *A. aspera* are scientifically proven.

Rhizomes of *Acorus calamus* is used in treatment of scorpion bites. Analgesic<sup>[36]</sup> anti-inflammatory<sup>[37]</sup> antipyretic<sup>[38]</sup> antioxidant<sup>[39]</sup>, antispasmodic<sup>[40]</sup> and properties of rhizome<sup>[41]</sup>, cardioprotective<sup>[42]</sup> property of entire plant and antihistaminic<sup>[43]</sup> effect of leaves of *A. calamus* are scientifically proven.

Rhizomes of *Alpinia galanga* is used in treatment of scorpion bites and its analgesic,<sup>[44]</sup> anti-inflammatory<sup>[45]</sup> and neuroprotective properties<sup>[46]</sup> have been scientifically proven.

Entire plant of *Amaranthus viridis* is used to treat scorpion bites. Analgesic, antipyretic<sup>[47]</sup> and cardioprotective<sup>[48]</sup> effects of entire plant and anti-inflammatory<sup>[49]</sup> and antioxidant<sup>[50]</sup> properties of leaves of *A. viridis* are scientifically proven.

Leaves of *Andrographis paniculata* is used to treat scorpion bites. Analgesic, antipyretic and anti-inflammatory effects of Andrographolide derivatives of *Andrographis paniculata*<sup>[51]</sup>, antihistaminic, and cardioprotective effects of leaves<sup>[52,53]</sup>, anti-scorpion activity <sup>[54]</sup> of aerial part of the plant are scientifically proven.

Fruits and old latex of *Artocarpus heterophyllus* are used to treat scorpion bites. Anti-inflammatory effects of phenolic compounds isolated

from the fruits of *A. heterophyllum* is proven scientifically.<sup>[55]</sup>

Leaves of *Benincasa hispida* are used to treat scorpion bites. Analgesic effect of *B. hispida* is proven scientifically.<sup>[56]</sup>

Leaves of *Cajanus cajan* are used in treatment of scorpion bites and its property of neuroprotection is proven scientifically.<sup>[57]</sup>

Leaves, roots, seeds, latex and entire plant of *Calotropis gigantea* are used in treatment of scorpion bites. Analgesic effect of flowers and latex<sup>[58,59]</sup>, anti-inflammatory and cardioprotective effects of latex<sup>[60,61]</sup>, antioxidant property of leaves and entire plant<sup>[62,63]</sup>, antipyretic and neuroprotective property of roots<sup>[64,65]</sup>, of *C. gigantea* are proven scientifically.

Leaves of *Carica papaya* are used in treatment of scorpion stinging and their analgesic<sup>[66]</sup>, anti-inflammatory<sup>[67]</sup>, antipyretic<sup>[68]</sup>, and antioxidant<sup>[69]</sup>, properties of leaves are proven scientifically.

Seeds of *Carum bulbocastanum* are used in treatment of scorpion bites and their analgesic effect is scientifically proven.<sup>[70]</sup>

Leaves of *Cassipourea* are used in treatment of scorpion bites and their anti-inflammatory<sup>[71]</sup> and neuroprotective<sup>[72]</sup> properties are scientifically proven.

Fruits of *Citrus medica* are used in treatment of scorpion bites. Analgesic,<sup>[73]</sup> antioxidant<sup>[74]</sup> and cardioprotective<sup>[75]</sup> properties of fruits of *C. medica* are scientifically proven.

Leaves of *Cleome viscosa* are used in treatment of scorpion bites and their analgesic<sup>[76]</sup> and antioxidant<sup>[77]</sup> properties of leaves are scientifically proven.

Roots of *Clitoria ternatea* are used in treatment of scorpion bites. Analgesic and anti-inflammatory,<sup>[78]</sup> antihistaminic, antipyretic, antioxidant, respiratory protective and neuroprotective<sup>[79]</sup> properties of roots are proven scientifically.

Stem of *Coscinium fenestratum* is used in treatment of scorpion bites. Analgesic,<sup>[80]</sup> anti-inflammatory<sup>[81]</sup> and antioxidant<sup>[82]</sup> activities of *C. fenestratum* are proven scientifically.

Rhizome of *Curcuma longa* is used in treatment of scorpion stinging. Analgesic,<sup>[83]</sup> anti-inflammatory,<sup>[84]</sup> anti spasmic<sup>[85]</sup> antioxidant,<sup>[86]</sup> cardioprotective<sup>[87]</sup> and neuroprotective<sup>[88]</sup> properties of rhizome of *C. longa* have been scientifically proven.

Tubers of *Cyperus rotundus* are used in treatment of scorpion stinging. Antihistaminic and antipyretic,<sup>[89]</sup> anti spasmic,<sup>[90]</sup> cardio protective<sup>[91]</sup>

and neuroprotective<sup>[92]</sup> properties of tubers of *C. Rotundus* have been scientifically proven.

Leaves of *Emilia sonchifolia* are used in treatment of scorpion bites. Analgesic, anti-inflammatory<sup>[93]</sup> and antioxidant<sup>[94]</sup> properties of leaves of *E. sonchifolia* are proven scientifically.

Leaves and seeds of *Erythrina variegata* are used in treatment of scorpion bites. Analgesic and anti-inflammatory<sup>[95]</sup> properties of leaves of *E. variegata* are scientifically proven.

Latex (gum oleoresin) of *Ferula foetida* is used to treat scorpion bites. Analgesic and anti-inflammatory,<sup>[96]</sup> antispasmodic,<sup>[97]</sup> antioxidant<sup>[98]</sup> and cardioprotective<sup>[99]</sup> properties of latex (gum oleoresin) of *F. foetida* are scientifically proven.

Leaves of *Fluggea leucopyrus* are used in treatment of scorpion stinging. Anti-inflammatory and antihistaminic<sup>[100]</sup> properties of aerial part of *F. leucopyrus* have been scientifically proven.

Leaves and roots of *Gossypium arboreum* (Kapu) are used in treatment of scorpion stings and anti-inflammatory<sup>[101]</sup> activity of leaves has been scientifically proven.

Seed oil of *Azadirachta indica* (Kohomba) is used in treatment of scorpion bites. Analgesic<sup>[102]</sup> and anti-inflammatory<sup>[103]</sup> properties of seed oil of *A. indica* are scientifically proven.

Entire plant of *Mimosa pudica* is used in treatment of scorpion bites. Analgesic and anti-inflammatory activities of leaves<sup>[104]</sup> and anti-scorpion-venom<sup>[105]</sup> are scientifically proven.

Stem bark of *Moringa oleifera* is used in treatment of scorpion bites. Cardioprotective<sup>[106]</sup> property of stem bark of *M. oleifera* is scientifically proven.

Leaves of *Ocimum sanctum* are used in treatment of scorpion bites. Anti-inflammatory<sup>[107]</sup> antioxidant<sup>[108]</sup> cardioprotective<sup>[109]</sup> properties of leaves of *O. sanctum* are proven scientifically.

Fruits of *Piper longum* are used in treatment of scorpion bites. Anti-inflammatory<sup>[110]</sup> antioxidant and cardioprotective<sup>[111]</sup> activities of fruit of *P. longum* have been proven scientifically

Entire plant of *Pogostemon heyneanus* is used in treatment of scorpion bites. Antioxidant<sup>[112]</sup> property of leaves of *P. heyneanus* has been proven scientifically.

Roots, leaves and seeds of *Pongamia pinnata* are used in treatment of scorpion bites. Analgesic and antipyretic,<sup>[113]</sup> anti-inflammatory and antihistaminic<sup>[114]</sup> and antioxidant<sup>[115]</sup> property of leaves are scientifically proven.

Stem bark of *Schleichera oleosa* is used in treatment of scorpion bites. Analgesic and anti-



inflammatory<sup>[116]</sup> and antihistaminic<sup>[117]</sup> properties of stem bark have been scientifically proven.

Pericarp of *Terminalia chebula* is used in treatment of scorpion bites. Analgesic and anti-inflammatory,<sup>[118]</sup> antioxidant,<sup>[119]</sup> cardioprotective,<sup>[120]</sup> and neuroprotective,<sup>[121]</sup> properties of the pericarp are scientifically proven.

Entire plant of *Tragia involucrate* Linn is used in treatment of scorpion stinging. Anti-inflammatory and analgesic activity of root,<sup>[122]</sup> antihistaminic and respiratory protective property<sup>[123]</sup> of leaves, Nephroprotective and antioxidant property<sup>[124]</sup> of aerial part of the plant are scientifically proven.

Rhizomes of *Zingiber officinale* (Inguru) are used in treatment of scorpion bites. Anti-inflammatory and analgesic,<sup>[125]</sup> antioxidant,<sup>[126]</sup> cardioprotective<sup>[127]</sup> properties of rhizome are scientifically proven.

Cow's urine and ghee are used in treatment of scorpion bites. Analgesic and antioxidant properties<sup>[128,129]</sup> of cow's urine are scientifically proven.

Analgesic, anti-inflammatory and antihistaminic properties of some of the ingredients used to treat scorpion stings are scientifically proven. Therefore, these ingredients help to reduce pain, redness and odema caused by scorpion bites.

Muscle spasm is one of the effects appearing near the sting site. *Achyranthes aspera*,<sup>[32]</sup> *Acorus calamus*,<sup>[40]</sup> *Curcuma longa*,<sup>[85]</sup> *Cyperus rotundus*,<sup>[89]</sup> *Ferula foetida*<sup>[97]</sup> also possess anti-spasmodic effect. External application of these plants may become helpful to overcome muscular spasms occurring at the site of scorpion sting.

Pyrexia may occur in scorpion bites. Antipyretic effects of some ingredients mentioned in this study are proven. Therefore, these medicines may be effective in controlling fever.

A study was carried out to evaluate the anti-scorpion venom (ASV) property of *Andrographis paniculata* (AP) in comparison with anti-red scorpion venom serum and this study was aimed to determine its combined effect with anti-red scorpion venom serum. It is concluded that both anti-scorpion venom (ASV) and ethanolic extract of AP, have scorpion venom neutralization property *in vivo* and *in vitro*, when used individually, but their combination provides maximum benefit in all the aspects of anti-venom property. However, further studies are required for elaboration of potentiating action of plant with ASV in humans. Considering the scorpion sting as acute medical emergency, further studies to prepare injectable form of extract of AP will be helpful. Larger *in vivo* doses may be more beneficial.<sup>[54]</sup>

*Mesobuthus tamulus* (red scorpion) is common in many parts of India. Acute toxicity of the scorpion venom and its neutralization by the *Mimosa pudica* plant extract in-vivo was done. In the acute toxicity and in-vivo neutralization by plant extract in the dose of 1gm/Kg and 2gm/Kg resulted in the mean survival time of 7 minutes and 9.83 minutes respectively. In combination of prazosin+ plant extract the mean survival time was increased to 20.83 minutes. This protective property of *Mimosa pudica* can be exploited in practice where a significant amount of time is lost while shifting the patient from the Primary Health Care Centre to the Tertiary Health Care Centre.<sup>[105]</sup>

Antioxidant property of many ingredients is scientifically proven. An antioxidant is a molecule that inhibits the oxidation of other molecules. Oxidation is a chemical reaction that can produce free radicals, leading to chain reactions that may damage cells. Antioxidants help to regenerate damaged cells. Dousset *et al.* (2005) hypothesized that scorpion toxins could induce the generation of high levels of free radicals responsible for membrane damage in organs targeted by venom action. They have shown that scorpion toxins cause considerable lipid peroxidation in most vital organs. They have concluded that free radical generation only plays a minor role in the toxicity of scorpion venom.<sup>[130]</sup> Even though, antioxidants may play a role in overcoming the toxicity of scorpion venom.

Leaves of *Andrographis paniculata* strongly protect the myocardium against isoproterenol-induced infarction and suggest that the cardioprotective effects could be related to antioxidant activities.<sup>[131]</sup> Deaths due to scorpion sting occur through cardiac failure. Therefore, *Andrographis paniculata* is beneficial in preventing cardiac failure in scorpion stings.

*Acalypha indica* leaves possess possibly active metabolites that protect the cardiac tissue from cardiovascular diseases.<sup>[28]</sup> Eliyas (2011) reported that entire plant of *Achyranthes aspera* has significant dose dependent protective effect on heart against isoproterenol induced myocardial infarction probably by improving endogenous antioxidant enzyme activities.<sup>[33]</sup>

Respiratory failure is another cause of death especially in patients with upper respiratory infections or related problems. *Achyranthes aspera* and *Clitoria ternatea* also possess anti-asthmatic effects.<sup>[32,79]</sup> *Andrographis paniculata* is effective in treatment of respiratory infections.<sup>[132]</sup> These herbs will help to prevent respiratory failure in scorpion sting.

Although it is well established that symptomatology, morbidity and death following scorpion envenomation are due to increases in neurotransmitter release secondary to toxins binding to voltage-sensitive sodium channels, the mechanism by which venom action is involved in damaging heart, liver, lungs and kidneys remains unclear<sup>[130]</sup>. Treatments mentioned in the present study to treat scorpion bites are beneficial because some of the ingredients have the neuroprotective property and act as a brain tonic.

## CONCLUSION

Ayurvedic and Sri Lankan traditional physicians are rich with knowledge on treatment of scorpion bites. They use various methods to administer medicines, internally and externally. The properties which are capable of overcoming effects of scorpion bites in some ingredients used for these treatments are scientifically proven.

## REFERENCE

1. Anonymous, 2017, Scorpion (Scorpiones) - Animals - A-Z Animals, Cited on 26 Sept 2017, Available from: <https://a-z-animals.com/animals/scorpion/>
2. Mullen, G.R., and Stockwell, S.A.(2002).Scorpions. Medial and veterinary entomology. Elsevier science. USA.411-423
3. Anonymous, 2016, List of scorpions of Sri Lanka, Cited on 26 Sept 2017, Available from: [http://en.wikipedia.org/wiki/List\\_of\\_scorpions\\_of\\_Sri\\_Lanka](http://en.wikipedia.org/wiki/List_of_scorpions_of_Sri_Lanka)
4. Sharma, P.V. (ed. and trans.) 2010, Sushruta Samhita, Chaukambha Visvabharati, Varanasi, India 87- 90.
5. Isbister, G.K.,Volschenk,E.S., Balitb,C.R.,Harveyd, M.S., (2003) Australian scorpion stings: a prospective study of definite stings. Toxicon 41. 877-883.
6. Jayathilaka, K. G. P., 1957, Visha Vidyava, publisher not mentioned, 129 -130.
7. Liyanaarchchi, S.K., 1952, Sithiyam Sahita Visha Vaidya Chinthamani, Nuwana Printers, Colombo, Sri Lanka 1:72-7.
8. Gunarathna, S., 1992, Visha Vidyawa hevath Sarpavisha Samharaya, Ratna publishers, Colombo, Sri Lanka, 24-25.
9. Gunasena, D., 1955, Visha Vaidya Prakaranaya, Modern printers, Nugegoda, Sri Lanka, 64-65.
10. Liyanaarchchi, S.K., 1972, Visha Veda Muthuhara, Diamond Printers, Colombo, Sri Lanka 23- 29.
11. Anonymous, 1986, Ath Beheth, Department of Ayurveda, Colombo, Sri Lanka,116.
12. Mudannayaka, C. S., 1962, Purana Sarpa Vedapotha, Gunasekara printers, Aluthgama, Sri Lanka, 27.
13. Ukwattage, S., 2003, Ayurveda Osu (Ath Beheth), Samayawardhana Printers, Colombo, Sri Lanka 41.
14. Deraniyagala, P. A. P., 1952, Es Vedakama, Ceylon National Manuscript series Vol 3, Government printers, Colombo, Sri Lanka, 31.
15. Jayasekara, D. C., 1948, Pavule Veda Potha, Sarasavi printers, Gampaha, Sri Lanka, pp.251.
16. Deraniyagala, P. A. P., 1956, Sarpa Vedakama, Ceylon National Manuscript series Vol 3, Government printers, Colombo, Sri Lanka, pp.11.
17. Karunarathna, S., year not mentioned, Nivase Ath Veda Potha, Vasana publishers, Negombo, Sri Lanka, 40 - 43.
18. Liyanaarchchi, S.K., 1977, Sarpa Visha Prathamadhara, Gunasena and Company, Anuradhapura, Sri Lanka pp.5.
19. Anonymous, 1993, Thalpathe Piliyam, Department of Ayurveda, Colombo, Sri Lanka, pp.74.
20. Senavirathna, L., 2005, Deshiya Ath Beheth, Chatura printers, Wellampitiya, Sri Lanka, pp. 69.
21. Kannangara, J.C., 1892, Vishaushadha Samgraha, Indika printers, Colombo, Sri Lanka, pp.18-19.
22. Rev. Samyadeva, S., 1988, Sarpa Visha Vinodaya, Gunasena Printers, Colombo, Sri Lanka, pp.44.
23. Bahekar, S., Kale, R. and Nagpure, S., 2012, A review on medicinal plants used in scorpion bite treatment in India, Mintage journal of Pharmaceutical & Medical Sciences; pp.1-6.
24. Al-Asmari, A., Manthiri, R., A., Abdo, N., Al-Duaiji, F. A. and Khan, H., A., 2016, Saudi medicinal plants for the treatment of scorpion sting envenomation, Saudi Journal of Biological Sciences pp.2-23.
25. Rahman, M.A., Bachar, S.C. and Rahmatullah, D., 2010, Analgesic and antiinflammatory activity of methanolic extract of *Acalypha indica* Linn, Pakistan journal of pharmaceutical sciences 23 (3): 256-258.
26. Vijayabhaskar, K., Venkateshwarlu, G., Bhaskar, J., Sravanprasad, M., Sathis Kumar, D., Suresh Kumar, P. and Somsuhara, G., 2011, Evaluation of effect of *Acalypha indica* Linn. leaves extract on bronchodilation and bronchial hyperreactivity in experimental animals, Journal of Pharmacy Research 4(7): 2250.
27. Gagan. S., Simran, S.P., Bharpur, J.A.S. and Mann, A.S. 2011, Antipyretic activity of *Acalypha indica*, Journal of Pharmacy 4(7): 2333.
28. Murugan, P.S., Ramprasath, T. and Selvam, G.S., 2014, Cardioprotective role of *Acalypha indica* extract on Isoproterenol induced myocardial infarction in rats, Journal of Pharmacy Research 4(7): 2129-2132.
29. Sanseera, D., Niwatananun, W., Liawruangrath, B., Liawruangrath, S. and Baramee, A., 2012, Antioxidant and anticancer activities from aerial parts of *Acalypha indica* Linn, Chiang Mai University Journal of Natural Sciences 11(2): 157-168.
30. Purwaningsih, E.H., Ibrahim, N. and Zain, H., 2010, The nerve protection and in vivo therapeutic effect

- of *Acalypha indica* extract in frogs, *Medical Journal of Indonesia* 19(2): 96-102.
31. Kumar, S.V., Sankar, P. and Varatharajan, R., 2009, Anti-inflammatory activity of roots of *Achyranthes aspera*, *Pharmaceutical Biology* 47(10): 973-975.
  32. Dey, A., 2011, *Achyranthes aspera*: Phytochemical and pharmacological aspects- Review article, *International Journal of Pharmaceutical Sciences Review and Research* 9(2): 72-82.
  33. Eliyas, S., 2017, Evaluation of cardioprotective effect of *Achyranthes aspera* on isoproterenol induced myocardial infarction in rats, *International Journal of Pharma and Bio Sciences* 8(4): 124-130.
  34. Goli, V., Macharl, S.P. A Gowrishankar, N.L., 2011, Anti-pyretic activity of *Achyranthes aspera* Linn, *An International Journal of Advances In Pharmaceutical Sciences* 2(2-3): 204-206.
  35. Abi Beulah, G., Sadiq A. M. and Santhi, J., 2011, Antioxidant and antibacterial activity of *Achyranthes aspera*: An invitro study, *Annals of Biological Research* 2 (5) :662-670.
  36. Jayaraman, R., Anitha, T. and Vishal D. Joshi, V.D., 2010, Analgesic and anticonvulsant effects of *Acorus calamus* roots in mice, *International Journal of PharmTech Research* 2(1):552-555.
  37. Shi1, Wang, B., Wu1,Q., Wang, T., Wang, C., Sun1, X., Zong, W., Yan, M., Zhao, Q., Chen, Y. and Zhang, W., 2014, Evaluation of the wound-healing activity and anti-inflammatory activity of aqueous extracts from *Acorus calamus* L., *Pakistan journal of pharmaceutical sciences* 27(1): 91-95.
  38. Daniel, J. A., Ragavee, A., Sabina, E. P. and Devi, S.A., 2014, Evaluation of analgesic, antipyretic and ulcerogenic activities of *Acorus calamus* rhizome extract in Swiss albino mice, *Research Journal of Pharmaceutical, Biological and Chemical Sciences* 5(6): 503-507.
  39. Prabha, S., Kumar, J. and Bapat, U.C., 2016, Evaluation of antioxidant activity of *Acorus calamus* Linn. and *Vitex negundo* L., *European Journal of Pharmaceutical and Medical Research* 3(12): 583-587.
  40. Gilani, A.H., Shah, A.J., Ahamad, M. and Shaheen, F., 2006, Antispasmodic effect of *Acorus calamus* Linn. In mediated through Calcium channel blockade, *Phytotherapy Research* 20: 1080 -1084.
  41. Shukla, P. K., Khanna, V. K. and Ali, M. M., 2006, Neuroprotective effect of *Acorus calamus* against middle cerebral artery occlusion-induced ischaemia in rat, *Human & Experimental Toxicology* 25(4):187-94.
  42. Kumar, M.S. and Hiremath, W.S.M.A., 2016, Cardioprotective effect of *Acorus calamus* against doxorubicin-induced myocardial toxicity in albino Wistar rats, *Indian Journal of Health Sciences* 9 (2): 225 - 234.
  43. Vijayapandia, p., Annabathinab, V., Srikanth, B. S.N., Manjunathb, V., Boggavarapub, P., Mohammed, P. A.K., Prasadb, K.R. and Kumarappanc, C. T., 2013, In vitro anticholinergic and antihistaminic activities of *Acorus calamus* Linn. leaves extracts, *African Journal of Traditional, Complementary and Alternative Medicines* 10(1):95-101.
  44. Acharya, S., Ullal, S., Padiyar, S. and Raj, V., 2011, Analgesic effect of extracts of *Alpinia galanga* rhizome in mice, *Journal of Chinese Integrative Medicine* 9(1): 100 - 104.
  45. Ghosh, A.K., Banerjee, M. and Bhattacharyya, 2011, Anti-inflammatory activity of root of *Alpinia galanga* willd, *Chronicles of Young Scientists* 2(3): 139-142.
  46. Singh J.C.H., Alagarsamy, V., Diwan, P.V., Kumar S.S., Nisha J. C. and Reddy, Y.N., 2011, Neuroprotective effect of *Alpinia galanga* (L.) fractions on  $\text{A}\beta$ (25-35) induced amnesia in mice, *Journal of Ethnopharmacology* 138(1):85-91.
  47. Kumar, B.S. A., Lakshman, K., Jayaveera, K.N., Shekar, D. S., and Vivek, C., 2010, Antinociceptive and antipyretic activities of *Amaranthus viridis* Linn. in different experimental models, *Archives of Biological Sciences* 62 (2): 397-402.
  48. Saravanan, G., Ponmurugan, P., Sathiyavathi, M., Vadivukkarasi, S. and Sengottuvelu, S., 2013, Cardioprotective activity of *Amaranthus viridis* Linn: effect on serum marker enzymes, cardiac troponin and antioxidant system in experimental myocardial infarcted rats, *International Journal of Cardiology* 165(3):494-498.
  49. Salvamani, S., Gunasekaran, B., Shukor, M.Y., Shaharuddin, N. A., Sabullah, M. K. and Ahmad, S. A., 2016, Anti-HMG-CoA reductase, antioxidant, and anti-inflammatory activities of *Amaranthus viridis* leaf extract as a potential treatment for hypercholesterolemia, *Evidence-Based Complementary and Alternative Medicine* 1-10.
  50. Jadhav, V. and Birada, S. D., 2016, Evaluation of antioxidant activity of *Amaranthus viridis* L. methanolic extract, *International Journal of Pharmacy and Biological Sciences* 6(3): 150-153.
  51. Suebsasana, S., Pongnaratorn, P., Sattayasai, J., Arkaravichien, T., Tiamkao, S. and Aromdee, C., 2009, Analgesic, antipyretic, anti-inflammatory and toxic effects of *Andrographolide* derivatives in experimental animals, *Archives of Pharmacal Research* 32(9): 1191-1200.
  52. Karpakavalli, M., Perumal, G. N., Christi, V. E.I., Dhanamani, M. and Rao, U. S., 2010, Anti-histamine effect of hydroalcoholic extract of *Andrographis paniculata* leaf (Burm. F), *Der Pharmacia Sinica* 1 (2): 17-23.
  53. Sah, D. K., and Nagarathana, P.K. M., 2016, Screening of cardioprotective activity of leaves of *Andrographis paniculata* against isoproterenol



- induced myocardial infarction in rats, International Journal of Pharmacological Research 6(1): 23 -28.
54. Kale, R. S., Bahekar, S. E., Nagpure, S. R. and Salwe, K., J., 2013, Anti-scorpion venom activity of *Andrographis paniculata*: A combined and comparative study with anti-scorpion serum in mice, Ancient Science of Life ; 32(3): 156-160.
55. Fang, S., Hsu, C. and Yen, G., 2008, Anti-inflammatory Effects of Phenolic Compounds Isolated from the Fruits of *Artocarpus heterophyllus*, Journal of Agricultural and Food Chemistry, 56 (12): 4463-4468.
56. Jahan, F.I., Hossain, M. S., Al Mamun, A., Hossain, T., Seraj, S. and Chowdhury, A.R., 2010, An evaluation of antinociceptive effect of methanol extracts of *Desmodium gangeticum* (L.) DC. stems and *Benincasa hispida* (Thunb.) Cogn. leaves on acetic acid-induced gastric pain in mice, Advances in Natural and Applied Sciences 4(3): 365-370.
57. Liu, Y., Shen, S., Xia, F., Chang, Q., Liu, X. and Pan, R., 2015, Neuroprotection of Stilbenes from Leaves of *Cajanus cajan* against Oxidative Damage Induced by Corticosterone and Glutamate in Differentiated PC12 Cells, Chinese Herbal Medicines 7(3):238-246.
58. Pathak, A. K. and Argal, A., 2007, Analgesic activity of *Calotropis gigantea* flower, Fitoterapia 78(1): 40-42.
59. Dewan, S., Sangraula, H. and Kumar, V. L., 2000, Preliminary studies on the analgesic activity of latex of *Calotropis procera*, Journal of Ethnopharmacology 7(1-2): 307-311.
60. Kumar, V. L. and Basu, N., 1994, Anti-inflammatory activity of the latex of *Calotropis procera*, Journal of Ethnopharmacology 44(2):123-125.
61. Ahmed, K.K.M., Rana, A. C. and V.K.Dixit, V. K., 2004, Effect of *Calotropis procera* latex on isoproterenol induced myocardial infarction in albino rats, Phytomedicine 11(4):327-330.
62. Singh, N., Jain, N. K., Kannoja, P., Garud, N., Pathak, A. K., and Mehta, S. C., 2010, In vitro antioxidant activity of *Calotropis gigantea* hydroalcoholic leaves extract, Der Pharmacia Lettre, 2010, 2(3): 95-100.
63. Amit, J., Namrata, S., Pathak A.K., Tailang M., 2010, Phytochemistry and evaluation of antioxidant activity of whole plant of *Calotropis gigantea* Linn, International Journal of Research in Ayurveda and Pharmacy 1 (1): 120-125.
64. Chitme, H. R., Chandra, R. and Kaushik, S., 2005, Evaluation of antipyretic activity of *Calotropis gigantea* (Asclepiadaceae) in experimental animals, Phytotherapy Research 19(5): 454-456.
65. Verma, S., Varma, R. K. and Singh, S., 2017, Nootropic activity of *Calotropis gigantea* (Linn) root against scopolamine induce amnesia in albino rats, International journal of Indigenous Herbs and Drugs 2(4): 13-17.
66. Hasimuna, P., Suwendara, Ernasaria, G.I., 2014, Analgetic Activity of Papaya (*Carica papaya* L.) leaves extract, Procedia Chemistry 13: 147 - 149.
67. Owoyele, B. V., Adebukola, O. M., Funmilayo, A. A. and Soladoye, A. O., 2008, Anti-inflammatory activities of ethanolic extract of *Caricapapaya* leaves Inflammopharmacology 16: 168-173
68. Swati, S., Chitra, G., John P.P., Faizal, A. and Bhawana, N., 2011, Antipyretic activity of leaf extract of *Carica papaya* Linn., Invention Rapid - Ethnopharmacology <http://invention.in/journal/article/rapid/3/9506/ethnopharmacology/pi>
69. Fatokun, O.T, Wojuola, T. E., Esievo, K. B. and Kunle, O.F., 2016, Medicinal plants used in the management of asthma: A review, European journal of Pharmaceutical and Medical Research 3(7): 82-92.
70. Kapoor, I.P., Singh, B., Singh, G., De Heluani, C.S., De Lampasona, M. P., Catalan, C. A., 2010, Chemistry and antioxidant activity of essential oil and oleoresins of black caraway (*Carumbulbocastanum*) fruits: Part 69, Journal of the Science of Food and Agriculture 90(3): 385-90.
71. Maity, T. K., Mandal, S. C., Mukherjee, p. k. and Saha, B. P., 1998, Studies on antiinflammatory effect of *Cassia tora* leaf extract (Fam. Leguminosae), Phytotherapy Research 12(3):221 - 223.
72. Ravi, S. K., Narasingappa, R. B., Joshi, C. G., Girish, T. K. and Vincent, B., 2017, Neuroprotective effects of *Cassia tora* against paraquat-induced neurodegeneration: relevance for Parkinson's disease, Natural Product Research 16:1-5.
73. Negi, S. A., Juyal, V. and Melkani, A. B., 2010, Analgesic activity of fruit decoction of *Citrus medica* Linn, Journal of Pharmacy Research.3(9) 2119-2121
74. Munwar, S., Roy, H. and Rahaman, S. A., 2015, Antioxidant and free radical scavenging activity of *Citrus medica*, International Journal of Pharma Research and Health Sciences 3 (4): 810-816.
75. Al-Yahya, M. A., Mothana, R. A., Al-Said, M. S., El-Tahir, K. E., Al-Sohaibani, M. and Rafatullah, S., 2013, *Citrus medica* "Otroj": Attenuates Oxidative Stress and Cardiac Dysrhythmia in Isoproterenol-Induced Cardiomyopathy in Rats, Nutrients 5(11): 4269-4283.
76. Parimaladevi, B., Boominathan, R. and Mandal, S.C., 2003, Studies on analgesic activity of *Cleome viscosa* in mice, Fitoterapia 74 (3): 262-266.
77. Sowunmi, L. I. and Afolayan, A. I., 2015, Phytochemical constituents and antioxidant properties of acetone extract of *Cleome gynandra* (L.) growing in the Eastern Cape, South Africa, African Journal of Traditional, Complementary and Alternative Medicines 12(3):1-8.
78. Devi, B. P., Boominathan, R. and Mandal, S. C., 2003, Anti-inflammatory, analgesic and antipyretic

- properties of *Clitoria ternatea* root, *Fitoterapia* 74(4):345-349.
79. Al-Snafi, A.E., 2016, Pharmacological importance of *Clitoria ternatea* – A review, *IOSR Journal of Pharmacy* 6(3): 68-83.
  80. Kekuda, T. R. P., Sudharshan, S. J., Chinmaya, A., Valleesha, N. C., Murthuza, S. and Achur, N. R., 2009, Central nervous system (CNS) depressant and Analgesic activity of methanolic extracts of *Nardostachys jatamansi* DC. and *Coscinium fenestratum* Colebr. in experimental animal model, *Journal of Pharmacy Research* 2(11): 1716-1719.
  81. Sudharshan, S. J., Kekuda, T. R. P. and Sujatha, M., 2010, Antiinflammatory activity of *Curcuma aromatica* Salisb and *Coscinium fenestratum* Colebr: A comparative study, *Journal of Pharmacy Research* 3(1) : 24-25.
  82. Goveas, S.W. and Abraham, A., 2013, Evaluation of antimicrobial and antioxidant activity of stem and leaf extracts of *Coscinium fenestratum*. *Asian Journal of Pharmaceutical and Clinical Research*, 6(3):218-221.
  83. Ibrionke, G. F. and Owemidu, I. O., 2014, Analgesic activity of the ethanol extract of *Curcumalonga* rhizome and its mechanism of action, *Nigerian journal of physiological sciences* 29(1):67-70.
  84. Chainani-Wu, N., 2003, Safety and anti-inflammatory activity of curcumin: a component of tumeric (*Curcuma longa*), *Journal of Alternative and Complementary Medicine* 9(1):161-168.
  85. Itthipanichpong, C., Ruangrunsi, N., Kemsri, W. and Sawasdipanich, A., 2003, Antispasmodic effects of curcuminoids on isolated guinea-pig ileum and rat uterus, *Journal of the Medical Association of Thailand* 2: S299-309.
  86. Tanvir, E.M., Hossen, S., Hossain, F., Afroz, R., Gan, S. H., Khalil, I. and Karim, N., 2017, Antioxidant Properties of Popular Turmeric (*Curcuma longa*) Varieties from Bangladesh, *Hindawi Journal of Food Quality* 1- 8.
  87. El-Sayed, E. M., Abd El-azeem, S. A., Afify, A. A., Shabana, M. H., and Ahmed, H. H., 2011, Cardioprotective effects of *Curcumalonga* L. extracts against doxorubicin-induced cardiotoxicity in rats, *Journal of Medicinal Plants Research* 5(17) : 4049-4058.
  88. Rajakrishnan, V., Viswanathan, P., Rajasekharan, K. N. and Menon, V. P., 1999, Neuroprotective role of curcumin from *Curcumalonga* on ethanol-induced brain damage, *Phytotherapy Research* 13(7):571-4.
  89. Jain, P. K. and Das, D., 2016, Ethno pharmacological study of *Cyperus rotundus* - A herb used by tribal community as a traditional medicine for treating various diseases, *Innovare Journal of Ayurvedic Science* 4(2): 5-8.
  90. Shamkuwar, P. B., Hoshamani, A. H. and Gonjari, I. D., 2012, Antispasmodic effect of *Cyperus rotundus* L. (Cyperaceae) in diarrhoea, *Der Pharmacia Lettre* 4 (2):522-524.
  91. Jahan, N., Khalil-ur-Rahman and Ali, S., 2012, Cardioprotective and antilipidemic potential of *Cyperus rotundus* in chemically induced cardiotoxicity, *International Journal of Agriculture and Biology* 14 (6):989-992.
  92. Kumar K. H., Tamatam, A., Pal, A. and Khanum, F., 2013, Neuroprotective effects of *Cyperus rotundus* on SIN-1 induced nitric oxide generation and protein nitration: ameliorative effect against apoptosis mediated neuronal cell damage, *Neurotoxicology* 34:150-9.
  93. Essien, G. E, Nwidi, L. L and Nwafor, P. A., 2009, Anti-Inflammatory and Analgesic Potential of Methanolic Extract of *Emilia Sonchifolia* (Compositae) Leaves in Rodents, *African Journal of biomedical Research* 12(3): 199-207.
  94. Sophia, D., Gomathy, M., Shebin, T., Ragavendran, P., Arulraj, C. and Gopalakrishnan, V. K., 2011, Effect of *Emilia sonchifolia* (Linn.)DC on alcohol- induced oxidative stress in pancreas of male albino rats, *Asian Pacific Journal of Tropical Medicine* 973-977.
  95. Murugalakshmi, M., Mari Selvi, J. and Thangapandian, V., 2014, Analgesic and Anti-Inflammatory Activities of *Erythrina variegata* Leaves Extracts, *Journal of Advanced Botany and Zoology* 2(2):1-5.
  96. Bagheri, S. M., Dashti-R, M. H. and Morshedi, A., 2014, Antinociceptive effect of *Ferula assa-foetida* oleo-gum-resin in mice, *Research in Pharmaceutical Sciences* 9(3): 207-212.
  97. Fatehi, M., Farifteh, F. and Fatehi-Hassanabad, Z., 2004, Antispasmodic and hypotensive effects of *Ferula asafoetida* gum extract, *Journal of Ethnopharmacology* 91 (2-3): 321-324.
  98. Kiasalari, Z., Khalili, M., Roghani, M., Heidari, H., and Azizi, Y., 2013, Antiepileptic and Antioxidant Effect of Hydroalcoholic Extract of *Ferula Assa Foetida* Gum on Pentylentetrazole- induced Kindling in Male Mice, *Basic and Clinical Neuroscience* 4(4): 299-306.
  99. Esmalidehaj, M., Kakoo, M., Rezvani, M. E. and Mosaddeghmehrjardi, M. H., 2014, Pretreatment with *Assafoetida* exerts dose-dependent dual effects on rat hearts, *Pharmacognosy Magazine* 10(38): 147-153.
  100. Gopal, T. K., Sheela, T., Chamundeeswari, D. and Reddy, C. U., 2013, Investigation of in-vitro antioxidant, anti-inflammatory and anti-arthritis activity of aerial parts of *Securinegaleucopyrus* (Willd.) Muell, *Indian Journal of Research in Pharmacy and Biotechnology* 1(3): 371-378.
  101. Osuntoki, A. A. and Olagundoye, O. R., 2007, A Mechanism for the Anti-inflammatory Activity of *Gossypium arboreum* Linn Leaves, *Nigerian Journal of Health and Biomedical Sciences* 6 (2): 30-32.

102. Kumar, S., Agrawal, D., Patnaik, J., and Patnaik, S., 2012, Analgesic effect of Neem (*Azadirachta indica*) seed oil on albino rats, *International Journal of Pharma and Bio Sciences* 3(2): 222-225.
103. Pillai, N. R. and Santhakumari, G., 1981, Anti-arthritis and anti-inflammatory actions of Nimbidin, *Journal of Medicinal Plant Research* 43(9): 59-63.
104. Vikram, P. K., Malvi, R. and Jain, D. K., 2012, Evaluation of analgesic and anti-inflammatory potential of *Mimosapudica* Linn, *International Journal of Current Pharmaceutical Research* 4(4): 47-50.
105. Sunil, M., John, P., Brahmane, R., Jayasree, T. and Naveen, A., 2015, Evaluation of anti-scorpion-venom property of ethanolic extract of *Mimosa pudica* in albino mice, *International Journal for Pharmaceutical Research Scholars* 4(1-2): 372-377.
106. Gunjal, Shah, M. A., Wakade, A. S., Juvekar, A. S. and Archana, R., 2010, Protective effect of aqueous extract of *Moringa oleifera* Lam. stem bark on serum lipids, marker enzymes and heart antioxidants parameters in isoproterenol-induced cardiotoxicity in Wistar rats, *Indian Journal of Natural Products and Resources* 1(4): 485-492.
107. Thakur, K. and Pitre, K.S., 2009, Anti-inflammatory activity of extracted eugenol from *Ocimum sanctum* L. leaves, *Rasayana Journal of Chemistry*.2(2): 472-474.
108. Shetty, S., Udupa, S. and Udupa, L., 2007, Evaluation of antioxidant and wound healing effects of alcoholic and aqueous extract of *Ocimum sanctum* Linn in rats, *Evidence-Based Complementary and Alternative Medicine* 5(1): 95-101.
109. Sharma, M., Kishore, K., Gupta, S.K., Joshi, S. and Arya, D.S., 2001, Cardioprotective potential of *Ocimum sanctum* in isoproterenol induced myocardial infarction in rats, *Molecular and cellular biochemistry*, 225(1-):75-83.
110. Kumar, A., Panghal, S., Mallapur, S.S., Kumar, M., Ram, V., and Singh, B. K., 2009, Antiinflammatory activity of *Piper longum* fruit oil, *Indian Journal of Pharmaceutical Sciences* 71(4): 454-456.
111. Wakade, A.S., Shah, A. S. Kulakarani, M., P. and Juvekar, A. R., 2008, Protective effect of *Piper longum* on oxidative stress induced injury and cellular abnormality on adriamycin induced cardiotoxicity in rats, *Indian journal of Experimental Biology* 46: 522-533.
112. Rai, V., Pai, V. R., Kedaliya, P., 2016, A preliminary evaluation of anticancer and antioxidant potential of two traditional medicinal plants from *Lamiaceae*-*Pogostemon heyneanus* and *Plectranthus amboinicus*, *Journal of Applied Pharmaceutical Science* Vol. 6 (08), 073 - 078.
113. Srinivasan, K., Muruganandan, S., Lal, J., Chandra, S., Tandan, S.K, Raviprakash, V, and Kumar, D., 2003, Antinociceptive and antipyretic activities of *Pongamia pinnata* leaves, *Phytotherapy Research*17(3):259-64.
114. Srinivasan, K., Muruganandan, S., Lal, J., Chandra, S., Tandan, S.K and Raviprakash, V, 2001, Evaluation of anti-inflammatory activity of *Pongamia pinnata* leaves in rats, *Journal of Ethnopharmacology* 78(2-3): 151-157.
115. Sangwan, S., Rao, D.V. and Sharma, R.A., 2010, A Review on *Pongamiapinnata* (L.) Pierre: A Great Versatile Leguminous Plant, *Nature and Science* 8(11): 130 -139.
116. Santha, M. L., Kanchana, P. and Supriya, C. H., 2015, Anti Nociceptive and Anti Inflammatory Activity of *Scheilcheroaleosa* (lour) Oken Bark, *American Journal of Pharmtech Research* 5(3): 187-196.
117. Khan, M. J., Saraf, S. and Saraf, S., 2017, Anti-inflammatory and associated analgesic activities of HPLC standardized alcoholic extract of known Ayurvedic plant *Scheilcheroaleosa*, *Journal of Ethnopharmacology*197: 257-265.
118. Jami, S. I., Sultana, Z., Ali, E., Begum, M. and Haque, M., 2014, Evaluation of analgesic and anti-inflammatory activities on ethanolic extract of *Terminalia chebula* fruits in experimental animal models, *American Journal of Plant Sciences*5: 63-69.
119. Suchalatha, Subramaniyan, D. and Shyamala, C.S., 2005, Antioxidant activity of ethanolic extract of *Terminaliachebula* fruit against isoproterenol-induced oxidative stress in rats, *Indian Journal of Biochemistry and Biophysics* 42 (4): 246-249.
120. Bag, A., Bhattacharyya, S. K. and Chattopadhyay, R.R., 2013, The development of *Terminalia chebula* Retz. (*Combretaceae*) in clinical research, *Asian Pacific Journal of Tropical Biomedicine* 3(3): 244-252.
121. Shen, Y., Juan, C., Lin, C., Chen, C. and Chang, C., 2017, Neuroprotective effect of *Terminalia chebula* extracts and Ellagic acid in PC12 cells, *African Journal of Traditional, Complementary and Alternative Medicine*14 (4): 22-30.
122. Dhara, A. K., Suba, V., Sen, T., Pal, S. and Chaudhuri, A.K.N., 2000, Preliminary studies on the anti-inflammatory and analgesic activity of the methanolic fraction of the root extract of *Tragia involucrata* Linn, *Journal of Ethnopharmacology* 7(1-2): 265-268
123. Yadav, S. A., Ramalingam, S., Raj, A. J. and Subban, R., 2015, Antihistamine from *Tragia involucrata* L. leaves, *Journal of Complementary and Integrative Medicine* 12(3): 217-226.
124. Palani, S., Kumar, S. N., Gokulan, R., Rajalingam, D., and Kumar, B. S., 2009, Evaluation of Nephroprotective and antioxidant potential of *Tragiainvolucrata*, *Drug Invention Today* 1(1): 55-60.
125. Raji, Y., Udoh, U. S., Oluwadara, O.O., Akinsomisoye, O. S., Awobajo, O. and Adeshoga, K., 2002, Anti-



- inflammatory and analgesic properties of the rhizome extract of *Zingiber officinale*, African Journal of Biomedical Research 5: 121 -124.
126. Ahmed, R.S., Seth, V. and Banerjee, B.D., 2000, Influence of dietary ginger (*Zingiber officinales* Rosc) on antioxidant defense system in rat: comparison with ascorbic acid, Indian journal of experimental biology, 38(6): 604-606.
127. Rohini, A., Agrawal, N., Chandrasekar, M. J. N. and Sara, U.V.S., 2013, Evaluation of Cardioprotective Effect of *Zingiber Officinale* in Experimental Animals, International Journal of Current Pharmaceutical Review and Research 4(1): 1-9.
128. Kaushik, R., Jain, J. and Rai, P., 2015, Therapeutic potentials of cow derived products- A review, International Journal of Pharmaceutical Sciences and Research 7(4): 1383-1390.
129. Jarald, E., Edwin, S., Tiwari, V., Garg, R. and Toppo, E., 2008, Antioxidant and Antimicrobial Activities of Cow Urine, Global Journal of Pharmacology 2 (2): 20-22.
130. Dousset, E., Carrega, I., Steinberg, J.G., Clot-Faybesse, O., Jouirou, B., Sauze, N., Devaux, C., Autier, Y., Jammes, Y., Martin-Eauclaire, M.F. and Guieu, R., 2005, Evidence that free radical generation occurs during scorpion envenomation, Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology 140 (2): 221-226.
131. Sah, D. K. and Nagarathana, P.K. M., 2016, Screening of cardioprotective activity of leaves of *Andrographis paniculata* against isoproterenol induced myocardial infarction in rats, International Journal of Pharmacological Research 6 (1): 23-27.
132. Sareer, O., Ahmad, S. and Umar, S., 2014, *Andrographis paniculata*: a critical appraisal of extraction, isolation and quantification of *Andrographolide* and other active constituents, Natural Product Research 28 (23): 2081-2101.

**Cite this article as:**

E.R.H.S.S. Ediriweera, W.D. Ratnasooriya, K. W. M. D. Kamathewatta. Ethnomedicinal Uses in Scorpion Poisoning (Gonusu Visha) in Sri Lankan Traditional Medicine. International Journal of Ayurveda and Pharma Research. 2018;6 (7):29-43.

**Source of support: Nil, Conflict of interest: None Declared**

**\*Address for correspondence**

**Dr E.R.H.S.S. Ediriweera**

Professor,  
Department of Nidana Chikithsa,  
Institute of Indigenous Medicine,  
University of Colombo, Rajagiriya,  
Sri Lanka.

Email:

[ayurvedadocsujatha@yahoo.com](mailto:ayurvedadocsujatha@yahoo.com)

Tel: +94714447648

Disclaimer: IJAPR is solely owned by Mahadev Publications- dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IJAPR cannot accept any responsibility or liability for the articles content which are published. The views expressed in articles by our contributing authors are not necessarily those of IJAPR editor or editorial board members.