



Review Article

THERAPEUTIC APPLICATION OF A SIDDHA FORMULATION PATAIC CANKĀRAN – A REVIEW

Navinkumar M¹, Nivetha G¹, Radha P^{2*}, Ahamed Mohideen M¹

¹Department of Sirappu Maruthuvam, Govt. Siddha Medical College and Hospital, Palayamkottai, Tirunelveli, Tamil Nadu, India.

^{2*}Department of Botany, Siddha Clinical Research Unit (CCRS), Govt. Siddha Medical College and Hospital campus, Palayamkottai, Tirunelveli, Tamil Nadu, India.

ABSTRACT

Padarthamarai is the fungal skin infections and are categorised under *Kutta Rogam* (*Padarthamarai -Pundareega Kutam*). Many siddha drugs are prescribed for these diseases among them, *Pataic Cankāran* is exclusively mentioned for these disease as an internal and external medicine in Siddha literature - *Siddha Vaidya Thirattu*. Two different preparations are mentioned under *Pataic Cankāran* and differences observed in the ingredients as in *Pataic Cankāran I* *Ankol* (*Alangium salvifolium* (L.f.) Wang.), *Marukkarai* (*Catunaregam spinosa* (Thunb.) Tirvengadumia) and *Elumitchai* (*Citrus aurantifolia* (Christm.) Swingle), where as in *Pataic Cankāran II*, *Thagarai* (*Senna tora* (L.) Roxb.), *Erruku* (*Calotropis procera* (Ait.) R.Br.) are used. Even though *Pataic Cankāran* is mentioned in literature exclusively for dermatophytosis which is no longer in uses; Clinical study and standardisation of this drug have not been done so far, but *In-vitro* anti-dermatophytotic activities of the individual ingredient of these formulations have already been proved. The present review article aimed to document the biological, pharmacological and therapeutic applications such as anti-microbial, anti-bacterial and anti-inflammatory activities of individual ingredients used in *Pataic Cankāran I* with reference to skin diseases. This study will help to understand the need of research study on this ancient Siddha formulation.

KEYWORDS: *Pataic Cankāran*, Siddha formulation, Skin diseases, Dermatophytes.

INTRODUCTION

Siddha, a traditional system of medicine is being followed by the South Indians especially Tamilians and has drawn a great attention world wide. A number of ancient texts of Siddha perceived treasure to cure various ailments. In Siddha, skin diseases are collectively known as “*Thol Noikal*”. *Padai* and *Padarthamarai* (dermatophytoses) are common skin diseases grouped under “*Kutta Rogam*” and probably Dermatophytosis.^[1,2] *Padai* - differentiated into *Erichalpada*, *Thothupadai*, *Pithapadai*, *Megapadai*, *Themalpadai* etc^[2,3]. *Padarthamarai* (*Pundareega Kutam*) is another type skin disease caused by Ring worm, *Tinea corporis* and other species of *Tinea*. Symptoms of these diseases are patches, black, white, and red colored skin with watery or dried that spread on the body ^[2,3]. Siddha drugs prescribed for dermatophytosis/other skin infections are; *Parangipattai* *Chooranam*, *Chinathamani Kuligai*, *Ilaikalli Pattru*, *Arugantailam*, *Nantimeluku*, *Cirratattailam*, *Pataic Cankāran* and *Vekarappodi* ^[1,2]. Among them, *Pataic Cankāran* is an important drug used for fungal skin infection both *Padai* and *Padarthamarai*. In Siddha literature two

types of preparations named under *Pataic Cankāran*, and prescribed exclusively for *Padai* and *Padarthamarai* (dermatophytosis). Ingredients of *Pataic Cankāran I* (Table 1). In the review article presented siddha perspective and pharmacological important of *Pataic Cankāran I* and its ingredients in brief with reference to the skin diseases.

MATERIALS AND METHODS

Literature survey was done using search engines such as pubmed, pubchem, Science direct, and siddha literature.^[1,4,5]

RESULTS AND DISCUSSION

Preparation of *Pataic Cankāran I* mentioned in Siddha Vaidya Thirattu as follows: Finely powdered *Ankol Verpattai*, *Marukkarai Verpattai* and *Marukkarai Vidhai* were ground with *Elumitchai* fruit juice till a waxy consistency obtains. Recommended dosage: as external application also can use internal administration with the dose of 0.5-1g oral.^[4] Ingredients of *Pataic Cankāran I* are given in Table 1 with botanical name and Tamil name.

Table 1: Details of ingredients used in Pataic Cankāran I^[1, 4]

Drugs used in Pataic Cankāran	Botanical name of the drug
Ankol verpattai (Rt.Bk.) Marukkarai verpattai (Rt. Bk.) Marukkarai vidhai (Sd.) Elumitchai (Fr. Juice)	<i>Alangium salvifolium</i> (L.f.) Wang. * <i>Catunaregam spinosa</i> (Thunb.) Tirvengadumia <i>Citrus aurantifolia</i> (Christm.) Swingle

**Randia dumetorum*; Rt. Bk. – Root Bark; Sd – Seed; Fr.-Fruit

Botanical description of the plants used in Pataic Cankāran I:

***Alangium salvifolium* (L.f.) Wang. (Family: Alangiaceae):** *Alangium salvifolium* (L.f.) Wang. subsp. *salvifolium* and *A. salvifolium* (L.f.) Wang. subsp. *hexapetalum* (Lam.) Wang. belongs to the family Alangiaceae, it is commonly known as *Alingi*, *Alingil* or *Ankolam* in Tamil and it is distributed in all districts of Tamil Nadu.

Small deciduous trees; bark greyish brown; branchlets tomentose; stems with leaf scars outside. Leaves simple, alternate, oblong-lanceolate, base oblique, apex acute, margin entire, glabrous above, puberulous beneath, 3-5 nerved at base, lateral nerves 4-7 pairs, 5-14×2-2.5cm. flowers 1.5cm across, white, fragrant, in axillary cymes; bracts ovate, 4 mm. Calyx 3mm, cupular, tomentose; lobes 10, triangular-ovate. Corolla white, lobes 10, 2.5×5cm, linear-oblong, tomentose. Stamens ca. 20; filaments 1 cm long, base villous, subconnate, anthers linear. Ovary turbinate, 1-celled, 1-ovuled, pendulous; style 2 mm long, glabrous; stigma capitate. Berry globose, ca. 2-3×1-1.5cm, red when ripen, crowned with calyx lobes. Seed 1, ovoid.

The medicinal properties of Alingil relieve *Vatha* and *Kapha* diseases, leprosy with pus and all types of poison bites.^[5] *Azhingi* or *Ankolam* is an important drug and all parts of these plants are used in Siddha and Ayurveda in the treatment of various ailments^[5] especially for skin diseases. The bitter tasted root bark of *Alangium* is an astringent, purgative, emetic, febrifuge in nature which has been used in the treatment of rheumatism, leprosy, piles, deworming, hypertension, diarrhoea, fever, back pain etc. Recently, anti-cancer activities of the *Alangium* root bark have also been proved.^[6-10] Other Siddha formulation in which *alangium* root is used are *Naalpattapunnukupugai*, *Ayapirunka Raja Karpam* and *Purayodum Viranathailam* which are prescribed medicines for skin diseases.^[1]

The active chemical constituents reported from the different parts of *Alangium* are: alangimarckine, deoxutubulosine, alangiside, β-sistosterols from leaves; iso-alangiside, demethylneoalngiside etc from fruit. Alkaloids reported from seeds are alangimarine, emetine, cephaeline and other phytoconstituents includes betulinic acid, lupeol, alangol, tannins. Alagine A,

Alangine B, ankoline, myricyl, Benzoquinolizidine, isalamarine, stigmasterol and alangicine, d-methylpsychotrine, marckine are reported from the stem and root bark.^[6-11]

Alangium is being used as an ethno medicine for various diseases such as anti-diabetic, rheumatoid arthritis; skin diseases like leprosy, scabies; fever, anti-tuberculosis etc. Number of review studies have been attempted on Pharmacological and phytochemicals to emphasis the anti-bacterial, anti-fungal and anti-microbial activities of this plant^[11-14]

The anti-bacterial, anti-fungal and anti-microbial activities are reported against the micro-floras such as *Aspergillus niger*, *A. flavus*, *Escherichia coli*, *Staphylococcus aureas*, *Pseudomonas aeruginosa*, *Klebsella pneumonia*, *Solmonella typhyrrium*, *Candida albicans*, *Shiegella spp.*, etc. Beyond the anti-microbial activity the root and root barks shows significant anti-oxidant, anti-inflammatory and anti-helminthic activities.^[15-17]

***Catunaregam spinosa* (Thunb.) Tirveng. (Rubiaceae) - Marukkarai**

Syn.: *Randia dumetorum* (Retz.) Lam.

The therapeutic character (*Gunam*) of the root is mentioned in *Pathartha Guna Cintamani* as follows; Root cures pain during dysentery, *Rakhtha Pitam*, eczema, diseases and *Maantha Ganam*. Fruits are emetic, diaphoretic and antispasmodic.^[5]

Large shrub to small armed trees; spines to 3.5cm, paired; branchlets pubescent. Leaves opposite, obovate, 4×2.5cm, puberulous above, hirsute on nerves below; petiole 0.8 cm long; stipule ovate, cuspidate. Flowers solitary, 2-5, terminal on lateral branches. Calyx tube 5mm long; lobes 5, obovate, 5 × 3.5mm. Corolla white turning yellow, 2 cm across; tube 5 mm, densely villous at the base inside; lobes 5, obovate, twisted, 9 × 7 mm, apex acute. Stamens 5, anthers sessile at mouth of corolla. Ovary 2-6 celled; ovules many; style stout, 12 mm long; stigma fusiform, ribbed. Berry ovoid, 4 × 3 mm, glabrous, with a crown of enlarged calyx lobes. Seeds many, embedded in pulp.

Mannitol, saponins, coumarin glycosides Scolpoletin, Saponins (Randialic and Randialic acid) Urosolic acid, Randianin from roots and Oleanolic acid-3-glucose from seeds and from ripen fruit, Randioside A. Randianin, Dumetoronin A,B,C,D,E Ripe

fruit contains glycosides, randioside A. are important phyto-chemical constituents reported so far. [18-19]

In Ayurveda, this emetic nut known as *Mainphal* and it shows various biological activities such as diaphoretic, anthelmintic, expectorant, anti-spasmodic and cure cold, cough, rhinitis, rheumatism, diarrhoea and dysentery. Seeds are appetizer and shows significant results in the treatment of skin diseases, gastrointestinal tract diseases and wounds.¹⁹⁻²¹ Number of studies were conducted on pharmacological and phytochemical characteristics of stem bark, root bark and fruits and it proves that *Marukkarai* shows anti-bacterial, anti-allergic, anti-inflammatory, analgesic and immune modulatory activities. [21-26]

The extract of *Randia* shows Anti-microbial activity against *Staphylococcus aureus*, *Streptococcus mutans*, *Aspergillus niger*, *Candida albicans*, *Escherichia coli* and oral pathogenic organisms such as *Streptococcus sobrinus* and *Lactobacillus acidophilus*. [22-26]

***Citrus aurantifolia* (Christm.) Swingle (Rutaceae) - Elumitchai**

Citrus aurantifolia (Christm.) Swingle and *Citrus limon* (L.) Osbeck and their cultivars are commonly available as *Elumitchai* in markets. Lemon is being used as a medicine for various ailments. The anti-bacterial and anti-inflammatory activities of lemon juice are well studied by various researchers. In Siddha medicines, lemon juice mostly used as an adjuvant. Lemon juice shows a significant action against the dermatophytes such as *Pseudomonas aeruginosa*, *Staphylococcus aureus* and other bacteria includes *Klebsiella spp.*, *Proteus spp.* and *Escherichia coli*. Flavonoids such as naringin, hesperidin and neohesperidin are showed antifungal activities in four fungi species like *Aspergillus parasiticus*, *A. flavus*, *Fusarium semitectum* and *Penicillium expansum*. The Hesperidin glucoside lurate and pruning decanoate is the most inhibiting flavonoid for *P. expansum* and *A. flavus* respectively [27]. Anti-fungal activities are reported against the different micro-flora such as *Kluyveromyces fragilis*, *Rhodotorula rubra*, *Candida albicans*, *Hanseniaspora guilliermondii*, *Acne vulgaris*, *Rhizoctonia solanii* and *Debaryomyces hansenii* (yeast) and the results were referenced against the anti-fungal agents such as nystatin, ketaconazole and clotrimazole from Citrus peel oil. [28-31]

Anti-microbial activities from juice of *Citrus limon*, *C. reticulata* and *C. grandis* against pathogenic bacteria such as *Staphylococcus aureus*, *E. coli*, *Bacillus subtilis*, *Vibrio spp.* and *Salmonella typhi* from the phytochemical components of linalyl acetate, linolool, lomonene, pinene, β -pinene, 1,8-cineole, carvacrol

and thymol.^[32-35] Citrus fruit peel oil is used to suppress the growth of the *Acinetobacter*^[36] also anti-fungal properties of *Citrus limon* is used to investigate the safety and efficacy in the treatment of oral thrush in HIV/AIDS patients.^[37] Hence, it is evident that, *Citrus limon* having an anti-bacterial, anti-fungal and anti-microbial activities and play a significant role in the Siddha System of medicine.

CONCLUSION

Ingredients of *Pataic Cankāran I & II*, are different. In *Pataic Cankāran II*, Root bark of velleruku – (*Calotropis procera* (Ait.) R.Br.) and seeds of thagarai – (*Senna tora* (L.) Roxb.) are the important ingredient and recommended medicine for skin diseases. The medicinal applications/biological role of these plants are reported in number of folk/traditional medicine studies and proved their pharmacological and phytochemical activities with reference to the treatment of skin diseases [14,38-44]. Most of the Senna species are used by the tribal and inhabitants for the various dermatological diseases. The other medicinal plants used in the treatment of Padai and padarthamarai traditionally are *Kovai-Coccinia grandis* (L.) Voigt, *Sittirapaladai- Euphorbia thymifolia* L., *Vembu-Azadirachta indica* A.Juss., *poovarasu-Thespesia populnea* (L.) Soland ex.Correa; *Tumbai- Leucas aspera* (Willd.) Link, *Karisalai-Eclipta prostrata* (L.) L., *Kuppeimeni- Acalypha indica* L, *Mookiratai -Boerhavia diffusa* L., *Karuvelam-Acacia nilotica* Willd. ex Del. subsp. *indica* (Benth) Brenan, *Brahma thandu-Argemone mexicana* L., etc., and their *in-vitro* anti-dermatophytic activities have been proved with significant results.^[45-47] While reviewing the case studies of skin diseases, dermatophytosis are treated with siddha drugs such as *Kandhaga Rasayanam*, *Parankipattai Chooranam* etc.^[48,49] Even though, individual ingredient of *Pataic Cankāran I* shows therapeutic efficacy towards skin diseases, it is no longer in use and Clinical studies and standardisation of these drugs has also not been attempted. Thus, we can understand that there is a need for the evaluation and standardisation of these formulations which will be highly useful for the mankind in future.

REFERENCES

1. Siddha Formulary of India-SFI. The Siddha Formulary of India. Part 1. First Edition. Govt. of India. Ministry of Health and Family Welfare, Delhi. 1992. pp 167.
2. Thas JJ. Siddha Medicine-background and principles and the application for skin diseases. Clinics in Dermatology. 2008; 26: 62-78.
3. Sudharsana S, Mohanapriya A. Documentation of traditional Siddha medicines for skin diseases from Katpadi taluk, Vellore District, Tamil Nadu, India.

- European Journal of Integrative Medicine, Volume 9, January 2017, Pages 52-62.
4. Kuppusamy Mudaliar KN, Uthamarayan KS. Siddha Vaithiya Thirattu. 3 rd edition. Indian Medicine and Homeopathy. 2009; pp. 310.
 5. Anandha AR, Pathartha Guna Cintamani. Department of Indian medicine and Homeopathy, Chennai. 2009; pp, 66,140,222.
 6. Venkateshwarlu R, Raju AB, Yarragunta VG. Phytochemistry and pharmacology of *Alangium salvifolium*: A review. Journal of Pharmacy Research. 2011; 4(5):1423-1425.
 7. Kaur J, Singh R, Kaur J, Kaur H. *Alangium salvifolium* (Linn. F) Wang: A Phytopharmaceutical Review. International Journal of Current Pharmaceutical Review and Research. 2017; 8(01), 45-48.
 8. Shrivya S, Vinod BN, Sunil C. Pharmacological and phytochemical studies of *Alangium salvifolium* Wang-A review. Bulletin of Faculty of Pharmacy, Cairo University. 2017, 55(2): 217-222.
 9. Tanwer SB, Vijayvergia R. Biological evaluation of *Alangium salviifolium* (L. F.)Wangerin. Journal of Chemical and Pharmaceutical Research. 2014; 6(12): 611-618.
 10. Panara K, Singh PK, Rawat P, Vivek Kumar, Marur M, Patel K, Ravikumar RK, Vipin Kumar. Importance of *Alangium salvifolium* and its Pharmacological Update. European Journal of Medicinal Plants. 2016; 12(4): 1-15.
 11. Jha AK, Yogesh G, Professor A. Some Wild Trees of Bihar and Their Ethnobotanical Study. IOSR Journal of Research & Method in Education Ver. II. 2015; 5(6):2320-7388.
 12. Madhu V, Yarra R. Investigation on ethnomedicinal plants used to cure skin diseases in Adilabad, Andhra Pradesh, India. International Journal of Pharmacy and Life Sciences. 2011; 2(5): 742-745.
 13. Kumar G, Sharmila Banu G, Rajarajan T, Sathishkumar G. Medicinal Flora of Palayapalayam, Namakkal District, Tamil Nadu. Indian Journal of Applied and Pure Biology. 2011; 26(1): 135-158.
 14. Pathak K, Das R. An Overview on Medicinal Herbs in the Treatment of Skin Diseases. World Journal of Pharmacy and Pharmaceutical Sciences. 2013; 2(2): 437-449.
 15. Khan MN. The evaluation of the in-vitro antimicrobial activity of *Alangium salvifolium* (Linn.). World News of Natural Sciences. 2018; 20: 176-181.
 16. Wuthi-udomlert M, Prathanturarug S, Wongkrajang Y. Antifungal activity and local toxicity study of *Alangium salviifolium* subsp hexapetalum. The Southeast Asian Journal of Tropical Medicine and Public Health. 2002; 33 Suppl 3: 152-154.
 17. Sharma AK, Agarwal V, Kumar R, Balasubramaniam A, Mishra A, Gupta R. Pharmacological studies on seeds of *Alangium salvifolium* Linn. Acta Poloniae Pharmaceutica - Drug Research. 2011; 68(6):897-904.
 18. Khare CP. Indian Medicinal plants – All illustrated Dictionary. Springer. 2007; pp.535-536.
 19. Ghaisas MM, Ninave P B, Ganu GP, Zope VS, Tanwar MB, Deshpande AD. Effect of *Randia dumetorum* Lam on clonidine and haloperidol-induced catalepsy in mice. Pharmacologyonline. 2008; 2: 42-50.
 20. Singh, NK, Mishra AK, Gupta JK, Jayalakshmi S. *Randia spinosa* (Poir.): Ethnobotany, phytochemistry and pharmacology -A review. International Journal of Pharmaceutical Sciences Review and Research. 2010; 4(1): 127-131.
 21. Jyothirmayi B, Tejaswini L, Latha GA, Swathi C, Nishitha G. Anthelmintic evaluation of *Randia dumetorum* fruit extracts. 2018; 12(6): 880-883.
 22. Ritesh PG, Pathak NL, Rathod JD, Patel LD, Bhatt Nayna M. Phytopharmacological properties of *Randia dumetorum* as a potential medicinal Tree: An overview. Journal of Applied Pharmaceutical Sciences. 2011; 1(10): 24-26.
 23. Kumar PV, Chauhan SN, Padh H, Rajani M. Search for antibacterial and antifungal agents from selected Indian medicinal plants. Journal of Ethnopharmacology, 2006; 107(2): 182-188.
 24. Dharmishtha M, Falguni G. Antibacterial activity of methanolic fruit extract of *Randia dumetorum* Lamk. International Journal of Pharm Tech Research. 2009; 1(3): 679-681.
 25. Tilloo SK, Bodhankar MM, Waikar SB, Pande VB. Antimicrobial activity of formulation containing bioactive extract of endocarp of *Randia spinosa*. International Journal of Advances in Pharmacy Biology and Chemistry. 2012; 1(2): 172-175.
 26. Unissa R, Saikumar M. Antibacterial Activity of Methanolic extract of *Randia dumetorum* Lamk against Oral Pathogens. International Journal of ChemTech Research. 2018; 11(1): 116-120.
 27. Salas MP, Céliz G, Geronazzo H, Daz M, Resnik SL. Antifungal activity of natural and enzymatically-modified flavonoids isolated from citrus species. Food Chemistry. 2011; 124: 1411-1415.
 28. Amadioha AC. Fungicidal activity of some plant extracts against rhizoctonia. Archives of Phytopathology and Plant Protection. 2001; 33(6): 509-517.
 29. Kirbaslar FG, Tavman A, Dulger B, Turker G. Antimicrobial activity of Turkish Citrus Peel Oils. Pak. J. Bot., 2009; 41(6): 3207-3212.
 30. Verma RK, Chaurasia L, Kumar M. Antifungal activity of essential oils against selected building fungi. Indian Journal of Natural Products and Resources. 2011; 2(4): 448-451.

31. Shinkafi SA, Ndanusa H, Antibacterial Activity of Citrus limon on Acne vulgaris (Pimples). International Journal of Science Inventions Today. 2013; 2(5): 397-403.
32. Tomotake H, Koga T, Yamato M, Kassu A, Ota F. Antibacterial Activity of Citrus Fruit Juices against Vibrio Species. Journal of Nutritional Science and Vitaminology. 2006; 52(2):157-160.
33. Soković M, Marin PD, Brkić D, Griensven LJLD Van. Chemical Composition and Antibacterial Activity of Essential Oils of Ten Aromatic Plants against Human Pathogenic Bacteria. Food. 2007; 1: 220-226.
34. Ehigbali IO, Ehimwenma SO, Faith EO Kelly O. Phytochemical, antimicrobial, and antioxidant activities of different citrus juice concentrates. Food Science & Nutrition. 2015; 4(1): 103-109.
35. Saeb S, Amin M, Gooybari SR, Aghel N. Evaluation of Antibacterial Activities of Citrus limon, Citrus reticulata, and Citrus grandis Against Pathogenic Bacteria. International Journal of Enteric Pathogens. 2017;4(4):11-15.
36. Felipe QSG, Mendes JM, Oliveira1 AW, Souza1 SF, Trajano VN, Coutinho HDM, Lima EO. Antibacterial activity of the essential oil of Citrus limon against multidrug resistant Acinetobacter strains. Rev. Bras. Farm., 2013; 94 (2): 142-147.
37. Wright SC, Maree JE, Sibanyoni M. Treatment of oral thrush in HIV/AIDS patients with lemon juice and lemongrass (Cymbopogon citratus) and gentian violet. Phytomedicine. 2009; 16:118-124.
38. Jain S, Patel UK, Phytochemical and pharmacological profile of Cassia tora Linn. - An Overview. Indian Journal of Natural Products and Resources. 2010; 1(4): 430-437.
39. Habib MR, Karim MR. Evaluation of antitumour activity of Calotropis gigantea L. root bark against Ehrlich ascites carcinoma in Swiss albino mice. Asian Pacific Journal of Tropical Medicine. 2011; 4(10), 786-790.
40. Singhal M, Kansara N. Cassia tora Linn Cream Inhibits Ultraviolet-B-Induced Psoriasis in Rats. ISRN Dermatology. 2012; 1-6.
41. Kori P, Alawa P. Antimicrobial activity and phytochemical analysis of Calotropis gigantea root, latex extracts. IOSR Journal of Pharmacy. 2014; 4(6): 07-11.
42. Aliyu RM, Abubakar MB, Kasarawa AB, Dabai YU, Lawal N, Bello MB, Fardami AY. Efficacy and Phytochemical Analysis of Aqueous Extract of Calotropis procera against selected Dermatophytes. Journal of Intercultural Ethnopharmacology. 2015; 4(4): 314-317.
43. Anoma GM, Ajantha Chethana Kumari and Anjana. Alcoholic extraction and phytochemical evaluation of Chakramarda Seeds (Cassia tora L.). International Journal of Research in Ayurveda & Pharmacy. 2017; 8(3): 157-161.
44. Meena R, Ramaswamy RS. Herbs for combatting dematophytosis - a review. IJP. 2014; 1(6):373-79.
45. Vaidyanathan D, Salaisenthilkumar MS, Basha MG, Studies on ethno medicinal plants used by malayaltribals in Kolli hills of Eastern Ghats, Tamil Nadu, India. Asian Journal of Plant Science and Research. 2013; 3(6): 29-45.
46. Xavier FT, Kannan M, Auxilia A. Traditional Medicinal Plants Used in the treatment of different skin diseases. Int.J.Curr.Microbiol. App. Sci. 2015; 4(5): 1043-1053.
47. Gupta DK, Gupta SG, Ghanshyam G, Gupta S. Endemic Use of Medicinal Plants for the Treatment of Skin Diseases in the Balod district. Journal of Pharmacy. 2018; 8(2): 18-24.
48. Meena R, Punithavathy PM, Ramaswamy RS. MIC of a Siddha formulation Kandhaga Rasayanam against Dermatophytosis. Int.J.Phar.Sci. Rev. 2018; 50(2):37-40.
49. Lekha GS, Kanagarajan A. Management of Dermatophytosis or Pardarthamarai in Siddha System- a case study. World Journal of Pharmaceutical Research. 2018; 7(19): 1264-1271.

Cite this article as:

Navinkumar M, Nivetha G, Radha P, Ahamed Mohideen M. Therapeutic Application of A Siddha Formulation Pataic Cankaran - A Review. International Journal of Ayurveda and Pharma Research. 2019;7(2):46-50.

Source of support: Nil, Conflict of interest: None Declared

***Address for correspondence**

Dr.P.Radha

Research Officer - Botany,
Department of Botany, Siddha
Clinical Research Unit (CCRS),
Govt. Siddha Medical College and
Hospital campus, Palaymkottai,
Tirunelveli, Tamil Nadu, India.
Email: randkr@rediffmail.com
Mobile: 9618948682

Disclaimer: IJAPR is solely owned by Mahadev Publications - dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IJAPR cannot accept any responsibility or liability for the articles content which are published. The views expressed in articles by our contributing authors are not necessarily those of IJAPR editor or editorial board members.