

## Limitation of Life-Sustaining Care in the Critically Ill: A Systematic Review of the Literature

Katie McPherson, MD<sup>1</sup>; W Graham Carlos III, MD<sup>2</sup>; Thomas W Emmett, MD, MLS<sup>3</sup>;  
James E Slaven, MS, MA<sup>4</sup>; Alexia M Torke, MD, MS<sup>5,6\*</sup>

<sup>1</sup>Division of Pulmonary and Critical Care Medicine, University of Colorado, Denver Colorado; <sup>2</sup>Division of Pulmonary, Critical Care, Sleep, and Occupational Medicine, Indiana University School of Medicine, Indianapolis, Indiana; <sup>3</sup>Ruth Lilly Medical Library at Indiana University School of Medicine, Indianapolis, Indiana; <sup>4</sup>Department of Biostatistics, Indiana University School of Medicine, Indianapolis, Indiana; <sup>5</sup>Indiana University Center for Aging Research, Indianapolis Indiana; <sup>6</sup>Daniel F. Evans Center for Spiritual and Religious Values in Healthcare and Fairbanks Center for Medical Ethics, Indiana University Health, Indianapolis, Indiana.

When life-sustaining treatments (LST) are no longer effective or consistent with patient preferences, limitations may be set so that LSTs are withdrawn or withheld from the patient. Many studies have examined the frequency of limitations of LST in intensive care unit (ICU) settings in the past 30 years. This systematic review describes variation and patient characteristics associated with limitations of LST in critically ill patients in all types of ICUs in the United States. A comprehensive search of the literature was performed by a medical librarian between December 2014 and April 2017. A total of 1,882 unique titles and abstracts were reviewed, 113 were selected for article review, and 36 studies were fully reviewed. Patient factors associated with an increased likelihood of limiting LST included white race, older age, female sex, poor preadmission functional

status, multiple comorbidities, and worse illness severity score. Based on several large, multicenter studies, there was a trend toward a higher frequency of limitation of LST over time. However, there is large variability between ICUs in the proportion of patients with limitations and on the proportion of deaths preceded by a limitation. Increases in the frequency of limitations of LST over time suggests changing attitudes about aggressive end-of-life-care. Limitations are more common for patients with worse premorbid health and greater ICU illness severity. While some differences in the frequency of limitations of LST may be explained by personal factors such as race, there is unexplained wide variability between units. *Journal of Hospital Medicine* 2019;14:303-310. Published online first February 20, 2019. © 2019 Society of Hospital Medicine

Access to life-sustaining treatment (LST) became a mainstay in hospitals across the United States in the 1970s. This has raised complex ethical questions surrounding the use of these therapies, particularly in the face of a poor prognosis or significant morbidity. The Society for Critical Care Medicine formed a consensus panel in 1989 to construct ethical guidelines regarding the initiation, continuation, and withdrawal of intensive care.<sup>1</sup> These guidelines emphasized that withdrawing and withholding are not only permissible but may be necessary to preserve the balance between quantity and quality of life. Nevertheless, an increasing number of Americans are dying after aggressive LST in the hospital, and greater than one in five deaths occur after admission to the ICU.<sup>2</sup> Understanding the factors associated with decisions to withhold or withdraw LST are important to policy makers, ethicists, and healthcare leaders because they affect

resources used at the end of life and the need for palliative care and hospice in the ICU setting.

Several studies have characterized the patient characteristics, incidence, and variability associated with limitation of LST in various populations of critically ill patients in the US. We are unaware of another systematic review of the literature that has examined data from these studies in order to understand the process and outcomes of LST limitations. We defined limitations of LST as decisions to withdraw or withhold cardiopulmonary resuscitation through Do Not Resuscitate (DNR) orders, mechanical ventilation, renal replacement therapy, intravenous blood pressure support, or artificial nutrition (enteric or intravenous).

### METHODS

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement was used for reporting. A comprehensive literature search was performed by a medical librarian (TWE) in Ovid MEDLINE, PubMed, Embase, the full Cochrane Library, CINAHL, PsycINFO, the Philosopher's Index, Scopus, Web of Science, and Google Scholar. PubMed was limited to non-MEDLINE records in order to complement the Ovid results. The Georgetown Bioethics Research Library at the Kennedy Institute (<https://bioethics.georgetown.edu/>) was also searched for any unpublished literature. Initial searches were


\*Corresponding Author: Alexia M Torke, MD, MS; E-mail: [atorke@iu.edu](mailto:atorke@iu.edu);  
Telephone: 317-274-9221; Twitter: @AlexiaMTorke

Additional Supporting Information may be found in the online version of this article.

Received: July 7, 2018; Revised: December 3, 2018;

Accepted: December 3, 2018

© 2019 Society of Hospital Medicine DOI 10.12788/jhm.3137



**FIG 1.** Percent of patients with limitations, by year, for studies that included all Intensive Care unit admissions (A) and studies that included patients who died in the ICU (B).

conducted in December 2014, and an update was performed in April 2017. All databases were searched from inception, and bibliographies of relevant studies were reviewed for additional references (Appendix 1).

Database-specific subject headings and keyword variants for each of the five main concepts—intensive care, end-of-life, decision making, limitation of treatment, and death—were identified and combined. Results were limited to the adult population and to the English language.

Two authors independently reviewed article titles and abstracts (KM, AMT). The full text of potentially eligible studies was then reviewed for inclusion. All disputes were discussed and resolved by consensus. The criteria for inclusion were reporting of patient-level data, critical care patients only (or reported separately from other unit types), US setting, and reporting of data on limitations of LST. The exclusion criteria were studies published only as research abstracts, surveys of physicians or families, organ donors, studies of brain death, surveys, patients less than 18 years old, and long-term intensive care settings (ie, long-term acute care hospitals, long-term respiratory units). Also excluded were studies in which an intervention was performed; as a result, all included studies were observational. Research abstracts were excluded because they lacked sufficient detail from which to abstract study quality or results. Studies of organ donation, brain death, and pediatrics were excluded due to differences in the decision-making context that would make it difficult to draw conclusions about adult ICU care. Studies which included an intervention were excluded to avoid affecting the rate of limitation of LST as a result of the intervention, since our goal was to quantify the number of limitations of LST in usual medical practice.

For each article, we abstracted the number of patients who experienced a limitation of LST out of the total population and factors associated with the limitation. If a multivariable analysis was performed, we reported only variables that remained significant in this analysis. We also reported the number of patients who died, and of those, the number of decedents who

underwent a limitation of LST before death. In some cases, this proportion was not reported in the manuscript but could be calculated based on the data presented. This number was calculated based on the number of deaths that were preceded by a limitation in life-sustaining care divided by the total number of deaths. Patients with brain death were not counted as having had a “limitation” if support was withdrawn after the declaration of brain death. We were unable to conduct a meta-analysis of the findings because of the wide variation in study populations and criteria used to define limitations of care.

To assess risk of bias in individual studies, the two raters independently made a yes/no determination regarding several quality metrics established at the outset of the review: clarity of the eligibility criteria for participant inclusion, whether a power or sample size calculation was done, adequacy of the description of the sampling approach and recruitment, and generalizability. Disagreements were resolved by consensus.

## RESULTS

### Study Selection

A total of 2,460 references were identified, and after removal of 578 duplicates, 1,882 unique titles and abstracts were reviewed. One hundred thirteen titles met the inclusion criteria. After review of complete texts, 83 were excluded based on the above criteria (Appendix). This led to a final number of 36 studies included for analysis.

Fifteen articles were prospective, observational studies. The rest were retrospective analyses of patient-level data. Seven were large, multicenter studies with greater than 20 centers involved (including Project IMPACT); six such studies included medical and surgical patients. The remaining large, multicenter study examined a surgical trauma cohort.

Fifteen of the studies addressed DNR as a limitation and 25 addressed other limitations such as withdrawing or withholding LST (several addressed both DNR and another limitation). Nine studies enrolled only patients who had died and the remaining 27 enrolled all ICU admissions.

## Historical Trends

Examination of the three studies that looked at >20 regionally diverse ICUs revealed a trend over time toward increased limitation prior to death (Figure). Jayes looked at the number of DNR orders preceding death from 1979 to 1980 then compared that to a cohort from 1988 to 1990; Prendergast included withholding/withdrawing of LST prior to death from 1994 to 1995; and Quill used the IMPACT database to examine limitations prior to death from 2001 to 2009.<sup>3-5</sup>

## Effect of Unit Specialty

Twelve studies were mixed (surgical/medical or medical/neuro) ICUs, 11 were medical/cardiac units, five were neurologic units, and six were surgical/trauma units only. Two studies did not report unit specialty. Four studies that compared surgical and medical ICUs found that surgical patients were more likely to die with full intervention.<sup>4-7</sup> In all of these studies, medical patients were more likely to have limitations of LST preceding death. Quill, et al. further detailed that emergency surgery was more likely to be associated with limitation than elective surgery.<sup>5</sup>

## Patient Factors

In 15 studies, older age was associated with an increased likelihood of limitations on LST.<sup>3,5-18</sup> In one study, advanced age was associated with early versus late withdrawal.<sup>19</sup> Poor performance status and multiple medical comorbidities were also associated with limitations of LST. The largest population-based study by Quill et al. found that being fully dependent on others upon admission to the ICU was associated with an increased likelihood of limiting LST.<sup>5</sup> Sise et al. found, in an analysis performed over 10 years in one trauma center, that increased age, comorbidities, and a fall as the reason for trauma admission were associated with limitation of LST.<sup>9</sup> Salottolo et al. found that if the reason for trauma admission was a fall, there was an increased odds ratio of DNR status.<sup>18</sup> Many studies found that having medical comorbidities prior to admission was associated with increased likelihood of limiting LST in both medical and surgical patients.<sup>3,7,9,13,15,18</sup>

Five studies found a statistically significant difference between women and men in the likelihood of limitation of LST,<sup>3,5,9,14,16</sup> and another study reported that women who were trauma patients had an increased odds ratio of changing to DNR code status.<sup>18</sup> Only one study found that males were associated with an increased likelihood of limiting aggressive treatment.<sup>20</sup>

White race was associated with increased limitation of LST in nine studies.<sup>4,5,10,11,14-16,21,22</sup> One study in neurocritical care patients found that both white and Hispanic races were correlated with a higher likelihood of limitations.<sup>23</sup> Muni et al. found that nonwhite patients had a statistically significantly lower likelihood of having comfort measures and DNR orders written prior to death, but discussion of prognosis was more likely to be documented in nonwhite patients.<sup>21</sup>

In summary, white race, female gender, and older age were the most frequent factors associated with a higher likelihood of limiting LST.

## Factors Related to Critical Illness

There were several illness severity indicators that were associated with limitations. The Acute Physiology and Chronic Health Evaluation (APACHE) scores were the most common for medical patients and Glasgow Coma Scale (GCS) was the most common for patients with neurologic injury. Eight studies reported that a higher APACHE score was associated with an increased likelihood of limitations.<sup>3,7,10,15,17,20,22,24</sup> Similar associations were found based on the Sepsis Related Organ Failure Assessment score in one study and a scoring system developed by the author in a second study.<sup>25,26</sup>

Seven studies, consisting of three neurologic, two medical-surgical, and two trauma cohorts, reported that a lower GCS score increased the likelihood that the patient would have limited LST.<sup>5,10,11,13,14,18,22</sup> Additionally, Geocadin and colleagues discussed the difficulty with neurological prognostication in clinical practice; they reported that the cortical evoked potential (CEP) was correlated with the time to withdraw LST if the CEP was malignant, and the time to withdraw LST was less in malignant than in benign CEP.<sup>27</sup>

## Mortality and End Effects of Limiting LST

Chen and colleagues used propensity scores to control for mortality differences between patients who had full interventions versus those with limitations and found that higher mortality correlated with the decision to withhold or withdraw LST.<sup>10</sup> Weimer and colleagues used modeling to predict the probable outcome of patients who experienced an intracranial hemorrhage who had limitation of LST. Based on this model, nearly all the patients in their study would have died or had severe disability at 12 months despite having maximal therapy; they concluded that withdrawal of LST may not have been a self-fulfilling prophecy as others have proposed.<sup>28</sup> Mulder and colleagues reported that in a small cohort of out-of-hospital cardiac arrest survivors admitted to the hospital, over one-third had good neurological outcomes after coding after 72 hours.<sup>29</sup> The study highlighted the importance of timing in neurological prognostication.

## Variation in Limitation Rates among Centers

In the 36 studies, we found an overall range of DNR orders from 5.4%<sup>7</sup> to 82.0%.<sup>30</sup> For other limitations, the rates ranged from 6.3%<sup>13</sup> to 80.4%.<sup>31</sup> Hart reported a low rate of limitations (4.8%) at the time of ICU admission.<sup>16</sup> Four large, multicenter studies drew attention to the large variability between critical care centers and the limitation of end-of-life care.<sup>3-5,14</sup> Jayes first described this phenomenon when examining the frequency of DNR orders from 1979 to 1980 and 1988 to 1990.<sup>3</sup> This study found a range from 1.5% to 22%. Later, in another large, multicenter study, Prendergast et al. looked at 131 ICUs at 110 different institutions in 38 states that participated in postgraduate training and found variability in CPR attempts prior to death between 4% and 79%.<sup>4</sup> In 2008, Nathens et al. reported significant variation in DNR rates across trauma centers; they found a higher incidence of DNR orders when there was an open ICU structure.<sup>14</sup>

TABLE. Main Results of the 36 Final Manuscripts

Article Year	Period of Data Collection	Study Population (Number, Description, Study)	Limitation Addressed in the Study	Type of Intensive Care Unit	Study Design	Number (%) with Limitation of LST	Of Those Who Died, What Number (%) Had Limitations	Patient Characteristics Associated with Limitation
Albaeni 2014 <sup>19</sup>	2004-2010	189 survivors of out-of-hospital cardiac arrest pts; 1 unit	Withdrawal of LST	Cardiac/medical	Retrospective	99/189 (52%)	In hospital death 94/147 (63.9%)	No associations were found
Brown 2016 <sup>30</sup>	2003-2008	829 pts who died in ICUs from ILD, COPD, or metastatic; 15 hospitals, 2 neuro ICUs, and 5 non-neuro ICUs	DNR, withdrawal of LST	Medical	Secondary analysis of randomized trial	DNR 681/829 (82.1%); Withdrawal LST 621/829 (74.9%)	DNR 681/829 (82.1%); Withdrawal LST 621/829 (74.9%)	Not reported
Chen 2008 <sup>10</sup>	2002-2005	2,211 consecutive ICU admissions; 1 unit	Withholding LST, excluded withdrawal of treatment	Medical	Retrospective	201/2,211 (9.1%)	Not reported	Age, GCS, APACHE II, race
Creutzfeldt 2015 <sup>36</sup>	2001-2008	200 admission to 2 neuro ICUs who met palliative care triggers, 1,909 admissions to 5 other ICUs in the same hospitals who met triggers (Project IMPACT)	DNR, withholding LST	Neuro vs other	Retrospective	Neuro ICU DNR 16/175 (9.1%); WD/WH 36/175 (20.6%); Non-neuro ICU DNR 212/1,711 (12.4%); WD/WH 195/1,711 (11.4%)	Not reported	Not reported
Diringer 2001 <sup>11</sup>	1994- 2000	2,109 pts treated with mechanical ventilation; 1 unit	Withdrawal of mechanical ventilation	Neurology/ Neurosurgery	Retrospective analysis of prospectively collected clinical database	284/2,109 (13.5%)	279/720 (38.8%)	Surgical pts, age, GCS, race
Geocadin 2006 <sup>27</sup>	Dates not given	58 comatose pts after cardiac arrest consulted on by the neurology service; 1 unit	Withholding and withdrawal of LST	Cardiac, medical	Prospective observational cohort	40/58 (70.0%)	40/48 (83.3%)	Not reported
Hamel 2002 <sup>25</sup>	1989- 1994	596 Non-traumatic coma pts (SUPPORT); 5 medical centers	Withholding CPR and ventilation	Unknown	Secondary analysis of prospective study	121/549 (22.0%)	Not reported	Higher risk assessment score
Hart 2015 <sup>16</sup>	2001-2008	277,693 ICU admissions (Project IMPACT ); 141 ICUs in 105 hospitals	Treatment limitation at the time of ICU admission	Mixed	Retrospective	13,405/277,693 (4.8%)	Not reported	Age, gender, race, pre-admission functional status
Huynh 2013 <sup>12</sup>	2005- 2006	322 ICU pts who died; 1 hospital	Withdrawal of mechanical ventilation	Not specified	Retrospective	159/322 (49.4%)	159/322 (49.4%)	Age, nonsurgical pts
Jayes 1993 <sup>3</sup>	1982; 1988-1990	7,265 ICU admissions 1979-1982 17,440 ICU admissions from 1988 -1990; 42 ICUs	DNR	Medical, surgical	Prospective inception cohort	1979-1982 393/7,265 (5.4%) 1988-1990 1,577/17,440 (9.0%)	60% in 1988-1990; 39% in 1979-1982	Age, sex, APACHE III score, pre-admission daily living activities, chronic disease
Kerlin 2015 <sup>43</sup>	2001-2008	270,742 ICU admissions (Project IMPACT); 143 ICUs	Limitation of LST	Mixed	Retrospective	Not reported	Not reported	Not reported
Kish Wallace <sup>44</sup>	1994-1996	270 admissions to ICU; 1 hospital	DNR	Medical ICU in cancer hospital	Matched pairs case control	41/270 (15.2%)	Unable to determine	Matched groups
Kowalski 2013 <sup>13</sup>	1991-2009	1,134 aneurysmal SAH pts; 2 units	Withdrawal of LST	Neurology/ neurosurgery	Prospective observational	72/1,134 (6.3%)	In hospital deaths 72/207 (34.7%)	GCS, age, comorbidities
Lissauer 2011 <sup>20</sup>	2008- 2010	151 surgical ICU pts who died; 1 unit	Withdrawal and withholding of LST	Surgical	Retrospective analysis of prospectively collected data	111/151 (73.5%)	111/151 (73.5%)	Sex, APACHE IV score

Continued on page 307

TABLE. Main Results of the 36 Final Manuscripts (continued)

Article Year	Period of Data Collection	Study Population (Number, Description, Study)	Limitation Addressed in the Study	Type of Intensive Care Unit	Study Design	Number (%) with Limitation of LST	Of Those Who Died, What Number (%) Had Limitations	Patient Characteristics Associated with Limitation
Mayer 1999 <sup>23</sup>	1994- 1997	105 neurocritical ICU pts who died; 1 unit	Withdrawal of ventilation	Neuro	Retrospective	50/105 (47.6%)	50/105 (47.6%)	Race
Mehter 2014 <sup>17</sup>	2000-2005	809 ICU pts with ARDS and documented code status; multicenter	DNR, withdrawal and withholding of LST	Medical	Retrospective observation of clinical trial subset	DNR 232/809 (28.7%); WD/WH 195/809 (24.1%)	90 day mortality DNR 225/284 (79.2%)	APACHE III score
Mulder 2013 <sup>37</sup> and 2014 <sup>29</sup>	2007-2011	154 comatose survivors out of hospital cardiac arrest survivors admitted to the ICU; 1 center	Withdrawal of LST	Medical	Prospective observational	63/154 (40.9%)	63/78 (80.8%)	Not reported
Muni 2011 <sup>21</sup>	2003- 2008	3,138 ICU pts who died; 15 ICUs	DNR, comfort care, RRT, MV, pressors, enteral feedings	Medical	Retrospective	2,274/3,121 (72.90%)	2,274/ 3,121 (72.90%)	Race
Naib 2015 <sup>38</sup>	2012	1,368 pts; 1 CICU	Withdrawal and withholding of LST	Cardiac	Retrospective	Not reported	ICU deaths 85/117 (72.6%)	Not reported
Nathens 2008 <sup>14</sup>	2001- 2002	6,765 trauma pts; 68 centers	DNR	Trauma, mixed	Prospective observational	464/6,765 (6.9%)	408/603 67.7%	Age, race, sex, global injury severity score, admission GCS
Plaisier 2002 <sup>31</sup>	1994- 1998	102 trauma pts who died; 1 unit	Withdrawal and withholding of LST	Trauma	Retrospective	82/102 (80.4%)	82/102 (80.4%)	Not reported
Prendergast 1997 <sup>39</sup>	1987-1988;1992-1993	1987-1988: 1,719 ICU admissions 1992-1993: 1,711 ICU admissions; 2 hospitals	Withdrawal and withholding of LST	Medical/surgical	Retrospective for 1987-1988, prospective for 92-93	1987-1988 114/1719 (6.6%); 1992-1993 177/1711 (10%)	1987-1988 114/224 (50.9%); 1992-1993 179/200 (89.5%)	Not reported
Prendergast 1998 <sup>4</sup>	1994- 1995	6,303 ICU pts who died; 131 ICUs	DNR, withdrawal and withholding of LST	Medical/surgical	Prospective observational	DNR 5,506/6,303 75.5%; LST 3,036/6,303 (48.2%)	DNR 5,506/6,303 75.5%; LST 3,036/6,303 (48.2%)	Not reported
Quill 2014 <sup>5</sup>	2001- 2009	269,002 full code ICU admissions; 153 units	Withdrawal and withholding of LST	Medical/surgical	Retrospective	31,408/269,002 (11.7%)	ICU deaths 18,460/21,758 (84.8%); Hospital deaths 23,469/33,910 (69.2%)	Surgery type, age, race, sex, GCS score, dependent pre-admission status
Reichner 2006 <sup>26</sup>	2002- 2004	47 lung cancer pts admitted to ICU/1 unit	DNR/terminal extubation	Medical	Retrospective	DNR on ICU admission 12/47 (25.5%); any DNR order 35/47 (74.5%); terminally extubated 5/47 (10.6%)	Not reported	Sorse SOFA score
Rubin 2014 <sup>22</sup>	2002- 2009	1885 pts with neurologic injury with a GCS < 9, and mechanical ventilation; 1 unit	Withdrawal of ventilation, DNR	Neuro	Prospective observational	529/1,885 (28%) withdrawal ventilation; 714/1,885 (37.9%) DNR	533/788 (67.6%)	Nonsurgical pts low GCS, high APACHE II white race
Salottolo 2015 <sup>18</sup>	2008- 2013	10,053 trauma pts; 1 ICU	DNR, withdrawal of LST noted for deaths only	Surgical trauma	Retrospective	1,523 (15.1%)	In hospital death or hospice discharge, DNR 336/455 (73.8%); withdraw LST 45/455 (9.9%)	Age, sex, GCS score, Charlson Comorbidity Index, fall/head injury
Sise 2012 <sup>9</sup>	2000-2009	698 trauma pts who died; 1 unit	Withdrawal or withholding of LST	Trauma	Retrospective	375/698 (53.7%)	375/698 (53.7%)	Age, sex, ISS score, GCS score, fall, Charlson Comorbidity Index, trauma consult
Smedira 1990 <sup>40</sup>	1987- 1988	1,719 ICU patient; 2 hospitals	Withdrawal or withholding of LST	Medical/surgical	Prospective observational	97/1,719 <sup>a</sup> (5.6%)	ICU deaths 71/109pts (62.1%)	Not reported

Continued on page 308

TABLE. Main Results of the 36 Final Manuscripts (continued)

Article Year	Period of Data Collection	Study Population (Number, Description, Study)	Limitation Addressed in the Study	Type of Intensive Care Unit	Study Design	Number (%) with Limitation of LST	Of Those Who Died, What Number (%) Had Limitations	Patient Characteristics Associated with Limitation
Turnbull 2014 <sup>6</sup>	2004- 2007	490 acute lung injury patient; 13 ICUs	Withdrawal or withholding of LST	Medical/surgical	Prospective observational	192/490 (39.2%)	ICU deaths 166/214 (77.60%)	Patient type, age
Van Scoy 2013 <sup>41</sup>	2006- 2008	100 pts who died after 72 hour stay in the ICU; 1 hospital	DNR	Medical/surgical	Retrospective	52/100 (52.0%)	52/100 (52.0%)	APACHE II score
Weimer 2016 <sup>28</sup>	2008- 2011	383 intracranial hemorrhage pts; 1 ICU	Withdrawal of LST	Neuro	Retrospective analysis of prospectively collected data	26/383 (6.8%)	(26/67) 38.8%	None
Weireter 2014 <sup>8</sup>	2008- 2012	536 trauma pts who died; 1 unit	Withdrawal of LST	Trauma	Retrospective analysis of prospectively collected data	158/536 (29.5%)	158/536 (29.5%)	Older age
White 2006 <sup>42</sup>	2003-2004	303 ICU pts; 1 hospital	DNR, limitation of LST	Medical	Prospective observational	DNR 13/49 (26.5%); Withdrawal LST 8/49 (16.3%)	DNR 9/13 (69.2%); Withdrawal of LST 8/13 (61.5%)	
Youngner 1985 <sup>15</sup>	1983 - 1983	506 MICU admissions; 1 unit	DNR	Medical	Prospective observational	71/506 (14%)	Hospital deaths 62/123 (50.4%)	Age, race, APACHE score, pre-admission health status
Zimmerman 1986 <sup>7</sup>	1979-1982	7265 ICU admissions; 13 units	DNR	Mixed	Prospective observational	393/7,265 (5.4%)	ICU deaths 237/611 (39%)	Surgery type, age, APACHE score, pre-admission health status

\*Pts who were brain dead were not counted as having life-sustaining treatments withdrawn or withheld.

Abbreviations: APACHE, acute physiology and chronic health evaluation; COPD: chronic obstructive pulmonary disease; DNR, do not resuscitate; GCS, Glasgow coma scale ICU, intensive care unit; IMPACT, informing the pathway of COPD treatment; LST, life-sustaining treatments; MV, mechanical ventilation; Pts, patients; RRT, rapid response team; SAH, subarachnoid hemorrhage; SUPPORT: study to understand prognoses and preferences for outcomes and risks of treatments; WD, withdraw; WH, withhold.

Overall, there was wide variation in the proportion of deaths preceded by limitation of LST, ranging from 29.5% in one study of trauma patients<sup>8</sup> to 92% in another study of trauma patients whose death occurred after 24 hours of care.<sup>9</sup> In the largest study to date by Quill and colleagues utilizing the IMPACT database, they found large variability in the number of deaths preceded by full intervention based on differences in practice patterns of critical care centers.<sup>5</sup>

## Bias

All studies indicated clear eligibility criteria for inclusion and described their sampling approach in adequate detail. All but one stated their method of participant recruitment, and the one remaining study was a secondary analysis and referenced the earlier manuscript.<sup>30</sup> No study provided a power or sample size calculation, and sample sizes varied widely. Generalizability was most affected by the fact that many studies were conducted in a single ICU.

## DISCUSSION

This systematic review of LST in US ICUs found several patient and illness factors that were associated with limitation of LST. The association of preadmission functional status and comorbidities with limitation of LST suggest that prior health is a factor in decision making. Further, ICU severity of illness, as measured by several commonly used indices, was associated with limitations.

Although variations in study design precluded meta-analysis, examination of the largest studies suggests that limitations are becoming more frequent over time. Also, early studies generally addressed DNR status, while later studies included withdrawal or withholding of LST, most commonly artificial ventilation. These findings reflect the current consensus in US medicine that it is ethically acceptable to limit LSTs in cases when they no longer benefit the patient or the patient would no longer want them.<sup>32,33</sup>

Some studies found variability by unit type, suggesting that decision making may differ among surgical, medical, and neurologic illness. Mayer and Kossoff concluded, in study of a cohort of neurocritical care ICU patients, that medical patients often have issues of physiologic futility and imminent death, whereas neurologic patients more often confront issues of quality of life. They also note that there is a difference in how patients with differing illnesses die; medical patients will have limitation of hemodialysis or vasopressors, whereas neurologic surrogate decision makers often confront decisions around terminal extubation.<sup>23</sup>

Some patient-level factors, such as race or ethnicity, may point to cultural differences in preferences for LST at the end of life. Other authors have documented that African American patients are more likely to choose end-of-life care for themselves or their family members, which may be due to cultural or religious factors as well as to a history of unequal access to

medical care.<sup>34</sup> Reasons for the finding that women are more likely to have limitations has not been as well described. Further research could explore whether this is due to differences in patient preferences by gender or to other factors.

Even when examining patient-level factors, illness severity and type of ICU, the wide variability in end-of-life care in critical care units across the country is still large. A worldwide review also found a high degree of variability, even within geographical regions.<sup>35</sup> More research is needed to understand the factors associated with this wide variability, as this seems to indicate that approaches to end-of-life care may vary based on the ICU as much as individual patient preferences or clinical factors.

These findings can inform clinicians about variables that are important in the decision-making process. Patient age and race are factors to consider in the likelihood of reaching a decision to set limitations. Information about patients' health status prior to critical illness, as well as ICU illness severity, are also important considerations.

The limitations of this review include the wide variety of LSTs assessed, including code status change, ventilator withdrawal, removal of pressors, and cessation of renal replacement therapy. Also, there was variation in sample size and the number of included units. There was also significant heterogeneity in the outcomes addressed and the variety of methods used in the included studies. We attempted to address this with an analysis of the quality of the studies, but given the wide variability, we were unable to account for all of the differences; unfortunately, this is a standard issue within studies that utilize systematic reviews, as well as similar concepts such as meta-analyses.

In conclusion, the increase in the frequency of limitations of LST in critically ill patients and a change in the nature of limitations from DNR order to withdrawal or withholding of LST suggests a trend toward growing acceptance of limiting treatments in critical illness. The wide variation in withdrawal of care in US ICUs does not seem fully explained by patient variables including preferences, illness type, or changes over time. Factors such as poor prefunctional status, a higher number of comorbid conditions prior to critical illness, and the severity of critical illness are likely important for surrogates and clinicians to consider during goals of care discussions. Further research is needed to explore why patients may receive very different types of care at the end of life depending the institution and ICU in which they receive their care.

Disclosures: The authors have no conflicts of interest to disclose. This work was performed at the Indiana University School of Medicine.

Funding: Financial support for Dr. Torke was provided by a Midcareer Investigator Award in Patient Oriented Research from the National Institute on Aging (K24AG053794). Dr. McPherson was supported by the Indiana University Department of Medicine.

## References

1. Sprung CL, Raphaely RC, Hynninen M, et al. Consensus report on the ethics of foregoing life-sustaining treatments in the critically ill. Task Force on Ethics of the Society of Critical Care Medicine. *Crit Care Med*. 1990;18(12):1435-1439.
2. Angus DC, Barnato AE, Linde-Zwirble WT, et al. Use of intensive care at the end of life in the United States: an epidemiologic study. *Crit Care Med*. 2004;32(3):638-643.
3. Jayes RL, Zimmerman JE, Wagner DP, Draper EA, Knaus WA. Do-not-resuscitate orders in intensive care units. Current practices and recent changes. *JAMA*. 1993;270(18):2213-2217. doi: 10.1001/jama.1993.03510180083039.
4. Prendergast TJ, Claessens MT, Luce JM. A national survey of end-of-life care for critically ill patients. *Am J Respir Crit Care Med*. 1998;158(4):1163-1167. doi: 10.1164/ajrccm.158.4.9801108.
5. Quill CM, Ratcliffe SJ, Harhay MO, Halpern SD. Variation in decisions to forgo life-sustaining therapies in US ICUs. *Chest*. 2014;146(3):573-582. doi: 10.1378/chest.13-2529.
6. Turnbull AE, Ruhl AP, Lau BM, Mendez-Tellez PA, Shanholtz CB, Needham DM. Timing of limitations in life support in acute lung injury patients: a multisite study. *Crit Care Med*. 2014;42(2):296-302. doi: 10.1097/CCM.0b013e3182a272db.
7. Zimmerman JE, Knaus WA, Sharpe SM, Anderson AS, Draper EA, Wagner DP. The use and implications of do not resuscitate orders in intensive care units. *JAMA*. 1986;255(3):351-356. doi: 10.1001/jama.1986.03370030071030.
8. Weireter LJ, Jr., Collins JN, Britt RC, Novosel TJ, Britt LD. Withdrawal of care in a trauma intensive care unit: the impact on mortality rate. *Am Surg*. 2014;80(8):764-767.
9. Sise MJ, Sise CB, Thorndike JF, Kahl JE, Calvo RY, Shackford SR. Withdrawal of care: A 10-year perspective at a Level I trauma center. *J Trauma Acute Care Surg*. 2012;72(5):1186-1191. doi: 10.1097/TA.0b013e31824d0e57.
10. Chen Y-Y, Connors AF, Jr., Garland A. Effect of decisions to withhold life support on prolonged survival. *Chest*. 2008;133(6):1312-1318. doi: 10.1378/chest.07-1500.
11. Diringer MN, Edwards DF, Aiyagari V, Hollingsworth H. Factors associated with withdrawal of mechanical ventilation in a neurology/neurosurgery intensive care unit. *Crit Care Med*. 2001;29(9):1792-1797.
12. Huynh TN, Walling AM, Le TX, Kleerup EC, Liu H, Wenger NS. Factors associated with palliative withdrawal of mechanical ventilation and time to death after withdrawal. *J Palliat Med*. 2013;16(11):1368-1374. doi: 10.1089/jpm.2013.0142.
13. Kowalski RG, Chang TR, Carhuapoma JR, Tamargo RJ, Naval NS. Withdrawal of technological life support following subarachnoid hemorrhage. *Neurocrit Care*. 2013;19:269-275. doi: 10.1007/s12028-013-9929-8.
14. Nathens AB, Rivara FP, Wang J, Mackenzie EJ, Jurkovich GJ. Variation in the rates of do not resuscitate orders after major trauma and the impact of intensive care unit environment. *J Trauma*. 2008;64(1):81-88;discussion 8-91. doi: 10.1097/TA.0b013e31815dd4d7.
15. Youngner SJ, Lewandowski W, McClish DK, Juknialis BW, Coulton C, Bartlett ET. 'Do not resuscitate' orders. Incidence and implications in a medical-intensive care unit. *JAMA*. 1985;253(1):54-57. doi: 10.1001/jama.1985.03350250062023.
16. Hart JL, Harhay MO, Gabler NB, Ratcliffe SJ, Quill CM, Halpern SD. Variability among US intensive care units in managing the care of patients admitted with preexisting limits on life-sustaining therapies. *JAMA Intern Med*. 2015;175(6):1019-1026. doi: 10.1001/jamainternmed.2015.0372.
17. Mehter HM, Wiener RS, Walkey AJ. "Do not resuscitate" decisions in acute respiratory distress syndrome: a secondary analysis of clinical trial data. *Ann Am Thorac Soc*. 2014;11(10):1592-1596. doi: 10.1513/AnnalsATS.201406-244BC.
18. Salottolo K, Offner PJ, Orlando A, et al. The epidemiology of do-not-resuscitate orders in patients with trauma: a community level one trauma center observational experience. *Scand J Trauma Resusc Emerg Med*. 2015;23(1):9. doi: 10.1186/s13049-015-0094-2.
19. Albaeni A, Chandra-Strobos N, Vaidya D, Eid SM. Predictors of early care withdrawal following out-of-hospital cardiac arrest. *Resuscitation*. 2014;85(11):1455-1461. doi: 10.1016/j.resuscitation.2014.08.030.
20. Lissauer ME, Naranjo LS, Kirchoffner J, Scalea TM, Johnson SB. Patient characteristics associated with end-of-life decision making in critically ill surgical patients. *J Am Coll Surg*. 2011;213(6):766-770. doi: 10.1016/j.jamcollsurg.2011.09.003.
21. Muni S, Engelberg RA, Treece PD, Dotolo D, Curtis JR. The influence of race/ethnicity and socioeconomic status on end-of-life care in the ICU. *Chest*. 2011;139(5):1025-1033. doi: 10.1378/chest.10-3011.
22. Rubin MA, Dhar R, Diringer MN. Racial differences in withdrawal of mechanical ventilation do not alter mortality in neurologically injured patients. *J Crit Care*. 2014;29(1):49-53. doi: 10.1016/j.jc.2013.08.023.
23. Mayer SA, Kossoff SB. Withdrawal of life support in the neurological intensive care unit. *Neurology*. 1999;52(8):1602-1609. doi: 10.1212/WNL.52.8.1602.

24. 2nd National Congress on Medicinal Plants. *Iranian J Pharm Res.* 2013;12:43.
25. Hamel MB, Phillips R, Teno J, et al. Cost effectiveness of aggressive care for patients with nontraumatic coma. *Crit Care Med.* 2002;30(6):1191-1196.
26. Reichner CA, Thompson JA, O'Brien S, Kuru T, Anderson ED. Outcome and code status of lung cancer patients admitted to the medical ICU. *Chest.* 2006;130(3):719-723. doi: 10.1378/chest.130.3.719.
27. Geocadin RG, Buitrago MM, Torbey MT, Chandra-Strobos N, Williams MA, Kaplan PW. Neurologic prognosis and withdrawal of life support after resuscitation from cardiac arrest. *Neurology.* 2006;67(1):105-108. doi: 10.1212/01.wnl.0000223335.86166.b4.
28. Weimer JM, Nowacki AS, Frontera JA. Withdrawal of life-sustaining therapy in patients with intracranial hemorrhage: self-fulfilling prophecy or accurate prediction of outcome? *Crit Care Med.* 2016;44(5):1161-1172. doi: 10.1097/CCM.0000000000001570.
29. Mulder M, Gibbs HG, Smith SW, et al. Awakening and withdrawal of life-sustaining treatment in cardiac arrest survivors treated with therapeutic hypothermia. *Crit Care Med.* 2014;42(12):2493-2499. doi: 10.1097/CCM.0000000000000540.
30. Brown CE, Engelberg RA, Nielsen EL, Curtis JR. Palliative care for patients dying in the intensive care unit with chronic lung disease compared with metastatic cancer. *Ann Am Thorac Soc.* 2016;13(5):684-689. doi: 10.1513/AnnalsATS.201510-667OC.
31. Plaisier BR, Blostein PA, Hurt KJ, Malangoni MA. Withholding/withdrawal of life support in trauma patients: is there an age bias? *Am Surg.* 2002;68(2):159-162.
32. Beauchamp, Childress JF. *Principles of Biomedical Ethics.* 13th ed. Oxford: Oxford University Press; 2013.
33. Jonson AR, Siegler M, Winslade WJ. *Clinical Ethics: A Practical Approach to Ethical Decisions in Clinical Medicine.* New York: McGraw Hill; 2015.
34. Johnson KS, Elbert Avila KI, Tulsy JA. The influence of spiritual beliefs and practices on the treatment preferences of African Americans: a review of the literature. *J Am Geriatr Soc.* 2005;53(4):711-719. doi: 10.1111/j.1532-5415.2005.53224.x.
35. Mark NM, Rayner SG, Lee NJ, Curtis JR. Global variability in withholding and withdrawal of life-sustaining treatment in the intensive care unit: a systematic review. *Intensive Care Med.* 2015;41(9):1572-1585. doi: 10.1007/s00134-015-3810-5.
36. Creutzfeldt CJ, Wunsch H, Curtis JR, Hua M. Prevalence and Outcomes of Patients Meeting Palliative Care Consultation Triggers in Neurological Intensive Care Units. *Neurocrit Care.* 2015;23:14-21.
37. Mulder M, Smith SW, Dhaliwal RS, Goodwin HE, Scott NL, Geocadin RG. Comatose survivors of cardiac arrest and therapeutic hypothermia: Time of awakening and withdrawal of life sustaining therapies. *Neurocrit Care.* 2013;19:S281.
38. Naib T, Lahewala S, Arora S, Gidwani U. Palliative care in the cardiac intensive care unit. *Am J Cardiol.* 2015;115:687-90.
39. Prendergast TJ, Luce JM. Increasing incidence of withholding and withdrawal of life support from the critically ill. *Am J Respir Crit Care Med.* 1997;155:15-20.
40. Smedira NG, Evans BH, Grais LS, et al. Withholding and withdrawal of life support from the critically ill. *N Engl J Med.* 1990;322:309-15.
41. Van Scoy LJ, Sherman M. Factors Affecting Code Status in a University Hospital Intensive Care Unit. *Death Stud.* 2013;37:768-81.
42. White DB, Curtis JR, Lo B, Luce JM. Decisions to limit life-sustaining treatment for critically ill patients who lack both decision-making capacity and surrogate decision-makers. *Crit Care Med.* 2006;34:2053-9.
43. Kerlin MP, Harhay MO, Kahn JM, Halpern SD. Nighttime intensivist staffing, mortality, and limits on life support; a retrospective cohort study. *Chest.* 2015;147(4):951-958.
44. Kish Wallace S, Martin CG, Shaw AD, Price KJ. Influence of an advance directive on the initiation of life support technology in critically ill cancer patients. *Crit Care Med.* 2001;29(12):2294-2298.