

Proceedings of the Institute of Mathematics and Mechanics,
National Academy of Sciences of Azerbaijan
Volume 43, Number 2, 2017, Pages 240–251

RIEMANN BOUNDARY VALUE PROBLEMS IN GENERALIZED WEIGHTED HARDY SPACES

BILAL T. BILALOV, ALI A. HUSEYNLI, AND FIDAN SH. SEYIDOVA

Abstract. Riemann boundary value problem of analytic function theory in weighted Hardy classes with variable summability index is considered in this work. The Fredholmness of this problem is investigated under certain conditions on coefficients and a weight. The general solution for homogeneous problem is obtained in weighted Hardy classes with variable summability index. In the case where the weight function satisfies the Muckenhoupt condition with variable summability index, the solvability of the non-homogeneous Riemann problem with the right side from the generalized weighted Lebesgue space is studied.

1. Introduction

The boundary value problems of analytic functions theory has a deep history. Well-known monographs like [12, 24, 13, 27, 20], have been dedicated to the boundary value problems of analytic functions. In the classical formulation these problems are well studied (see [24, 13, 27, 20]). In terms of L_p -metrics in Hardy and Smirnov classes, the theory of boundary value problems has been developed by various mathematicians and has been illuminated in monographs I.I. Danilyuk [12], G.S.Litvinchuk [20], etc. Since recently, there arose an interest in the study of the Riemann boundary value problems in different spaces of analytic functions (see e.g. [17, 22, 15, 21, 14, 23, 8, 9, 2, 3, 4, 5, 6, 7, 10], etc). All the above-cited works dealing with weighted spaces of analytic functions consider the spaces with power weights.

In the present paper, we consider the Riemann boundary value problem in the weighted Hardy space with general weight and piecewise Hölder coefficient. We will find a sufficient condition on the weight under which this problem is Noetherian. We will also calculate the index of the problem. The general solution for homogeneous problem is obtained in weighted Hardy classes with variable summability index. In the case where the weight function satisfies the Muckenhoupt condition with variable summability index, the solvability of the non-homogeneous Riemann problem with the right side from the generalized weighted Lebesgue space is studied. It should be noted that in [25] the Noetherness of Riemann problem is studied in generalized weighted Hardy classes with power weight.

2010 *Mathematics Subject Classification.* 30E25, 30D55, 30B60, 42C15, 46A35.

Key words and phrases. Riemann problem, variable exponent, Hardy classes.

2. Needful information

Let C be the complex plane, $\omega \equiv \{z \in C : |z| < 1\}$ be the unit circle, and $\partial\omega \equiv \{z \in C : |z| = 1\}$ be the unit circumference. We will need some notion from the theory of generalized Lebesgue spaces.

Let $p : [-\pi, \pi] \rightarrow [1, +\infty)$ be some Lebesgue-measurable function. By \mathcal{L}_0 we denote the class of all functions measurable on $[-\pi, \pi]$ with respect to Lebesgue measure. Denote

$$I_p(f) \stackrel{def}{=} \int_{-\pi}^{\pi} |f(t)|^{p(t)} dt.$$

Let

$$\mathcal{L} \equiv \{f \in \mathcal{L}_0 : I_p(f) < +\infty\}.$$

With respect to the usual linear operations of addition and multiplication by a number \mathcal{L} is a linear space as $p^+ = \sup_{[-\pi, \pi]} p(t) < +\infty$. With respect to the norm

$$\|f\|_{p(\cdot)} \stackrel{def}{=} \inf \left\{ \lambda > 0 : I_p\left(\frac{f}{\lambda}\right) \leq 1 \right\},$$

\mathcal{L} is a Banach space, and we denote it by $L_{p(\cdot)}$. Let

$$WL \stackrel{def}{=} \left\{ p : p(-\pi) = p(\pi); \exists C > 0, \forall t_1, t_2 \in [-\pi, \pi] : |t_1 - t_2| \leq \frac{1}{2} \Rightarrow |p(t_1) - p(t_2)| \leq \frac{C}{-\ln|t_1 - t_2|} \right\}.$$

Throughout this paper, $q(t)$ will denote the conjugate of a function $p(t)$: $\frac{1}{p(t)} + \frac{1}{q(t)} \equiv 1$. Denote $p^- = \inf_{[-\pi, \pi]} p(t)$. The following generalized Hölder inequality is true

$$\int_{-\pi}^{\pi} |f(t)g(t)| dt \leq c(p^-, p^+) \|f\|_{p(\cdot)} \|g\|_{q(\cdot)},$$

where $c(p^-, p^+) = 1 + \frac{1}{p^-} - \frac{1}{p^+}$. Directly from the definition we get the property, which will be used in the sequel.

Property 2.1. *If $|f(t)| \leq |g(t)|$ a.e. on $(-\pi, \pi)$, then $\|f\|_{p(\cdot)} \leq \|g\|_{p(\cdot)}$.*

We will need the following easy-to-prove statement.

Statement 2.1. *Let $p \in WL$, $p(t) > 0, \forall t \in [-\pi, \pi]; \{\alpha_i\}_1^m \subset R$. The weighted function $\rho(t) = \prod_{i=1}^m |t - \tau_i|^{\alpha_i}$, belongs to the space $L_{p(\cdot)}$, if $\alpha_i > -\frac{1}{p(\tau_i)}, \forall i = \overline{1, m}$; where $-\pi = \tau_1 < \tau_2 < \dots < \tau_m = \pi$.*

By S we denote the singular integral

$$Sf = \frac{1}{2\pi i} \int_{\partial\omega} \frac{f(\tau)}{\tau - t} d\tau, \quad t \in \partial\omega,$$

Let $\rho : [-\pi, \pi] \rightarrow (0, +\infty)$ be some weight function. Define the weighted classes

$$L_{p(\cdot),\rho(\cdot)} : L_{p(\cdot),\rho(\cdot)} \stackrel{def}{=} \{f : \rho f \in L_{p(\cdot)}\},$$

with a norm $\|f\|_{p(\cdot),\rho(\cdot)} \stackrel{def}{=} \|\rho f\|_{p(\cdot)}$. The validity of the following statement is established in [16].

Statement 2.2. [16] *Let $p \in WL$, $1 < p^-$. Then, singular operator S is acting boundedly from $L_{p(\cdot),\rho(\cdot)}$ to $L_{p(\cdot),\rho(\cdot)}$ if and only if*

$$-\frac{1}{p(\tau_k)} < \alpha_k < \frac{1}{q(\tau_k)}, \quad k = \overline{1, m}. \tag{2.1}$$

We also will use the following variable Muckenhoupt condition.

Definition 2.1. Given $p(\cdot) : [-\pi, \pi] \rightarrow [1, +\infty]$, and a weight function $\omega(\cdot)$. We say that $\omega \in A_{p(\cdot)}$ if

$$\sup_{I \subset [-\pi, \pi]} |I|^{-1} \|\omega(\cdot) \chi_I(\cdot)\|_{p(\cdot)} \|\omega^{-1}(\cdot) \chi_I(\cdot)\|_{q(\cdot)} < +\infty.$$

It should be noted that for more details on these and other facts one can see works [11, 19, 28, 26].

We will also need the weighted Hardy classes. By H_p^+ we denote the usual Hardy class of analytical functions in ω . Let \mathcal{A} be the σ -algebra of Borel sets in $[-\pi, \pi]$ and ρ be a σ -finite measure on \mathcal{A} . $L_{p(\cdot);d\rho} \equiv L_{p(\cdot);d\rho}(-\pi, \pi)$ will denote the Banach space \mathcal{A} of measurable functions on $[-\pi, \pi]$ furnished with the norm

$$\|f\|_{p(\cdot)} \stackrel{def}{=} \inf \left\{ \lambda > 0 : I_{p(\cdot);d\rho} \left(\frac{f}{\lambda} \right) \leq 1 \right\},$$

where

$$I_{p(\cdot);d\rho}(f) = \int_{-\pi}^{\pi} |f(t)|^{p(t)} d\rho(t).$$

Assume

$$\tilde{H} \equiv \{f \in H_1^+ : f^+ \in L_{p(\cdot);d\rho}\},$$

where $f^+(e^{it})$ are non-tangential boundary values of function f on $\partial\omega$. The norm in space \tilde{H} is defined by

$$\|f\|_{\tilde{H}} = \|f^+(e^{it})\|_{p(\cdot);d\rho}. \tag{2.2}$$

Let us show that if the continuous inclusion $L_{p(\cdot);d\rho} \subset L_1$ is true, then \tilde{H} is a Banach space with respect to the norm (2.2), where $L_1 \equiv L_1(-\pi, \pi)$ is a Lebesgue space of summable functions on $(-\pi, \pi)$. Let $\{f_n\}_{n \in \mathbb{N}} \subset \tilde{H}$ be some fundamental sequence, i.e.

$$\|f_n^+(e^{it}) - f_m^+(e^{it})\|_{p(\cdot);d\rho} \rightarrow 0, \quad n, m \rightarrow \infty,$$

where $f_n^+ = f_n|_{\partial\omega}$ are non-tangential boundary values of function f_n on $\partial\omega$. From completeness of $L_{p(\cdot);d\rho}$ it follows that

$$\exists \varphi \in L_{p(\cdot);d\rho} : f_n^+(e^{it}) \rightarrow \varphi(t), \quad n \rightarrow \infty, \text{ in } L_{p(\cdot);d\rho}.$$

We have

$$\begin{aligned} \|f_n - f_m\|_{H_1^+} &= \|f_n^+(e^{it}) - f_m^+(e^{it})\|_{L_1} \leq \\ &\leq M \|f_n^+(e^{it}) - f_m^+(e^{it})\|_{p(\cdot);d\rho} \rightarrow 0, \quad n, m \rightarrow \infty, \end{aligned}$$

where M is a constant depending only on $p(\cdot)$ and $\rho(\cdot)$. This immediately implies that $\{f_n\}_{n \in \mathbb{N}}$ is fundamental in H_1^+ and, consequently

$$\exists f \in H_1^+ : f_n \rightarrow f, \quad n \rightarrow \infty, \quad \text{in } H_1^+.$$

Thus

$$f_n^+(e^{it}) \rightarrow f^+(e^{it}), \quad n \rightarrow \infty, \quad \text{in } L_1, \quad \text{where } f^+ = f/\partial\omega.$$

On the other hand, from

$$\|f_n^+(e^{it}) - \varphi(t)\|_{L_1} \leq M \|f_n^+(e^{it}) - \varphi(t)\|_{p(\cdot);d\rho} \rightarrow 0, \quad n \rightarrow \infty,$$

it follows that, $f_n^+(e^{it}) \rightarrow \varphi(t)$, $n \rightarrow \infty$, in L_1 . Then it is absolutely clear that $f^+(e^{it}) = \varphi(t)$ a.e. on $(-\pi, \pi)$, and, as a result, $f_n \rightarrow f$, $n \rightarrow \infty$, in \tilde{H} , i.e. \tilde{H} is a Banach space, and we denote it by $H_{p(\cdot);d\rho}^+$. Thus, we get the validity of

Theorem 2.1. *Let the embedding $L_{p(\cdot);d\rho} \subset L_1$, $1 \leq p < +\infty$, be continuous. Then $H_{p(\cdot);d\rho}^+$ is a Banach space.*

Let us define the class ${}_m H_{p(\cdot)}^-$, where $m \in \mathbb{Z}$ is some integer. So, let $\Phi(z)$ be an analytic function outside ω , which has finite order at infinity, i.e. let

$$\Phi(z) = \sum_{n=-\infty}^{n=k} a_n z^n, \quad k < +\infty,$$

be the Laurent series in a neighborhood of the infinitely remote point of the function $\Phi(z)$.

Under the order of the function $\Phi(z)$ at the infinitely remote point we will mean the largest number $n = k$ of these expansions, for which $a_n \neq 0$. In the case of $k = 0$ the function $\Phi(z)$ is bounded and different from zero at the point $z = \infty$; in the case of $k > 0$ it has a pole of order k ; and in the case of $k < 0$ —has a zero order $(-k)$. Thus, $\Phi(\cdot)$ has a form $\Phi(z) = \Phi_0(z) + \Phi_1(z)$, where $\Phi_0(z)$ —is a regular part, and $\Phi_1(z)$ is a main part of Laurent series of $\Phi(z)$ at the infinitely remote point. If in this case $k \leq m$ and the function $\overline{\Phi_0\left(\frac{1}{z}\right)}$ belongs to the class $H_{p(\cdot)}^+$, then we will say that the function $\Phi(\cdot)$ belongs to Hardy class ${}_m H_{p(\cdot)}^-$ outside ω . Denote

$$\tilde{H}^- \equiv \{f \in {}_m H_1^- : f^-(e^{it}) \in L_{p(\cdot);d\rho}\},$$

where $f^-(e^{it}) = f/\partial\omega$ are non-tangential boundary values of f on $\partial\omega$ from the outside of ω . The norm in \tilde{H}^- is defined as follows

$$\|f\|_{\tilde{H}^-} \equiv \|f^-(e^{it})\|_{p(\cdot);d\rho}, \quad \forall f \in \tilde{H}^-. \tag{2.3}$$

Similarly to the previous case, we can prove that for $L_{p(\cdot);d\rho} \subset L_1$, the class \tilde{H}^- is a Banach space with respect to the norm (2.3), and we denote it by ${}_m H_{p(\cdot);d\rho}^-$. Thus, we have

Theorem 2.2. *Let $L_{p(\cdot);d\rho} \subset L_1$, $1 \leq p < +\infty$, be true. Then the space defined above ${}_m H_{p(\cdot);d\rho}^-$ is a Banach space.*

3. General solution of homogeneous problem

In the sequel, we will assume that the measure $\rho(\cdot)$ is absolutely continuous with respect to a Lebesgue measure on $[-\pi, \pi]$ and $d\rho(x) = \vartheta^{p(x)}(x) dx$. Corresponding spaces are denoted by $L_{p(\cdot);\vartheta}$ and $H_{p(\cdot);\vartheta}^\pm$, respectively. Consider the following homogeneous Riemann problem

$$F^+(\tau) - G(\tau)F^-(\tau) = 0, \quad \tau \in \partial\omega, \tag{3.1}$$

where $G : \partial\omega \rightarrow C$ is some complex-valued function (coefficient of the problem). By solution of the problem (3.1) we mean a pair of analytic functions

$$(F^+(z); F^-(z)) \in H_{p(\cdot);\vartheta}^+ \times_m H_{p(\cdot);\vartheta}^-$$

whose non-tangential boundary values on $\partial\omega$ a.e. satisfy the relation (3.1), where $p(\cdot) \in WL \wedge p^- > 1$ and $m < 0$ is some fixed integer. In the sequel, we will assume that G satisfies the following conditions:

- $\alpha)$ $G^{\pm 1} \in L_\infty(\partial\omega)$;
- $\beta)$ $\theta(t) \equiv \arg G(e^{it})$ is a piecewise Hölder function on $[-\pi, \pi]$ and $-\pi < s_1 < \dots < s_r < \pi$ are the corresponding points of discontinuity.

Consider the following functions:

$$X_1(z) = \exp \left\{ \frac{1}{4\pi} \int_{-\pi}^{\pi} \ln |G(e^{it})| \frac{e^{it} + z}{e^{it} - z} dt \right\},$$

$$X_2(z) = \exp \left\{ \frac{i}{4\pi} \int_{-\pi}^{\pi} \theta(t) \frac{e^{it} + z}{e^{it} - z} dt \right\},$$

which are analytic inside and outside ω , respectively. Let

$$Z_k(z) \equiv \begin{cases} X_k(z), & |z| < 1, \\ [X_k(z)]^{-1}, & |z| > 1, \quad k = 1, 2. \end{cases}$$

Denote

$$Z(z) \equiv Z_1(z) Z_2(z).$$

As usual, by $Z^\pm(\tau)$ we denote non-tangential boundary values of $Z(z)$ on $\partial\omega$ from inside and outside ω , respectively. Using Sochockii-Plemelj formulas, we directly obtain

$$|G(e^{it})| = \frac{Z_1^+(e^{it})}{Z_1^-(e^{it})}, \quad \exp i\theta(t) = \frac{Z_2^+(e^{it})}{Z_2^-(e^{it})}.$$

Consequently, we have

$$G(e^{it}) = \frac{Z^+(e^{it})}{Z^-(e^{it})}, \text{ a.e. on } (-\pi, \pi). \tag{3.2}$$

$Z(z)$ will be called the canonical solution of the problem (3.1). Taking into account the relation (3.2), from (3.1) we get

$$\frac{F^+(\tau)}{Z^+(\tau)} = \frac{F^-(\tau)}{Z^-(\tau)}, \text{ a.e. } \tau \in \partial\omega.$$

Assume

$$\Phi(z) \equiv \frac{F(z)}{Z(z)}.$$

Due to the fact that $Z(z)$ has neither zeros and nor poles at $z \notin \partial\omega$, it is clear that the functions $\Phi(z)$ and $F(z)$ have the same order at infinity. By definition of solution, we have $F \in H_1^\pm$. Let us express the function $\theta(t)$ as the sum

$$\theta(t) = \theta_0(t) + \theta_1(t),$$

where θ_0 is a continuous part, and θ_1 is a jump function defined by

$$\theta_1(-\pi + 0) = 0, \theta_1(t) = \sum_{-\pi < s_k < t} h_k + [\theta(t) - \theta(t - 0)],$$

where

$$h_k = \theta(s_k + 0) - \theta(s_k - 0), k = \overline{1, r},$$

are jumps of the function $\theta(\cdot)$ at s_k . Following [12], we denote

$$h_0 = h_0^{(1)} - h_0^{(0)},$$

where

$$h_0^{(1)} = \theta_1(-\pi + 0) - \theta_1(\pi - 0), h_0^{(0)} = \theta_0(-\pi - 0) - \theta_0(\pi - 0).$$

Let

$$U_0(t) \equiv \left\{ \sin \left| \frac{t - \pi}{2} \right| \right\}^{-\frac{h_0^{(0)}}{2\pi}} \exp \left\{ -\frac{1}{4\pi} \int_{-\pi}^{\pi} \theta_0(s) \operatorname{ctg} \frac{t - s}{2} ds \right\}.$$

As shown in [12], from condition $\alpha)$ it follows that the following inequality is true

$$\|Z_1^-(e^{it})\|_\infty^{\pm 1} < +\infty,$$

where $\|\cdot\|_\infty$ denotes the norm of $L_\infty(-\pi, \pi)$. Using the results of [12] again, we find that the boundary values of $Z_2(z)$ can be represented as

$$|Z_2^-(e^{it})| = U_0(t) U(t) \left| \sin \frac{t - \pi}{2} \right|^{-\frac{h_0}{2\pi}},$$

where

$$U(t) = \prod_{k=1}^r \left| \sin \frac{t - s_k}{2} \right|^{-\frac{h_k}{2\pi}}.$$

Using this notation, we can write the boundary values of $|Z(\tau)|$ as follows

$$|Z^-(e^{it})| = U_0(t) |Z_1^-(e^{it})| \left| \sin \frac{t - \pi}{2} \right|^{-\frac{h_0}{2\pi}} U(t).$$

According to the results of [12], the functions $U_0^{\pm 1}(t)$ belong to $L_p(-\pi, \pi)$, $\forall p \in (0, +\infty)$, and, moreover, for sufficiently small values of $\delta > 0$ we have $Z^{\pm 1}(z) \in H_\delta^+$. As a result, we obtain that the function $\Phi(z)$ belongs to the Hardy class H_μ^+ for some $\mu > 0$. It is absolutely obvious that if $\Phi^+(e^{it}) \in L_1(-\pi, \pi)$, then $\Phi \in H_1^+$. Since we have $\Phi^+(\tau) = \Phi^-(\tau)$ a.e. $\tau \in \partial\omega$, it is clear that it suffices to prove that $\Phi^-(e^{it})$ belongs to $L_1(-\pi, \pi)$. We have

$$|\Phi^-(e^{it})| = |F^-(e^{it})| |Z^-(e^{it})|^{-1}, \quad \text{a.e. } t \in (-\pi, \pi).$$

By definition of solution, $F^-(e^{it}) \in L_{p(\cdot); \vartheta}$. Therefore, if

$$|Z^-(e^{it})|^{-1} \in L_{q(\cdot); \vartheta^{-1}},$$

then $\Phi^-(\tau) \in L_1$, which follows directly from the generalized Hölder’s inequality in $L_{p(\cdot)}$. So assume that the following inequality is fulfilled

$$\int_{-\pi}^{\pi} |w(t) \vartheta^{-1}(t)|^{q(t)} dt < +\infty, \tag{3.3}$$

where

$$w(t) = \left| \sin \frac{t - \pi}{2} \right|^{\frac{h_0}{2\pi}} \prod_{k=1}^r \left| \sin \frac{t - s_k}{2} \right|^{\frac{h_k}{2\pi}}. \tag{3.4}$$

If the condition (3.3) holds, then $|Z^-(\tau)|^{-1}$ belongs to $L_{q(\cdot); \vartheta^{-1}}$, and, as a result, $\Phi^-(\tau) \in L_1$, so $F^- \in L_{p(\cdot); \vartheta}$ and $\Phi^-(e^{it}) = F^-(e^{it}) (Z^-(e^{it}))^{-1}$. Then, by Smirnov theorem, $\Phi \in H_1^\pm$, and, as a result, it follows from the uniqueness theorem that $\Phi(z)$ is a polynomial of order $k \leq m$ (because $\Phi^+(\tau) = \Phi^-(\tau)$ a.e. $\tau \in \partial\omega$), i.e. $\Phi(z) \equiv P_k(z)$, where $P_k(z)$ is a polynomial of order $k \leq m$. Thus

$$F(z) \equiv Z(z) P_k(z). \tag{3.5}$$

Assume that the inequalities

$$h_k < 2\pi, \quad k = \overline{0, r}, \tag{3.6}$$

are fulfilled. Then we have $Z^\pm(e^{it}) \in L_1(-\pi, \pi)$ and, again by Smirnov theorem, we obtain $Z(z) \in H_1^\pm$. So, $Z(\cdot)$ has a zero order at infinity, then it follows directly from (3.5) that $F \in H_1^\pm$. Let the following inequality also be fulfilled

$$\int_{-\pi}^{\pi} |w^{-1}(t) \vartheta(t)|^{p(t)} dt < +\infty. \tag{3.7}$$

Then from the expression for boundary values $Z^-(e^{it})$ it follows $Z^-(e^{it}) \in L_{p(\cdot); \vartheta}$, which means that $F^-(e^{it}) \in L_{p(\cdot); \vartheta}$. As a result, we obtain that if $m \geq 0$ and $k \leq m$, then it is true

$$(F^+(z); F^-(z)) \in H_{p(\cdot); \vartheta}^+ \times_m H_{p(\cdot); \vartheta}^-.$$

Thus, if $m \geq 0$, then the expression (3.5) is the general solution of the homogeneous problem (3.1) in the classes $H_{p(\cdot); \vartheta}^+ \times_m H_{p(\cdot); \vartheta}^-$, where $P_m(\cdot)$ is an arbitrary polynomial of order $k \leq m$. So the following theorem is true.

Theorem 3.1. *Let $p(\cdot) \in WL \wedge p^- > 1$ and the coefficient $G(\tau)$ of the problem (3.1) satisfy the conditions α) and β). Suppose that the jumps of functions $\theta(t) \equiv \arg G(e^{it})$ satisfy the relations (3.3), (3.6), (3.7), where $w(t)$ is defined by (3.4). Then, under $m \geq 0$, the general solution of the homogeneous problem (3.1) in classes $H_{p(\cdot); \vartheta}^+ \times_m H_{p(\cdot); \vartheta}^-$ can be represented in the form of (3.5), where $Z(z)$ is a canonical solution, and $P_k(z)$ is an arbitrary polynomial of order $k \leq m$. If $m \leq -1$, then under the conditions (3.3), (3.6), (3.7), the problem (3.1) in $H_{p(\cdot); \vartheta}^+ \times_m H_{p(\cdot); \vartheta}^-$ has only trivial, i.e. zero solution.*

4. General solution of nonhomogeneous problem

Consider an nonhomogeneous Riemann problem

$$F^+(\tau) - G(\tau) F^-(\tau) = g(\tau), \tau \in \partial\omega, \tag{4.1}$$

where $g \in L_{p(\cdot); \vartheta}$ is a given function. By the solution of problem (4.1) we mean a pair

$$(F^+(z); F^-(z)) \in H_{p(\cdot); \vartheta}^+ \times_m H_{p(\cdot); \vartheta}^-,$$

for which the boundary values $F^\pm(\tau)$ satisfy the relation (4.1) a.e. on $\partial\omega$. It is clear that the general solution of (4.1) can be represented as

$$F(z) = F_0(z) + F_1(z),$$

where $F_0(z)$ is a general solution of the corresponding homogeneous problem (3.1), and $F_1(z)$ is some particular solution of the nonhomogeneous problem (4.1). We will construct a particular solution of (4.1). Let $Z(z)$ be a canonical solution of (3.1). Consider the piecewise analytic function

$$F_1(z) \equiv \frac{Z(z)}{2\pi} \int_{-\pi}^{\pi} \frac{g(e^{i\sigma})}{Z^+(e^{i\sigma})} \frac{d\sigma}{1 - ze^{-i\sigma}}. \tag{4.2}$$

Applying Sokhotskii-Plemelj formulas to (4.2), we obtain that the boundary values $F_1^\pm(\tau)$ satisfy (4.1) a.e. on $\partial\omega$. We have

$$F_1^+(e^{it}) = \frac{1}{2}g(e^{it}) + \frac{Z^+(e^{it})}{2\pi} \int_{-\pi}^{\pi} \frac{g(e^{i\sigma})}{Z^+(e^{i\sigma})} \frac{d\sigma}{1 - e^{i(t-\sigma)}}.$$

Let

$$f(\sigma) = g(e^{i\sigma}) [Z^+(e^{i\sigma})]^{-1}.$$

Consequently

$$\tilde{F}_1^+(t) = \frac{1}{2}f(t) + \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{f(\sigma) d\sigma}{1 - e^{i(t-\sigma)}},$$

where

$$\tilde{F}_1(t) \equiv F_1^+(e^{it}) [Z^+(e^{it})]^{-1}.$$

It is absolutely clear that $f(\cdot)$ belongs to the weighted space $L_{p(\cdot); \nu}$, where

$$\nu(t) = w^{-1}(t) \vartheta(t).$$

It is known that the singular operator

$$S(f) = \int_{-\pi}^{\pi} \frac{f(\sigma) d\sigma}{1 - e^{i(t-\sigma)}},$$

is bounded in $L_{p(\cdot); \nu}$, $1 < p^- \leq p^+ < +\infty$, $p \in WL$, if and only if the weight $\nu(\cdot)$ satisfies the Muckenhoupt condition $A_{p(\cdot)}$, i.e, $\nu \in A_{p(\cdot)}$ (see, for example, [11, 16]). Consequently, if $\nu \in A_{p(\cdot)}$, then $\tilde{F}_1 \in L_{p(\cdot); \nu}$. It is not difficult to see that

$$\tilde{F}_1 \in L_{p(\cdot); \nu} \Leftrightarrow F_1 \in L_{p(\cdot); \vartheta}.$$

The condition $\nu \in A_{p(\cdot)}$ implies $\nu \in L_1$ and $\nu^{-1} \in L_{q(\cdot)}$. Then from Hölder's inequality it follows

$$\int_{-\pi}^{\pi} |f(t)| dt = \int_{-\pi}^{\pi} |f(t)| \nu(t) \nu^{-1}(t) dt \leq c_{p(\cdot)} \| |f(\cdot)| \nu(\cdot) \|_{p(\cdot)} \| \nu^{-1}(\cdot) \|_{q(\cdot)}.$$

Thus, f belongs to L_1 , and, as a result, the Cauchy-Lebesgue type integral

$$\int_{-\pi}^{\pi} \frac{f(\sigma) d\sigma}{1 - ze^{-i\sigma}},$$

belongs to the Hardy class $H_{p_1}^{\pm}$, for $\forall p_1 : 0 < p_1 < 1$, (see, for example, [18]). From expression for $F_1(z)$ it directly follows that it belongs to the space H_{μ}^+ for sufficiently small $\mu > 0$. As we have already established, $F_1^+(e^{it}) \in L_{p(\cdot); \vartheta}$. Then it is evident that if $\vartheta^{-1} \in L_{q(\cdot)}$, then $F_1^+(e^{it}) \in L_1$. Then, by Smirnov theorem [18], the function $F_1^+(z)$ belongs to the class H_1^+ . As a result, we obtain $F_1^+(z) \in H_{p(\cdot); \vartheta}^+$. Consider $F_1^-(z)$, and let all the above conditions be satisfied. Similar reasoning yields $F_1^-(z) \in_{-1} H_{p(\cdot); \vartheta}^-$. It is absolutely clear that $F_1^-(\infty) = 0$. Therefore

$$F_1^-(z) \in_m H_{p(\cdot); \vartheta}^-, \forall m \geq -1.$$

Let $m < -1$. In this case, in order for the inclusion $F_1^-(z) \in_m H_{p(\cdot); \vartheta}^-$, to be valid, it is necessary that the following orthogonality conditions hold

$$\int_{-\pi}^{\pi} \frac{g(e^{i\sigma})}{Z^+(e^{i\sigma})} e^{in\sigma} d\sigma = 0, \quad n = \overline{1, -m}. \tag{4.3}$$

Thus, under these conditions, the expression (4.2) is a particular solution of (4.1) in classes $H_{p(\cdot); \vartheta}^+ \times_m H_{p(\cdot); \vartheta}^-$. If the conditions (3.3), (3.6) and (3.7) are fulfilled, then the general solution of the corresponding homogeneous problem has the form (3.5), where $Z(z)$ is a canonical solution, and $P_m(z)$ is an arbitrary polynomial of order $\leq m$. For $m \leq -1$ it is clear that $P_m(z) \equiv 0$. In this case, the problem (4.2) is uniquely solvable. So we have the following

Theorem 4.1. *Let the coefficient $G(\tau)$ of the problem (4.1) satisfy the conditions $\alpha); \beta)$. Assume that the jumps of the function $\theta(t) \equiv \arg G(e^{it})$ satisfy the relations (3.3), (3.6), (3.7), where the weight function $w(t)$ is defined by the expression (3.4). Let $\nu \in A_{p(\cdot)}, p(\cdot) \in WL \wedge p^- > 1$, where $\nu(t) = w^{-1}(t)\vartheta(t)$ and $\vartheta^{-1} \in L_{q(\cdot)}$. Then the nonhomogeneous Riemann problem (4.1) is solvable in classes $H_{p(\cdot); \vartheta}^+ \times_m H_{p(\cdot); \vartheta}^-$, if the orthogonality conditions (4.3) hold. For $m \geq 0$ the general solution of (4.1) can be represented as*

$$F(z) = Z(z)P_m(z) + F_1(z), \tag{4.4}$$

where $Z(z)$ is a canonical solution of the homogeneous problem, $P_m(z)$ is an arbitrary polynomial of order $\leq m$, and $F_1(z)$ is defined by the expression (4.2). Moreover, for $m \leq -1$ the problem (4.1) is uniquely solvable, and for $m = -1$ it has a solution for $\forall g \in L_{p(\cdot); \vartheta}$.

5. Special case

Consider the special case

$$\vartheta(t) = \prod_{k=1}^m |t - t_k|^{\alpha_k}, \tag{5.1}$$

where $\{t_k\}_1^m \subset (-\pi, \pi)$ are different points. For simplicity, we assume

$$\{s_k\}_1^r \cap \{t_k\}_1^m = \emptyset. \tag{5.2}$$

In this case, the weight $\nu(t)$ has the following form

$$\nu(t) \equiv \left| \sin \frac{t - \pi}{2} \right|^{-\frac{h_0}{2\pi}} \prod_{k=1}^r \left| \sin \frac{t - s_k}{2} \right|^{-\frac{h_k}{2\pi}} \prod_{k=1}^m |t - t_k|^{\alpha_k}.$$

It is known that $\nu \in A_{p(\cdot)}$ if and only if (see, for example [16])

$$\begin{aligned} -\frac{1}{p(s_k)} < -\frac{h_k}{2\pi} < \frac{1}{q(s_k)}, \quad k = \overline{0, r}; \\ -\frac{1}{p(t_k)} < \alpha_k < \frac{1}{q(t_k)}, \quad k = \overline{1, m}. \end{aligned} \tag{5.3}$$

Consequently

$$-\frac{1}{q(s_k)} < \frac{h_k}{2\pi} < \frac{1}{p(s_k)}, \quad k = \overline{0, r}. \tag{5.4}$$

From this relation it directly follows that the conditions (3.3), (3.6), and (3.7) are fulfilled. It is obvious that the relation $\vartheta \in L_1$ also holds. As a result, from Theorem 4.1 we obtain the following

Corollary 5.1. *Let $p(\cdot) \in WL \wedge p^- > 1$ and the coefficient $G(\tau)$ satisfy the conditions $\alpha); \beta)$ and let (5.2)-(5.4) hold. Then the general solution of the Riemann problem (4.1) in classes $H_{p(\cdot); \vartheta}^+ \times_m H_{p(\cdot); \vartheta}^-$ is given by (4.4), where the weight ϑ has the form (5.1). For $m < 0$, the orthogonality conditions (4.3) are necessary for solvability. Then the problem is uniquely solvable and $P_m(z) \equiv 0$. For $m \geq 0$, the*

homogeneous problem has m linearly independent solutions, while the nonhomogeneous problem is solvable for $\forall g \in L_{p(\cdot); \vartheta}$. For $m = -1$, the nonhomogeneous problem is uniquely solvable for $\forall g \in L_{p(\cdot); \vartheta}$.

Acknowledgement

This research was supported by the Azerbaijan National Academy of Sciences under the program "Approximation by neural networks and some problems of frames".

References

- [1] K.I. Babenko, On conjugated functions, *DAN SSSR*, **62(2)** (1948), 157–160.
- [2] B.T. Bilalov, Basicity of Some Systems of Exponents, Cosines and Sines, *Diff. Uravneniya*, **26(1)** (1990), 10–16.
- [3] B.T. Bilalov, The basis properties of some systems of exponential functions, cosines, and sines, *Sibirsk. Mat. Zh.*, **45(2)** (2004), 264–273.
- [4] B.T. Bilalov, The basis properties of power systems in L_p , *Sibirsk. Mat. Zh.*, **47(1)** (2006), 25–36.
- [5] B.T. Bilalov, A system of exponential functions with shift and the Kostyuchenko problem, *Siberian Mathematical Journal*, **50(2)** (2009), 279–288.
- [6] B.T. Bilalov, On solution of the Kostyuchenko problem, *Siberian Mathematical Journal*, **53(3)** (2012), 509–526.
- [7] B.T. Bilalov and F.A. Guliyeva, A completeness criterion for a double system of powers with degenerate coefficients, *Siberian Mathematical Journal*, **54(3)** (2013), 536–543.
- [8] B.T. Bilalov and Z.G. Huseynov, Basicity of a system of exponents with a piece-wise linear phase in variable spaces, *Mediterr. J. Math.*, **9(3)** (2012), 487–498.
- [9] B.T. Bilalov and Z.G. Huseynov, On the basicity from exponents in Lebesgue spaces with variable exponent, *TWMS J. Pure Appl. Math.*, **1(1)** (2010), 14–23.
- [10] B.T. Bilalov, F.I. Mamedov and R.A. Bandaliev, On classes of harmonic functions with variable exponent, *Dokl NAN Azerb.*, **LXIII(5)** (2007), 16–21.
- [11] D.V. Cruz-Uribe and A. Fiorenza, *Variable Lebesgue Spaces, Foundations and Harmonic Analysis*, Springer, 2013.
- [12] I.I. Danilyuk, *Irregular Boundary Value Problems on the Plane*, Nauka, Moscow, 1975.
- [13] F.D. Gakhov, *Boundary Value Problems*, (Russian), 3rd Ed. Moscow, Nauka, 1977. (Transl. of 2nd Ed. Oxford, Pergamon Press, 1966, 561 p.).
- [14] S.B. Klimentov, Riemann and Hilbert boundary value problems in BMO classes for holomorphic functions, *Vladikavkaz. Mat. Zh.*, **12(4)** (2010), 28–38.
- [15] V. Kokilashvili and V. Paatashvili, The Riemann-Hilbert problem in a domain with piecewise smooth boundaries in weight classes of Cauchy type integrals with a density from variable exponent Lebesgue spaces, *Georgian Math. J.*, **16(4)** (2009), 737–755.
- [16] V. Kokilashvili and S. Samko, Singular integrals in weighted Lebesgue spaces with variable exponent, *Georgian Math. J.*, **10(1)** (2003), 145–156.
- [17] V. Kokilashvili, V. Paatashvili and S. Samko, Boundary value problems for analytic functions in the class of Cauchy type integrals with density in $L^{p(\cdot)}$, *Bound. Value Probl.*, **1-2** (2005), 43–71.
- [18] P. Koosis, *Introduction to H^p spaces*, M. Mir, 364 p. (in Russian)

- [19] O. Kovacik and J. Rakosnik, On spaces $L^{p(\cdot)}$ and $W^{k,p(\cdot)}$, *Czechoslovak Math.I.*, **41(116)** (1991), 592–618
- [20] G.S. Litvinchuk, *Boundary value problems and singular integral equations with shift*, Nauka, Moscow, 1977.
- [21] G. Manjavidze and N. Manjavidze, Boundary value problems for analytic and generalized analytic functions, *Journal of Mathematical Sciences*, **160(6)** (2009), 745–821.
- [22] Z. Meshveliani, The Riemann-Hilbert problem in weighted Smirnov classes of analytic functions, *Proc. A. Razmadze Math. Inst.*, **137** (2005), 65–86.
- [23] T.R. Muradov, On Riemann Boundary Value Problem in Hardy Classes with Variable Summability Exponent, *Int. Journal of Math. Analysis*, **615** (2012), 743–751.
- [24] N.I. Muskhelishvili, *Singular Integral Equations*, 2nd ed. Groningen, Noordhoff, 1968.
- [25] T.I. Najafov and N.P. Nasibova, On the Noetherness of the Riemann Problem in Generalized Weighted Hardy Classes, *Azerbaijan Journal of Mathematics*, **5(2)** (2015), 109–125.
- [26] I.I. Sharapudinov, On the topology of space $L^{p(x)}([0,1])$, *Math. Notes*, **26(4)** (1979), 613 – 632
- [27] N.P. Vekua, *Singular integrals, singular equations and their application*, 2nd Izd., Mosco, 1970.
- [28] F. Xianling and Z. Dun, On the spaces $L^{p(x)}(\Omega)$ and $L^{m,p(\cdot)}(\Omega)$, *Journal of Math. Anal. and Appl.*, **263** (2001), 424–446.

Bilal T. Bilalov

Institute of Mathematics and Mechanics of NAS of Azerbaijan, B.Vahabzadeh str. 9, AZ1141, Baku, Azerbaijan

E-mail address: b.bilalov@mail.ru

Ali A. Huseynli

Department of Mathematics, Khazar University, Mehseti str. 41, AZ1096, Baku, Azerbaijan

Institute of Mathematics and Mechanics of NAS of Azerbaijan, B.Vahabzadeh str. 9, AZ1141, Baku, Azerbaijan.

E-mail address: alihuseynli@gmail.com

Fidan Sh. Seyidova

Ganja State University

E-mail address: fidanseidova207@gmail.com

Received: March 11, 2017; Revised: July 2, 2017; Accepted: July 20, 2017