



LJMU Research Online

Uusimaa, J, Evans, J, Smith, C, Butterworth, A, Craig, K, Ashley, N, Liao, C, Carver, J, Diot, A, Macleod, L, Hargreaves, IP, Al-Hussaini, A, Faqeih, E, Asery, A, Al Balwi, M, Eyaid, W, Al-Sunaid, A, Kelly, D, van Mourik, I, Ball, S, Jarvis, J, Mulay, A, Hadzic, N, Samyn, M, Baker, A, Rahman, S, Stewart, H, Morris, AAM, Seller, A, Fratter, C, Taylor, RW and Poulton, J

Clinical, biochemical, cellular and molecular characterization of mitochondrial DNA depletion syndrome due to novel mutations in the MPV17 gene

<http://researchonline.ljmu.ac.uk/id/eprint/13683/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Uusimaa, J, Evans, J, Smith, C, Butterworth, A, Craig, K, Ashley, N, Liao, C, Carver, J, Diot, A, Macleod, L, Hargreaves, IP, Al-Hussaini, A, Faqeih, E, Asery, A, Al Balwi, M, Eyaid, W, Al-Sunaid, A, Kelly, D, van Mourik, I, Ball, S, Jarvis, J, Mulay, A, Hadzic, N, Samyn, M, Baker, A, Rahman, S, Stewart, H.

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

<http://researchonline.ljmu.ac.uk/>

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

ARTICLE

Clinical, biochemical, cellular and molecular characterization of mitochondrial DNA depletion syndrome due to novel mutations in the *MPV17* gene

Johanna Uusimaa^{1,2,17}, Julie Evans^{3,17}, Conrad Smith³, Anna Butterworth⁴, Kate Craig⁴, Neil Ashley¹, Chunyan Liao¹, Janet Carver¹, Alan Diot¹, Lorna Macleod¹, Iain Hargreaves⁵, Abdulrahman Al-Hussaini⁶, Eissa Faqih⁶, Ali Asery⁶, Mohammed Al Balwi⁷, Wafaa Eyaid⁸, Areej Al-Sunaid⁸, Deirdre Kelly⁹, Indra van Mourik⁹, Sarah Ball¹⁰, Joanna Jarvis¹¹, Arundhati Mulay¹², Nedim Hadzic¹³, Marianne Samyn¹³, Alastair Baker¹³, Shamima Rahman¹⁴, Helen Stewart¹⁵, Andrew AM Morris¹⁶, Anneke Seller³, Carl Fratter³, Robert W Taylor⁴ and Joanna Poulton^{*,1}

Mitochondrial DNA (mtDNA) depletion syndromes (MDS) are severe autosomal recessive disorders associated with decreased mtDNA copy number in clinically affected tissues. The hepatocerebral form (mtDNA depletion in liver and brain) has been associated with mutations in the *POLG*, *PEO1* (Twinkle), *DGUOK* and *MPV17* genes, the latter encoding a mitochondrial inner membrane protein of unknown function. The aims of this study were to clarify further the clinical, biochemical, cellular and molecular genetic features associated with MDS due to *MPV17* gene mutations. We identified 12 pathogenic mutations in the *MPV17* gene, of which 11 are novel, in 17 patients from 12 families. All patients manifested liver disease. Poor feeding, hypoglycaemia, raised serum lactate, hypotonia and faltering growth were common presenting features. mtDNA depletion in liver was demonstrated in all seven cases where liver tissue was available. Mosaic mtDNA depletion was found in primary fibroblasts by PicoGreen staining. These results confirm that *MPV17* mutations are an important cause of hepatocerebral mtDNA depletion syndrome, and provide the first demonstration of mosaic mtDNA depletion in human *MPV17* mutant fibroblast cultures. We found that a severe clinical phenotype was associated with profound tissue-specific mtDNA depletion in liver, and, in some cases, mosaic mtDNA depletion in fibroblasts.

European Journal of Human Genetics (2014) 22, 184–191; doi:10.1038/ejhg.2013.112; published online 29 May 2013

Keywords: *MPV17*; mitochondrial DNA depletion; hepatocerebral disease; neonatal liver disease

INTRODUCTION

During the last decade, an increasing number of nuclear genetic defects have been identified causing mitochondrial dysfunction either through the accumulation of mitochondrial DNA (mtDNA) deletions (multiple deletions) or through a reduction in mtDNA copy number causing mtDNA depletion syndromes (MDS), the latter of which are autosomal recessive diseases characterized by a severe, tissue-specific decrease of mtDNA copy number. At least 12 nuclear genes have been found to be involved in mtDNA maintenance, including *POLG*, *POLG2*, *PEO1*, *SLC25A4*, *TYMP*, *DGUOK*, *TK2*, *SUCLA2*, *SUCLG1*, *MPV17*, *OPA1* and *RRM2B* genes.¹ MDS are caused by recessive defects in proteins directly involved in mtDNA replication or in proteins that affect the availability of deoxyribonucleoside

triphosphates for mtDNA synthesis. Clinical presentations of MDS include early-onset hepatocerebral disease overlapping with Alpers-Huttenlocher syndrome (henceforth 'Alpers'), isolated myopathy, encephalomyopathy and mitochondrial neurogastrointestinal encephalomyopathy syndrome. MDS are classified as myopathic, encephalomyopathic or hepatocerebral forms, of which the latter group has been associated with mutations in *POLG*, *PEO1* (Twinkle), *DGUOK* and *MPV17* genes.²

The human *MPV17* gene is located on chromosome 2p21-23, comprising eight exons encoding 176 amino acids. It is expressed in human pancreas, kidney, muscle, liver, lung, placenta, brain and heart. *MPV17* encodes a mitochondrial inner membrane protein, which has a so far largely unknown role in mtDNA maintenance. Human

¹Nuffield Department of Obstetrics and Gynaecology, University of Oxford The Women's Centre, Oxford, UK; ²Institute of Clinical Medicine/Paediatrics, University of Oulu, Clinical Research Center, Oulu University Hospital, Oulu, Finland; ³Oxford Medical Genetics Laboratories, Oxford University Hospitals NHS Trust, Oxford, UK; ⁴Wellcome Trust Centre for Mitochondrial Research, Newcastle University, UK; ⁵Neurometabolic Unit, National Hospital for Neurology and Neurosurgery, London, UK; ⁶Department of Pediatrics, King Fahad Medical City, Riyadh, Saudi Arabia; ⁷Department of Pathology and Laboratory Medicine, King Abdulaziz Medical City, Riyadh, Saudi Arabia; ⁸Department of Pediatrics, King Abdulaziz Medical City, King Saud Bin Abdulaziz University for Health & Science, Riyadh, Saudi Arabia; ⁹Paediatric Liver Unit, Birmingham Children's Hospital, Birmingham, UK; ¹⁰Department of Newborn Screening and Biochemical Genetics, Birmingham Children's Hospital, Birmingham, UK; ¹¹Clinical Genetics Unit, Birmingham Women's Hospital, Birmingham, UK; ¹²Maternity Services, Manor Hospital, Walsall, UK; ¹³Paediatric Liver Centre, King's College Hospital, London, UK; ¹⁴Mitochondrial Research Group, UCL Institute of Child Health, London, UK; ¹⁵Department of Clinical Genetics, Oxford University Hospitals NHS Trust, Oxford, UK; ¹⁶Willink Biochemical Genetics Unit, St Mary's Hospital, Manchester, UK

¹⁷These authors contributed equally to this work.

*Correspondence: Professor J Poulton, Nuffield Department of Obstetrics and Gynaecology, The Women's Centre, John Radcliffe Hospital, Headington, Oxford OX3 9DU, UK. Tel: +44 01865 221067; Fax +44 01865 769141; E-mail: Joanna.Poulton@obs-gyn.ox.ac.uk

Received 9 November 2012; revised 15 April 2013; accepted 17 April 2013; published online 29 May 2013

MPV17 is the orthologue of the mouse kidney disease gene, *Mpv17*. Loss of function has been shown to cause hepatocerebral MDS with oxidative phosphorylation failure and mtDNA depletion both in affected individuals and in *Mpv17*^{-/-} mice.³ To date, MDS caused by *MPV17* mutations has been reported in 32 patients with the clinical manifestations including early progressive liver failure, neurological abnormalities, hypoglycaemia and raised blood lactate.^{4–15} Recently, *MPV17* mutations have also been associated with autosomal recessive adult-onset neuropathy and leukoencephalopathy with multiple mtDNA deletions in skeletal muscle.¹⁶ Thus, as for *POLG*, *RRM2B* and *TK2*, *MPV17* mutations can lead to recessive MDS or recessive multiple mtDNA deletion disorders.

The aim of this study was to clarify further the clinical, biochemical, cellular and molecular features associated with MDS due to *MPV17* gene mutations. We report 17 cases from 12 families with 11 novel *MPV17* mutations. All the patients presented with early-onset liver disease but only some patients manifested neurological dysfunction. Furthermore, we demonstrate that primary fibroblast cultures from patients with *MPV17* mutations may exhibit a mosaic mtDNA depletion.

SUBJECTS AND METHODS

We studied 70 unrelated probands with suspected hepatocerebral MDS who had been referred to Mitochondrial Diagnostic Centres at Oxford, Newcastle or London for clinical assessment, histological, biochemical and/or molecular genetic analyses. This study was approved and performed under the ethical guidelines issued by each institution for clinical studies, with written informed consent obtained for all subjects.

Molecular genetics

Total genomic DNA was isolated from blood leukocytes, skeletal muscle and liver samples by standard methods. Long-range PCR amplification of 13.8 kb mtDNA was undertaken to screen for mtDNA deletions in the available muscle and liver DNA samples, using primers previously described.¹⁷ mtDNA copy number relative to nDNA levels in muscle and/or liver DNA was estimated and compared with age-matched normal controls^{18,19} by real-time quantitative PCR as described previously,²⁰ except the nuclear probe was labelled with Vic at the 5' end and the assays were carried out simultaneously using a PE7500 real-time PCR instrument (Applied Biosystems, Life Technologies, Carlsbad, CA, USA). mtDNA copy number <30% compared with age-matched controls was classed as mtDNA depletion, 30–50% as borderline low, and >50% as normal. The entire coding and flanking intronic regions of *MPV17* were amplified by PCR from genomic DNA and sequenced by fluorescent dideoxy sequencing (Applied Biosystems Big Dye Terminator v3.1 kit) and capillary electrophoresis (Applied Biosystems 3730). Results were compared with Genbank reference sequences NM_002437.4 and NG_008075.1, and mutations described in accordance with HGVS nomenclature guidelines. *MPV17* exon copy number (exons 1–8) was assessed by MLPA (kit P089-A1; MRC-Holland, Amsterdam, The Netherlands) in patients 16 and 17 (family 12) to confirm the deletion of multiple exons. Where available, parental blood DNA samples were analysed for the familial mutation(s) by sequencing of the appropriate exon, or by MLPA for family 12.

Muscle histology, histochemistry and biochemistry

Tissue samples from muscle and liver were obtained according to standard procedures. Muscle and liver histology and histochemistry were performed by standard methods. The activities of mitochondrial respiratory chain enzyme complexes were determined from skeletal muscle and liver biopsy samples and cultured skin fibroblasts, as previously described.^{21,22}

Characterization of the cellular mtDNA depletion phenotype by PicoGreen staining of fibroblasts

Cell cultures were analysed using the fluorescent stains PicoGreen (Molecular Probes, Eugene, OR, USA) (dsDNA intercalator) and tetramethylrhodamine

methyl ester (TMRM; reflects mitochondrial membrane potential). Staining with PicoGreen is a simple, semi-quantitative method for visualizing mtDNA *in situ* within living cells as described.²³ mtDNA nucleoids were further quantified using the IN Cell 1000 analyser (GE Healthcare Life Science, Little Chalfont, UK). Control and patients' fibroblasts were stained with PicoGreen and 30 nM TMRM as above in clear Optimum media (Gibco, Life Technologies). Raw images were acquired and binarized to quantify mtDNA nucleoids per cell. Nucleoids were defined as PicoGreen punctae present in the cytoplasm and colocalizing with mitochondria.

RESULTS

Analysis of the *MPV17* gene

We identified pathogenic mutations in the *MPV17* gene in 12 out of 70 probands screened representing 17% of our undiagnosed cohort of children with suspected hepatocerebral MDS. Ten probands harboured homozygous mutations, and two had compound heterozygous mutations. A familial homozygous mutation was also identified in further affected siblings from families 4, 5, 9 and 12, resulting in a total of 17 patients from 12 families (Table 1). Eleven of the 12 mutations we identified are previously unpublished, and as such considered to be novel: c.62T>G (p.Leu21Arg), c.67G>C (p.Ala23-Pro), c.107A>C (p.Gln36Pro), c.121C>T (p.Arg41Trp), c.130C>T (p.Gln44*), c.135delA (p.Glu45Aspfs*8), c.191C>G (p.Pro64Arg), c.278A>C (p.Gln93Pro), c.279 + 1G>T, c.461 + 1G>C and a deletion including exons 3–8 of *MPV17* (Figures 1 and 2).

Five of the 11 novel mutations are predicted to be protein truncating: c.279 + 1G>T and c.461 + 1G>C alter a consensus splice site; c.130C>T is a nonsense mutation (p.Gln44*); c.135delA results in a frameshift (p.Glu45Aspfs*8) and one mutation is a large deletion of exons 3–8. The remaining six novel mutations are missense changes, which are scattered throughout the protein (Figure 2), and alter amino acids, which are highly conserved across vertebrates (Supplementary Figure 1).

Homozygosity for the *MPV17* mutation in patients from families 1, 3–5, 7–10 and 12 was confirmed by parental testing, which demonstrated all parents to be heterozygous carriers. Parental samples were not available for family 11. For patients 2 (family 2) and 9 (family 6), at least one parent was found to be a heterozygous carrier in each case, thereby confirming compound heterozygosity.

Molecular genetic studies on mtDNA from muscle and tissue samples

Molecular genetic analyses did not reveal any significant large-scale mtDNA rearrangements in any of the available muscle or liver DNA samples. mtDNA copy number was compared as previously^{18,19} with controls matched for age but not ethnicity. Severe mtDNA depletion in liver was demonstrated for five out of seven cases (5–14% of control mtDNA) where liver tissue was available (patients 2, 3, 11, 14 and 16). Patient 9 had clear but less severe mtDNA depletion in liver (21% of controls), whereas mtDNA copy number was borderline low (40% of controls) in the remaining case (patient 4). Mean mtDNA copy number of all seven cases was 16%. In contrast, mtDNA copy number was highly variable across the six cases where muscle tissue was available (10–100% of controls, mean 46%; Table 1).

Clinical and laboratory features of patients with *MPV17* mutations

The 17 individuals in our cohort came from 12 families, 6 being male and 11 female, from different ethnic populations including Asian, Middle-Eastern and Caucasian families. All patients manifested with liver disease with abnormal liver function tests. Poor feeding, hypoglycaemia, raised plasma lactate, hypotonia and faltering growth

Table 1 Clinical, biochemical, cellular and molecular findings in 17 patients with MPV17 mutations

Family	Sex	Consanguinity (family history)	Clinical information	Age at onset	Age at referral	Deceased/alive, age at death/cure	MPV17 mutations	mtDNA copy number in liver	mtDNA copy number in muscle	PicoGreen analysis of fibroblasts	Muscle/liver histology	Muscle/liver/fibroblasts respiratory chain activities or histochemistry
1	F	Y (N)	Hypotonia, progressive liver disease, coagulopathy, hypoglycaemia, developmental delay, raised blood lactate	4.5 mo	5 mo	Decd, 6.5 mo	c.62T>G (p.Leu21Arg) hom	ND	ND	ND	ND	ND
2	F	N (N)	Poor feeding, FG, progressive liver disease, hypoglycaemia, raised blood lactate, delayed development, vomiting	Birth	2 mo	Decd, 12 mo	c.67G>C (p.Ala23Pro) & c.135delA (p.Glu45Aspfs*8)	14%	51%	Normal	Mus: Normal; Liv: micro-and macro-vesicular change in most hepatocytes, fibrosis, inflammation	Mus: Normal; Liv: CIV ↓
3	F	Y (Y ^a)	IUGR, feeding problems, hypoglycaemia, raised tyrosine within days of birth, raised blood lactate, FG, hypotonia, progressive liver failure, encephalopathy	Birth	2 mo	Decd, 4 mo	c.107A>C (p.Gln36Pro) hom	5%	100%	Mosaic depletion	Liv: severe periportal loss of hepatocytes and stromal collapse	Mus: Normal; Fib: Normal
4	F	Y (Y, sibs P5 & P6)	Progressive neurological deterioration with peripheral neuropathy, progressive liver disease	5 yr	5 yr	Alive, 8.5 yr	c.121C>T (p.Arg41Trp) hom	40%	46%	Normal/minor changes	Mus: Normal; Liv: microvesicular steatosis and focal fibrosis	Mus: Normal; Liv: Normal
5	F	Y (Y, sibs P4 & P6)	Progressive liver failure, FG, corneal scarring, progressive neurological deterioration	Birth	9 mo	Decd, 4 yr 3 mo	c.121C>T (p.Arg41Trp) hom	ND	38%	N.D.	Mus: Normal Liv: cirrhosis	ND
6	M	Y (Y, sibs P4 & P5)	Progressive liver failure, encephalopathy, liver transplant at 15 mo, died 15 mo later following neurological and renal deterioration.	Birth	9 mo	Decd, 2 yr 6 mo	ND	ND	ND	ND	Liv: Giant cell hepatitis, haemorrhagic necrosis, focal fatty change	ND
7	M	Y (Y ^b , sib P8)	Poor feeding, hypoglycaemia, FG, raised blood lactate, encephalopathy, progressive liver failure	3.5 mo	3.5 mo	Decd, 4 mo	c.130C>T (p.Gln44*) hom	ND	ND	ND	Mus: Fatty infiltration; Liv: fatty infiltration	Mus: CI ↓, CII + III ↓, CIV ↓
8	F	Y (Y ^b , sib P7)	Tachycnea, hypoglycaemia, raised blood lactate within days of birth, poor weight gain, progressive liver failure	Birth	Birth	Decd, 7.5 mo	c.130C>T (p.Gln44*) hom	ND	ND	Normal/minor changes	ND	ND
9	M	N (N)	FG, liver disease, liver transplant at 3 yr, progressive demyelinating peripheral neuropathy, hypoparathyroidism, severe growth hormone deficiency,	5 mo	5 mo	Alive, 11.5 yr	c.191C>G (p.Pro64Arg) & c.293C>T (p.Pro98Leu)	21%	80%	Normal	Mus: minor fatty changes; Liv: steatosis, abundant mitochondria with mild pleomorphism	Mus: CI ↓, CIV ↓ Liv: CIV ↓
10	M	Y (N)	Gastrostomy Neonatal jaundice, FG, hypotonia, microcephaly, progressive liver disease, raised blood lactate, retinal pigmentation	2 mo	2 mo	Decd, 5 mo	c.278A>C (p.Gln93Pro) hom	ND	ND	ND	Liv: distension of hepatocytes, few periportal glycogen nuclei, fibrosis, microvesicular steatosis and portal inflammation	ND
11	F	Y (Y ^c)	Poor feeding, mild hypotonia, jaundice; liver dysfunction, coagulopathy, raised blood lactate	2 mo	3.5 mo	Decd, <16 mo	c.278A>C (p.Gln93Pro) hom	11%	ND	Mosaic depletion	Liv: cholestasis with scattered giant cell hepatocytes and extensive portal - portal bridging fibrosis	ND
12	F	Y (Y, sib P13)	Poor feeding, FG, hypotonia, subtle facial dysmorphism, hypoglycaemia, cholestatic jaundice, central hypocortisolism, progressive liver disease	2 mo	5 mo	Decd, 8 mo	c.278A>C (p.Gln93Pro) hom	ND	10%	ND	ND	Mus: CI ↓, CIII ↓, CIV ↓
13	F	Y (Y, sib P12)	Neonatal jaundice, FG, hypotonia, recurrent vomiting, progressive liver disease, raised blood lactate, retinal pigmentation	Birth	5 mo	Alive, 6 mo	c.278A>C (p.Gln93Pro) hom	ND	ND	ND	Liv: large droplet steatosis and microsteatosis, mild iron deposition in Kupffer cells	ND
14	M	Y (Y ^d)	FG, (lev delay, hypotonia, myopathy, liver disease, cholestasis; raised blood lactate	4 mo	5 mo	Alive, 14 mo	c.279+1G>T hom	5%	ND	ND	Liv: micro- and macro-vesicular steatosis with canalicular cholestasis, portal fibrosis and septae formation	ND
15	F	Y (N)	Conjugated jaundice, hypothyroidism, hypotonia, FG, developmental delay and dysmorphism	2.5 mo	13 mo	Decd, 3 yr	c.461+1G>C hom	ND	25%	Mosaic depletion	Liv: Patchy hepatocellular oncocytosis and single-cell necrosis	Mus: CI ↓, CII + III ↓, CIV ↓
16	M	Y (N)	Poor weight gain, hypoglycaemia, FG, hypotonia, subtle dysmorphic features, progressive liver failure, encephalomyopathy	2 mo	3 mo	Decd, 4 mo	Del exons 3-8 hom	14%	17%	ND	Mus: Generalized fibre atrophy, small angular fibres; Liv: cholestasis and steatosis	ND
17	F	Y (Y, sib P16)	Raised γGT at 6 weeks, progressive liver failure from 4 mo with rapid deterioration	Birth	Birth	Alive, 5 mo	Del exons 3-8 hom	ND	ND	Mosaic depletion	ND	ND

Abbreviations: CI, CII, CIII, CIV, mitochondrial respiratory chain enzyme complexes I, II, III and IV, respectively; Decd, deceased; F, female; FG, faltering growth; Fib, fibroblasts; γGT, gamma-glutamyltransferase; IUGR, intrauterine growth retardation;

Liv, liver; M, male; mo, months; Mus, muscle; N, no; ND, not determined; Y, yes; yr, years.

^aTwo relatives died in infancy due to liver disease (the mother's sister's two children whose parents were also consanguineous).^bLiver disease in two family members, two premature deaths in infancy of two other family members.^cTwo cousins died with liver disease.^dTwo previous siblings died in infancy with similar presentation.

Histopathological features

The histological assessment of a diagnostic muscle biopsy was performed in six patients. Histopathological changes varied from normal findings (patients 2, 4 and 5) to fatty infiltration (patients 7 and 9), to a generalized fibre atrophy with small angular fibres (patient 16).

Liver histology was available from 13 of the 17 patients. Histopathological features included: fatty infiltration (patients 2, 4, 6, 7, 9, 10, 13, 14 and 16), fibrosis/cirrhosis (patients 2, 4, 5, 10 and 14), severe panlobular loss of hepatocytes and stromal collapse (patient 3), giant cell hepatitis with haemorrhagic necrosis (patient 6), non-specific changes (patient 11), cholestasis (patients 14 and 16), patchy hepatocellular oncocytosis and single-cell necrosis (patient 15), abundant mitochondria with mild pleomorphism (patient 9), marked distension of hepatocytes and few periportal glycogen deposits (patient 10) and mild iron deposition in Kupffer cells (patient 13).

Biochemical and histochemical analyses

Table 1 Muscle tissue, liver tissue and/or skin fibroblasts were available for the assessment of mitochondrial respiratory chain function from 7 out of 12 probands. Respiratory chain analysis in muscle at or soon after initial clinical presentation suggested a combined deficiency of respiratory chain complex activities in four patients (7, 9, 10 and 15), whereas the enzyme activities were normal in muscle samples from patients 2, 3 and 4 as well as in liver from patient 4 (who had the highest mtDNA copy number) and fibroblasts from patient 3. Unfortunately, biochemical analyses of liver from patients 3, 11 and 15 with mosaic mtDNA depletion in fibroblasts were not available. Sequential COX-SDH histochemistry revealed a severe, near-global COX deficiency in the liver from patient 2. Initial muscle histology at the age of 28 months from patient 9 was within normal limits, concurrent respiratory enzyme assays showing a borderline complex I deficiency. A repeat muscle biopsy from patient 9 at 3 years revealed slightly decreased respiratory chain activities of both complex I and complex IV (cytochrome c oxidase; 82% of control values). Complex IV activity was 27% of control in the explanted liver tissue. These findings are also summarized in Table 1.

Characterization of the cellular mtDNA depletion phenotype by PicoGreen staining of fibroblasts

Fibroblast cultures were available from 8 of the 17 patients in our cohort. mtDNA was visualized in these cultures using PicoGreen fluorescence microscopy and the membrane potential-dependent mitochondrial probe, TMRM. PicoGreen staining provides a quantitative measure of mtDNA content in a cell, and is more sensitive to small localized depletion than real-time PCR. mtDNA is arranged into punctate DNA/protein structures, termed nucleoids *in vivo*, consisting of several mtDNA genomes complexed to nucleoid proteins. Mosaic mtDNA depletion was clearly observed in fibroblasts from four *MPV17* patients (patients 3, 11, 15 and 17; Figures 3c and e and Supplementary Information 2). In two patients, intermediate mtDNA depletion was observed with fibroblast mtDNA staining being generally paler (patients 4 and 8) whereas PicoGreen staining was normal in fibroblasts from patients 2 and 9. Cells showing the most marked mtDNA depletion exhibited a decreased mitochondrial membrane potential following TMRM staining. This was reflected by real-time PCR (Supplementary Information 2), which showed that the average mtDNA copy number in the cell lines with mosaic depletion was 70% (range 35–100%) of expected compared with 89% (range 43–150%) in the other patient lines (four controls, mean 100%, range 70–136%). Real-time PCR was supported by

quantitation of nucleoid numbers using IN Cell 1000 analyser (Supplementary Information 2).

DISCUSSION

We have performed clinical, biochemical, immunocytochemical and molecular genetic studies of 17 patients with MDS associated with *MPV17* mutations, demonstrating a loose relationship between the clinical phenotype and mutational genotype, with patients with the most severe mtDNA depletion in liver tending to present and die at an earlier age. We have also demonstrated a mosaic mtDNA depletion in fibroblasts from some of the more severely affected patients.

Previously, *MPV17* mutations have been reported in 32 patients with the hepatocerebral form of MDS in which the most common clinical manifestations included early progressive liver failure, hypoglycaemia and raised plasma lactate concentrations.^{4–15} Some of these patients manifested with neuromuscular abnormalities including hypotonia, nystagmus, encephalopathy, developmental delay/mental retardation, microcephaly, seizures, myoclonus, myopathy, ataxia and/or peripheral neuropathy (Navajo neurohepatopathy). White matter changes and abnormalities within the reticular formation of the lower brain stem and within the reticulospinal tracts at the cervicocranial junction on MR imaging have been reported.^{13,15}

Clinical findings and differential diagnoses

All our 17 patients presented with liver disease; other common features included raised serum lactate concentrations, hypotonia and faltering growth. Rarer features included endocrine abnormalities (hypoparathyroidism in patient 9, pituitary dysfunction in patient 12 and hypothyroidism in patient 15), retinal pigmentation (patients 10 and 13), corneal scarring²⁴ (patient 5) and progressive demyelinating peripheral neuropathy (patients 4 and 9).^{7,11} In addition, three patients were found with minor dysmorphic features (patients 12, 15 and 16), which have not been described previously. Thus, the clinical findings in our cohort of MDS patients with *MPV17* mutations are broadly similar to those previously reported.

The infantile phenotypes of hepatocerebral forms of MDS associated with mutations in *POLG*, *PEO1* (Twinkle), *DGUOK* or *MPV17* have similarities, presenting with liver failure along with various neurological and endocrinological manifestations. Significant hypoglycaemia, that is glucose levels that are more difficult to maintain with oral supplementation than would be expected from the degree of liver failure, was commonly present in patients with *MPV17* mutations as previously reported.⁴ We also observed raised levels of tyrosine in blood and/or urine in three patients, as can occur in patients with *DGUOK* mutations.²⁴ We have also noted a tendency for the liver failure to precede severe neurological problems, whereas in our experience the sequence tends to be the other way round in patients with classical Alpers syndrome due to *POLG* mutations: in only 1 out of 24 of our series did liver dysfunction clearly precede neurological involvement.²⁰ For this reason, patient 6 underwent a liver transplant at 15 months of age, but died 15 months later, following neurological and renal deterioration. In summary, presentations of patients with *MPV17* mutations were most like those of patients with *DGUOK*-associated MDS, who typically have neonatal liver failure and hypotonia.²⁵

MPV17 mutations

Twelve different mutations were identified in our cohort of 17 patients (12 probands), including 11 novel mutations (6 missense, 5 truncating) in the *MPV17* gene (Figure 1). This adds significantly to the 21 previously reported mutations.^{4–16} The seven missense




Figure 3 PicoGreen staining of fibroblast cultures. Healthy control fibroblasts exhibit typical bright punctate nucleoid staining when stained with the fluorescent DNA dye PicoGreen (a). Co-staining these cells with the potential sensitive mitochondrial stain, TMRM demonstrated a normal polarized mitochondrial network in all cells (b). Fibroblasts from patient 3 mostly exhibit typical bright nucleoids staining, but some cells (arrow) lack puncta (green in on-line version) because they are depleted of mtDNA, termed mosaic mtDNA depletion (c). Co-staining of these cells with TMRM demonstrates that some of the mtDNA-depleted cells have a depolarized mitochondrial network (d). Fibroblasts grown from patient 17 umbilical cord also show mosaic mtDNA depletion (e), most cells having normal TMRM signal (f) but one depolarized cell is shown (arrow) consistent with mosaic mtDNA depletion. Scale bars = 50 μ m.

mutations (six novel and one previously reported) identified in our cohort are scattered throughout the protein (Figure 2), with no significant clustering. This is in contrast to El-Hattab *et al*,¹¹ who reported clustering of missense mutations in the region of the putative protein kinase C phosphorylation site.

The majority of patients in our cohort presented in the first few weeks of life and died before 2 years of age (although three patients currently aged 5–14 months are living and so their clinical course cannot be predicted), and these patients had either truncating or missense mutations in *MPV17* that presumably led to complete loss of functional protein. However, patients from families 4 and 6 (patients 4, 5, 6 and 9) appear to have a milder disease with better prognosis. These patients are/were either homozygous or compound heterozygous for missense mutations, namely p.Arg41Trp, p.Pro64Arg and p.Pro98Leu, suggesting that these mutant *MPV17* proteins retain some residual function. This is similar to the findings of Karadimas *et al*,⁵ who reported patients homozygous for p.Arg50Gln or compound heterozygous for p.Gly94Arg and p.Pro98Leu, with a relatively mild form of hepatocerebral MDS, namely Navajo

neurohepatopathy. It is interesting to note that both p.Arg41Trp and p.Arg50Gln are located in the same protein loop in the intermembrane space (Figure 2).

As a result of identifying pathogenic *MPV17* mutations, prenatal diagnosis and/or preimplantation genetic diagnosis is now available to these 12 families. Indeed, the parents of patient 16 opted to have amniocentesis and *MPV17* analysis in two subsequent pregnancies, who were accurately identified as an unaffected carrier and an affected girl (patient 17).

Mitochondrial respiratory chain analyses and mtDNA depletion in tissue samples

Respiratory chain deficiency and mtDNA depletion are usually present in the muscle of MDS patients, but not in all patients. In cases with isolated liver involvement, respiratory chain enzyme activity and mtDNA copy number have been normal in muscle and decreased only in liver tissue. For example, many patients with autosomal recessive mutations in the *POLG* gene causing Alpers syndrome (intractable epilepsy, hepatopathy) have normal respiratory chain enzyme activity

and mtDNA copy number in skeletal muscle.^{20,26} The characterization of tissues from previously described patients with *MPV17* mutations has revealed a severe mtDNA depletion that was most pronounced in liver, followed by a less severe, but still significant depletion in skeletal muscle.^{6,15} In our cohort, respiratory chain analysis in muscle revealed a combined deficiency of respiratory chain complexes in four out of seven muscle biopsy samples available for biochemical studies, and mtDNA depletion was identified in three out of eight muscle samples with borderline low copy number in two samples. Therefore, muscle from three out of eight patients showed no evidence of mtDNA depletion. Interestingly, there was no clear correlation between the level of mtDNA depletion in muscle, disease severity or mutation location.

As predicted, respiratory chain enzyme activities and mtDNA copy number in liver tissue correlated more closely with disease severity. Liver was available for mtDNA copy number analysis in seven patients. Severe mtDNA depletion in liver (<20% compared with age-matched controls) was associated with early onset and severe disease course (Table 1). In addition, when measured in both tissues (five patients), the mtDNA content was significantly less in the liver than in the muscle ($P=0.04$, one-tailed paired sample *T*-test). Both of these features are also apparent in patients with Alpers syndrome due to *POLG* mutations.²⁰ Although liver respiratory chain enzyme activities or histochemistry were only available for three patients, this also showed some correlation with disease severity, as the only sample with normal activity was from patient 4 with the mildest disease.

PicoGreen stainings of fibroblasts

We identified mosaic mtDNA depletion in fibroblasts by PicoGreen staining in three cases (patients 3, 11 and 15) with an early-onset liver disease (birth–2.5 months), two of whom had severe mtDNA depletion in liver (5% in patient 3 and 11% in patient 11); fibroblasts grown from the umbilical cord of patient 17 (now 5 months old and deteriorating rapidly with liver failure, and whose older sibling, patient 16, was severely affected and had severe liver mtDNA depletion) also showed mosaic mtDNA depletion. In four other patients (2, 4, 8 and 9) from whom fibroblasts were available, there was no clear evidence of mosaic mtDNA depletion even after extensive passaging, and these patients had more variable onset (birth–5 years) and mtDNA copy number in liver (14–40% of controls). Muscle mtDNA copy number was highly variable both in patients with mosaic mtDNA depletion (25–100% of controls) and in patients without mosaic mtDNA depletion (46–80% of controls). Thus, mosaic mtDNA depletion was more apparent in patients with severe clinical phenotype and low mtDNA content in liver, but did not appear to correlate with muscle mtDNA content. A similar association between the cellular phenotype and severity of the clinical phenotype has previously been seen in patients with the most severe *POLG* mutations.^{20,23} Mosaic mtDNA depletion has also been detected in serum-deprived mouse embryonic fibroblasts lacking *MPV17* protein, using antiDNA antibodies,³ but has not previously been observed in patients with *MPV17* mutations. Finding mosaic depletion in fibroblasts, post mortem may provide a useful clue to aetiology in cases where fibroblasts are the only material available for diagnosis.²⁷

CONCLUSION

Our data confirm previous reports describing *MPV17* mutations as an important cause of mtDNA maintenance disorders, specifically hepatocerebral MDS. We describe 17 patients with 11 novel *MPV17* mutations and provide the first description of recessive *MPV17*

mutations associated with mosaic mtDNA depletion in fibroblasts. We present evidence that *MPV17* mutations lead to tissue selective impairment of mtDNA replication and to a mosaic defect pattern in fibroblasts, which is associated at least in some patients with the severity of clinical phenotype.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank the patients and their families for participating. This study was supported by grants from the MRC, the Wellcome Trust and the Angus Memorial Mitochondrial Fund (JP), the Foundation for Paediatric Research (JU), the Sigrid Juselius Foundation (JU), and the Academy of Finland (JU, Decision number: 138566). RWT is supported by a Wellcome Trust Strategic Award (906919). SR is supported by Great Ormond Street Hospital Children's Charity. We also thank the UK NHS Specialist Commissioners who fund the 'Rare Mitochondrial Disorders of Adults and Children' Diagnostic Service (<http://www.mitochondrialncg.nhs.uk>) in Oxford, London and Newcastle upon Tyne.

- Copeland WC: Inherited mitochondrial diseases of DNA replication. *Annu Rev Med* 2008; **59**: 131–146.
- Spinazzola A, Invernizzi F, Carrara F *et al*: Clinical and molecular features of mitochondrial DNA depletion syndromes. *J Inherit Metab Dis* 2009; **32**: 143–158.
- Viscomi C, Spinazzola A, Maggioni M *et al*: Early-onset liver mtDNA depletion and late-onset proteinuric nephropathy in *Mpv17* knockout mice. *Hum Mol Genet* 2009; **18**: 12–26.
- Spinazzola A, Viscomi C, Fernandez-Vizarrá E *et al*: *MPV17* encodes an inner mitochondrial membrane protein and is mutated in infantile hepatic mitochondrial DNA depletion. *Nat Genet* 2006; **38**: 570–575.
- Karadimas CL, Vu TH, Holve SA *et al*: Navajo neurohepatopathy is caused by a mutation in the *MPV17* gene. *Am J Hum Genet* 2006; **79**: 544–548.
- Wong LJ, Brunetti-Pierri N, Zhang Q *et al*: Mutations in the *MPV17* gene are responsible for rapidly progressive liver failure in infancy. *Hepatology* 2007; **46**: 1218–1227.
- Navarro-Sastre A, Martín-Hernández E, Campos Y *et al*: Lethal hepatopathy and leukodystrophy caused by a novel mutation in *MPV17* gene: description of an alternative *MPV17* spliced form. *Mol Genet Metab* 2008; **94**: 234–239.
- Spinazzola A, Santer R, Akman OH *et al*: Hepatocerebral form of mitochondrial DNA depletion syndrome: novel *MPV17* mutations. *Arch Neurol* 2008; **65**: 1108–1113.
- Kaji S, Murayama K, Nagata I *et al*: Fluctuating liver functions in siblings with *MPV17* mutations and possible improvement associated with dietary and pharmaceutical treatments targeting respiratory chain complex II. *Mol Genet Metab* 2009; **97**: 292–296.
- Parini R, Furlan F, Notarangelo L *et al*: Glucose metabolism and diet-based prevention of liver dysfunction in *MPV17* mutant patients. *J Hepatol* 2009; **50**: 2105–2110.
- El-Hattab AW, Li FY, Schmitt E, Zhang S, Craigen WJ, Wong LJ: *MPV17*-associated hepatocerebral mitochondrial DNA depletion syndrome: new patients and novel mutations. *Mol Genet Metab* 2010; **99**: 300–308.
- Navarro-Sastre A, García-Silva MT, Martín-Hernández E, Lluch M, Briones P, Ribes A: Functional splicing assay supporting that c.70+5G>A mutation in the *MPV17* gene is disease causing. *J Inherit Metab Dis* 2010; **33**: doi:10.1007/s10545-010-9155-x.
- Merkle AN, Nascene DR, McKinney AM: MR imaging findings in the reticular formation in siblings with *MPV17*-related mitochondrial depletion syndrome. *AJNR Am J Neuroradiol* 2012; **33**: E34–E51.
- AlSaman A, Tomoum H, Invernizzi F, Zeviani M: Hepatocerebral form of mitochondrial DNA depletion syndrome due to mutation in *MPV17* gene. *Saudi J Gastroenterol* 2012; **18**: 285–289.
- El-Hattab AW, Scaglia F, Craigen WJ, Wong LJC: *MPV17*-Related Hepatocerebral Mitochondrial DNA Depletion Syndrome; in Pagon RA, Bird TD, Dolan CR, Stephens K, Adam MP (eds) *GeneReviews™ [Internet]*. Seattle (WA): University of Washington 1993–2012.
- Blakely EL, Butterworth A, Hadden RD *et al*: *MPV17* mutation causes neuropathy and leukoencephalopathy with multiple mtDNA deletions in muscle. *Neuromuscul Disord* 2012; **22**: 587–591.
- Li YY, Hengstenberg C, Maisch B: Whole mitochondrial genome amplification reveals basal level multiple deletions in mtDNA of patients with dilated cardiomyopathy. *Biochem Biophys Res Commun* 1995; **210**: 211–218.
- Poulton J, Sewry C, Potter CG *et al*: Variation in mitochondrial DNA levels in muscle from normal controls. Is depletion of mtDNA in patients with mitochondrial myopathy a distinct clinical syndrome? *J Inherit Metab Dis* 1995; **18**: 4–20.

- 19 Morten KJ, Ashley N, Wijburg F *et al*: Liver mtDNA content increases during development: a comparison of methods and the importance of age- and tissue-specific controls for the diagnosis of mtDNA depletion. *Mitochondrion* 2007; **7**: 386–395.
- 20 Ashley N, O'Rourke A, Smith C *et al*: Depletion of mitochondrial DNA in fibroblast cultures from patients with POLG1 mutations is a consequence of catalytic mutations. *Hum Mol Genet* 2008; **17**: 2496–2506.
- 21 Kirby DM, Thorburn DR, Turnbull DM, Taylor RW: Biochemical assays of respiratory chain complex activity. *Methods Cell Biol* 2007; **80**: 93–119.
- 22 Hargreaves IP, Heales SJ, Land JM: Mitochondrial respiratory chain defects are not accompanied by an increase in the activities of lactate dehydrogenase or manganese superoxide dismutase in paediatric skeletal muscle biopsies. *J Inher Metab Dis* 1999; **22**: 925–931.
- 23 Ashley N, Harris D, Poulton J: Detection of mitochondrial DNA depletion in living human cells using PicoGreen staining. *Exp Cell Res* 2005; **303**: 432–446.
- 24 Johnsen SD, Johnson PC, Stein SR: Familial sensory autonomic neuropathy with arthropathy in Navajo children. *Neurology* 1993; **43**: 1120–1125.
- 25 Dimmock DP, Zhang Q, Dionisi-Vici C *et al*: Clinical and molecular features of mitochondrial DNA depletion due to mutations in deoxyguanosine kinase. *Hum Mutat* 2008; **29**: 330–331.
- 26 Nguyen KV, Ostergaard E, Holst Ravn S *et al*: POLG mutations in Alpers syndrome. *Neurology* 2005; **65**: 1493–1495.
- 27 Rahman S, Poulton J: Diagnosis of mitochondrial DNA depletion syndromes. *Arch Dis Child* 2009; **94**: 3–5.



This work is licensed under a Creative Commons Attribution 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by/3.0/>

Supplementary Information accompanies this paper on European Journal of Human Genetics website (<http://www.nature.com/ejhg>)