



Contents lists available at ScienceDirect

Journal of Diabetes and Its Complications

journal homepage: WWW.JDCJOURNAL.COM

A systematic review considering risk factors for mortality of patients discharged from hospital with a diagnosis of diabetes

Teesta Mukherjee^{a,1}, Tim Robbins^{a,b,1}, Sarah N. Lim Choi Keung^a, Sailesh Sankar^{b,c}, Harpal Randeva^{b,c}, Theodoros N. Arvanitis^{b,*}

^a Institute of Digital Healthcare, WMG, University of Warwick, Coventry CV4 7AL, United Kingdom

^b University Hospitals Coventry & Warwickshire NHS Trust, Clifford Bridge Road, Coventry CV2 2DX, United Kingdom

^c Warwick Medical School, University of Warwick, Coventry CV4 7AL, United Kingdom

ARTICLE INFO

Article history:

Received 15 May 2020

Received in revised form 16 July 2020

Accepted 30 July 2020

Available online xxxxx

Keywords:

Diabetes

Mortality

Inpatient care

Systematic review

ABSTRACT

Aim: To identify known risk factors for mortality for adult patients, discharged from hospital with diabetes.

Method: The systematic review was based on the PRISMA protocol. Studies were identified through EMBASE & MEDLINE databases. The inclusion criteria were papers that were published over the last 6 years, in English language, and focused on risk factors of mortality in adult patients with diabetes, after they were discharged from hospitals. This was followed by data extraction "with quality assessment and semi-quantitative synthesis according to PRISMA guidelines".

Results: There were 35 studies identified, considering risk factors relating to mortality for patients, discharged from hospital with diabetes. These studies are distributed internationally. 48 distinct statistically significant risk factors for mortality can be identified. Risk factors can be grouped into the following categories; demographic, socioeconomic, lifestyle, patient medical, inpatient stay, medication related, laboratory results, and glycaemic status. These risk factors can be further divided into risk factors identified in generalized populations of patients with diabetes, compared to specific sub-populations of people with diabetes.

Conclusion: A relatively small number of studies have considered risk factors relating to mortality for patients, discharged from hospital with a diagnosis of diabetes. Mortality is an important outcome, when considering discharge from hospital with diabetes. However, there has only been limited consideration within the research literature.

© 2020 The Authors. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Contents

1.	Introduction	0
2.	Methods	0
2.1.	Search strategy	0
2.2.	Study selection	0
2.3.	Data extraction	0
2.4.	Quality assessment	0
2.5.	Data synthesis	0
3.	Results	0
3.1.	Search results	0
3.2.	Study characteristics	0
3.3.	Risk factor identification	0
3.4.	Study quality	0
4.	Discussion	0
5.	Conclusions	0

* Corresponding author.

E-mail address: t.arvanitis@warwick.ac.uk (T.N. Arvanitis).

¹Authors contributed equally.

<https://doi.org/10.1016/j.jdiacomp.2020.107705>

1056-8727/© 2020 The Authors. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article as: T. Mukherjee, T. Robbins, S.N. Lim Choi Keung, et al., A systematic review considering risk factors for mortality of patients discharged from hospital with..., Journal of Diabetes and Its Complications, <https://doi.org/10.1016/j.jdiacomp.2020.107705>

Funding	0
Acknowledgment	0
References.	0

1. Introduction

Patients with diabetes are known to be at an increased risk of complications and mortality during their inpatient stay.^{1,2} Significant research and resource has been invested into understanding how to reduce this burden of inpatient mortality for patients with diabetes.^{3,4} There has been much less focus on the risk of mortality, following discharge from hospital with diabetes. Data available on post discharge outcomes, in this patient population, is not abundant and is ambiguous.⁵ There is evidence of an increased risk of mortality for patients with diabetes; however, tools for predicting mortality in such patients are not easily available.⁶ This study, therefore, aims to systematically identify the known risk factors for mortality in patients with diabetes, after they are discharged from hospitals.

This work builds on a previous published work, considering risk factors for readmission of patients discharged from hospital with diabetes.⁷ Readmission is a commonly considered measure of care quality, when patients are discharged from hospital, and, therefore, has been more closely studied than mortality following hospital discharge, arguably due to financial incentives and costs associated with readmission rates.⁸ Both readmission and mortality are, however, important outcome measures, when considering the discharge of patients with diabetes and the subsequent development of risk prediction models. The consideration of which patients are at a greatest risk of mortality, following discharge, will help allocated limited resources to supporting this population.

The identification of risk factors, in this manner, has the potential to support research that enables us to predict mortality in patients with

diabetes, following hospital discharge. Once risk factors are known and understood, they can be identified in the patient's individual Electronic Health Record (EHR). Understanding risk factors, relevant to patients discharged from hospital with diabetes, is important to patients, carers, healthcare practitioners and researchers. It supports the delivery and development of individualised medicine, based on each patient's underlying risks; supports our understanding of regional variations in readmission risk; and, supports development of evidence based interventions, targeted at reducing readmission risks. Interestingly, the paucity of research in this area, for diabetes, is in direct contrast to other medical conditions, such as heart failure.⁹

2. Methods

2.1. Search strategy

The systematic review was performed according to the PRISMA protocol.¹⁰ The PRISMA flowchart is shown in Fig. 1.

2.2. Study selection

The literature search was conducted in the MEDLINE, EMBASE & Engineering Village databases. Search terms used to identify relevant medical literature were ((“diabetes” AND “discharge”)) AND (“mortality” or “death” or “died”). Filters were applied to identify studies that were published, in English language, over the last 6 years (March 2014 to February 2020), to include the most recent clinical evidence available in this area of research. Additionally, filters were applied to set participants'




Fig. 1. PRISMA flow diagram of study selection.

Table 1

*Comprising of: 10 retrospective cohort, 4 retrospective analysis, 2 retrospective case control, 1 retrospective observational 1 retrospective registry and 1 retrospective risk prediction modeling.

Study design	Number of papers	Percentage of papers (%)
Retrospective study*	19	54
Prospective study	10	29
Registry based	5	14
Post hoc analysis	1	3

age as 18 years and above, as this research looked at only adult patients. Any study that focused on in-hospital mortality or diabetes, as a risk factor for other clinical conditions, was excluded.

2.3. Data extraction

Data was extracted to a pre-determined, data-extraction proforma. Data was extracted based on the following variables; year of publication, type of study, location of research, sample size, sub population of people with diabetes, statistically significant risk factors and statistical tests for analysis. Additionally, risk factors not meeting significance criteria, but had impact were also extracted. Data extraction was performed independently by two authors, with any discrepancies resolved by a third author.

2.4. Quality assessment

All studies included in the review were assessed for study quality. Data was extracted using Keshav's 5Cs (category, context, correctness, contribution and clarity),¹¹ alongside sample size, justification of their participant selection and statistical tests applied. Papers were not excluded based on producing a negative outcome or being a low quality study.

2.5. Data synthesis

The diversity among the identified studies, in terms of their sample sizes and patient population with various co-morbidities apart from diabetes, did not allow a meta-analysis to be conducted. Instead, both a semi-quantitative and narrative summary of identified prognostic risk factors were performed. Individual risk factors have then been grouped under broader categories, as and when necessary.

Table 2

Distribution of study geographies.

Study location	# studies	Study location	# studies	Study location	# studies
United States	3	Canada	1	Spain	1
Australia	2	China	1	Middle East	1
Brazil	2	Croatia	1	Romania & Germany	1
Greece	2	Finland	1	USA & EU	1
Italy	2	Germany	1		
Taiwan	2	Israel	1		
United Kingdom	2	Latvia	1		

3. Results

3.1. Search results

3436 studies were identified from the literature search, representing 2891 studies following removal of duplicates. Title and abstract based screening resulted in 2679 articles being removed, which left 212 for full text assessment. Full text evaluation resulted in 35 articles being shortlisted for the review (Fig. 1).

3.2. Study characteristics

From the 35 articles identified, 29 were full text articles (83%) and 6 (17%) were only in abstract form, typically from conference presentations (Table 1).

The study designs noted within the review were:

The distribution of studies over time is described in Fig. 2:

(11%) studies were based on international datasets, 12 studies (34%) were conducted based on national data, 9 studies (26%) used regional data, 10 (29%) were single-centre studies. The distribution of study geographies is shown in Table 2:

3.3. Risk factor identification

The majority of studies analysed (24 studies, 69%) identified one or more statistically significant risk factors for mortality, following discharge from hospital with a co-morbid diagnosis of diabetes. There were 11 studies (31%), which identified no statistically significant risk factors. The distribution of risk factors is demonstrated in more detail in Table 3.

The follow up period, during which mortality was looked for in the studies, was highly variable. The most common follow up period was




Fig. 2. Studies over time.

Table 3
Study characteristics.

Ref	Year	Data collection	Sample size	Sub-population	Significant risk factors	No. of significant risk factors	Follow up period (years)
¹²	2017	Regional	22,473	Post Intensive Care Unit (ICU)	Hospital, age, sex, ethnicity, APACHE score, co-morbidity	6	5.1
¹³	2016	National	1613	Post Myocardial Infarction (MI)	Admission creatinine, employment, age, Haemoglobin (Hb), Left Ventricular Systolic Dysfunction (LVSD), co-morbidity, activity, in hospital revascularisation, BMI, insulin, fasting blood glucose, angina frequency	12	5
¹⁴	2017	Regional	59,412	All diabetes	Not specific	0	Until death
¹⁵	2018	Single	312	Age over 60	Gait speed, calf circumference (BMI)	2	NA
¹⁶	2015	Single	4607	All diabetes	Age, Gender, Cardiovascular diseases and infectious diseases (co-morbidity)	4	337 days
¹⁷	2014	National	1082	Heart failure	Age, co-morbidity, sodium (biochemistry), anaemia (haematology), medication, BMI, NHYA (severity scoring)	7	1
¹⁸	2015	National	2904	Chronic Kidney Disease (CKD) and acute MI admission	0	0	1
¹⁹	2015	National	4054	Admission for acute MI	Statins prior to acute MI, number of medications	2	3
²⁰	2018	Single	207	Diabetic Ketoacidosis (DKA) diagnosed	Age, duration of diabetes, number of hospitalisations	3	5
²¹	2016	Single	409	Admitted with pneumonia	Hospitalization for pneumonia (admission diagnosis) Age, haemodialysis (procedure) Charlson co-morbidity index (score) renal insufficiency, pleural effusion, malnutrition (co-morbidity) pH < 7.35 (biochemistry)	8	1
²²	2017	Single	761	Admitted with poor glucose control	Estimated Glomerular Filtration Rate (eGFR), Albuminuria (biochemistry)	2	6.6
²³	2016	International	5005	Patients admitted with acute decompensated heart failure with diabetes	BMI (underweight, severely obese)	2	1
²⁴	2018	Single	304	All diabetes	0	0	1
²⁵	2019	Regional	104,525	Patients hospitalised and dispensed prescription for insulin and/or OHA within 8 days of discharge. Age > 66 years	new insulin use (medication)	1	30 days
²⁶	2018	Single	130	Type 2 Diabetes Mellitus (T2DM)	0	0	30 days
²⁷	2017	Single	304	T2DM patients with first ever noncardioembolic acute ischemic stroke	Age, Stroke severity (NIHSS scale), clopidogrel compared to aspirin (medication) co-morbidity	4	10
²⁸	2016	National	386	DKA admitted to Intensive Treatment Unit (ITU)	Age, APACHEII score, mechanical ventilation, number of organs supported, DKA severity, creatinine, bilirubin, pCO2 (biochemistry), lowest Glasgow Coma Score (GCS)	9	5
²⁹	2016	Regional	214,991	All diabetes	Age, sex, Charlson index	3	2
³⁰	2018	National	17,186	Diabetes and heart failure	0	0	NA
³¹	2016	National	1743	Diabetes and Acute Coronary Syndrome (ACS)	0	0	6 months
³²	2019	National	843,978	All diabetes	0	0	6 months
³³	2017	National	71,640	Diabetes and cancer	0	0	3
³⁴	2017	Regional	28,353	All diabetes	Median glucose during admission & glucose variability	2	6
³⁵	2015	International	221	T2DM admitted with ACS	0	0	1
³⁶	2014	International	1998	Patients hospitalised with worsening heart failure and an ejection fraction below 40%	0	0	9.9 months
³⁷	2017	Regional	218	Haematocrit, EF, ACEi. Red blood cell distribution	0	0	1
³⁸	2017	Single	202	All diabetes	hba1c, insulin resistance	2	2.5
³⁹	2020	International	974	ACS	Fibrinolysis	1	1
⁴⁰	2019	Single	306	All diabetes	Sarcopenia	1	2
⁴¹	2019	Regional	10,542	All diabetes	Heat wave intensity	1	NA
⁴²	2019	National	13,113	Heart failure	Diabetes and heart failure	1	1
⁴³	2019	National	41,776	Heart failure	0	0	1
⁴⁴	2019	Regional	NA	All diabetes	Pharmacy follow up	1	90 days
⁴⁵	2019	National	1221	Acute MI	Insulin prescription at discharge	1	5
⁴⁶	2019	Regional	100,000	T2DM over 40 years	Lower limb amputation, sex	2	10

12 months in 8 studies (23%). The shortest follow up period was 1 month and the longest follow up period was 10 years.

The extracted studies varied according to whether they considered risk in all patients with diabetes (6 studies, 17%) or a specific sub population. Those sub populations were defined either by the type of diabetes, or the characteristics relevant to the inpatient stay, and are demonstrated in more detail in Table 3.

In total, there were 48 distinct risk factors identified. These can be grouped into risk factor categories, alongside grouping according to whether they were identified for all patients with diabetes. The results

of this grouping is shown in Table 4, which represents a complete list of all statistically significant risk factors identified within the published research literature, alongside the number (and references) of studies, which identified these risk factors as significant.

3.4. Study quality

The sample size of extracted studies varied between 130 participants and 843,978 participants. The mean average number of participants per study was 44,484, with a median of 1743 participants.

Table 4
Risk factors.

Risk factors for specific diabetes subpopulations		Risk factors in general diabetes populations	
Risk factor	Number of studies & Ref	Risk Factor	No.& Ref
Demographic			
Age	7 ^{12,13,17,20,21,24,27,28}	Age	2 ^{16,29}
Gender	1 ^{12,46}	Gender	2 ^{16,29}
Race	1 ¹²		
Socioeconomic status			
Employment status	1 ¹³		
Lifestyle			
Leisure time activity	1 ¹³		
External factors			
		Weather conditions	1 ⁴¹
Patient medical factors			
Co-morbidity	6 ^{12,13,17,25,27,42}	Co-morbidity	2 ^{16,29,40}
Malnutrition	1 ²¹		
Duration of diabetes	1 ²⁰		
Severity score	4 ^{12,17,27,28}		
DKA Severity	1 ⁴⁷		
Body mass index	3 ^{13,17,23}		
Gait speed	1 ¹⁵		
Calf circumference	1 ¹⁵		
Angina frequency	1 ¹³		
Inpatient stay factors			
Procedure	2 ^{13,21,46}		
Admission diagnosis	1 ²¹		
Length of stay	1 ²⁴		
Glasgow coma score (GCS)	1 ²⁸		
Mechanical ventilation	1 ²⁸		
Number of organs supported	1 ²⁸		
No. of hospitalisations	1 ²⁰		
Which hospital admitted to	1 ¹²		
Medication related			
Beta blocker	1 ¹⁷	Pharmacy follow up	1 ⁴⁴
ACEi/ARB blocker	1 ¹⁷		
Statins prior to AMI	1 ¹⁹		
No. of medications at discharge	1 ¹⁹		
Insulin use	1 ^{25,45}		
Clopidogrel	1 ²⁷		
Laboratory results			
Admission creatinine	2 ^{13,28}	Hba1c	1 ³⁸
Fasting glucose	1 ¹³		
Sodium level	1 ¹⁷		
pH below 7.35	1 ²¹		
eGFR	1 ²²		
Albuminuria	1 ²²		
Bilirubin	1 ²⁸		
PCO2 (on blood gas)	1 ²⁸		
Admission haemoglobin	1 ¹³		
Anaemia	1 ²⁷		
Fibrinolysis	1 ³⁹		
Glycaemic Status (not including Hba1c above)			
		Glycaemic variability	1 ³⁴
		Mean capillary blood glucose	1 ³⁸
		Insulin resistance	1 ³⁸

All the papers used statistical tests to assess the impact of potential risk factors and evaluate their significance. Statistical significance was set at a standard of p -value less than 0.05. The majority of the papers, which was 14 in number (40%), used Cox proportional hazards model.

7 papers (20%) had not defined their statistical test, 5 papers (14%) used univariate and multivariate Cox regression analysis, 3 studies (8%) utilised univariate and multivariate analysis, 2 papers (6%) used multivariate analysis, 1 study (3%) used multivariate binary logistic regression, 1 study used Kaplan Meir curves (3%), 1 study used sensitivity analysis (3%) and finally 1 additional study (3%) used parametric and non-parametric tests. No studies considered the application of standardized effect size measures; there was very limited calculation of power sizes, in advance, to identify and appropriate population size for the study.

4. Discussion

The identification of risk factors, following discharge from hospital is an important consideration for patients, clinicians, managers and policy makers. This review represents the first systematic collection and identification of risk factors for mortality in the published research literature. We identify a total of 48 distinct, statistically significant risk factors across 9 categories reported in 35 studies. The most commonly reported risk factor was age, followed by co-morbidity burden. Thirty-seven of the risk factors were only identified as statistically significant in a single research paper. We would argue that this represents a research literature that is at a relatively early stage of maturity, in considering this important topic area.

The quality of studies identified was relatively high, with acceptable sample sizes, albeit frequently lacking a power-calculation for the results reported. The statistical tests used were acceptable, with the majority using Cox proportional hazard models. The small number of papers not defining the statistical test used was however concerning. A large proportion of the extracted articles represented conference abstracts rather than full research papers. These conference abstracts are typically shorter, lack full methodological detail and often do not undergo such rigorous peer review. This again supports the argument that the research literature in this area is at an early stage of maturity.

The results presented in this paper can be compared to a similar review that considered risk factors, specifically for *readmission*, when patients with diabetes are discharged from hospital.⁷ In the review, considering *readmission*, a significantly larger number of studies were reported (82 studies compared to 35) and 72 distinct risk factors were identified, in comparison to the 48 statistically significant risk factors identified here. There is a higher proportion of conference abstracts within the mortality articles, as compared to the *readmission* articles, and the mortality articles included fewer reports of prospective studies, compared to the *readmission* papers. These comparisons, therefore, contrast the early state of the research literature for mortality risk factors with a more complete picture seen relating to risk factors, for *readmission*. *Readmission* and *post-discharge mortality* are both important negative outcomes, in the context of discharge from hospitals. Clinicians, managers and policy makers are motivated to reduce both outcomes. However, it is more typically avoiding *readmission* that is associated with financial incentivisation for clinical providers. This incentivisation, alongside *readmission* typically happening sooner than mortality events, may have led researchers more towards conducting *readmission* rather than mortality studies. However, mortality is undoubtedly a vitally important consideration for all those involved with the discharge process and it is important that adequate research resource is directed towards this outcome.

The strengths of this research include the PRISMA approach to reporting, the use of Keshav's 5Cs approach to quality assessment and the level of detail in data collection. There are however a number of limitations that should be considered. The research only considers papers published in the English Language; there may be other risk factors identified in other studies that are published in different languages. Furthermore, the study only considers papers published over the last 6 years.

This ensures that all risk factors identified are relevant to current medical practice, although it is possible that previously identified risk factors may be also identified. Finally, the search term used to identify discharge from hospital was only “discharge,” this is a relatively universally accepted term for when a patient leaves hospital following an inpatient admission, however it is possible that a very small number of articles used an alternative descriptor and therefore the results should be interpreted with awareness of this.

There is a lot of scope and future work that can be recommended, based on this systematic review. The identified significant risk factors can be stratified according to their effectiveness in predicting mortality in type 2 diabetes patients, following their discharge from hospitals. These will highlight patients who are most vulnerable and need special attention. Additionally, available EHR based predictive tools, such as “deep learning” models that use machine learning techniques and could potentially be used for risk prediction of mortality in such patients, could be found out. Following this step, the feasibility of the chosen predictive tools can be examined and validated. Such evaluated tools for risk prediction can be integrated within the EHR system, to predict the risk of mortality post hospital discharge tailored for each patient who has type 2 diabetes.

5. Conclusions

Mortality, following hospital discharge, is necessarily an important outcome of relevance to patients, carers, clinicians and policymakers. This review is the first systematic collection of risk factors identified for mortality for patients being discharged from hospital with diabetes. The review demonstrates that the literature is at an early stage of maturity, particularly in comparison to the literature considering risks factors for readmission following discharge. This research is particularly important for the future development of risk prediction models that can support the better identification of patients at most risk, and the tailoring of support strategies to individual patient needs in an evidence-based manner.

Funding

Dr. Tim Robbins receives funding through a PhD Studentship at WMG, University of Warwick.

Professor Theodoros N. Arvanitis is partially funded by HDR UK. HDR UK is funded by the UK Medical Research Council, Engineering and Physical Sciences Research Council, Economic and Social Research Council, Department of Health and Social Care (England), Chief Scientist Office of the Scottish Government Health and Social Care Directorates, Health and Social Care Research and Development Division (Welsh Government), Public Health Agency (Northern Ireland), British Heart Foundation and Wellcome Trust. HDR-3001.

Declaration of competing interest

This is to declare that all authors have no conflicts of interest to declare and all funding has been noted in the manuscript.

Acknowledgment

The authors would like to thank WMG, University of Warwick for Dr. T. Robbins' PhD studentship and HDR UK for partially funding Prof. T. N. Arvanitis. HDR UK is funded by the UK Medical Research Council, Engineering and Physical Sciences Research Council, Economic and Social Research Council, Department of Health and Social Care (England), Chief Scientist Office of the Scottish Government Health and Social Care Directorates, Health and Social Care Research and Development Division (Welsh Government), Public Health Agency (Northern Ireland), British Heart Foundation and Wellcome Trust.

References

- Valent F, Tonutti L, Grimaldi F. Does diabetes mellitus comorbidity affect in-hospital mortality and length of stay? Analysis of administrative data in an Italian Academic Hospital. *Acta Diabetol* 2017;54:1081-90.
- Nirantharakumar K, Marshall T, Kennedy A, Narendran P, Hemming K, Coleman J. Hypoglycaemia is associated with increased length of stay and mortality in people with diabetes who are hospitalized. *Diabet Med* 2012;29:e445-8.
- Savage M, Dhatariya K, Kilvert A, Rayman G, Rees J, Courtney C, et al. Diabetes UK position statements and care recommendations: joint British Diabetes Societies guideline for the management of diabetic ketoacidosis. *Diabet Med* 2011;28:508-15.
- Akiboye F, Adderley N, Martin J, Gokhale K, Rudge G, Marshall T, et al. Impact of the diabetes inpatient care and education (DICE) project on length of stay and mortality. *Diabetic Medicine* 2019;37(2):277-85.
- Sarma S, Mentz RJ, Kwasny MJ, Fought AJ, Huffman M, Subacius H, et al. Association between diabetes mellitus and post-discharge outcomes in patients hospitalized with heart failure: findings from the EVEREST trial. *Eur J Heart Fail* 2013;15:194-202.
- Wells BJ, Jain A, Arrigain S, Yu C, Rosenkrans WA, Kattan MW. Predicting 6-year mortality risk in patients with type 2 diabetes. *Diabetes Care* 2008;31:2301-6.
- Robbins TD, Keung SLC, Sankar S, Randeve H, Arvanitis T. Risk factors for readmission of inpatients with diabetes: a systematic review. *J Diabetes Complications* 2019;33(5):398-405.
- Soeken KL, Prescott PA, Herron DG, Creasia J. Predictors of hospital readmission: a meta-analysis. *Eval Health Prof* 1991;14:262-81.
- Philbin EF, DiSalvo TG. Prediction of hospital readmission for heart failure: development of a simple risk score based on administrative data. *J Am Coll Cardiol* 1999;33:1560-6.
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009;151:264-9.
- Keshav S. How to read a paper. *ACM SIGCOMM Computer Communication Review* 2007;37:83-4.
- Abdelhamid YA, Plummer MP, Finnis ME, Biradar V, Bihari S, Kar P, et al. Long-term mortality of critically ill patients with diabetes who survive admission to the intensive care unit. *Crit Care Resusc* 2017;19:303.
- Arnold SV, Spertus JA, Jones PG, McGuire DK, Lipska KJ, Xu Y, et al. Predicting adverse outcomes after myocardial infarction among patients with diabetes mellitus. *Circ Cardiovasc Qual Outcomes* 2016;9:372-9.
- Barengo NC, Teuschl Y, Moltchanov V, Laatikainen T, Jousilahti P, Tuomilehto J. Coronary heart disease incidence and mortality, and all-cause mortality among diabetic and non-diabetic people according to their smoking behavior in Finland. *Tobacco induced diseases* 2017;15:12.
- Beretta MV, Feldman JV, Rodrigues T. Predictor of mortality after hospital discharge in the elderly with type 2 diabetes mellitus—follow-up of two years. *Am Diabetes Assoc* 2018;67(Supplement 1).
- Buchs AE, Braverman M, Rapoport MJ. Hyperglycemia in diabetic patients in general medicine wards: no independent correlation with in-and out-of-hospital mortality. *Sat* 2015;15:19.
- Carrasco-Sanchez EJ, Gomez-Huelgas R, Formiga F, Conde-Martel A, Trullas JC, Bettencourt P, et al. Association between type-2 diabetes mellitus and post-discharge outcomes in heart failure patients: findings from the RICA registry. *Diabetes Res Clin Pract* 2014;104:410-9.
- Chen D-Y, Wang S-H, Mao C-T, Tsai M-L, Lin Y-S, Chou C-C, et al. Sitagliptin and cardiovascular outcomes in diabetic patients with chronic kidney disease and acute myocardial infarction: a nationwide cohort study. *Int J Cardiol* 2015;181:200-6.
- Colombo MG, Meisinger C, Amann U, Heier M, von Scheidt W, Kuch B, et al. Association of obesity and long-term mortality in patients with acute myocardial infarction with and without diabetes mellitus: results from the MONICA/KORA myocardial infarction registry. *Cardiovasc Diabetol* 2015;14:24.
- Vencio S, Caiado A, Morgental D, Bufaiçal N, Carneiro R, Vencio RC. *21st Brazilian diabetes society congress*. 2018.
- Falcone M, Tiseo G, Russo A, Giordo L, Manzini E, Bertazzoni G, et al. Hospitalization for pneumonia is associated with decreased 1-year survival in patients with type 2 diabetes: results from a prospective cohort study. *Medicine* 2016;95.
- Hsieh Y-M, Lee W-J, Sheu WH-H, Li Y-H, Lin S-Y. Inpatient screening for albuminuria and retinopathy to predict long-term mortality in type 2 diabetic patients: a retrospective cohort study. *Diabetol Metab Syndr* 2017;9:29.
- Abi Khalil C, Suleiman K, Singh R, Asaad N, Jayyousi A, Al Suwaidi J, et al. Severe obesity, but not overweight, is associated with short-and long-term decreased mortality in patients with type 2 diabetes hospitalized for acute decompensated heart failure: findings from the gulf-care. *Circulation* 2016;134. p. A12149-A12149.
- Lim W, Goh S-Y, Bee YM, Chan TCE, Tan XHA, Wee Z, et al. High one-year mortality following hospitalization for severe hypoglycemia among patients with diabetes mellitus: findings of a retrospective cohort study at an acute tertiary care hospital in Singapore. *Curr Med Res Opin* 2019;35:631-5.
- Lysz Z, Fung K, Giannakeas V, Fischer HD, Bell CM, Lipscombe LL. The association between insulin initiation and adverse outcomes after hospital discharge in older adults: a population-based cohort study. *J Gen Intern Med* 2019;34:575-82.
- Marušić S, Meliš P, Lucijanić M, Grgurević I, Turčić P, Neto PRO, et al. Impact of pharmacotherapeutic education on medication adherence and adverse outcomes in patients with type 2 diabetes mellitus: a prospective, randomized study. *Croat Med J* 2018;59:290.
- Milionis H, Ntaios G, Papavasileiou V, Spengos K, Manios E, Elisaf M, et al. Aspirin versus clopidogrel for type 2 diabetic patients with first-ever noncardioembolic acute ischemic stroke: ten-year survival data from the athens stroke outcome project. *J Stroke Cerebrovasc Dis* 2017;26:2769-77.

28. Ramaesh A. Incidence and long-term outcomes of adult patients with diabetic ketoacidosis admitted to intensive care: a retrospective cohort study. *Journal of the Intensive Care Society* 2016;17:222-33.
29. Seghieri G, Policardo L, Profili F, Francesconi P, Anichini R, Del Prato S. Hospital incidental diagnosis of diabetes: a population study. *J Diabetes Complications* 2016;30:457-61.
30. Sharma A, Wu J, Hernandez AF, Felker G, Al-Khatib S, Green J, et al. Comparative effectiveness of primary prevention implantable cardioverter-defibrillators in older adults with diabetes mellitus: insights from the get with the guidelines-heart failure registry. *Circulation: Cardiovascular Quality and Outcomes* 2018;11. p. A122-A122.
31. Snir A, Dabin B, Hyun K, Yamen E, Ryan M, Aliprandi-Costa B, et al. Glycosylated haemoglobin assessment in diabetic patients with acute coronary syndromes. *Intern Med J* 2016;46:574-82.
32. Spanakis EK, Umpierrez GE, Siddiqui T, Zhan M, Snitker S, Fink JC, et al. Association of glucose concentrations at hospital discharge with readmissions and mortality: a nationwide cohort study. *The Journal of Clinical Endocrinology & Metabolism* 2019;104(9):3679-91.
33. Strele I, Pildava S, Repsa I, Kojalo U, Vilmanis J, Brigis G. Pre-existing diabetes mellitus and all-cause mortality in cancer patients: a register-based study in Latvia. *Acta Oncol* 2018;57:973-82.
34. Timmons JG, Cunningham SG, Sainsbury CA, Jones GC. Inpatient glycaemic variability and long-term mortality in hospitalized patients with type 2 diabetes. *J Diabetes Complications* 2017;31:479-82.
35. Trutz J, Babeş A, Babeş K. The importance of the treatment strategy changes in the long-term evolution of type 2 diabetic patients with sub-optimal Glycaemic control after acute coronary syndrome. *Romanian Journal of Diabetes Nutrition and Metabolic Diseases* 2015;22:29-37.
36. Vaduganathan M, Dei Cas A, Mentz RJ, Greene SJ, Khan S, Subacius HP, et al. Mineralocorticoid receptor antagonist use in hospitalized patients with heart failure, reduced ejection fraction, and diabetes mellitus (from the EVEREST trial). *Am J Cardiol* 2014;114:743-50.
37. Xanthopoulos A, Giamouzis G, Melidonis A, Kitai T, Paraskevopoulou E, Paraskevopoulou P, et al. Red blood cell distribution width as a prognostic marker in patients with heart failure and diabetes mellitus. *Cardiovasc Diabetol* 2017;16:81.
38. Xing Z, Zhang L, Liu Z, He P, Yang Y, Wulashan M. Prognostic value of glucose metabolism for non-ST-segment elevation infarction patients with diabetes mellitus and single concomitant chronic total occlusion following primary percutaneous coronary intervention. *Medicine* 2017;96(45).
39. Sumaya W, Wallentin L, James SK, Siegbahn A, Gabrysck K, Himmelmann A, et al. Impaired fibrinolysis predicts adverse outcome in acute coronary syndrome patients with diabetes: a PLATO sub-study. *Thromb Haemost* 2020;23.
40. Beretta MV, Dantas Filho FF, Freiberg RE, Feldman JV, Nery C, Rodrigues TC. Sarcopenia and Type 2 diabetes mellitus as predictors of 2-year mortality after hospital discharge in a cohort of hospitalized older adults. *Diabetes Research and Clinical Practice* 2020;159. (no pagination).
41. Xu Z, Tong S, Cheng J, Crooks JL, Xiang H, Li X, et al. Heatwaves and diabetes in Brisbane, Australia: a population-based retrospective cohort study. *Int J Epidemiol* 2019;48:1091-100.
42. Edmonston DL, Wu J, Matsouaka RA, Yancy C, Heidenreich P, Pina IL, et al. Association of post-discharge specialty outpatient visits with readmissions and mortality in high-risk heart failure patients. *Am Heart J* 2019;212:101-12.
43. Echouffo-Tcheugui JB, Sheng S, DeVore AD, Matsouaka RA, Hernandez AF, Yancy CW, et al. Glycated hemoglobin and outcomes of heart failure (from get with the guidelines-heart failure). *American Journal of Cardiology* 2019;123:618-26.
44. Wright EA, Graham JH, Maeng D, Tusing L, Zaleski L, Martin R, et al. Reductions in 30-day readmission, mortality, and costs with inpatient-to-community pharmacist follow-up. *J Am Pharm Assoc* 2019;59:178-86.
45. Bataille V, Ferrieres J, Danchin N, Puymirat E, Zeller M, Simon T, et al. Increased mortality risk in diabetic patients discharged from hospital with insulin therapy after an acute myocardial infarction: data from the FAST-MI 2005 registry. *Eur Heart J Acute Cardiovasc Care* 2019;8:218-30.
46. Lopez-De-Andres A, Jimenez-Garcia R, Esteban-Vasallo MD, Hernandez-Barrera V, Aragon-Sanchez J, Jimenez-Trujillo I, et al. Time trends in the incidence of long-term mortality in T2DM patients who have undergone a lower extremity amputation. Results of a descriptive and retrospective cohort study. *J Clin Med* 2019;8.
47. Azevedo LC, Choi H, Simmonds K, Davidow J, Bagshaw SM. Incidence and long-term outcomes of critically ill adult patients with moderate-to-severe diabetic ketoacidosis: retrospective matched cohort study. *J Crit Care* 2014;29:971-7.