



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Transcriptional analyses identify genes that modulate bovine macrophage response to *Toxoplasma* infection and immune stimulation

Citation for published version:

Gossner, A & Hassan, M 2020, 'Transcriptional analyses identify genes that modulate bovine macrophage response to *Toxoplasma* infection and immune stimulation', *Frontiers in Cellular and Infection Microbiology*. <https://doi.org/10.3389/fcimb.2020.00437>

Digital Object Identifier (DOI):

[10.3389/fcimb.2020.00437](https://doi.org/10.3389/fcimb.2020.00437)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Frontiers in Cellular and Infection Microbiology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Transcriptional Analyses Identify Genes That Modulate Bovine Macrophage Response to *Toxoplasma* Infection and Immune Stimulation

Anton Gossner¹ and Musa A. Hassan^{1,2*}

¹ Division of Infection and Immunity, The Roslin Institute, The University of Edinburgh, Edinburgh, United Kingdom, ² Centre for Tropical Livestock Genetics and Health, The University of Edinburgh, Edinburgh, United Kingdom

OPEN ACCESS

Edited by:

Prajwal Gurung,
The University of Iowa, United States

Reviewed by:

David Arranz-Solis,
University of California, Davis,
United States
Xiaoyu Hu,
University of Texas Southwestern
Medical Center, United States

*Correspondence:

Musa A. Hassan
musa.hassan@roslin.ed.ac.uk

Specialty section:

This article was submitted to
Microbes and Innate Immunity,
a section of the journal
Frontiers in Cellular and Infection
Microbiology

Received: 27 April 2020

Accepted: 16 July 2020

Published: 20 August 2020

Citation:

Gossner A and Hassan MA (2020)
Transcriptional Analyses Identify
Genes That Modulate Bovine
Macrophage Response to
Toxoplasma Infection and Immune
Stimulation.
Front. Cell. Infect. Microbiol. 10:437.
doi: 10.3389/fcimb.2020.00437

The obligate intracellular parasite, *Toxoplasma gondii*, is highly prevalent among livestock species. Although cattle are generally resistant to *Toxoplasma* strains circulating in Europe and North America, the underlying mechanisms are largely unknown. Here, we report that bovine bone marrow-derived macrophage (BMDM) pre-stimulated with interferon gamma (IFN γ) restricts intracellular *Toxoplasma* growth independently of nitric oxide. While *Toxoplasma* promoted the expression of genes associated with alternative macrophage activation and lipid metabolism, IFN γ abrogated parasite-induced transcriptional responses and promoted the expression of genes linked to the classical macrophage activation phenotype. Additionally, several chemokines, including *CCL22*, that are linked to parasite-induced activation of the *Wnt*/ β -catenin signaling were highly expressed in *Toxoplasma*-exposed naïve BMDMs. A chemical *Wnt*/ β -catenin signaling pathway antagonist (IWR-1-endo) significantly reduced intracellular parasite burden in naïve BMDMs, suggesting that *Toxoplasma* activates this pathway to evade bovine macrophage anti-parasitic responses. Congruently, intracellular burden of a mutant *Toxoplasma* strain (RH Δ ASP5) that does not secrete dense granule proteins into the host cell, which is an essential requirement for parasite-induced activation of the *Wnt*/ β -catenin pathway, was significantly reduced in naïve BMDMs. However, both the *Wnt*/ β -catenin antagonist and RHASP Δ 5 did not abolish parasite burden differences in naïve and IFN γ -stimulated BMDMs. Finally, we observed that parasites infecting IFN γ -stimulated BMDMs largely express genes associated with the slow dividing bradyzoite stage. Overall, this study provides novel insights into bovine macrophage transcriptional response to *Toxoplasma*. It establishes a foundation for a mechanistic analysis IFN γ -induced bovine anti-*Toxoplasma* responses and the counteracting *Toxoplasma* survival strategies.

Keywords: bovine toxoplasmosis, macrophages, RNA-sequencing, *Toxoplasma*, chemokine, *Wnt*/ β -catenin signaling

INTRODUCTION

Toxoplasma is a zoonotic protozoan parasite that infects virtually all warm-blooded vertebrates and is perhaps the most successful and widespread human pathogen. It is the leading cause of encephalitis and death in HIV/AIDS patients (Basavaraju, 2016) and is ranked 4th among foodborne parasites with the greatest global impact (FAO/WHO, 2014) and contributors to years lived with disability and disability-adjusted life years per million persons (Gkogka et al., 2011). *Toxoplasma* undergoes sexual reproduction exclusively in cats while asexual reproduction can occur in any intermediate host, including humans and farmed animals. Definitive-to-intermediate host parasite transmission occurs via ingestion of oocysts from infected cat feces, while transmission between intermediate hosts can occur through the consumption of contaminated food products or vertically from mother to fetus (Hill and Dubey, 2002). Although *Toxoplasma* infections in healthy individuals are mostly asymptomatic, severe disease or even death, often caused by damage to the brain or other organs, can occur in immunocompromised or congenitally infected individuals (Hill and Dubey, 2002). *Toxoplasma* typically establishes lifelong chronic infection in healthy intermediate hosts by encysting, after an initial phase of rapid intracellular proliferation and cell–cell spread, in the central nervous system and muscle tissues (Hill and Dubey, 2002). To do this, the parasite must carefully regulate immune activation and host cell anti-*Toxoplasma* effector mechanisms.

Toxoplasma invades host cells mostly through an active process powered by parasite-generated actin motor activity to form a specialized non-fusogenic parasitophorous vacuole (PV) (Fleckenstein et al., 2012) that helps the parasite avoid host cell immune surveillance (Morisaki et al., 1995). In phagocytic cells, such as macrophages, the parasite can also be taken up by phagocytosis and, sometimes the phagocytosed parasites can exit the phagosome to form a PV (Zhao et al., 2014). Besides hiding within the PV, the parasite also evades host cell immunity by sequentially discharging several effector proteins that modulate a variety of host immune and metabolic processes, including the inflammatory pathway (Hunter and Sibley, 2012). Generally, effective host anti-*Toxoplasma* responses are dependent on the production of interleukin (IL)-12 (Gazzinelli et al., 1994) by macrophages and dendritic cells (Gazzinelli et al., 1994). IL-12 in turn activates natural killer (NK) and T cells to secrete interferon gamma (IFN γ) (Gazzinelli et al., 1993, 1994), a pro-inflammatory cytokine that activates several anti-*Toxoplasma* effector mechanisms such as the interferon-regulated GTPases (IRGs) in mice (Zhao et al., 2009), reactive nitrogen/oxygen intermediates (Scharton-Kersten et al., 1997), tryptophan degradation and cell death in human cells (Pfefferkorn, 1984; Niedelman et al., 2013), and inflammasome activation (Cirelli et al., 2014). In return, *Toxoplasma* has evolved several mechanisms to counteract the IFN γ -induced host defenses, including the secretion of effector proteins from specialized apical organelles that co-opt host transcription and signaling pathways to control host cell responsiveness to inflammatory signals (Jensen et al., 2011; Koshy et al., 2012; Bougdour et al., 2013; He et al., 2018).

Besides producing the IL12 that primes IFN γ secretion by NK and T cells, macrophages are also the preferred intracellular niche for the fast-dividing parasite stage (Jensen et al., 2011). Thus, the innate defenses triggered by monocytes/macrophages are key to toxoplasmosis pathogenesis in humans and rodents (Channon et al., 2000; Dunay and Sibley, 2010; Gregg et al., 2013; Tosh et al., 2016; Song et al., 2017). Besides killing microbes, including *Toxoplasma*, macrophages can also initiate adaptive immune responses (van de Vosse et al., 2009; Thi et al., 2012). Human and mouse models show that upon the engagement of surface signaling receptors or pattern recognition receptors (PRRs) such as, toll-like (TLRs), RIG-I-like (RLRs), and the cytosolic NOD-like (NLRs) receptors by conserved pathogen-associated molecular patterns (PAMPs) such as, lipopolysaccharide (LPS) or immune factors, including cytokines, macrophages assume different activation phenotypes. The most extreme classical [M1, M(IFN γ)] and the alternative [M2, M (IL-4)] phenotypes are separated by several intermediate activation states (Murray et al., 2014). The M1 phenotype, which can be induced by IFN γ , is highly microbicidal and characterized by the production of reactive oxygen and nitrogen intermediates such as nitric oxide (NO), and a range of pro-inflammatory cytokines and chemokines, such as Tumor necrosis factor (TNF) alpha (De Paoli et al., 2014; Murray et al., 2014). In contrast, the M2 phenotype, which is induced by IL-4 and IL-13 and is important for regulating inflammation, is characterized by the production of anti-inflammatory cytokines and growth factors (Sindrilaru and Scharffetter-Kochanek, 2013). The general hypothesis is that macrophage activation phenotypes, which are underpinned by discrete transcriptional programs (Hassan et al., 2015), provide a high degree of plasticity that is exploited by some intracellular pathogens, including *Toxoplasma* and *Mycobacteria*, to turn this potentially hostile host cell into a favorable replication niche (Price and Vance, 2014). Although effective host response to *Toxoplasma* require the induction of inflammation, characterized by classical macrophage activation, *Toxoplasma* promotes its survival in macrophages by secreting effector proteins to dampen inflammatory responses and favor alternative macrophage activation (Jensen et al., 2011). Indeed, *Toxoplasma* strain differences in virulence in mice is partly due to strain differences in inducing alternative macrophage activation (Jensen et al., 2011), while host differences in susceptibility to *Toxoplasma* is due in part to differences in macrophage activation phenotypes after infection (Jensen et al., 2011, 2013). Therefore, to devise strategies to improve the early defense against *Toxoplasma* and a variety of intracellular pathogens, it is important to understand the molecular mechanisms underpinning macrophage response to *Toxoplasma* and/or immune effector proteins, such as cytokines.

Toxoplasma is common in many species of livestock, including cattle. Compared to other livestock species, such as sheep and pigs, cattle are highly resistant to *Toxoplasma* and rarely transmit the parasite to other intermediate hosts (Dubey, 1986; Esteban-Redondo and Innes, 1997). Unlike sheep and pigs, natural *Toxoplasma* infection in cattle is mostly asymptomatic and does not appear to result in abortion. However, the molecular factors and mechanisms that modulate bovine-*Toxoplasma*

interactions, which can be exploited to enhance resistance in other ruminants, are ambiguous. Although IFN γ is central to anti-*Toxoplasma* responses in virtually all vertebrates, the role of this cytokine, and the mechanism underpinning, bovine resistance to *Toxoplasma* is equivocal. In the present study we sought to determine the role of IFN γ , and host genes associated with effective bovine macrophage response to *Toxoplasma*. We performed RNA-sequencing on naïve or IFN γ -stimulated bovine bone marrow derived macrophages (BMDMs) that were either unexposed or exposed to a *Toxoplasma* strain that is highly virulent in laboratory inbred mice (RH) for 24 h. Analysis of the datasets provides novel insights into the *Toxoplasma*-induced transcriptional responses in naïve and pre-stimulated BMDMs. We report that IFN γ enhances bovine BMDMs anti-*Toxoplasma* responses and that, despite producing large amounts of nitric oxide, bovine macrophages restrict *Toxoplasma* independently of nitric oxide. *Toxoplasma* induces the Wnt/ β -catenin signaling pathway and the expression of several anti-inflammatory chemokines and arginine metabolism in naïve BMDMs. On the other hand, to survive in IFN γ -primed BMDMs, the parasite expresses mostly genes that are associated with its slow dividing bradyzoite stage.

MATERIALS AND METHODS

Parasites

The type I *Toxoplasma* strain (RH) engineered to express green fluorescent protein (GFP) and firefly luciferase has previously been described (Jensen et al., 2013). The RH Δ ASP5 (Hammoudi et al., 2015) was a generous gift from Dr. Mohamed-Ali Hakimi (INSERM). All parasite strains were maintained by serial passage on confluent human foreskin fibroblast (HFF) monolayer.

Primary Bone Marrow Derived Macrophages (BMDMs)

Marrow cells were flushed from the ribs of three ($n = 3$) \sim 2 year old calves using phosphate buffered saline (PBS, Invitrogen). The cells were centrifuged at $500 \times g$ for 5 min at 4°C and re-suspended in red cell lysis buffer (Sigma) and incubated on ice for 5 min. Next, the cells were passed through a 70 μ m cell strainer (BD Biosciences) and centrifuged at $500 \times g$ for 5 min at 4°C. Cells from each calf were subsequently differentiated into macrophages in 10 cm non-tissue culture petri dishes (Corning) in RPMI 1640 (Sigma-Aldrich) supplemented with heat-inactivated 20% fetal bovine serum (FBS, Thermo Fisher Scientific), GlutaMAX (Thermo Fisher Scientific), penicillin/streptomycin (Thermo Fisher Scientific), and recombinant human CSF1 (10^4 U/ml; a gift from Chiron, Emeryville, CA) for 10 days as previously described (Young et al., 2018).

In vitro Measurements

BMDMs from each calf were detached using a cell scraper, washed, counted, and seeded separately in triplicates in 96-well plates at a density of 10^5 cells/well. The BMDMs were left unstimulated (naïve) or stimulated with: recombinant bovine IFN γ (100 ng/mL), lipopolysaccharide (LPS) from *Salmonella*

enterica serotype Minnesota Re 595 (Sigma-Aldrich, 100 ng/mL), or a combination of IFN γ (100 ng/mL) and LPS (100 ng/mL) with or without aminoguanidine (final concentration of 500 μ M) and incubated at 37°C in 5% CO $_2$. 24 h post-stimulation, cell free supernatants were collected for Griess reagent-based nitric oxide assay, as previously described (Young et al., 2018), and cell viability, using the CellTiter 96[®] Aqueous One Solution Cell Proliferation kit (Promega) according to the manufacturer's recommendations. Freshly lysed parasites (by sequential passage through 25G and 27G needles) were passed through a 5 μ m filter to remove HFFs, counted, diluted in RPMI, and added to the BMDMs at a multiplicity of infection (MOI) 1 for 24 h before luciferase activity was measured using a luciferase assay kit (Promega) according to the manufacturer recommendations.

RNA Sequencing and Analysis

2×10^6 BMDMs from each calf were seeded in 6-well plates and left unstimulated or stimulated with IFN γ (100 ng/mL) for \sim 18 h at 37°C in 5% CO $_2$. Next, the media was replaced with fresh media containing *Toxoplasma* at a MOI 1 and incubated further for \sim 18 h. Fresh media was added to the non-infected control BMDMs. At the end of the incubation period, total RNA was isolated from each BMDM sample using QIAzol[®] Lysis Reagent and miRNeasy Mini Kit (Qiagen) according to the manufacturer recommendations. RNA quality and integrity were assessed on the Agilent 2200 TapeStation using an Agilent RNA ScreenTape and quantified using a Qubit RNA BR Assay Kit and Fluorometer. RNA-seq libraries were generated and sequenced by Edinburgh Genomics. All libraries were prepared using the Illumina TruSeq Stranded library protocol for total RNA libraries (Part: 15031048, Revision E). Briefly, polyA-tail enrichment (Dynabeads mRNA Purification Kit; Invitrogen) was performed on the total RNA and the mRNA fragmented into 200–400 base-pairs, and reverse transcribed into cDNA before Illumina sequencing adapters were added to each end. Twelve barcoded libraries were multiplexed and sequenced on a single S2 sequencing lane on the Illumina NovaSeq 6000 machine to yield \sim 60 million 50 bp high quality strand-specific paired reads per sample. Reads were pseudoaligned to the Ensembl bovine genome (ARS-UCD1.2) using Kallisto v.0.46.0 (Bray et al., 2016) with 100 bootstraps to generate transcript-level expression estimates as transcripts per million (TPM) as previously described (Young et al., 2018). Gene-level differential expression analysis was performed in sleuth as previously described (Pimentel et al., 2017). TPM values derived from pseudoalignment of RNA-seq reads to the GT1 genome (ToxoDB.org) were used as input in sleuth to identify differentially expressed *Toxoplasma* genes.

Quantitative Real-Time PCR

One microgram of total RNA from each BMDMs sample was reverse transcribed using Superscript III (Thermo Fisher Scientific), 10 mM dNTP mix (Thermo Fisher Scientific) and Oligo (dT) $_{12-18}$ Primer (Thermo Fisher Scientific) according to the manufacturer's instructions. Gene-specific primers (Table 1) were designed using Primer-BLAST (Ye et al., 2012). All primers were commercially synthesized by Thermo Fischer Scientific. Quantitative real-time RT-PCR (qPCR) was performed in a

TABLE 1 | Primer sets for qPCR validation of representative differentially expressed genes.

Symbol	Primer 1 forward	Primer 1 reverse	Amplicon length
<i>C1R</i>	GGTGCAGGATCAAGGACTGC	GTGTGCATCTTGTAAGGGCT	136
<i>ISG15</i>	GACCTGACGGTGAAGATGCTA	ATCTTCTGGGCGATGAAGTGC	98
<i>PSMB8</i>	TGTCATATGTACCACAT GAAGGAG	CACCATCACTGACTGGCCTC	102
<i>SGK1</i>	GCCAAGGATGACTTTATGGAGA	AGGATCAAAGTGTGCGCAGG	138
<i>IRF1</i>	AGGACATCATGAAGCTCTTGGGA	GCTCCTCCTTGCAACTGAAGT	129
<i>GBP5</i>	CCAGGAAAGGAATACAGGCTGA	TTCCATTGCTGTGAGAGCCAG	107
<i>RSAD2</i>	GTGGTTCCAGAAGTACGGTGA	CTTCTTTCCTTGACCACGGC	103
<i>CFB</i>	CTTGCAAAGGTGATTCTGGTGG	CGCTTGCAAACATCCACGAC	100
<i>GBP1</i>	CTCTCAAATGCAGGAACAGTC	TGCTTTGGATAAGAGT GACCAG	175
<i>CXCL10</i>	TCCTCGAACACGGAAAGA GGCATA	AGCTGATGGTGACTGG CTTGGT	164
<i>CCL22</i>	CGGGACTACATCCGTTACCC	CAGCACAGATCTCTCGGTCC	121
<i>CCl24</i>	GCAGGAGTGATCTTACCACC	TAGCGGAGGCTTTCTTCTGC	115
<i>SDHA</i>	ACCTGATGCTTTGTCTCTGC	CCTGGATGGGCTTGGAGTAA	126
<i>GAPDH</i>	GGTGATGCTGGTCTGAGTA	TCATAAGTCCCTCCACGATG	265

Rotor-Gene Q real-time PCR cyler (Qiagen) using FastStart Universal SYBR Green Master (Rox) in final volume of 20 μ L. The linearity and efficiency of qPCR amplification was determined for each primer pair using a standard curve generated by a serial dilution of cDNA pooled from all the samples. All reactions were performed in duplicate and “no template” controls included for each gene. Agarose gel electrophoresis was used to confirm product sizes and melt curve analysis confirmed specificity of amplification. Data were analyzed using the $2^{-\Delta\Delta C_t}$ method and statistical analyzed performed on the ΔC_t values. qPCR was performed on parasite DNA as previously described (Li et al., 2000).

RESULTS

IFN γ Induces Restriction of *Toxoplasma* Growth in Bovine Macrophages Independently of Nitric Oxide


To determine whether IFN γ is essential for the control of *Toxoplasma* growth in bovine macrophages, we exposed naïve or IFN γ -stimulated bovine bone marrow-derived macrophages (BMDMs) to a luciferase-expressing type I *Toxoplasma* strain (RH) for 24 h and measured luciferase activity, a proxy for parasite burden (Hassan et al., 2015). In murine macrophages, IFN γ is known to require a second stimulant, such as LPS or TNE, to effectively restrict *Toxoplasma* (Sibley et al., 1991; Hassan et al., 2015). Therefore, we included in our experiments, BMDMs that were pre-stimulated with IFN γ +LPS. As expected, there was reduced intracellular parasite burden in BMDMs pre-stimulated with IFN γ or IFN γ +LPS, relative to naïve BMDMs (Figure 1A). However, there were no significant differences in parasite burden between IFN γ - and IFN γ +LPS-stimulated BMDMs, suggesting that IFN γ alone can sufficiently induce anti-*Toxoplasma* activities in bovine BMDMs. A quantitative

real-time polymerase chain reaction (qRT-PCR) analysis of parasite DNA (Li et al., 2000) confirmed the intracellular parasite burden differences between naïve and IFN γ -stimulated BMDMs (Figure 1B).

Previously, others and we reported that Nitric oxide (NO), a product of the inducible nitric oxide synthase (iNOS or NOS2)-catalyzed L-arginine metabolism (MacMicking et al., 1997), is a major IFN γ -induced effector against intracellular *Toxoplasma* growth in murine BMDMs (Hassan et al., 2015). In addition, bovine BMDMs stimulated with LPS and/or IFN γ are reported to produce significantly more NO than ovine or equine BMDMs (Denis et al., 2005; Young et al., 2018), which are susceptible to *Toxoplasma*. Thus, we postulated that NO inhibits *Toxoplasma* growth in bovine IFN γ -stimulated BMDMs. Consequently, we determined whether IFN γ alone can induce NO production in bovine BMDMs and whether the levels of the induced NO correlate with intracellular parasite burden in the pre-stimulated BMDMs. Unlike IFN γ , IFN γ +LPS induced a 34-fold increase in NO, relative to unstimulated BMDMs (Figure 1C). Thus, unlike murine BMDMs, NO is dispensable for IFN γ -induced *Toxoplasma* growth restriction in bovine BMDMs.

Toxoplasma Induces a Robust Bovine Macrophage Transcriptional Response

To gain mechanistic insight into bovine innate immune response to *Toxoplasma*, we leveraged high throughput RNA-sequencing to profile the transcriptional landscape of bovine BMDMs that were left unstimulated (naïve) or pre-stimulated with IFN γ before being exposed to *Toxoplasma* for 24 h. Transcriptional analysis was also performed on uninfected controls. Downstream analysis was restricted to genes that were differentially expressed by more than 2-fold change and had at least 10 reads that uniquely aligned to the genome in at least two samples when compared the uninfected naïve BMDMs. In total, 1,349 unique genes were differentially expressed in at least one condition, of which 887 were differentially expressed in a condition-specific manner (Figure 2A and Supplementary Table 1). Unlike IFN γ that interacts mainly with the IFN γ receptors to induce the expression of several IFN γ -specific genes (ISGs), *Toxoplasma* is likely to interact with many, yet to be defined, bovine BMDM pattern recognition receptors to induce bovine gene expression. Indeed, there were more differentially expressed genes that were unique to naïve BMDMs exposed to *Toxoplasma* (RH-specific) than in the uninfected IFN γ -stimulated BMDMs, (657 vs. 134, respectively; Hypergeometric P -value ≤ 0.05). Interestingly, only 96 genes were differentially expressed exclusively in IFN γ -stimulated BMDMs exposed to *Toxoplasma* (IRH-specific), suggesting that pre-stimulation with IFN γ abrogates a majority of *Toxoplasma*-induced transcriptional changes in the BMDMs (Figure 2A). The variable expression of most genes (579/657, $\sim 88\%$), including immunoregulatory genes such as Krüppel-like factor 4 (*KLF4*) and Ornithine decarboxylase (*ODCI*), in the RH BMDMs was due to increased transcript abundance in the *Toxoplasma*-exposed BMDMs, rather than a downregulation of innately expressed genes, as revealed by hierarchical clustering




(Figure 2B), suggesting that effective response to the parasite is inducible. Three hundred and forty genes, including several inflammatory cytokines such C-X-C motif chemokine ligand 10 (*CXCL10*) that were differentially expressed in unexposed IFN γ -stimulated BMDMs, remained dysregulated by a similar fold-change magnitude when the IFN γ -stimulated BMDMs were exposed to *Toxoplasma* (IRH), suggesting an inability by the parasite to overcome most of the IFN γ -induced transcriptional changes. Arginase 2 (*ARG2*) was among 13 genes that were highly expressed in RH but downregulated in IFN γ -stimulated BMDMs. Similarly, 60 genes, including the C-C motif chemokine 22 (*CCL22*), were upregulated in RH but downregulated by a difference of more than 2-fold change between in RH and IRH BMDMs.

To gain greater insights into the biological processes underpinning bovine BMDMs transcriptional response to *Toxoplasma*, and to determine whether the transcriptional changes in the different BMDM conditions was due to differences in the induction of distinct gene sets or the magnitude of induction of the same gene sets, we performed a pre-ranked gene set enrichment analysis (GSEA) (Subramanian et al., 2005) on the differentially expressed genes in IFN γ -stimulated, IRH, and RH BMDMs using the curated “HALLMARK” gene set database (Liberzon et al., 2015). Several “HALLMARK” gene sets, including the interferon gamma response, inflammatory response, and the regulation of cytokine production, were significantly (*FDR* \leq 0.05) enriched in at least two BMDM conditions. Some gene sets, including hypoxia, which is not only a well-established *Toxoplasma*-induced host cell response, but also supports intracellular parasite growth (Wiley et al., 2010), were significantly enriched in RH but absent in IRH (Figure 2C). Thus, infection of naïve and IFN γ -stimulated

BMDMs modulated by differences in the induction of unique and common gene sets.

IFN γ Reverses *Toxoplasma*-Induced Dysregulation of Genes Linked to Inflammatory and Metabolic Pathways to Restrict Parasite Replication in Bovine Macrophages


As demonstrated above, *Toxoplasma* significantly dysregulate several bovine genes, potentially to support its intracellular survival. We reasoned that since IFN γ -stimulated BMDMs are refractory to the parasite, *Toxoplasma*-induced genes that support the intracellular parasite lifestyle are likely to be targeted and reversed when BMDMs are pre-stimulated with IFN γ prior to infection. Several inflammatory genes were dysregulated in IRH but not in RH BMDMs. For example, *CXCL9*, *10*, and *11*, were 22-, 10-, and 12-fold, respectively, upregulated in IRH, but not dysregulated in RH, BMDMs. Markers for the *Toxoplasma*-refractory classically activated macrophages, such as *CD180*, *CD74*, and *CD1D* were highly expressed in IRH but not in RH BMDMs. Similarly, c-type lectins including *CLEC6A*, which are associated with classically activated macrophages (Jensen et al., 2011; Murray et al., 2014), were upregulated in IRH BMDMs. Conversely, calcium signaling genes that are reported to play a significant role in intracellular parasite survival, such as *CAMK2G*, and histone deacetylase (*HDAC*) 5, and 7, were downregulated in RH, but remained unchanged in IRH BMDMs. A functional analysis on genes upregulated or unchanged in IRH BMDMs revealed an enrichment for, among others, calcium signaling pathway, phospholipase D signaling, defense response



to virus, and positive regulation of I- κ B kinase/NF κ B signaling (Figure 3A).

Genes implicated in chemotaxis or anti-inflammatory responses were differentially expressed by a margin of more than 2-fold change between RH and IRH BMDMs. For example, chemotaxis-related *CCL22* and *CCL24* chemokines were upregulated by 20- and 12-fold in RH but upregulated by 14- and 8-fold in IRH BMDMs. Similarly, the chemokine receptor (*CCR7*) was upregulated by 26-fold in RH but not differentially expressed in IRH BMDMs. The anti-inflammatory-related suppressor of cytokine signaling (*SOCS* 3), and TNF receptor-associated factor 1 (*TRAF1*) were also upregulated in RH but downregulated in IRH BMDMs. Mitogen-activated protein kinases (MAPK), including *MAP3K14* and *MAP4K1*, which induce non-canonical (Jin et al., 2014) and suppresses canonical (Brenner et al., 2005) NF- κ B signaling pathways, respectively, were upregulated in RH but downregulated in IRH BMDMs. Studies in model animals, including mice, have shown that *Toxoplasma* induces alternative macrophage activation to favor its intracellular lifestyle (Melo et al., 2011). Arginine metabolism, via arginase, is key to alternative macrophage activation. Congruently, Arginase 2 (*ARG2*) was 4-fold upregulated in RH but virtually not expressed in IRH BMDMs. Additional genes

associated with alternative macrophage activation, such as *KLF4* and *IL34* were upregulated in RH but downregulated in IRH BMDMs. The expression of protein arginine methyltransferase 5 (*PRMT5*), which is necessary in c-MYC-mediated alternative macrophage differentiation, was slightly increased in RH BMDMs. Similarly, genes encoding glucose transporters were upregulated in RH but not IRH BMDMs. Solute carriers (*SLC*), including *SLC2A1* and *SLC2A3* that encode glucose transporters, were downregulated in IRH but upregulated in RH BMDMs. Other solute carriers, including *SLC7A1* and *SLC7A5* that function in the L-arginine transport pathway, and *ODC1*, the rate limiting enzyme in polyamine biosynthesis and a negative regulator of macrophage inflammation (Hardbower et al., 2017), were upregulated in RH BMDMs. *Toxoplasma* reportedly rely on host lipid droplets, which increase in abundance in *Toxoplasma*-infected cells, to sustain rapid intracellular replication (Gomes et al., 2014). The expression of pentraxin 3 (*PTX3*), which is induced by lipid accumulation (Liu et al., 2014), was increased 8-fold in RH but unexpressed in IRH BMDMs. Similarly, the expression of genes involved in fatty acid elongation, such as *ELOVL6*, were upregulated in RH but not expressed in IRH BMDMs. Genes upregulated in RH but not IRH BMDMs were significantly ($FDR \leq 0.05$) linked to purine



metabolism, NIK/NF κ B (non-canonical NF κ B) signaling, and regulation of apoptotic signaling pathway (Figure 3B), all of which are expected to play a role in regulating *Toxoplasma*'s intracellular fate (Yamamoto et al., 2011; Blume and Seeber, 2018). Quantitative RT-PCR (qPCR) confirmed the differential expression of genes, including *CXCL10*, *CCL22*, and *CCL24* that are reportedly upregulated in *Toxoplasma*-infected murine macrophages (Melo et al., 2013), in the different macrophage conditions (Figure 3C and Supplementary Figure 1). Thus, IFN γ targets both the host cell inflammatory and metabolic pathways to revoke intracellular parasite replication.

IFN γ Abrogates *Toxoplasma*-Induced *Wnt*/ β -Catenin Signaling in Bovine Macrophages

Recent studies have documented *Toxoplasma*-induced expression of several chemokines, including *CCL17*, *CCL22*, and *CCL24* (He et al., 2018; Majumdar et al., 2019) via the activation of the *Wnt*/ β -catenin signaling pathway (He et al., 2018). *Wnt*/ β -catenin-signaling has an important role in various cellular processes, including proliferation (Reya et al., 2003) and immunity (Staal et al., 2008). The variable expression of *CCL22* and *CCL24*, in RH and IRH BMDMs, prompted us to investigate the expression of *Wnt*-target genes to establish a role of *Wnt*/ β -catenin signaling in bovine BMDM response to *Toxoplasma*. However, apart from the downregulation of delta like canonical Notch ligand 1 (*DLL1*) in IFN γ -stimulated BMDMs, none of the

canonical *Wnt*-target genes were differentially expressed in RH and/or IRH BMDMs.

Unlike *Toxoplasma*, IFN γ is known to suppress *Wnt*/ β -catenin signaling (Nava et al., 2014; Bai et al., 2017). To determine whether *Toxoplasma* growth restriction in IFN γ -stimulated bovine BMDMs is linked to *Wnt*/ β -catenin signaling, we evaluated parasite burden in naïve or IFN γ -stimulated BMDMs cultured in the presence or absence of a *Wnt*/ β -catenin antagonist (IWR-1-endo). IWR-1-endo reduced parasite burden in naïve BMDMs in a dose-dependent manner without abolishing the differences in parasite burden between naïve and IFN γ -stimulated BMDMs (Figure 4A). *Toxoplasma*-induced accumulation of β -catenin in the host cell nucleus, and the dysregulation of *CCL22* is reportedly induced by a dense granule (GRA) protein (GRA18) that is secreted beyond the parasites' parasitophorous vacuole membrane (PVM) into the host cell during infection (He et al., 2018). Thus, to overcome the potential effect of IWR-1-endo on unrelated host cell functions and establish a direct link between infection and *Wnt*/ β -catenin activation, we infected naïve or IFN γ -stimulated BMDMs with a mutant *Toxoplasma* strain (RH Δ ASP5) and quantified parasite burden using qPCR. The RH Δ ASP5 mutant strain lacks the ASP5 enzyme that is essential for the processing of GRA proteins, including GRA18, that are destined for secretion into the host cell (Coffey et al., 2015; Hammoudi et al., 2015; Curt-Varesano et al., 2016; He et al., 2018). We reasoned that if GRAs that are secreted in the host cell interact with β -catenin to induce chemokine secretion and support intracellular parasite survival

and replication in bovine BMDMs, then the RH Δ ASP5 strain should not be able to replicate efficiently in naïve BMDMs. Indeed, parasite burden was significantly reduced in naïve BMDMs infected with RH Δ ASP5, relative to the wildtype (Figure 4B). Unlike wildtype parasites, IWR-1-endo did not significantly alter parasite burden in naïve or IFN γ -stimulated BMDMs infected with RH Δ ASP5 (Figure 4C). Neither did IWR-1-endo abrogate intracellular RH Δ ASP5 burden differences between naïve and IFN γ -stimulated BMDMs. Finally, when infected with RH Δ ASP5, but not the wildtype parental strain, naïve BMDMs exhibited high abundance for *CCL22* transcripts (Figure 4D). Together, these results indicate a role for *Wnt*/ β -catenin signaling and parasite secreted GRAs that translocate to the host cell in bovine BMDMs anti-*Toxoplasma* responses.

Parasites in IFN γ -Stimulated Bovine Macrophages Transcribe Mostly Bradyzoite-Related Genes

To evaluate whether the infection of bovine BMDMs is underpinned by distinct parasite expression signatures, we used the RNA-sequencing reads uniquely aligning to the parasite genome (GT1 v.46) to evaluate stochastic changes in *Toxoplasma* transcript abundance in naïve and IFN γ -stimulated BMDMs. We focused our analysis on genes that were modulated by more than 3-fold change and had more than 10 uniquely aligned RNA-sequencing reads in at least two samples when comparing parasites infecting naïve and IFN γ -stimulated BMDMs. One hundred and eighty one genes were differentially expressed, of which 94 corresponded to genes upregulated in IFN γ -stimulated BMDMs (Figure 5 and Supplementary Table 1). Parasite genes that were highly expressed when the parasite is in IFN γ -stimulated BMDMs included DnaK-tetratricopeptide repeat (*DnaK-TPR*) and Cyclic AMP-Dependent Protein Kinase Subunit 3 (*cAMPK3*), which are associated with or involved in stress-induced *Toxoplasma* stage conversion from the fast dividing tachyzoite to the semi-dominant bradyzoite parasite stage (Ueno et al., 2011; Sugi et al., 2016). Thus, consistent with the well-defined immune pressure-induced parasite stage conversion, *Toxoplasma* potentially responds to the IFN γ -induced immune pressure by slowing down its rate of replication.


DISCUSSION

The outcome of *Toxoplasma* infections in cattle is fundamentally different from that of rodents, small ruminants, and pigs: clinical disease in naturally infected cattle are rare and many large-scale studies have failed to detect viable parasites in bovine tissues (Stelzer et al., 2019), suggesting that, like humans, cattle are largely an end-stage host for the parasite. However, the mechanisms underlying bovine resistance to *Toxoplasma* are largely unknown. In this study, we performed high throughput functional genomics and parasitological assays to investigate the molecular factors that modulate *Toxoplasma* interactions with bovine bone marrow-derived macrophages (BMDMs). Similar to observations in rodent and human macrophages, interferon gamma (IFN γ) significantly enhanced intracellular

parasite growth restriction in bovine BMDMs. Interestingly, despite the ability to restrict intracellular parasite growth, IFN γ on its own was not able to induce nitric oxide (NO) secretion in bovine BMDMs, indicating that, unlike murine BMDMs, IFN γ -induced toxoplasmatostatic activities in bovine BMDMs are independent of NO. Additionally, we found that secreted parasite dense granule proteins potentially induce the expression of chemokines, including *CCL22*, to enhance parasite growth in naïve bovine BMDMs.


Understanding the mechanisms of innate recognition and response to *Toxoplasma* in mice has been invaluable in advancing knowledge on response to *Toxoplasma* in other vertebrates, including cattle. The IFN γ -induced murine anti-*Toxoplasma* effectors, which are largely dependent on the activation of immunity-related GTPases (IRGs) and guanylate binding proteins (GBPs), are well-characterized. However, cattle, like humans, lack functional toll-like receptor (TLR) 11 and 12 (O'Neill et al., 2013) that facilitate the recognition of *Toxoplasma*-derived pathogen-associated molecular patterns (PAMPs) in murine cells (Melo et al., 2010). Although bovine BMDMs express a functional TLR5, unlike murine TLR5, it does not recognize *Toxoplasma* antigens (Tombácz et al., 2018). Thus, the mechanisms by which bovine cells recognize and respond to *Toxoplasma* are equivocal and potentially different from those defined in the murine system. Previous studies, including our own (Jensen et al., 2013; Hassan et al., 2015), show that reactive oxygen and nitrogen species, including NO, are central to *Toxoplasma* growth inhibition in murine macrophages synergistically activated with IFN γ and LPS or TNF. Bovine macrophages are known to secrete significantly more NO in response to LPS than sheep and pigs, which are susceptible to *Toxoplasma*, LPS (Jungi et al., 1996; Young et al., 2018), suggesting that species-specific differences in susceptibility to *Toxoplasma* is due to differences in NO production. Although parasite growth was significantly restricted in stimulating bovine BMDMs with IFN γ -stimulated bovine BMDMs, IFN γ , on its own, did not induce NO in bovine BMDMs, which is consistent with previous observations (Denis et al., 2005; Young et al., 2018; Imrie and Williams, 2019). In fact, intracellular parasite burden in IFN γ - and IFN γ +LPS-stimulated BMDMs were similar, despite the latter secreting significantly more NO. Combined, these results suggest that NO is dispensable in IFN γ -induced bovine BMDM toxoplasmatostatic activities. A similar non-essential role for NO in macrophage response to *Toxoplasma* has previously been observed in human macrophages. However, human macrophages are distinct from bovine BMDMs, since NO cannot be induced by conventional activating regimes *in vitro* in human macrophages (Schneemann and Schoedon, 2002).

Recent studies have documented a role for *Wnt*/ β -catenin signaling in host responses to *Toxoplasma*. Enhanced intracellular *Toxoplasma* growth was found to co-occur with increased β -catenin (*CTNNB1*) gene expression in human cells (Majumdar et al., 2019), while a parasite secreted GRA protein is reported to stabilize β -catenin in murine BMDMs leading to an increased expression of chemokines such as *CCL22* (He et al., 2018; Majumdar et al., 2019). Consistent with IFN γ -induced inhibition of the *Wnt*/ β -catenin signaling



pathway (Nava et al., 2014; Bai et al., 2017), we observed a downregulation of several *Wnt*/β-catenin signaling-associated genes, including *CCL17*, *CCL22*, and *CCL24* (He et al., 2018),

in IFN γ -stimulated BMDMs. Congruently, a *Wnt*/β-catenin signaling chemical inhibitor (IWR-1-endo) reduced intracellular parasite burden in naïve BMDMs in a dose-dependent manner.



However, IWR-1-endo did not abolish intracellular parasite burden differences between naïve and IFN γ -stimulated BMDMs, probably because IFN γ suppresses *Wnt*/ β -catenin signaling to a level below which further inhibition produces no discernible effect. Considering that we used a single IFN γ concentration (100 ng/mL), it is plausible that IWR-1-endo would have an effect at lower IFN γ concentrations. A more plausible explanation for the lack of IWR-1-endo efficacy in IFN γ -stimulated cells is that besides the *Wnt*/ β -catenin signaling, IFN γ induces additional effector mechanisms against the parasite. Although GRA18 is reported to induce β -catenin accumulation in the host cell nucleus, this was not discernible when GRA18 was expressed at physiological levels (He et al., 2018), which may partly explain the lack of β -catenin nuclear accumulation in the present study. Nevertheless, intracellular parasite burden in bovine BMDMs infected with a mutant *Toxoplasma* strain that does not secrete GRA18 into the host cell cytoplasm (RH Δ ASP5) (He et al., 2018), was significantly

different from cells infected the wildtype parasites. Similar to IWR-1-endo, infection with RH Δ ASP5 did not abolish intracellular parasite burden differences between naïve and IFN γ -stimulated BMDMs.

The mechanisms regulating the impact of *Wnt*/ β -catenin on intracellular *Toxoplasma* replication are largely equivocal. Recent studies indicate that β -catenin alters intracellular parasite growth dynamics by interacting with indoleamine 2,3-dioxygenase 1 (IDO1): IDO1 promoter activity is supported by β -catenin. IDO1 is an IFN γ -inducible protein that degrades tryptophan to kynurenine and is known to impede *Toxoplasma* growth in human fibroblasts (Pfefferkorn, 1984), epithelial, and endothelial cells (MacKenzie et al., 2007) *in vitro*. Because IDO1 degrades tryptophan, which supports intracellular *Toxoplasma* replication, increased IDO1 promoter activity in the presence of β -catenin may appear contradictory. However, IDO1 transcript and protein levels are often discordant in *Toxoplasma*-infected cells: *Toxoplasma* promotes the degradation of IDO1 protein

(Majumdar et al., 2019). Although we cannot conclude, from this study, that tryptophan plays an essential role in bovine BMDM response to *Toxoplasma*, *IDO1* was significantly downregulated in naïve BMDMs exposed to *Toxoplasma*. In the absence of *IDO1*, tryptophan is catabolized to melatonin, which scavenges reactive oxygen species (ROS) and promotes cell survival (Dolado and Nebreda, 2008), both of which are beneficial for intracellular *Toxoplasma* survival. However, considering that NO and ROS are dispensable for *Toxoplasma* growth restriction in bovine BMDMs, scavenging of ROS is unlikely to be a viable mechanism by which β -catenin can affect parasite replication. Rather, it is plausible that melatonin promotes cell survival, which in turn promotes intracellular parasite survival. Indeed, we did not observe significant changes in cell viability in *Toxoplasma*-infected, relative to uninfected, BMDMs. Alternatively, *Wnt*/ β -catenin-mediated anti-*Toxoplasma* mechanisms may involve the induction of alternative macrophage activation, which is supportive of intracellular *Toxoplasma* survival and growth (Melo et al., 2011). Besides, the expression of *Wnt*/ β -catenin-associated chemokines, we observed increased expression of genes associated with alternatively activated or foamy macrophages, such as *KLF4*, *ARG2*, *ODC1*, and *PTX3*, in naïve, but not IFN γ -stimulated, BMDMs. Combined, with the increased expression of glucose transporters in naïve BMDMs, which is reversed in IFN γ -stimulated BMDMs, we conclude that *Toxoplasma* exploits the bovine BMDMs metabolic pathways to enhance intracellular survival and replication. It is worth noting that we have tested only one clonal parasite strain (RH) at a single MOI and that different *Toxoplasma* strains, including the genetically distinct atypical strains such as GUYDOS, and higher MOIs may induce significantly different responses in the bovine BMDMs.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are publicly available. This data can be found here: <https://www.ncbi.nlm.nih.gov/PRJNA646376>.

REFERENCES

- Bai, M., Li, W., Yu, N., Zhang, H., Long, F., and Zeng, A. (2017). The crosstalk between β -catenin signaling and type I, type II and type III interferons in lung cancer cells. *Am. J. Transl. Res.* 9, 2788–2797.
- Basavaraju, A. (2016). Toxoplasmosis in HIV infection: an overview. *Trop. Parasitol.* 6, 129–135. doi: 10.4103/2229-5070.190817
- Blume, M., and Seeber, F. (2018). Metabolic interactions between *Toxoplasma gondii* and its host. *F1000Res.* 7:F1000. doi: 10.12688/f1000research.16021.1
- Bougourd, A., Durandau, E., Brenier-Pinchart, M. P., Ortet, P., Barakat, M., Kieffer, S., et al. (2013). Host cell subversion by *Toxoplasma* GRA16, an exported dense granule protein that targets the host cell nucleus and alters gene expression. *Cell Host Microbe* 13, 489–500. doi: 10.1016/j.chom.2013.03.002
- Bray, N. L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic RNA-seq quantification. *Nat. Biotechnol.* 34, 525–527. doi: 10.1038/nbt.3519

ETHICS STATEMENT

The animal study was reviewed and approved by the Protocols and Ethics Committees of The Roslin Institute, The University of Edinburgh, and the Royal (Dick) School of Veterinary Medicine. In accordance with the United Kingdom Animal (Scientific Procedures) Act 1986, this study did not require a Home Office project license as no regulated procedures were carried out.

AUTHOR CONTRIBUTIONS

MH conceived and designed the experiments and wrote the manuscript. MH and AG performed the experiments and analyzed the data. All authors contributed to the article and approved the submitted version.

FUNDING

MH was funded by a University of Edinburgh Chancellor's Fellowship and a Bill and Melinda Gates Foundation award to the Centre for Tropical Livestock Genetics and Health (OPP1127286). The Roslin Institute receives strategic investment funding from the Biotechnology and Biological Sciences.

ACKNOWLEDGMENTS

The authors thank the University of Edinburgh Genomics Core for providing facilities and services.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fcimb.2020.00437/full#supplementary-material>

Supplementary Figure 1 | Quantitative real-time polymerase chain reaction (qPCR) of some differentially expressed genes in naïve infected (RH; black bars), IFN γ -stimulated and infected (IRH; blue bars), and IFN γ -stimulated (red bars), unstimulated-uninfected control (open bars) BMDMs. Data are average value \pm s.d. of three replicates. Data are representative of two independent experiments.

- Brenner, D., Golks, A., Kiefer, F., Krammer, P. H., and Arnold, R. (2005). Activation or suppression of NF κ B by HPK1 determines sensitivity to activation-induced cell death. *EMBO J.* 24, 4279–4290. doi: 10.1038/sj.emboj.7600894
- Channon, J. Y., Seguin, R. M., and Kasper, L. H. (2000). Differential infectivity and division of *Toxoplasma gondii* in human peripheral blood leukocytes. *Infect. Immun.* 68, 4822–4826. doi: 10.1128/IAI.68.8.4822-4826.2000
- Cirelli, K. M., Gorf, G., Hassan, M. A., Printz, M., Crown, D., Leppla, S. H., et al. (2014). Inflammasome sensor NLRP1 controls rat macrophage susceptibility to *Toxoplasma gondii*. *PLoS Pathog.* 10:e1003927. doi: 10.1371/journal.ppat.1003927
- Coffey, M. J., Sleeb, B. E., Uboldi, A. D., Garnham, A., Franco, M., Marino, N. D., et al. (2015). An aspartyl protease defines a novel pathway for export of *Toxoplasma* proteins into the host cell. *elife* 4:e10809. doi: 10.7554/eLife.10809.025
- Curt-Varesano, A., Braun, L., Ranquet, C., Hakimi, M. A., and Bougourd, A. (2016). The aspartyl protease TgASP5 mediates the export of the *Toxoplasma*

- GRA16 and GRA24 effectors into host cells. *Cell Microbiol.* 18, 151–167. doi: 10.1111/cmi.12498
- De Paoli, F., Stael, B., and Chinetti-Gbaguidi, G. (2014). Macrophage phenotypes and their modulation in atherosclerosis. *Circ. J.* 78, 1775–1781. doi: 10.1253/circj.CJ-14-0621
- Denis, M., Wedlock, D. N., and Buddle, B. M. (2005). IFN- γ enhances bovine macrophage responsiveness to *Mycobacterium bovis*: impact on bacterial replication, cytokine release and macrophage apoptosis. *Immunol. Cell Biol.* 83, 643–650. doi: 10.1111/j.1440-1711.2005.01386.x
- Dolado, I., and Nebreda, A. R. (2008). AKT and oxidative stress team up to kill cancer cells. *Cancer Cell* 14, 427–429. doi: 10.1016/j.ccr.2008.11.006
- Dubey, J. P. (1986). A review of toxoplasmosis in cattle. *Vet. Parasitol.* 22, 177–202. doi: 10.1016/0304-4017(86)90106-8
- Dunay, I. R., and Sibley, L. D. (2010). Monocytes mediate mucosal immunity to *Toxoplasma gondii*. *Curr. Opin. Immunol.* 22, 461–466. doi: 10.1016/j.coi.2010.04.008
- Esteban-Redondo, I., and Innes, E. A. (1997). *Toxoplasma gondii* infection in sheep and cattle. *Comp. Immunol. Microbiol. Infect. Dis.* 20, 191–196. doi: 10.1016/S.0147-9571(96)00039-2
- FAO/WHO (2014). *Multicriteria-Based Ranking for Risk Management of Food-Borne Parasites*. Available online at: <http://www.fao.org/3/a-i3649e.pdf> %5Cnfiles/540/a-i3649e.pdf (accessed August 9, 2020).
- Fleckenstein, M. C., Reese, M. L., Konen-Waisman, S., Boothroyd, J. C., Howard, J. C., and Steinfield, T. (2012). A *Toxoplasma gondii* pseudokinase inhibits host IRG resistance proteins. *PLoS Biol.* 10:e1001358. doi: 10.1371/journal.pbio.1001358
- Gazzinelli, R. T., Hieny, S., Wynn, T. A., Wolf, S., and Sher, A. (1993). Interleukin 12 is required for the T-lymphocyte-independent induction of interferon γ by an intracellular parasite and induces resistance in T-cell-deficient hosts. *Proc. Natl. Acad. Sci. U.S.A.* 90, 6115–6119. doi: 10.1073/pnas.90.13.6115
- Gazzinelli, R. T., Wysocka, M., Hayashi, S., Denkers, E. Y., Hieny, S., Caspar, P., et al. (1994). Parasite-induced IL-12 stimulates early IFN- γ synthesis and resistance during acute infection with *Toxoplasma gondii*. *J. Immunol.* 153, 2533–2543.
- Gkogka, E., Reij, M. W., Havelaar, A. H., Zwietering, M. H., and Gorris, L. G. M. (2011). Risk-based estimate of effect of foodborne diseases on public health, Greece. *Emerg. Infect. Dis.* 17, 1581–1590. doi: 10.3201/eid1709.101766
- Gomes, A. F., Magalhães, K. G., Rodrigues, R. M., De Carvalho, L., Molinaro, R., Bozza, P. T., et al. (2014). *Toxoplasma gondii*-skeletal muscle cells interaction increases lipid droplet biogenesis and positively modulates the production of IL-12, IFN- γ and PGE 2. *Parasit. Vectors* 7:47. doi: 10.1186/1756-3305-7-47
- Gregg, B., Taylor, B. C., John, B., Tait-Wojno, E. D., Girgis, N. M., Miller, N., et al. (2013). Replication and distribution of *Toxoplasma gondii* in the small intestine after oral infection with tissue cysts. *Infect. Immun.* 81, 1635–1643. doi: 10.1128/IAI.01126-12
- Hammoudi, P. M., Jacot, D., Mueller, C., Di Cristina, M., Dogga, S. K., Marq, J. B., et al. (2015). Fundamental roles of the golgi-associated *Toxoplasma* aspartyl protease, ASP5, at the host-parasite interface. *PLoS Pathog.* 11:e1005211. doi: 10.1371/journal.ppat.1005211
- Hardbower, D. M., Asim, M., Luis, P. B., Singh, K., Barry, D. P., Yang, C., et al. (2017). Ornithine decarboxylase regulates M1 macrophage activation and mucosal inflammation via histone modifications. *Proc. Natl. Acad. Sci. U.S.A.* 114, E751–E760. doi: 10.1073/pnas.1614958114
- Hassan, M. A., Jensen, K. D., Butty, V., Hu, K., Boedec, E., Prins, P., et al. (2015). Transcriptional and linkage analyses identify loci that mediate the differential macrophage response to inflammatory stimuli and infection. *PLoS Genet.* 11:e1005619. doi: 10.1371/journal.pgen.1005619
- He, H., Brenier-Pinchart, M.-P. P., Braun, L., Kraut, A., Touquet, B., Couté, Y., et al. (2018). Characterization of a *Toxoplasma* effector uncovers an alternative GSK3 β -catenin-regulatory pathway of inflammation. *elife* 7:e39887. doi: 10.7554/eLife.39887.027
- Hill, D., and Dubey, J. P. (2002). *Toxoplasma gondii*: transmission, diagnosis, and prevention. *Clin. Microbiol. Infect.* 8, 634–640. doi: 10.1046/j.1469-0691.2002.00485.x
- Hunter, C. A., and Sibley, L. D. (2012). Modulation of innate immunity by *Toxoplasma gondii* virulence effectors. *Nat. Rev. Microbiol.* 10, 766–778. doi: 10.1038/nrmicro2858
- Imrie, H., and Williams, D. J. L. (2019). Stimulation of bovine monocyte-derived macrophages with lipopolysaccharide, interferon- γ , Interleukin-4 or Interleukin-13 does not induce detectable changes in nitric oxide or arginase activity. *BMC Vet. Res.* 15:45. doi: 10.1186/s12917-019-1785-0
- Jensen, K., Kenneth, H., Whitmarsh, R., Hassan, M. A., Julien, L., Lu, D., et al. (2013). *Toxoplasma gondii* rhoptry 16 kinase promotes host resistance to oral infection and intestinal inflammation only in the context of the dense granule protein GRA15. *Infect. Immun.* 81, 2156–2167. doi: 10.1128/IAI.01185-12
- Jensen, K. D. C., Wang, Y., Wojno, E. D. T., Shastri, A. J., Hu, K., Cornel, L., et al. (2011). *Toxoplasma* polymorphic effectors determine macrophage polarization and intestinal inflammation. *Cell Host Microbe* 9, 472–483. doi: 10.1016/j.chom.2011.04.015
- Jin, J., Hu, H., Li, H. S., Yu, J., Xiao, Y., Brittain, G. C., et al. (2014). Noncanonical NF- κ B pathway controls the production of type I interferons in antiviral innate immunity. *Immunity* 40, 342–354. doi: 10.1016/j.immuni.2014.02.006
- Jungi, T. W., Adler, H., Adler, B., Thöny, M., Krampe, M., and Peterhans, E. (1996). Inducible nitric oxide synthase of macrophages. Present knowledge and evidence for species-specific regulation. *Vet. Immunol. Immunopathol.* 54, 323–330. doi: 10.1016/S0165-2427(96)05690-5
- Koshy, A. A., Dietrich, H. K., Christian, D. A., Melehani, J. H., Shastri, A. J., Hunter, C. A., et al. (2012). *Toxoplasma* co-opts host cells it does not invade. *PLoS Pathog.* 8:e1002825. doi: 10.1371/journal.ppat.1002825
- Li, M. H., Che, T. C., Kuo, T. T., Tseng, C. C. P., and Tseng, C. C. P. (2000). Real-time PCR for quantitative detection of *Toxoplasma gondii*. *J. Clin. Microbiol.* 38, 4121–4125. doi: 10.1128/JCM.38.11.4121-4125.2000
- Liberzon, A., Birger, C., Thorvaldsdóttir, H., Ghandi, M., Mesirov, J. P., and Tamayo, P. (2015). The molecular signatures database hallmark gene set collection. *Cell Syst.* 1, 417–425. doi: 10.1016/j.cels.2015.12.004
- Liu, W., Jiang, J., Yan, D., Li, D., Li, W., Ma, Y., et al. (2014). Pentraxin 3 promotes oxLDL uptake and inhibits cholesterol efflux from macrophage-derived foam cells. *Exp. Mol. Pathol.* 96, 292–299. doi: 10.1016/j.yexmp.2014.03.007
- MacKenzie, C., Hessler, K., Muller, A., and Daubener, W. (2007). Role of indoleamine 2,3-dioxygenase in antimicrobial defence and immunoregulation: tryptophan depletion versus production of toxic kynurenines. *Curr. Drug Metab.* 8, 237–244. doi: 10.2174/138920007780362518
- MacMicking, J., Xie, Q. W., and Nathan, C. (1997). Nitric oxide and macrophage function. *Annu. Rev. Immunol.* 15, 323–350. doi: 10.1146/annurev.immunol.15.1.323
- Majumdar, T., Sharma, S., Kumar, M., Hussain, M. A., Chauhan, N., Kalia, I., et al. (2019). Tryptophan-kynurenine pathway attenuates β -catenin-dependent pro-parasitic role of STING-TICAM2-IRF3-IDO1 signalosome in *Toxoplasma gondii* infection. *Cell Death Dis.* 10:161. doi: 10.1038/s41419-019-1420-9
- Melo, M., Nguyen, Q., Cordeiro, C., Hassan, M., Yang, N., McKell, R., et al. (2013). Transcriptional analysis of murine macrophages infected with different *Toxoplasma* strains identifies novel regulation of host signaling pathways. *PLoS Pathog.* 9:e1003779. doi: 10.1371/journal.ppat.1003779
- Melo, M. B., Jensen, K. D., and Saeij, J. P. (2011). *Toxoplasma gondii* effectors are master regulators of the inflammatory response. *Trends Parasitol.* 27, 487–495. doi: 10.1016/j.pt.2011.08.001
- Melo, M. B., Kasperkovitz, P., Cerny, A., Konen-Waisman, S., Kurt-Jones, E. A., Lien, E., et al. (2010). UNC93B1 mediates host resistance to infection with *Toxoplasma gondii*. *PLoS Pathog.* 6:e1001071. doi: 10.1371/journal.ppat.1001071
- Morisaki, J. H., Heuser, J. E., and Sibley, L. D. (1995). Invasion of *Toxoplasma gondii* occurs by active penetration of the host cell. *J. Cell Sci.* 108, 2457–2464.
- Murray, P. J., Allen, J. E., Biswas, S. K., Fisher, E. A., Gilroy, D. W., Goerdt, S., et al. (2014). Macrophage activation and polarization: nomenclature and experimental guidelines. *Immunity* 41, 14–20. doi: 10.1016/j.immuni.2014.06.008
- Nava, P., Kamekura, R., Quirós, M., Medina-Contreras, O., Hamilton, R. W., Kolegraf, K. N., et al. (2014). IFN γ -induced suppression of β -catenin signaling: evidence for roles of Akt and 14.3.3 ζ . *Mol. Biol. Cell* 25, 2894–2904. doi: 10.1091/mbc.e13-09-0512
- Niedelman, W., Sprockholt, J. K., Clough, B., Frickel, E. M., and Saeij, J. P. (2013). Cell death of gamma interferon-stimulated human fibroblasts upon *Toxoplasma gondii* infection induces early parasite egress and limits parasite replication. *Infect. Immun.* 81, 4341–4349. doi: 10.1128/IAI.00416-13

- O'Neill, L. A. J., Golenbock, D., and Bowie, A. G. (2013). The history of toll-like receptors—redefining innate immunity. *Nat. Rev. Immunol.* 13, 453–460. doi: 10.1038/nri3446
- Pfefferkorn, E. R. (1984). Interferon gamma blocks the growth of *Toxoplasma gondii* in human fibroblasts by inducing the host cells to degrade tryptophan. *Proc. Natl. Acad. Sci. U.S.A.* 81, 908–912. doi: 10.1073/pnas.81.3.908
- Pimentel, H., Bray, N. L., Puente, S., Melsted, P., and Pachter, L. (2017). Differential analysis of RNA-seq incorporating quantification uncertainty. *Nat. Methods* 14, 687–690. doi: 10.1038/nmeth.4324
- Price, J. V., and Vance, R. E. (2014). The macrophage paradox. *Immunity* 41, 685–693. doi: 10.1016/j.immuni.2014.10.015
- Reya, T., Duncan, A. W., Ailles, L., Domen, J., Scherer, D. C., Willert, K., et al. (2003). A role for wnt signalling in self-renewal of haematopoietic stem cells. *Nature* 423, 409–415. doi: 10.1038/nature01593
- Scharton-Kersten, T. M., Yap, G., Magram, J., and Sher, A. (1997). Inducible nitric oxide is essential for host control of persistent but not acute infection with the intracellular pathogen *Toxoplasma gondii*. *J. Exp. Med.* 185, 1261–1273. doi: 10.1084/jem.185.7.1261
- Schneemann, M., and Schoedon, G. (2002). Species differences in macrophage NO production are important. *Nat. Immunol.* 3:102. doi: 10.1038/ni0202-102a
- Sibley, L. D., Adams, L. B., Fukutomi, Y., and Krahenbuhl, J. L. (1991). Tumor necrosis factor-alpha triggers antitoxoplasmal activity of IFN-gamma primed macrophages. *J. Immunol.* 147, 2340–2345.
- Sindrilaru, A., and Scharfetter-Kochanek, K. (2013). Disclosure of the culprits: macrophages—versatile regulators of wound healing. *Adv. Wound Care* 2, 357–368. doi: 10.1089/wound.2012.0407
- Song, H. B., Jun, H. O., Kim, J. H. J. H., Lee, Y. H., Choi, M. H., and Kim, J. H. J. H. (2017). Disruption of outer blood-retinal barrier by *Toxoplasma gondii*-infected monocytes is mediated by paracrine activated FAK signaling. *PLoS ONE* 12:e0175159. doi: 10.1371/journal.pone.0175159
- Staal, F. J. T., Luis, T. C., and Tiemessen, M. M. (2008). WNT signalling in the immune system: WNT is spreading its wings. *Nat. Rev. Immunol.* 8, 581–593. doi: 10.1038/nri2360
- Stelzer, S., Basso, W., Benavides Silván, J., Ortega-Mora, L. M., Maksimov, P., Gethmann, J., et al. (2019). *Toxoplasma gondii* infection and toxoplasmosis in farm animals: risk factors and economic impact. *Food Waterborne Parasitol.* 15:e00037. doi: 10.1016/j.fawpar.2019.e00037
- Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M. A., et al. (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U.S.A.* 102, 15545–15550. doi: 10.1073/pnas.0506580102
- Sugi, T., Ma, Y. F., Tomita, T., Murakoshi, F., Eaton, M. S., Yakubu, R., et al. (2016). *Toxoplasma gondii* cyclic AMP-dependent protein kinase subunit 3 is involved in the switch from tachyzoite to bradyzoite development. *mBio* 7, e 00755–16. doi: 10.1128/mBio.00755-16
- Thi, E. P., Lambert, U., and Reiner, N. E. (2012). Sleeping with the enemy: how intracellular pathogens cope with a macrophage lifestyle. *PLoS Pathog.* 8:e1002551. doi: 10.1371/journal.ppat.1002551
- Tombácz, K., Burgess, G., Holder, A., Werners, A., and Werling, D. (2018). *Toxoplasma gondii* profilin does not stimulate an innate immune response through bovine or human TLR5. *Innate Immun.* 24, 422–429. doi: 10.1177/1753425918798450
- Tosh, K. W., Mittereder, L., Bonne-Annee, S., Hieny, S., Nutman, T. B., Singer, S. M., et al. (2016). The IL-12 response of primary human dendritic cells and monocytes to *Toxoplasma gondii* is stimulated by phagocytosis of live parasites rather than host cell invasion. *J. Immunol.* 196, 345–356. doi: 10.4049/jimmunol.1501558
- Ueno, A., Dautu, G., Haga, K., Munyaka, B., Carmen, G., Kobayashi, Y., et al. (2011). *Toxoplasma gondii*: a bradyzoite-specific DnaK-tetratricopeptide repeat (DnaK-TPR) protein interacts with p23 co-chaperone protein. *Exp. Parasitol.* 127, 795–803. doi: 10.1016/j.exppara.2011.01.015
- van de Vosse, E., van Dissel, J. T., and Ottenhoff, T. H. (2009). Genetic deficiencies of innate immune signalling in human infectious disease. *Lancet Infect. Dis.* 9, 688–698. doi: 10.1016/S1473-3099(09)70255-5
- Wiley, M., Sweeney, K. R., Chan, D. A., Brown, K. M., McMurtrey, C., Howard, E. W., et al. (2010). *Toxoplasma gondii* activates hypoxia-inducible factor (HIF) by stabilizing the HIF-1 α subunit via type I activin-like receptor kinase receptor signaling. *J. Biol. Chem.* 285, 26852–26860. doi: 10.1074/jbc.M110.147041
- Yamamoto, M., Ma, J. S., Mueller, C., Kamiyama, N., Saiga, H., Kubo, E., et al. (2011). ATF6 β is a host cellular target of the *Toxoplasma gondii* virulence factor ROP18. *J. Exp. Med.* 208, 1533–1546. doi: 10.1084/jem.20101660
- Ye, J., Coulouris, G., Zaretskaya, I., Cutcutache, I., Rozen, S., and Madden, T. L. (2012). Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction. *BMC Bioinf.* 13:134. doi: 10.1186/1471-2105-13-134
- Young, R., Bush, S. J., Lefevre, L., McCulloch, M. E. B., Lisowski, Z. M., Muriuki, C., et al. (2018). Species-specific transcriptional regulation of genes involved in nitric oxide production and arginine metabolism in macrophages. *Immunohorizons* 2, 27–37. doi: 10.4049/immunohorizons.1700073
- Zhao, Y., Marple, A. H., Ferguson, D. J. P., Bzik, D. J., and Yap, G. S. (2014). Avirulent strains of *Toxoplasma gondii* infect macrophages by active invasion from the phagosome. *Proc. Natl. Acad. Sci. U.S.A.* 111, 6437–6442. doi: 10.1073/pnas.1316841111
- Zhao, Y. O., Khaminets, A., Hunn, J. P., and Howard, J. C. (2009). Disruption of the *Toxoplasma gondii* parasitophorous vacuole by IFN γ -inducible immunity-related GTPases (IRG proteins) triggers necrotic cell death. *PLoS Pathog.* 5:e1000288. doi: 10.1371/journal.ppat.1000288

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Gossner and Hassan. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.