



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Anatomy of a crosslinker

Citation for published version:

Belsom, A & Rappsilber, J 2020, 'Anatomy of a crosslinker', *Current opinion in chemical biology*, vol. 60, pp. 39-46. <https://doi.org/10.1016/j.cbpa.2020.07.008>

Digital Object Identifier (DOI):

[10.1016/j.cbpa.2020.07.008](https://doi.org/10.1016/j.cbpa.2020.07.008)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Current opinion in chemical biology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Anatomy of a crosslinker

Adam Belsom¹ and Juri Rappsilber^{1,2}

Abstract

Crosslinking mass spectrometry has become a core technology in structural biology and is expanding its reach towards systems biology. Its appeal lies in a rapid workflow, high sensitivity and the ability to provide data on proteins in complex systems, even in whole cells. The technology depends heavily on crosslinking reagents. The anatomy of crosslinkers can be modular, sometimes comprising combinations of functional groups. These groups are defined by concepts including: reaction selectivity to increase information density, enrichability to improve detection, cleavability to enhance the identification process and isotope-labelling for quantification. Here, we argue that both concepts and functional groups need more thorough experimental evaluation, so that we can show exactly how and where they are useful when applied to crosslinkers. Crosslinker design should be driven by data, not only concepts. We focus on two crosslinker concepts with large consequences for the technology, namely reactive group reaction kinetics and enrichment groups.

Addresses

¹ Bioanalytics, Institute of Biotechnology, Technische Universität Berlin, 13355, Berlin, Germany

² Wellcome Centre for Cell Biology, University of Edinburgh, Edinburgh, EH9 3BF, UK

Corresponding author: Rappsilber, Juri (Juri.Rappsilber@tu-berlin.de)

Current Opinion in Chemical Biology 2021, 60:39–46

This review comes from a themed issue on Omics

Edited by Nichollas Scott and Laura Edgington-Mitchell

For a complete overview see the [Issue](#) and the [Editorial](#)

<https://doi.org/10.1016/j.cbpa.2020.07.008>

1367-5931/© 2020 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Keywords

Crosslinking mass spectrometry, Proteins structure, Protein–protein interactions, Chemical reagents, Design of chemical reagents.

Introduction

Crosslinking mass spectrometry (MS) technology and its applications have been reviewed extensively [1–6], which includes subdisciplines providing high-density crosslinking data [7,8] and quantitative crosslinking data [9]. Here, we would like to turn the spotlight back


onto the crosslinking reagent, focussing on the multiple aspects of chemical structure and reactivity that these applications depend upon.

Protein crosslinking predates the electrospray-ionisation MS that is used routinely nowadays for its detection, by 30 years. The first identification of a crosslink (via paper chromatography) in 1958 by Zahn et al. was made following the use of a Sanger's reagent derivative 1,5-difluoro-2,4-dinitrobenzene (Figure 1) on insulin [10,11]. Since this time, there has been an explosion of protein crosslinking reagents available and something of an evolution/revolution in the chemicals used specifically for crosslinking-MS [12].

The range of available reagents and the chemistry and diverse functionality that can be incorporated within them is so great that it becomes impossible to review them on a case-by-case basis (to date, Sigma–Aldrich lists 268 crosslinkers, Thermo Fisher Scientific lists 104 crosslinkers, crosslinkers are also not a part of standardisation efforts in crosslinking-MS [<https://arxiv.org/abs/2007.00383>]). Instead, the crosslinking reagent can rather be dissected, understood and discussed in terms of its constituent parts. Note that 'zero-length' crosslinkers [13,14], reactive oxygen species [15] and metal-induced redox reactions [16], and photo-activatable noncanonical amino acids [17] can lead to crosslinks. However, the largest individual group of crosslinking reagents, which is also the most used for crosslinking-MS today, comprises molecules with functional groups, whereby at least two of the groups are reactive and capable of conjugation, which are themselves separated by a group functioning as a spacer or distance restraint (Figure 2). The simplest examples of this type include disuccinimidyl suberate (DSS) and bis(sulfosuccinimidyl)suberate (BS3) (NHS and sulfo-NHS esters, respectively), where the spacer constitutes a straightforward alkyl chain (Figure 1).

Despite the large selection of available crosslinkers, and their predating crosslinking-MS by nearly two decades, DSS [18] and BS3 [19] have become the go-to reagents of choice [6]. This was also evidenced in a recent interlaboratory study where the majority of participating labs chose DSS/BS3 (35/57 data sets) [20]. The choice is understandable considering the huge range of applications and successes using DSS/BS3, including integrative structural biology in purified complexes [21],

Figure 1




Evolution of crosslinkers. Reactive groups are highlighted in magenta. (a) 1,5-difluoro-2,4-dinitrobenzene (DFDNB), the first crosslinker applied to protein analysis. (b) Bis(sulfosuccinimidyl)suberate (BS3), which together with its nonsulfonated form disuccinimidyl suberate (DSS), are currently the most commonly used crosslinkers. (c) Protein interaction reporter (PIR), a large, multifunctional crosslinker. This crosslinker contains an MS-cleavable Rink group (highlighted in grey, cleavable bonds indicated by black dashed lines) and an enrichable biotin group (highlighted in cyan). MS, mass spectrometry.

organelles [22,23] and in cells [24], and both labelled [25,26] and label-free [27] quantitation for analysing protein structural changes [9,28].

The simplicity of DSS/BS3, reaction specificity, ease-of-use, reaction product stability and lack of reaction by-products, all contribute to the tremendous success of these reagents. However, an overreliance on this specific reactive group chemistry significantly limits the

Figure 2



Crosslinking reagent anatomy. Requisite features are shown in red. Optional features are highlighted in grey. Many different reactive groups and spacer scaffolds, and multiple different cleavable, releasable and enrichable groups are in use. Even more can be imagined or will be developed in the future. The resulting combinatorial complexity has already resulted in a large number of crosslinkers and allows for an even larger number of future crosslinkers. Reagents are typically only demonstrated in proof-of-principle, although some also in biological applications. However, the actual benefit of most of these reagents for crosslinking-MS has not been assessed rigorously. MS, mass spectrometry.

information density of crosslinking and may mislead structural analysis as we will discuss in the following context. Furthermore, there is significantly more to crosslinker design than only protein reactive groups.

In this review, we give a general overview of the array of functional groups found in crosslinking reagents. We propose that crosslinker design has been led by concepts, including reactive group specificity, cleavability to enhance identification and enrichability to improve detection. We argue that currently the enthusiasm for concepts pushes the design of new reagents. As appealing as concepts can be, they still must be underpinned by data. Ultimately, it is the data that reveal, where further improvements are required. After all, it is data that crosslinking-MS feeds, into structural and systems biology.

Reactive groups

The NHS-ester (and derivatives) is found ubiquitously in crosslinking reagents. Its reaction specificity in proteins is well characterised, preferentially reacting with amines from lysine side chains and protein N-termini but also with hydroxyl groups from the side chains of serine, threonine and tyrosine, at physiological conditions [29,30]. This confined specificity, predominantly towards pairs of proximal lysine residues, limits the numbers of distance restraints that can be introduced into proteins; hence data are sparse and low-resolution. Data density can be increased by expanding reactivity to target other amino acid residues,

including employment of aromatic glyoxal groups for arginine-targeting [31] and dihydrazides/diazo-derivatives targeting carboxylic acids [32–34].

Expanding reactivity even further, to all amino acids, has led to the greatest leap towards increasing data density so far (Figure 3), via crosslinkers containing photo-activatable reactive groups including diazirine (sulfo-SDA/SDA) [35] and benzophenone (sulfo-SBP/SBP) [36]. Upon activation by ultraviolet irradiation, the resulting reactive radical intermediates react promiscuously and readily with any proximal N–H and C–H bonds. These all-amino-acid-targeting photoactivatable groups are combined with highly amino acid-specific NHS-ester groups into heterobifunctional, semispecific crosslinkers. Protein crosslinking becomes a two-stage reaction: First, the NHS-ester groups decorate proteins specifically on nucleophiles (as previously described). Second, photoactivation completes crosslinking, to any part of a protein within a crosslinkable distance. Similar to the reaction products of homobifunctional, NHS-ester crosslinkers BS3/DSS, the beauty of the reaction products of the diazirine in sulfo-SDA, lies in their simplicity. The spacer group introduced is a short, 4-carbon, alkyl chain (3.9 Å spacer length, compared with 11.4 Å for BS3/DSS). The crosslinked products from use of photoactivatable benzophenone on the other hand, retain the aromatic groups. Interestingly, there is evidence of complementarity between crosslinked products with diazirine and benzophenone [36], despite involving the same anchoring residues. The influence of crosslinker spacer composition and conformational preference is an aspect of reagents that is being actively studied [37,38].

Semispecific, heterobifunctional crosslinkers can increase data density (not crosslinks per protein molecule) by orders of magnitude, compared with amino acid-specific, homobifunctional crosslinkers (a cross-laboratory study reported 44 crosslinks for monomeric BSA summed from three liquid chromatography (LC)-MS runs, as an average of the participating labs, whereas sulfo-SDA yielded 311 in a single LC-MS run [20]). The work carried out by our laboratory demonstrated the implication of this on one of the original application fields of crosslinking-MS, resolving detailed tertiary protein structure [8,35]. Indeed, high-density data using semispecific crosslinkers can improve protein-structure predictions [35] and routine implementation is intensely investigated [7,8,39–42]. This work inspired a number of other works in pursuit of this same ambition [41–44]. Further studies continue to demonstrate the benefits of semispecific crosslinkers in protein complexes [45]. The costs involved with the use of semispecific crosslinkers are the increased needs for high-quality MS and MS/MS data, and robust database search and respective false-discovery rate [46–49] determination pipelines [28,50]. The same is also true

wherever the scope of crosslinked products is expanded, including proteome-wide analyses [51–53].


An oft-mentioned concern in crosslinking-MS is the occurrence of artefacts, especially for photocrosslinkers [2,54]. Large numbers of crosslinks per protein molecule supposedly lead to catching, and stabilising, rare or artificial protein conformations and interactions, by ‘zippering’ (Figure 3). However, even at very high reagent concentrations, homobifunctional NHS-ester crosslinking reagents perturb protein structure only locally, with global folding relatively unscathed [55,56]. On the other hand, crosslinking with semispecific photoactivatable reagents is typically carried out with lower reagent concentrations than normal homobifunctional NHS-ester crosslinking, with high-density information resulting from the aggregated data of many individual protein molecules [28,35]. Zippering instead appears to be linked to reaction kinetics, rather than the number of crosslinks per molecule.

The reaction kinetics of crosslinker reactive groups is a concept that has recently been garnering more attention with regards to protein crosslinking reactions [55,57]. In short, the two relevant key factors of the reactive group are reaction rate and half-life. The NHS-ester group is highly reactive with a long half-life, consequently waiting for protein to come close enough for capture. This makes homobifunctional NHS-crosslinkers susceptible to zippering. The diazirine photoactivatable group however, produces an extraordinarily reactive (order of nanoseconds) carbene radical upon photoactivation, with an almost nonexistent half-life [58]. This indiscriminate and fast reaction therefore accurately captures protein states and interactions [54,59]. Taking reactivity in the other direction, the sulfur fluoride exchange (SuFEx) clickable group [60] has a much lower reactivity than NHS-ester, thus requiring longer-lived contacts and therefore not catching short-lived protein states [61]. Considering reaction kinetics and not only the reaction specificity of reactive groups, suggests a strong future for reactive groups complementary to the NHS-ester and the importance of data-driven crosslinker design.

Spacer functionality

The spacer can act as a scaffold for functionalities that address the low abundance and complexity of analytes, characteristics of crosslinking products that challenge analysis. Functionality includes MS-cleavable groups, isotope-coding, enrichment handles and related capture and release groups. Modular design approaches applied to crosslinker synthesis can be used to create reagents combining multiple functional groups [62]. This is where the range of applicable chemistry also greatly expands. This is demonstrated in the PIR crosslinkers [63] (Figure 1), azide/alkyne-A-DSBSO [64], CBDPS

Figure 3



Consequences of reactive group chemistry in crosslinkers. **(a)** Semi-specific photocrosslinkers can increase crosslink data density (the protein sequence is represented by a grey circular line; the crosslinks are represented as internal lines) by expanding reactivity to all amino acid residues. Photocrosslinkers such as SDA contain lysine (and S/T/Y) reactive NHS-esters on one side (blue circle). The photoactivatable reactive group (red circle) on the other side can react with any amino acid. The reaction specificity of standard homobifunctional NHS-ester-based

[65] and Leiker [66], which have seen application success including proteome-wide analyses [64,67,68].

MS-cleavable groups include multiple classes of bonds, which cleave upon collisional activation in the mass spectrometer and have been incorporated into numerous crosslinkers. MS-cleavable crosslinkers have been reviewed extensively [2,3,69]. The concept of MS-cleavability is proposed as important for proteome-wide analyses [52,70–72]. Other comparable proteome-wide studies however, have relied simply on DSS/BS3 [22,23,28]. This leads to the question of where the real gains of MS-cleavable groups can be achieved versus the conceptual benefits. Contributing to this, it is still unclear to what extent available reagents realise the concept. MS-cleavable reagents will not cleave with the same efficiency between all crosslinked peptides, an aspect that original proof-of-concept studies did not investigate and that urgently demands to be quantified. Assuming perfect behaviour when designing acquisition or search strategies will come at the cost of excluding a potentially large number of crosslinked peptides that cleave with lower efficiency than selected examples. Furthermore, the data obtained using noncleavable reagents can be computationally dissected into contributions of the individual peptides [73]. This computational ‘cleavage’ blurs the distinction between MS-cleavable and other crosslinkers and increases the need for more rigorous experimental validation.

Enrichable groups have also been used in crosslinkers (sometimes in combination with MS-cleavable groups), for proteome-wide analyses [64,67,68]. The predominant enrichment group used in the spacer of crosslinkers is biotin [62–66,74–80], which exploits the remarkably high affinity of the avidin–biotin complex. Biotinylated crosslinked peptides can be isolated on a solid-supported avidin derivative (including monomeric avidin, streptavidin and neutravidin), with the streptavidin–biotin interaction being the strongest noncovalent biological interaction known ($K_d \sim 10^{-15}$ M). This extremely high affinity presents a problem however, when release from the solid support is required [81]. It is unclear what is the extent of recovery from streptavidin using organic solvent elution alone [74,77,79]. Solid-supported monomeric avidin, with a far lower binding affinity with biotin ($K_d \sim 10^{-7}$ M), represents a compromise [63,65,75,76,78]. Other attempts to overcome the problem of release from the solid support have been to include release groups between the biotin and the rest of the crosslinker backbone [62,64,66,80]. These include polyethylene glycol (PEG, which is hydroxylamine-cleavable) [62], pinacol esters (acid-cleavable) [64], azobenzene (cleaved by sodium dithionite reduction) [66], disulphide (reductive-cleavage) [80] and photocleavable groups [66]. Click chemistry groups have been proposed as both capture groups for coupling a biotin enrichment group

crosslinkers, targets lysines (and S/T/Y residues to a lesser extent) only. This limited specificity also limits the density of resulting crosslink networks. **(b)** Rare protein conformations and interactions are over-represented by homobifunctional NHS-ester-based crosslinkers. The long half-life yet still fast reaction rate of the NHS-ester reactive group, allows capturing rare conformations (on the timescale of typical protein dynamics) by apparent zippering. Photocrosslinkers generate reactive radicals with vanishingly short half-life and consequently do not “wait” until protein is nearby. Therefore protein populations of mixed conformations are captured accurately. **(c)** The relative reaction rates and half-lives of the diazirine (red circle), NHS-ester (blue circle) and SuFEx (orange circle) reactive groups.

after crosslinking [64,78], and as enrichment groups in their own right [64,82,83], with reversibility achieved via an acid-cleavable acetal [64,83] or disulphide [82]. Recently, a different enrichment group, a negatively charged phosphonic acid, was developed into a crosslinker reagent [84]. This simple, small and short crosslinker is a refreshing exception to the biotin/Click chemistry conundrum and the trend towards functional crosslinker expansion. Paradoxically though, the negative charge prevents the crossing of cell membranes for in-cell crosslinking, a major application goal that thirsts for enrichment.

The impressive array of chemistry applied to crosslinking reagents in the quest for crosslinker enrichment shows that synthesis of even protracted functionality is possible. What is not yet clear is whether increasing functionality guarantees or translates into improved identification of crosslinked analytes, that is, greater information content which might be reached by increased number of crosslinks (quantity) at maintained confidence (quality) but also different types of crosslinks (kinetics, selectivity). A concern is that increasing and combining functional chemistry into larger crosslinking reagents could end up burdening analysis, owing to issues such as increased hydrophobicity, unexpected side reactions, inefficient reactions, complicated fragmentation spectra and other related causes of analyte or information losses.

Conclusion

The power and potential of crosslinking-MS has been thoroughly established. Extensive claims regarding conceptual gains, through the use of crosslinker chemistry and consequent functional groups, have been made repeatedly. Crosslinker anatomical concepts including MS-cleavability and enrichment have potential, but the benefits must be demonstrated clearly. This must be done in order to move forward with developing better

performing crosslinkers and optimal combinations of functionality.

Functional groups applied in crosslinkers need to be thoroughly tested, characterised and proven specifically in the context of crosslinking-MS. A larger focus on data is urgently needed, to enable data-driven crosslinker design. An important element of this is the inclusion of statistical data as part of proof-of-concept studies to complement cherry-picked examples, which would also include providing open and stable access to the raw data. One way this could be done is through synthetic peptide libraries [50].

The increased appeal of crosslinking-MS, means that many scientists now want to apply crosslinking-MS as a tool to answer their own structural biology questions. However, they must first navigate concepts when deciding which crosslinker reagent to choose for their application. It is possible that an ideal crosslinker already exists, despite crosslinker developers continuing to churn out new variants. But without proving concepts through data, crosslinker developers are unable to make concrete suggestions or convince crosslinking MS users to apply their crosslinker reagents with confidence. A consequence is that researchers will err on the side of caution, opting for the simplicity of homobifunctional NHS-ester based crosslinkers such as BS3 and DSS.

Crosslinking reagents are of course essential but still only one of the critical pieces of the crosslinking MS machinery, which also relies on MS, software engineering, data processing and analysis, data visualisation and integration with structural biology. Ultimately the true test of crosslinker chemistry will be through the applications that are achieved and the discoveries that will be made.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The study is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC 2008–390540038 – UniSysCat and 329673113. This work was also supported by the Wellcome Trust through a Senior Research Fellowship to JR [103139]. The Wellcome Centre for Cell Biology is supported by core funding from the Wellcome Trust [203149].

References

Papers of particular interest, published within the period of review, have been highlighted as:

- * of special interest
- ** of outstanding interest

1. Leitner A, Faini M, Stengel F, Aebersold R: **Crosslinking and mass spectrometry: an integrated technology to understand the structure and function of molecular machines.** *Trends Biochem Sci* 2016, **41**:20–32.
2. Sinz A: **Cross-linking/mass spectrometry for studying protein structures and protein-protein interactions: where are we now and where should we go from here?** *Angew Chem Int Ed Engl* 2018, **57**:6390–6396.
3. Yu C, Huang L: **Cross-linking mass spectrometry: an emerging technology for interactomics and structural biology.** *Anal Chem* 2018, **90**:144–165.
4. O'Reilly FJ, Rappsilber J: **Cross-linking mass spectrometry: methods and applications in structural, molecular and systems biology.** *Nat Struct Mol Biol* 2018, **25**:1000–1008.
5. Chavez JD, Bruce JE: **Chemical cross-linking with mass spectrometry: a tool for systems structural biology.** *Curr Opin Chem Biol* 2019, **48**:8–18.
6. Steigenberger B, Albanese P, Heck AJR, Scheltema RA: **To cleave or not to cleave in XL-MS?** *J Am Soc Mass Spectrom* 2020, **31**:196–206.
7. Belsom A, Schneider M, Brock O, Rappsilber J: **Blind evaluation of hybrid protein structure analysis methods based on cross-linking.** *Trends Biochem Sci* 2016, **41**:564–567.
8. Schneider M, Belsom A, Rappsilber J: **Protein tertiary structure by crosslinking/mass spectrometry.** *Trends Biochem Sci* 2018, **43**:157–169.
9. Chen ZA, Rappsilber J: **Protein dynamics in solution by quantitative crosslinking/mass spectrometry.** *Trends Biochem Sci* 2018, **43**:908–920.
10. Zahn VH, Meienhofer J: **Reaktionen von 1,5-difluor-2,4-dinitrobenzol mit insulin 1. Mitt. Synthese von modellverbindungen.** *Die Makromol Chem* 1958, **26**:126–152.
11. Zahn VH, Meienhofer J: **Reaktionen von 1,5-difluor-2,4-dinitrobenzol mit insulin 2. Mitt. Versuche mit insulin.** *Die Makromol Chem* 1958, **26**:153–166.
12. Hermanson GT: *Bioconjugate techniques.* Academic Press; 2013.
13. Staros JV, Wright RW, Swingle DM: **Enhancement by N-hydroxysulfosuccinimide of water-soluble carbodiimide-mediated coupling reactions.** *Anal Biochem* 1986, **156**:220–222.
14. Grabarek Z, Gergely J: **Zero-length crosslinking procedure with the use of active esters.** *Anal Biochem* 1990, **185**:131–135.
15. Aeschbach R, Amadò R, Neukom H: **Formation of dityrosine cross-links in proteins by oxidation of tyrosine residues.** *Biochim Biophys Acta* 1976, **439**:292–301.
16. Fancy DA, Kodadek T: **Chemistry for the analysis of protein-protein interactions: rapid and efficient cross-linking triggered by long wavelength light.** *Proc Natl Acad Sci U S A* 1999, **96**:6020–6024.
17. Suchanek M, Radzikowska A, Thiele C: **Photo-leucine and photo-methionine allow identification of protein-protein interactions in living cells.** *Nat Methods* 2005, **2**:261–267.
18. Pilch PF, Czech MP: **Interaction of cross-linking agents with the insulin effector system of isolated fat cells. Covalent linkage of 125I-insulin to a plasma membrane receptor protein of 140,000 daltons.** *J Biol Chem* 1979, **254**:3375–3381.
19. Staros JV: **N-hydroxysulfosuccinimide active esters: bis(N-hydroxysulfosuccinimide) esters of two dicarboxylic acids are hydrophilic, membrane-impermeant, protein cross-linkers.** *Biochemistry* 1982, **21**:3950–3955.
20. Iacobucci C, Piotrowski C, Aebersold R, Amaral BC, Andrews P, Bernfur K, Borchers C, Brodie NI, Bruce JE, Cao Y, et al.: **First community-wide, comparative cross-linking mass spectrometry study.** *Anal Chem* 2019, **91**:6953–6961.
21. Chen ZA, Jawhari A, Fischer L, Buchen C, Tahir S, Kamenski T, Rasmussen M, Larivière L, Bukowski-Wills J-C, Nilges M, et al.: **Architecture of the RNA polymerase II-TFIIF complex revealed by cross-linking and mass spectrometry.** *EMBO J* 2010, **29**:717–726.
22. Ryl PSJ, Bohlke-Schneider M, Lenz S, Fischer L, Budzinski L, Stuiver M, Mendes MML, Sinn L, O'Reilly FJ, Rappsilber J: **In Situ structural restraints from cross-linking mass spectrometry in human mitochondria.** *J Proteome Res* 2020, **19**:327–336.
 First proteome-wide Crosslinking MS study using a standard cross-linker, i.e. one that is not designed to be MS-cleavable. MS-cleavable crosslinkers are believed to be essential for such large-scale studies.
23. Linden A, Deckers M, Parfentev I, Pflanz R, Homberg B, Neumann P, Ficner R, Rehling P, Urlaub H: **A cross-linking mass spectrometry approach defines protein interactions in yeast mitochondria.** *Mol Cell Proteomics* 2020, <https://doi.org/10.1074/mcp.RA120.002028>.
24. O'Reilly FJ, Xue L, Graziadei A, Sinn L, Lenz S: **In-cell architecture of an actively transcribing-translating expressome.** *bioRxiv* 2020, <https://doi.org/10.1126/science.abb3758>.
25. Fischer L, Chen ZA, Rappsilber J: **Quantitative cross-linking/mass spectrometry using isotope-labelled cross-linkers.** *J Proteomics* 2013, **88**:120–128.
26. Chen ZA, Fischer L, Cox J, Rappsilber J: **Quantitative cross-linking/mass spectrometry using isotope-labeled cross-linkers and MaxQuant.** *Mol Cell Proteomics* 2016, **15**:2769–2778.
27. Müller F, Fischer L, Chen ZA, Auchynnikava T, Rappsilber J: **On the reproducibility of label-free quantitative cross-linking/mass spectrometry.** *J Am Soc Mass Spectrom* 2018, **29**:405–412.
28. Mendes ML, Fischer L, Chen ZA, Barbon M, O'Reilly FJ, Giese SH, Bohlke-Schneider M, Belsom A, Dau T, Combe CW, et al.: **An integrated workflow for crosslinking mass spectrometry.** *Mol Syst Biol* 2019, **15**: e8994.
29. Kalkhof S, Sinz A: **Chances and pitfalls of chemical cross-linking with amine-reactive N-hydroxysuccinimide esters.** *Anal Bioanal Chem* 2008, **392**:305–312.
30. Tomko Jr RJ, Taylor DW, Chen ZA, Wang H-W, Rappsilber J, Hochstrasser M: **A single α helix drives extensive remodeling of the proteasome lid and completion of regulatory particle assembly.** *Cell* 2015, **163**:432–444.
31. Jones AX, Cao Y, Tang Y-L, Wang J-H, Ding Y-H, Tan H, Chen Z-L, Fang R-Q, Yin J, Chen R-C, et al.: **Improving mass spectrometry analysis of protein structures with arginine-selective chemical cross-linkers.** *Nat Commun* 2019, **10**:3911.
 Expands the reactive groups used in crosslinking to aromatic glyoxal groups for arginine-targeting and OPA for lysine-targeting, also departing from the omnipresent use of N-hydroxysuccinimide esters for lysine-targeting.
32. Novák P, Kruppa GH: **Intra-Molecular cross-linking of acidic residues for protein structure studies.** *Eur J Mass Spectrom* 2008, **14**:355–365.
33. Leitner A, Joachimiak LA, Unverdorben P, Walzthoeni T, Frydman J, Förster F, Aebersold R: **Chemical cross-linking/mass spectrometry targeting acidic residues in proteins and protein complexes.** *Proc Natl Acad Sci USA* 2014, **111**:9455–9460.
34. Zhang X, Wang J-H, Tan D, Li Q, Li M, Gong Z, Tang C, Liu Z, Dong M-Q, Lei X: **Carboxylate-selective chemical cross-linkers for mass spectrometric analysis of protein structures.** *Anal Chem* 2018, **90**:1195–1201.
35. Belsom A, Schneider M, Fischer L, Brock O, Rappsilber J: **Serum Albumin Domain structures in human Blood Serum by mass spectrometry and computational biology.** *Mol Cell Proteomics* 2016, **15**:1105–1116.
 Introduces high-data densities to crosslinking by use of a photo-activatable diazine and shows that crosslinking data can guide modelling in solving protein structures de novo.
36. Belsom A, Mudd G, Giese S, Auer M, Rappsilber J: **Complementary benzophenone cross-linking/mass spectrometry photochemistry.** *Anal Chem* 2017, **89**:5319–5324.
 Shows that the crosslinker, as a molecule and not just its reactive groups, influences which residues are captured.

37. Yu C, Novitsky EJ, Cheng NW, Rychnovsky SD, Huang L: **Exploring spacer arm structures for designs of asymmetric sulfoxide-containing MS-cleavable cross-linkers**. *Anal Chem* 2020, **92**:6026–6033.
38. Gong Z, Ye S-X, Nie Z-F, Tang C: **The conformational preference of chemical crosslinkers determines the crosslinking probability of reactive protein residues**. *J Phys Chem B* 2020: 4446–4453, <https://doi.org/10.1021/acs.jpcc.0c02522>.
39. Belsom A, Schneider M, Fischer L, Mabrouk M, Stahl K, Brock O, Rappsilber J: **Blind testing cross-linking/mass spectrometry under the auspices of the 11th critical assessment of methods of protein structure prediction (CASP11)**. *Wellcome Open Res* 2016, **1**:24.
40. Schneider M, Belsom A, Rappsilber J, Brock O: **Blind testing of cross-linking/mass spectrometry hybrid methods in CASP11**. *Proteins* 2016, **84**(Suppl 1):152–163.
41. Ogorzalek TL, Hura GL, Belsom A, Burnett KH, Kryshtafyovych A, Tainer JA, Rappsilber J, Tsutakawa SE, Fidelis K: **Small angle X-ray scattering and cross-linking for data assisted protein structure prediction in CASP 12 with prospects for improved accuracy**. *Proteins* 2018, **86**(Suppl 1):202–214.
42. Fajardo JE, Shrestha R, Gil N, Belsom A, Crivelli SN, Czaplewski C, Fidelis K, Grudin S, Karasikov M, Karczyńska AS, et al.: **Assessment of chemical-crosslink-assisted protein structure modeling in CASP13**. *Proteins* 2020, **88**:540.
43. Brodie NI, Popov KI, Petrotchenko EV, Dokholyan NV, Borchers CH: **Solving protein structures using short-distance cross-linking constraints as a guide for discrete molecular dynamics simulations**. *Sci Adv* 2017, **3**. e1700479.
44. Mintseris J, Gygi SP: **High-density chemical cross-linking for modeling protein interactions**. *Proc Natl Acad Sci U S A* 2020, **117**:93–102.
45. Higashi TL, Eickhoff P, Simoes JS, Locke J, Nans A, Flynn HR, Snijders AP, Papageorgiou G, O'Reilly N, Chen ZA, et al.: **A structure-based mechanism for DNA entry into the cohesin ring**. *Mol Cell* 2020. In Press, Corrected Proof, <https://doi.org/10.1016/j.molcel.2020.07.013>.
46. Maiolica A, Cittaro D, Borsotti D, Sennels L, Ciferri C, Tarricone C, Musacchio A, Rappsilber J: **Structural analysis of multiprotein complexes by cross-linking, mass spectrometry, and database searching**. *Mol Cell Proteomics* 2007, **6**:2200–2211.
47. Walzthoeni T, Claassen M, Leitner A, Herzog F, Bohn S, Förster F, Beck M, Aebersold R: **False discovery rate estimation for cross-linked peptides identified by mass spectrometry**. *Nat Methods* 2012, **9**:901–903.
48. Fischer L, Rappsilber J: **Quirks of error estimation in cross-linking/mass spectrometry**. *Anal Chem* 2017, **89**:3829–3833.
49. Fischer L, Rappsilber J: **False discovery rate estimation and heterobifunctional cross-linkers**. *PLoS One* 2018, **13**. e0196672.
50. Beveridge R, Stadlmann J, Penninger JM, Mechtler K: **A synthetic peptide library for benchmarking crosslinking-mass spectrometry search engines for proteins and protein complexes**. *Nat Commun* 2020, **11**:742.
- Assesses search algorithms and error estimation procedures by means of a sample with known composition; a synthetic peptide library.
51. Keller A, Chavez JD, Felt KC, Bruce JE: **Prediction of an upper limit for the fraction of interprotein cross-links in large-scale in vivo cross-linking studies**. *J Proteome Res* 2019, **18**: 3077–3085.
52. Yugandhar K, Wang T-Y, Leung AK-Y, Lanz MC, Motorykin I, Liang J, Shayhidin EE, Smolka MB, Zhang S, Yu H: **MaxLinker: proteome-wide cross-link identifications with high specificity and sensitivity**. *Mol Cell Proteomics* 2020, **19**:554–568.
53. Lenz S, Sinn LR, O'Reilly FJ, Fischer L, Wegner F, Rappsilber J: **Reliable identification of protein-protein interactions by crosslinking mass spectrometry**. *bioRxiv* 2020, <https://doi.org/10.1101/2020.05.25.114256>.
- Proteome-wide Crosslinking MS study demonstrating the possibility and importance of controlled false discovery rate estimation for reliable identification of PPIs.
54. Ziemianowicz DS, Ng D, Schryvers AB, Schriemer DC: **Photo-cross-linking mass spectrometry and integrative modeling enables rapid screening of antigen interactions involving bacterial Transferrin receptors**. *J Proteome Res* 2019, **18**: 934–946.
- Demonstrates that the fast photoactivatable diazirine-containing SDA crosslinker provides more accurate reflection of the in-solution state of a protein complex than N-hydroxysuccinimide esters.
55. Rozbeský D, Rosůlek M, Kukačka Z, Chmelík J, Man P, Novák P: **Impact of chemical cross-linking on protein structure and function**. *Anal Chem* 2018, **90**:1104–1113.
56. Xia L, Ma Z, Tong J, Tang Y, Li S, Qin S, Lou R, Zhao S, Lei X, Shui W: **Evaluation of chemical cross-linkers for in-depth structural analysis of G protein-coupled receptors through cross-linking mass spectrometry**. *Anal Chim Acta* 2020, **1102**: 53–62.
57. Fürsch J, Kammer K-M, Kreft SG, Beck M, Stengel F: **Proteome-wide structural probing of low-abundant protein interactions by cross-linking mass spectrometry**. *Anal Chem* 2020, **92**: 4016–4022.
58. Wang J, Burdzinski G, Kubicki J, Platz MS, Moss RA, Fu X, Piotrowiak P, Myahkostupov M: **Ultrafast spectroscopic study of the photochemistry and photophysics of arylhalodiazirines: direct observation of carbene and zwitterion formation**. *J Am Chem Soc* 2006, **128**:16446–16447.
59. Li Z, Hao P, Li L, Tan CYJ, Cheng X, Chen GYJ, Sze SK, Shen H-M, Yao SQ: **Design and synthesis of minimalist terminal Alkyne-Containing diazirine photo-crosslinkers and their incorporation into kinase inhibitors for cell-and tissue-based proteome profiling**. *Angew Chem Int Ed* 2013, **52**:8551–8556.
60. Barrow AS, Smedley CJ, Zheng Q, Li S, Dong J, Moses JE: **The growing applications of SuFEx click chemistry**. *Chem Soc Rev* 2019, **48**:4731–4758.
61. Yang B, Wu H, Schnier PD, Liu Y, Liu J, Wang N, DeGrado WF, Wang L: **Proximity-enhanced SuFEx chemical cross-linker for specific and multitargeting cross-linking mass spectrometry**. *Proc Natl Acad Sci U S A* 2018, **115**:11162–11167.
- Introduces slow crosslinking as a method by using aryl sulfonyl fluorides that target His, Ser, Thr, Tyr, and Lys side chains in such a slow manner that rapid, rare conformers are not captured.
62. Trester-Zedlitz M, Kamada K, Burley SK, Fenyő D, Chait BT, Muir TW: **A modular cross-linking approach for exploring protein interactions**. *J Am Chem Soc* 2003, **125**:2416–2425.
63. Zhang H, Tang X, Munske GR, Tolic N, Anderson GA, Bruce JE: **Identification of protein-protein interactions and topologies in living cells with chemical cross-linking and mass spectrometry**. *Mol Cell Proteomics* 2009, **8**:409–420.
64. Kaake RM, Wang X, Burke A, Yu C, Kandur W, Yang Y, Novitsky EJ, Second T, Duan J, Kao A, et al.: **A new in vivo cross-linking mass spectrometry platform to define protein-protein interactions in living cells**. *Mol Cell Proteomics* 2014, **13**:3533–3543.
65. Petrotchenko EV, Serpa JJ, Borchers CH: **An isotopically coded CID-cleavable biotinylated cross-linker for structural proteomics**. *Mol Cell Proteomics* 2011, **10**. M110.001420.
66. Tan D, Li Q, Zhang M-J, Liu C, Ma C, Zhang P, Ding Y-H, Fan S-B, Tao L, Yang B, et al.: **Trifunctional cross-linker for mapping protein-protein interaction networks and comparing protein conformational states**. *Elife* 2016, **5**. e12509.
67. Schweppe DK, Chavez JD, Lee CF, Caudal A, Kruse SE, Stuppard R, Marcinek DJ, Shadel GS, Tian R, Bruce JE: **Mitochondrial protein interactome elucidated by chemical cross-linking mass spectrometry**. *Proc Natl Acad Sci U S A* 2017, **114**:1732–1737.

68. Makepeace KAT, Mohammed Y, Rudashevskaya EL, Petrochenko EV, Vögtle F-N, Meisinger C, Sickmann A, Borchers CH: **Improving identification of in-organello protein-protein interactions using an affinity-enrichable, isotopically coded, and mass spectrometry-cleavable chemical cross-linker.** *Mol Cell Proteomics* 2020, **19**:624–639.
69. Sinz A: **Divide and conquer: cleavable cross-linkers to study protein conformation and protein-protein interactions.** *Anal Bioanal Chem* 2017, **409**:33–44.
70. Liu F, Rijkers DTS, Post H, Heck AJR: **Proteome-wide profiling of protein assemblies by cross-linking mass spectrometry.** *Nat Methods* 2015, **12**:1179–1184.
- First proteome-wide Crosslinking MS study using an MS-cleavable crosslinker.
71. Liu F, Lössl P, Scheltema R, Viner R, Heck AJR: **Optimized fragmentation schemes and data analysis strategies for proteome-wide cross-link identification.** *Nat Commun* 2017, **8**:15473.
72. Götze M, Iacobucci C, Ihling CH, Sinz A: **A simple cross-linking/mass spectrometry workflow for studying system-wide protein interactions.** *Anal Chem* 2019, **91**:10236–10244.
73. Giese SH, Fischer L, Rappsilber J: **A study into the collision-induced dissociation (CID) behavior of cross-linked peptides.** *Mol Cell Proteomics* 2016, **15**:1094–1104.
74. Fujii N, Jacobsen RB, Wood NL, Schoeniger JS, Guy RK: **A novel protein crosslinking reagent for the determination of moderate resolution protein structures by mass spectrometry (MS3-D).** *Bioorg Med Chem Lett* 2004, **14**:427–429.
75. Sinz A, Kalkhof S, Ihling C: **Mapping protein interfaces by a trifunctional cross-linker combined with MALDI-TOF and ESI-FTICR mass spectrometry.** *J Am Soc Mass Spectrom* 2005, **16**:1921–1931.
76. Chu F, Mahrus S, Craik CS, Burlingame AL: **Isotope-coded and affinity-tagged cross-linking (ICATXL): an efficient strategy to probe protein interaction surfaces.** *J Am Chem Soc* 2006, **128**:10362–10363.
77. Kang S, Mou L, Lanman J, Velu S, Brouillette WJ, Prevelige Jr PE: **Synthesis of biotin-tagged chemical cross-linkers and their applications for mass spectrometry.** *Rapid Commun Mass Spectrom* 2009, **23**:1719–1726.
78. Chowdhury SM, Du X, Tolić N, Wu S, Moore RJ, Mayer MU, Smith RD, Adkins JN: **Identification of cross-linked peptides after click-based enrichment using sequential collision-induced dissociation and electron transfer dissociation tandem mass spectrometry.** *Anal Chem* 2009, **81**:5524–5532.
79. Rey M, Dupré M, Lopez-Neira I, Duchateau M, Chamot-Rooke J: **eXL-MS: an enhanced cross-linking mass spectrometry workflow to study protein complexes.** *Anal Chem* 2018, **90**:10707–10714.
80. Stadlmeier M, Runtsch LS, Streshnev F, Wühr M, Carell T: **A click-chemistry-based enrichable crosslinker for structural and protein interaction analysis by mass spectrometry.** *ChemBiochem* 2020, **21**:103–107.
81. Oda Y, Nagasu T, Chait BT: **Enrichment analysis of phosphorylated proteins as a tool for probing the phosphoproteome.** *Nat Biotechnol* 2001, **19**:379–382.
82. Buncherd H, Nessen MA, Nouse N, Stelder SK, Roseboom W, Dekker HL, Arents JC, Smeenk LE, Wanner MJ, van Maarseveen JH, et al.: **Selective enrichment and identification of cross-linked peptides to study 3-D structures of protein complexes by mass spectrometry.** *J Proteomics* 2012, **75**:2205–2215.
83. Matzinger M, Kandollner W, Doppler P, Heiss EH, Mechtler K: **Fast and highly efficient affinity enrichment of azide-A-DSBSO cross-linked peptides.** *J Proteome Res* 2020, **19**:2071–2079.
84. Steigenberger B, Pieters RJ, Heck AJR, Scheltema RA: **PhoX: an IMAC-enrichable cross-linking reagent.** *ACS Cent Sci* 2019, **5**:1514–1522.
- Departs from nearly two decades of field-focus on biotin as an enrichment group and introduces the phosphonic acid moiety for this purpose, converting the enrichment challenge of crosslinks onto the already solved challenge of enriching phosphorylated peptides.