

OPERATORS ON ANTI-DUAL PAIRS: GENERALIZED KREIN–VON NEUMANN EXTENSION

ZSIGMOND TARCSAY AND TAMÁS TITKOS

ABSTRACT. The main aim of this paper is to generalize the classical concept of positive operator, and to develop a general extension theory, which overcomes not only the lack of a Hilbert space structure, but also the lack of a normable topology. The concept of anti-duality carries an adequate structure to define positivity in a natural way, and is still general enough to cover numerous important areas where the Hilbert space theory cannot be applied. Our running example – illustrating the applicability of the general setting to spaces bearing poor geometrical features – comes from noncommutative integration theory. Namely, representable extension of linear functionals of involutive algebras will be governed by their induced operators. The main theorem, to which the vast majority of the results is built, gives a complete and constructive characterization of those operators that admit a continuous positive extension to the whole space. Various properties such as commutation, or minimality and maximality of special extensions will be studied in detail.

1. INTRODUCTION

One of the most natural questions that arises when someone dealing with partially defined objects in mathematics is whether there exists an extension sharing some prescribed properties. A great many authors have studied abstract extension problems for operators on Hilbert spaces, that go at least back to M. G. Krein [17] and J. von Neumann [19]. (For various different developments of their groundbreaking work see e.g. [4–6, 8, 12, 13, 22, 26], and the references therein.) The following extension problem was posed by P. R. Halmos [11]: Assume that a positive operator $A : D \rightarrow H$ is given, where D is a linear subspace of the complex Hilbert space H . Positivity here means that $(Ax | x) \geq 0$ for all $x \in D$. The question is: under what conditions can we guarantee the existence of an everywhere defined bounded positive extension \tilde{A} of A ? Of course, if there is any then A itself must be bounded. Hence, extending it to the closure by continuity, we may suppose that D is closed. Consider the matrix representation of A with respect to the orthogonal

2010 *Mathematics Subject Classification.* Primary 47A20, 47B65, 46A20 Secondary 46K10, 46A22.

Key words and phrases. Positive operator, anti-duality, factorization, operator extension, *-algebra, positive functional, operator kernel.

The corresponding author Zs. Tarcsay (tarcsay@cs.elte.hu) was supported by the “For the Young Talents of the Nation” scholarship program (NTP-NFTÖ-17) of the Hungarian Ministry of Human Capacities.

T. Titkos was supported by the Hungarian National Research, Development and Innovation Office – NKFIH (PD128374), and by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences.

decomposition $H = D \oplus D^\perp$

$$[A] = \begin{pmatrix} A_{11} & * \\ A_{21} & * \end{pmatrix},$$

where $A_{11} : D \rightarrow D$ and $A_{21} : D \rightarrow D^\perp$ arise in the usual way, whereas the second column (of symbols $*$) waits to be filled to obtain a positive operator. It is easy to see that every positive extension of A has representation of the form

$$(1.1) \quad [\tilde{A}] = \begin{pmatrix} A_{11} & A_{21}^* \\ A_{21} & X \end{pmatrix},$$

where $A_{11} : D \rightarrow D$ and $X : D^\perp \rightarrow D^\perp$ are positive. So, extending A to a positive operator \tilde{A} is equivalent to find $X \geq 0$ such that the operator matrix (1.1) is positive. (For a more general completion problem for block operators see [7].) This form helps us to demonstrate also that such an extension need not exist even in the most simple case. Indeed, assume that H is of the form $H = K \oplus K$ with a complex Hilbert space K and assume that A_{11} is the zero operator, while $A_{21} = A_{21}^*$ is any positive but nonzero bounded operator on K . Then an elementary calculation shows that there is no positive completion at all.

The main aim of this paper is to develop a general extension theory which overcomes the problem of not having orthogonal decomposition when we drop the Hilbert space structure. This level of generality is indeed necessary in our considerations, because we intend to investigate extendibility of “positive” mappings acting on spaces without inner product. In order to introduce the appropriate analogues of standard operator classes, that cover the original Hilbert space setting, and is general enough to be applicable for objects like operator kernels and representable functionals, we are going to consider anti-dual pairs. That is, two appropriately chosen complex linear spaces intertwined by a sesquilinear and separating map, called anti-duality. This is a slight modification of the well known dual pair setting (see [21, Chapter IV]), the only difference we make is the conjugate linearity in the second variable. Having an anti-duality at hand also allows us to introduce various topologies, and hence continuity and boundedness of maps acting between the spaces in question.

The organization of the paper is as follows. Section 2 contains a short overview of concepts that are needed in later sections of the paper. In particular, we are going to introduce the notions of positivity and symmetry of operators in context of anti-dual pairs. It will turn out that these operators are weakly continuous, which suggests that the most adequate topologies for our investigations are the weak topologies.

In Section 3 we present a quite general extension theorem, which can be considered as the main result of the paper. In fact, this result will serve as the base of our further investigations throughout. Roughly speaking, we are going to characterize (in both topological and algebraic ways) operators possessing positive extension to the whole space. In addition, the construction which is based on the ingenious paper of Z. Sebestyén [24] has some useful theoretical consequences, including an explicit formula for the obtained positive extension, as well as for its “quadratic form”. Also, it will turn out that this extension is minimal in some sense, thus we shall call it Krein-von Neumann extension, in accordance with the classical literature. We emphasize that in this paper we restrict ourselves to continuous extensions, and thus the operators we are going to deal with are typically not densely defined.

Hence, the maximal (Friedrichs) extension, apart from trivial cases, does not exist. Nevertheless, we will show that the set of positive extensions bounded by a fixed positive operator always possesses a maximal element. Closing that chapter we will also show how the abstract result can be used for positive definite operator functions. The special case when the anti-duality is the evaluation on the pair of a fixed Banach space and its conjugate topological dual will be considered in Section 4. On the one hand, we will obtain a strengthening of the main result of [26], on the other hand, we will conclude that Halmos' original result follows indeed as an immediate consequence of our main theorem. Finally, we apply our results to obtain representable positive extensions of linear functionals given on a left ideal of an involutive algebra.

2. PRELIMINARIES

In this section we collect all the necessary ingredients from the theory of topological vector spaces, anti-dual pairs, and their special linear operators which we are going to use throughout.

2.1. Anti-dual pairs and related topologies. We start by recalling the notion of anti-duality, which is just a slight (and technical) modification of dual pairing. Although there is no crucial difference between these two notions, we choose anti-duality in order to stay formally as close as possible to the Hilbert space case. Let E, F be complex vector spaces. A function $\langle \cdot, \cdot \rangle : F \times E \rightarrow \mathbb{C}$ is called anti-duality if it is sesquilinear (that is to say, linear in the first argument and conjugate linear in the second one) and $\langle \cdot, \cdot \rangle$ separates the points of F and E (that is to say, if $\langle f, x \rangle = 0$ for all $x \in E$ then $f = 0_F$ and if $\langle f, x \rangle = 0$ for all $f \in F$ then $x = 0_E$). The triple $((E, F), \langle \cdot, \cdot \rangle)$ is called an anti-dual pair, and it is denoted shortly by $\langle F, E \rangle$. Observe that if $\langle F, E \rangle$ is an anti-dual pair then $\langle E, F \rangle'$ is also an anti-dual pair where $\langle \cdot, \cdot \rangle' : E \times F \rightarrow \mathbb{C}$ is given by

$$(2.1) \quad \langle x, f \rangle' := \overline{\langle f, x \rangle}, \quad x \in E, f \in F.$$

The most natural anti dual pair is a linear space and a linear subspace of its conjugate algebraic dual, intertwined by the evaluation as anti-duality. In fact, every anti-dual pair can be written in the above form. Indeed, if $\langle F, E \rangle$ is an anti-dual pair, then due to the identification

$$x \mapsto \varphi_x; \quad \varphi_x(f) := \langle f, x \rangle \quad \text{for all } f \in F,$$

E may be regarded as a linear subspace of F^* , the algebraic dual of F . Similarly, due to the mapping

$$f \mapsto \psi_f; \quad \psi_f(x) := \langle f, x \rangle \quad \text{for all } x \in E,$$

F can be identified as a linear subspace of \bar{E}^* , the algebraic anti-dual space of E . Our prototype of anti-dual pairs is the system $((H, H), (\cdot | \cdot))$ where H is a Hilbert space with inner product $(\cdot | \cdot)$. This particular anti-dual pair has the useful feature that H can be identified with its topological dual along the maps $x \mapsto \varphi_x$ and $f \mapsto \psi_f$, according to the Riesz representation theorem. A similar feature is obtained in the general setting if we endow E and F with appropriate topologies. For this purpose the most natural at hand are the *weak topologies* $\sigma(E, F)$ and $\sigma(F, E)$ on E and F , respectively: $\sigma(E, F)$ is the smallest topology making φ_x continuous for all $x \in E$, and similarly, $\sigma(F, E)$ is the smallest topology on F such

that all the functionals of the form ψ_f ($f \in F$) are continuous. Both $(E, \sigma(E, F))$ and $(F, \sigma(F, E))$ are locally convex Hausdorff spaces such that $\bar{E}' = F$ and $F' = E$, where \bar{E}' and F' refer to the topological anti-dual and dual spaces of E and F , respectively. We call an anti dual pair $\langle F, E \rangle$ w^* -sequentially complete, if F is $\sigma(F, E)$ sequentially complete. In order to demonstrate that being w^* -sequentially complete does not mainly depend on the spaces but on the way of how to intertwine them, we cite an example of Wilansky.

Example 2.1. Let us denote by c and ℓ^1 the spaces of convergent and absolute summable complex sequences, respectively. Define an anti-duality by

$$\langle x, y \rangle_1 := \sum_{n=1}^{\infty} x(n) \overline{y(n)}, \quad x \in \ell^1, \quad y \in c.$$

Clearly, $\langle \ell^1, c \rangle_1$ is an anti-dual pair, which is not w^* -sequentially complete, as the sequence of Kronecker delta's ($\delta^n(n) = 1$ and $\delta^n(k) = 0$ if $k \neq n$) is a w^* -Cauchy but not w^* -convergent sequence in ℓ^1 . On the other hand, the anti-dual pair $\langle \ell^1, c \rangle_2$ with

$$\langle x, y \rangle_2 := x(1) \cdot \overline{\lim_{n \rightarrow \infty} y(n)} + \sum_{n=1}^{\infty} x(n+1) \overline{y(n)}$$

is w^* -sequentially complete according to the Banach-Steinhaus theorem (for the details see Example 7 of 8-2 and Problem 108-109 of 8-2 in [30]).

We remark also here that a linear space and its algebraic conjugate dual intertwined by evaluation is always w^* -(sequentially) complete. A Banach space and its topological conjugate dual form also a w^* -sequentially complete anti-dual pair, according to the Banach-Steinhaus theorem. More generally, a barrelled space and its topological conjugate dual form a quasi-complete, and hence a w^* -sequentially complete anti-dual pair. Nevertheless we mention that the a Banach space is w^* -complete if and only if the space is finite dimensional.

Besides the weak topology, the strong topology will be particularly important for our investigations. Recall that if $\langle F, E \rangle$ is an anti-dual pair, then the polar of $A \subseteq F$ is the absolutely convex set

$$A^\circ = \{x \in E \mid \forall a \in A : |\langle a, x \rangle| \leq 1\}.$$

The strong topology $\beta(E, F)$ on E is obtained as the linear topology induced by polars of $\sigma(F, E)$ -bounded subsets of F . Similarly, the strong topology $\beta(F, E)$ on F is induced by polars of $\sigma(E, F)$ -bounded subsets of E . Now we turn to investigate special linear operators acting between two sides of anti-dual pairs.

2.2. Positive- and symmetric operators on anti-dual pairs. If an anti-dual pair $\langle F, E \rangle$ is given, we will use the short notation $A : E \supseteq \text{dom } A \rightarrow F$ for *linear* operators acting on a subspace $\text{dom } A$ of E with values in F . We prefer this setting instead of considering duality with conjugate linear operators as in [1]. An operator $A : E \supseteq \text{dom } A \rightarrow F$ is called *positive* if it satisfies

$$\langle Ax, x \rangle \geq 0 \quad \text{for all } x \in \text{dom } A,$$

and *symmetric* if

$$\langle Ax, y \rangle = \overline{\langle Ay, x \rangle} \quad \text{for all } x, y \in \text{dom } A.$$

It is obvious that these are direct generalizations of the well-known notions of Hilbert space theory. The main advantage is that this setting allows us to handle structures without Hilbert space structure analogously. As the next example will illustrate, operators on anti-dual pairs appear very naturally for example in noncommutative integration theory.

Example 2.2. Let \mathcal{A} be a $*$ -algebra with algebraic conjugate-dual $\overline{\mathcal{A}}^*$, and let $\mathcal{I} \subseteq \mathcal{A}$ be a left-ideal. Then $\langle \overline{\mathcal{A}}^*, \mathcal{A} \rangle$ is an anti-dual pair with $\langle f, a \rangle := f(a)$. If a positive linear functional $f : \mathcal{I} \rightarrow \mathbb{C}$ is given, we can associate a positive operator $A_f : \mathcal{I} \rightarrow \overline{\mathcal{A}}^*$ as $\langle A_f a, b \rangle := f(b^* a)$. It will turn out later that positive extendibility of f to the whole algebra can be characterized by means of A_f . Furthermore, the canonical extension of f itself will be gained from the canonical extension of A_f .

Recall one of the main advantages of weak topology, namely that a linear operator T acting on a topological vector space (V, \mathcal{T}_V) with values in $(F, \sigma(F, E))$ is continuous if and only if the linear functionals

$$(2.2) \quad \vartheta_x : V \rightarrow \mathbb{C}; \quad \vartheta_x(h) := \langle Th, x \rangle$$

are continuous for each $x \in E$. For the sake of simplicity we introduce the following terminology: if two anti-dual pairs $\langle F_1, E_1 \rangle_1$ and $\langle F_2, E_2 \rangle_2$ are given, we will call a map $T : E_1 \rightarrow E_2$ *weakly continuous* if T is $\sigma(E_1, F_1)$ - $\sigma(E_2, F_2)$ continuous. The set of weakly continuous linear operators $T : E_1 \rightarrow E_2$ is denoted by $\mathcal{L}(E_1; E_2)$. By replacing $\langle F_2, E_2 \rangle_2$ with $\langle E_2, F_2 \rangle_2'$ (see (2.1)) one can characterize weak continuity of an operator $T : E_1 \rightarrow F_2$. Indeed, according to (2.2), T is weakly continuous if and only if for all $x_2 \in E_2$ there exists $f_1 \in F_1$ such that

$$(2.3) \quad \langle Tx_1, x_2 \rangle_2 = \overline{\langle f_1, x_1 \rangle_1} (= \langle x_1, f_1 \rangle_1') \quad \text{for all } x_1 \in E_1.$$

The (necessarily weakly continuous) operator $T^* : E_2 \rightarrow F_1$ satisfying

$$\langle Tx_1, x_2 \rangle_2 = \overline{\langle T^* x_2, x_1 \rangle_1}, \quad x_1 \in E_1, x_2 \in E_2.$$

is called the adjoint of T . In particular, the adjoint A^* of an operator $A \in \mathcal{L}(E; F)$ belongs again to $\mathcal{L}(E; F)$ and satisfies $\langle Ax, y \rangle = \overline{\langle A^* y, x \rangle}$ for all $x, y \in E$. Hence it makes sense to speak about self-adjointness $A = A^*$ of an operator $A \in \mathcal{L}(E; F)$. An everywhere defined symmetric operator (that is, an operator $S : E \rightarrow F$ such that $\langle Sx, y \rangle = \overline{\langle Sy, x \rangle}, x, y \in E$) is automatically weakly continuous, and hence self-adjoint. If $A : E \supseteq \text{dom } A \rightarrow F$ is an operator such that $\langle Ax, x \rangle$ is real for all $x \in \text{dom } A$ then the sesquilinear form $\mathfrak{t}_A(x, y) := \langle Ax, y \rangle$ ($x, y \in \text{dom } A$) is hermitian, thus A is symmetric. Indeed, $\langle Ax, y \rangle = \mathfrak{t}_A(x, y) = \overline{\mathfrak{t}_A(y, x)} = \overline{\langle Ay, x \rangle}$ holds for all $x, y \in \text{dom } A$. Let us summarize these observations in the following proposition.

Proposition 2.1. *Let $\langle F, E \rangle$ be an anti-dual pair.*

- (a) *Every symmetric operator $A : E \rightarrow F$ is weakly continuous and self-adjoint.*
- (b) *An operator $A : E \rightarrow F$ is symmetric if and only if $\langle Ax, x \rangle$ is real for each $x \in E$.*
- (c) *Every positive operator $A : E \rightarrow F$ is weakly continuous and self-adjoint.*

As our main interest in this paper lies in positive extensions, we present a prototype of positive operators. In rest of the paper we are mainly interested in positive operators on an anti-dual pair. In the next example we present the prototype of such positive operators.

Example 2.3. Let $\langle F, E \rangle$ be an anti-dual pair and H a Hilbert space. If $T : E \rightarrow H$ is a $\sigma(E, F)$ - $\sigma(H, H)$ continuous linear operator then its adjoint $T^* : H \rightarrow F$ is $\sigma(H, H)$ - $\sigma(F, E)$ continuous so that $T^*T \in \mathcal{L}(E; F)$ is positive:

$$\langle T^*Tx, x \rangle = (Tx | Tx) \geq 0, \quad x \in E.$$

We will see later that, under some natural conditions on F , each positive operator $A \in \mathcal{L}(E; F)$ possesses a factorization of the form $A = T^*T$ with a suitable T and H of Example 2.3.

3. MAIN THEOREM – EXTENSIONS OF POSITIVE OPERATORS

The central problem of this section is to provide necessary and sufficient conditions under which a linear operator $A : E \supseteq \text{dom } A \rightarrow F$ possesses a positive extension to the whole E . The following set associated to A will play a key role in our treatment:

$$(3.1) \quad W(A) := \{Ax \mid x \in \text{dom } A, \langle Ax, x \rangle \leq 1\} \subseteq F.$$

The construction below is motivated by the work of Sebestyén [24]. Here we emphasize that no topological conditions on $\text{dom } A$ are assumed.

Theorem 3.1. *Let $\langle F, E \rangle$ be a w^* -sequentially complete anti-dual pair and let $A : E \supseteq \text{dom } A \rightarrow F$ be a linear operator with domain $\text{dom } A$, which is assumed to be only a linear subspace. Then the following statements are equivalent.*

- (i) *There is a positive operator $\tilde{A} \in \mathcal{L}(E; F)$ extending A ,*
- (ii) *$W(A)$ is $\beta(F, E)$ -bounded in F ,*
- (iii) *$W(A)$ is $\sigma(F, E)$ -bounded in F ,*
- (iv) *To any y in E there is $M_y \geq 0$ such that*

$$(3.2) \quad |\langle Ax, y \rangle|^2 \leq M_y \langle Ax, x \rangle \quad \text{for all } x \in \text{dom } A.$$

If one (and hence all) of the above conditions is satisfied then there exists a distinguished extension called the Krein-von Neumann extension which is minimal in the following sense: $A_N \leq \tilde{A}$ holds for any positive extension $\tilde{A} \in \mathcal{L}(E; F)$ of A .

Proof. We start by proving that (iv) implies (i). Let us consider the following inner product on the range space $\text{ran } A$ of A :

$$(3.3) \quad (Ax \mid Ax')_A := \langle Ax, x' \rangle, \quad x, x' \in \text{dom } A.$$

First of all note that $(\cdot \mid \cdot)_A$ is well defined: if $Ax = Ax'$ and $Ay = Ay'$, then

$$\langle Ax, y \rangle = \langle Ax', y \rangle = \overline{\langle Ay, x' \rangle} = \overline{\langle Ay', x' \rangle} = \langle Ax', y' \rangle.$$

Furthermore, if $\langle Ax, x \rangle = 0$ for some $x \in \text{dom } A$ then (iv) implies $\langle Ax, y \rangle = 0$ for all $y \in E$, hence $Ax = 0$, which means that $(\text{ran } A, (\cdot \mid \cdot)_A)$ is a pre-Hilbert space. Consider now the completion H_A of $\text{ran } A$ equipped with that inner product. Let us consider the densely defined canonical embedding operator

$$(3.4) \quad J : H_A \supseteq \text{ran } A \rightarrow F; \quad J(Ax) := Ax, \quad x \in \text{dom } A.$$

For any y in E , we see by (iv) that

$$|\langle Ax, y \rangle|^2 \leq M_y (Ax \mid Ax)_A, \quad x \in \text{dom } A,$$

hence the linear functional

$$H_A \supseteq \text{ran } A \ni Ax \mapsto \langle Ax, y \rangle,$$

is continuous with respect to the norm induced by $(\cdot | \cdot)_A$. That means that J is $\sigma(\text{ran } A, H_A)$ - $\sigma(F, E)$ continuous and thus it admits a unique continuous extension to H_A by w^* -sequentially completeness of F . As it will not cause ambiguity, this extension will be denoted by the same symbol $J \in \mathcal{L}(H_A; F)$. The adjoint operator $J^* \in \mathcal{L}(E; H_A)$ fulfills

$$(Ax' | J^*x)_A = \langle J(Ax'), x \rangle = \langle Ax', x \rangle = (Ax' | Ax)_A$$

for every $x, x' \in \text{dom } A$, whence we gain the useful identity

$$(3.5) \quad J^*x = Ax, \quad x \in \text{dom } A.$$

And here we have arrived: for any x in $\text{dom } A$ we find that $JJ^*x = J(Ax) = Ax$, hence $JJ^* \in \mathcal{L}(E; F)$ is a positive extension of A . The extension obtained by this construction will be denoted by A_N .




FIGURE 1. Factorization of the Krein-von Neumann extension.

If $A \in \mathcal{L}(E; F)$ is any positive extension of A then clearly $W(A) \subseteq W(\tilde{A})$. Hence, to see that (i) implies (ii) we may assume that A is everywhere defined, i.e., $A \in \mathcal{L}(E; F)$. We have to show that $W(A)$ is bounded in the $\beta(F, E)$ topology, that is, B° absorbs $W(A)$ for each $\sigma(E, F)$ -bounded subset B of E . Note that we have $A = JJ^*$ and that $W(A) \subseteq J\langle B_A \rangle$, where B_A refers to the closed unit ball in H_A . Since B_A is weakly compact, $J\langle B_A \rangle$ is absorbed by the barrel B° , according to the absorbing lemma. Hence B° absorbs $W(A)$ as well, proving that (i) \Rightarrow (ii).

As the topology $\beta(F, E)$ is stronger than $\sigma(F, E)$, (ii) implies (iii) obviously.

Assume now that $W(A)$ is weak- $*$ bounded in F and take y from E . By continuity we see that $\langle \cdot, y \rangle$ is bounded on $W(A)$ whence we conclude that

$$M_y := \sup\{|\langle Ax, y \rangle|^2 \mid x \in \text{dom } A, \langle Ax, x \rangle \leq 1\} < +\infty,$$

which clearly implies (iv). The equivalence of (i)-(iv) has been proved.

In order to prove minimality of A_N , we are going to show first that

$$(3.6) \quad \langle A_N y, y \rangle = \sup\{|\langle Ax, y \rangle|^2 \mid x \in \text{dom } A, \langle Ax, x \rangle \leq 1\}, \quad y \in E.$$

Indeed, using density of $\text{ran } A$ in H_A , we conclude for all $y \in E$ that

$$\begin{aligned} \langle JJ^*y, y \rangle &= (J^*y | J^*y)_A = \{|\langle Ax | J^*y \rangle_A|^2 \mid x \in \text{dom } A, (Ax | Ax)_A \leq 1\} \\ &= \sup\{|\langle Ax, y \rangle|^2 \mid x \in \text{dom } A, \langle Ax, x \rangle \leq 1\}. \end{aligned}$$

Consider now a positive extension $\tilde{A} \in \mathcal{L}(E; F)$ of A . Then

$$\begin{aligned} \langle JJ^*y, y \rangle &= \sup\{|\langle Ax, y \rangle|^2 \mid x \in \text{dom } A, \langle Ax, x \rangle \leq 1\} \\ &\leq \sup\{|\langle \tilde{A}x, y \rangle|^2 \mid x \in E, \langle \tilde{A}x, x \rangle \leq 1\} \\ &\leq \sup\{\langle \tilde{A}x, x \rangle \langle \tilde{A}y, y \rangle \mid x \in E, \langle \tilde{A}x, x \rangle \leq 1\} \\ &= \langle \tilde{A}y, y \rangle, \end{aligned}$$

holds for any $y \in E$. Hence $JJ^* \leq \tilde{A}$, indeed. \square

For the sake of brevity and clarity we introduce now a short term to express that $\langle F, E \rangle$ is a given anti-dual pair where the topological space $(F, \sigma(F, E))$ is sequentially complete, $\text{dom } A$ is a linear subspace of E , and $A : \text{dom } A \rightarrow F$ is a linear operator that satisfies one (and hence all) of the properties (i)-(iv) in Theorem 3.1. Following Halmos' terminology we will say shortly that A is a *subpositive operator on the w^* -sequentially complete anti-dual pair $\langle F, E \rangle$* . Observe that (3.6) provides a useful explicit formula for the ‘‘quadratic form’’ of A_N . From the density of $\text{ran } A$ in H_A we can establish another one, namely,

$$\begin{aligned} 0 &= \inf\{(J^*y - Ax | J^*y - Ax)_A \mid x \in \text{dom } A\} \\ &= \langle JJ^*y, y \rangle + \inf\{-\langle Ax, y \rangle - \overline{\langle Ax, y \rangle} + \langle Ax, x \rangle \mid x \in \text{dom } A\} \end{aligned}$$

for every $y \in E$, which gives us

$$(3.7) \quad \langle A_N y, y \rangle = \sup\{\langle Ax, y \rangle + \overline{\langle Ax, y \rangle} - \langle Ax, x \rangle \mid x \in \text{dom } A\}, \quad y \in E,$$

after rearrangement.

The simple example below demonstrates that w^* -sequentially completeness of F was really essential in the main theorem.

Example 3.1. Let H be a Hilbert space and let A be an unbounded positive self-adjoint operator in H , with (dense) domain $\text{dom } A$. Let E denote the domain of $A^{1/2}$, i.e., $E := \text{dom } A^{1/2}$. Then clearly, $\langle H, E \rangle$ forms an anti-dual pair with respect to the duality induced by the inner product, but H is not sequentially complete with respect to the corresponding weak topology. It is readily seen that $A : \text{dom } A \rightarrow H$ fulfills condition (iv) of Theorem 3.1: indeed, for $y \in E$,

$$|(Ax | y)|^2 = |(A^{1/2}x | A^{1/2}y)|^2 \leq \|A^{1/2}y\|^2 (Ax | x), \quad x \in \text{dom } A.$$

Although condition (iv) is satisfied, the implication of Theorem 3.1 does not remain true because of the lack of w^* -sequentially completeness. Indeed, assume that A extends to a positive operator $\tilde{A} : E \rightarrow H$. Recall that a self-adjoint operator may not have any proper symmetric extension, hence $A = \tilde{A}$ and, in particular, $\text{dom } A = \text{dom } A^{1/2}$. But this is impossible because $\text{dom } A \subsetneq \text{dom } A^{1/2}$ whenever A is unbounded (see [28, Corollary 2.4]).

Notice that the set of positive extensions of a given positive operator A has no maximal element (unless $\text{dom } A$ is dense): for example, in the trivial case when $\text{dom } A = \{0\}$, every positive operator is an extension of A . However, the next theorem says that we will get a maximum among positive extensions, bounded by a positive operator B .

Theorem 3.2. *Let A be a subpositive operator on the w^* -sequentially complete anti-dual pair $\langle F, E \rangle$. Let $B \in \mathcal{L}(E; F)$ be a positive operator such that $A_N \leq B$, then there exists a positive operator $A_{max}^B \in \mathcal{L}(E; F)$, $A_{max}^B \leq B$ such that for every positive extension $\tilde{A} \in \mathcal{L}(E; F)$ of A , $0 \leq \tilde{A} \leq B$, one has $\tilde{A} \leq A_{max}^B$. In other words,*

$$A_{max}^B = \max\{\tilde{A} \in \mathcal{L}(E; F) \mid 0 \leq \tilde{A} \leq B, A \subset \tilde{A}\}.$$

Furthermore, a positive operator $0 \leq \tilde{A} \leq B$ is an extension of A if and only if $A_N \leq \tilde{A} \leq A_{max}^B$:

$$(3.8) \quad [A_N, A_{max}^B] = \{\tilde{A} \in \mathcal{L}(E; F) \mid 0 \leq \tilde{A} \leq B, A \subset \tilde{A}\}$$

Proof. By assumption, $B - A_N \geq 0$ is a positive extension of $B - A$, hence $(B - A)_N$ exists. Set

$$A_{max}^B := B - (B - A)_N.$$

We have apparently $0 \leq A_{max}^B \leq B$, and $A \subset A_{max}^B$. If $0 \leq \tilde{A} \leq B$ extends A then $B - A \subset B - \tilde{A}$ implies $(B - A)_N \leq B - \tilde{A}$ and hence $\tilde{A} = B - (B - \tilde{A}) \leq A_{max}^B$. In order to prove (3.8) let $A_N \leq \tilde{A} \leq A_{max}^B$. Take $x \in \text{dom } A$, then for every $y \in E$ we have

$$\begin{aligned} |\langle \tilde{A}x - Ax, y \rangle|^2 &= |\langle (\tilde{A} - A_N)x, y \rangle|^2 \\ &\leq \langle (\tilde{A} - A_N)x, x \rangle \langle (\tilde{A} - A_N)y, y \rangle \\ &\leq \langle (A_{max}^B - A_N)x, x \rangle \langle (\tilde{A} - A_N)y, y \rangle = 0, \end{aligned}$$

because $A_N x = A_{max}^B x = Ax$. Hence $\tilde{A}x = Ax$, that is, $A \subset \tilde{A}$. \square

The next theorem tells us that the Krein-von Neumann extension preserves certain commutation properties as well. We remark in advance that the proof is a bit unusual: although the claim concerns continuous operators, the proof itself applies unbounded operator techniques.

Theorem 3.3. *Let $A : \text{dom } A \rightarrow F$ be a subpositive operator on the w^* -sequentially complete anti-dual pair $\langle F, E \rangle$. Suppose that there are two weakly continuous operators $B, C \in \mathcal{L}(E; E)$ leaving $\text{dom } A$ invariant, and that the spectrum of BC restricted to $\text{dom } A$ is bounded. Assume in addition that B and C satisfy*

$$(3.9) \quad C^*A \subset AB, \quad \text{and} \quad B^*A \subset AC,$$

then the Krein-von Neumann extension of A satisfies

$$C^*A_N = A_N B, \quad \text{and} \quad B^*A_N = A_N C.$$

Proof. Let us introduce two linear operators \hat{B} and \hat{C} on the dense linear subspace $\text{ran } A \subseteq H_A$ by letting

$$\hat{B}(Ax) := ABx, \quad \hat{C}(Ax) := ACx, \quad x \in \text{dom } A.$$

Observe that \hat{B} and \hat{C} satisfy

$$(\hat{B}(Ax) | Ay)_A = (Ax | \hat{C}(Ay))_A, \quad x, y \in \text{dom } A$$

because from (3.9) it follows that

$$(ABx | Ay)_A = \langle ABx, y \rangle = \langle Ax, Cy \rangle = (Ax | ACy)_A.$$

We see therefore that \hat{B} and \hat{C} are closable operators and their closures satisfy

$$(3.10) \quad (\hat{B}^{**}h | k)_A = (h | \hat{C}^{**}k)_A, \quad h \in \text{dom}(\hat{B}^{**}), k \in \text{dom}(\hat{C}^{**}).$$

We are going to prove first that \hat{B}^{**} and \hat{C}^{**} are adjoints of each other, i.e.,

$$\hat{C}^* = \hat{B}^{**}, \quad \hat{B}^* = \hat{C}^{**}.$$

By [25, Theorem 3.1] that will be shown once we prove that the operator matrix

$$\Pi := \begin{pmatrix} tI_A & -\hat{B}^{**} \\ \hat{C}^{**} & tI_A \end{pmatrix}$$

has full range, i.e., $\text{ran } \Pi = H_A \times H_A$, for some $t \in \mathbb{R}$, $t \neq 0$ (here I_A stands for the identity operator of H_A). From (3.10) it is easy to deduce that

$$\|\Pi(h, k)\|^2 \geq t^2 \|(h, k)\|^2, \quad (h, k) \in \text{dom } \Pi,$$

and since Π is a closed operator, its range is closed. On the other hand, with

$$\Pi_0 := \begin{pmatrix} tI_A & -\widehat{B} \\ \widehat{C} & tI_A \end{pmatrix},$$

we have $\Pi = \Pi_0^{**}$, hence it suffices to show that Π_0 has dense range. To do so observe that

$$\begin{pmatrix} tI_A & -\widehat{B} \\ \widehat{C} & tI_A \end{pmatrix} \begin{pmatrix} tI_A & \widehat{B} \\ -\widehat{C} & tI_A \end{pmatrix} = \begin{pmatrix} t^2I_A + \widehat{B}\widehat{C} & 0 \\ 0 & t^2I_A + \widehat{C}\widehat{B} \end{pmatrix},$$

and here the latter one has range

$$(3.11) \quad \{(A(t^2I + BC)x, A(t^2I + CB)y) \mid x, y \in \text{dom } A\} \subset H_A \times H_A.$$

By assumption, we have that the restriction $t^2I + BC|_{\text{dom } A} : \text{dom } A \rightarrow \text{dom } A$ is weakly continuously invertible whenever t is large enough. The same holds true for $t^2I + CB$ because of the Jacobson lemma. Hence we conclude that the set in (3.11) is identical with $\text{ran } A \times \text{ran } A$, and therefore $\text{ran } A \times \text{ran } A \subseteq \text{ran } \Pi_0$, which proves our claim.

Next we show that $\text{ran } J^* \subseteq \text{dom}(\widehat{B}^{**})$ and $J^*B = \widehat{B}^{**}J^*$. Let $y \in E$ and $x \in \text{dom } A$, then $(\widehat{C}(Ax) \mid J^*y)_A = \langle ACx, y \rangle = \langle Ax, By \rangle = (Ax \mid J^*By)_A$, whence we have $J^*y \in \text{dom}(\widehat{C}^*)$ and $\widehat{C}^*J^*y = J^*By$. Since we have $\widehat{C}^* = \widehat{B}^{**}$, this proves $J^*B = \widehat{B}^{**}J^*$. Similarly, $J^*C = \widehat{C}J^*$. And now we have arrived:

$$A_N B = J J^* B = J \widehat{B}^{**} J^* = (J \widehat{B}^* J^*)^* = (J \widehat{C}^{**} J^*)^* = (A_N C)^* = C^* A_N,$$

and similarly, $A_N C = B^* A_N$. The proof is complete. \square

We have proved above that the auxiliary operators \widehat{B} and \widehat{C} (more precisely, their closures) are adjoint of each other, but we do not know whether they are bounded or not. If the underlying anti-dual pair comes from a Banach space E and its conjugate dual space $F = E'$ then one can imitate the proof of [14, Theorem 2.2] to show that \widehat{B} , \widehat{C} are bounded operators with common norm bound $\sqrt{r(BC)}$, the spectral radius of BC .

We close this section by a quick application of our main theorem. Namely, we are going to deal with so called positive definite operator functions see e.g. [3, 27] for the Hilbert space case, and [15, 18] for the Banach space setting. As these frameworks were too restrictive for various applications, many authors dealt with the case of functions taking values in a locally convex topological vector spaces in duality (see for example [4] and the references therein). Our setting is as follows: let $\langle F, E \rangle$ be a w^* -sequentially complete anti-dual pair and Z be a non-void set. Let us denote by \mathcal{F} the vector space of functions $u : Z \rightarrow E$ which have finite support, and denote by \mathcal{F}^* the algebraic conjugate dual of \mathcal{F} . A function $K : Z \times Z \rightarrow \mathcal{L}(E; F)$ is called a positive definite function (or shortly, a *kernel*) if it satisfies

$$(3.12) \quad \sum_{s, t \in Z} \langle K(s, t)u(s), u(t) \rangle \geq 0, \quad \text{for all } u \in \mathcal{F}.$$

We can naturally associate a positive operator $A_K : \mathcal{F} \rightarrow \tilde{\mathcal{F}}^*$ with K by setting

$$\langle A_K u, v \rangle := \sum_{s,t \in Z} \langle K(s,t)u(s), u(t) \rangle, \quad u, v \in \mathcal{F}.$$

For given two kernels K, L we write $K \preceq L$ if their associated positive operators satisfy $A_K \leq A_L$. Using this correspondence one can consider a kernel K as subspace of $\mathcal{F} \times \tilde{\mathcal{F}}^*$ of the form $\{(u, A_K u) \mid u \in \mathcal{F}\}$, say the graph of K . In the following theorem we will characterize whether a subspace implemented by an operator $A : \mathcal{F}_0 \rightarrow \tilde{\mathcal{F}}^*$ can be extended to be a graph of a kernel.

Theorem 3.4. *Let \mathcal{F}_0 be a linear subspace of \mathcal{F} , and assume that a positive operator $A : \mathcal{F}_0 \rightarrow \tilde{\mathcal{F}}^*$ is given. Then the following assertions are equivalent:*

- (i) *There exists a kernel $K : Z \times Z \rightarrow \mathcal{L}(E; F)$ so that $A \subseteq A_K$,*
- (ii) *For every $v \in \mathcal{F}$ there is $M_v \geq 0$ such that*

$$|\langle Au, v \rangle|^2 \leq M_v \cdot \langle Au, u \rangle \quad \text{for all } u \in \mathcal{F}_0.$$

Moreover, there is a K that is smallest in the sense that $K \preceq L$ for any kernel L with $A \subseteq A_L$.

Proof. It is obvious that (i) implies (ii). For the converse observe that \mathcal{F} and $\tilde{\mathcal{F}}^*$ intertwined by the evaluation form a w^* -sequentially complete anti-dual pair, thus we can apply Theorem 3.1. Therefore the smallest (Krein-von Neumann) extension A_N of A exists. Our only duty is to show that A_N is induced by a suitable kernel K . To this aim, for any fixed $s \in Z$ and $x \in E$ introduce the function $u_{s,x} \in \mathcal{F}$ by setting

$$u_{s,x}(s') := \delta_s(s') \cdot x, \quad s' \in Z,$$

where δ_s is the usual Dirac-function concentrated at s . For $s, t \in Z$ define a linear operator $K(s, t) : E \rightarrow F$ by

$$\langle K(s, t)x, y \rangle := \langle A_N u_{s,x}, u_{t,y} \rangle, \quad x, y \in E.$$

By the symmetry of A_N we conclude that

$$\langle K(s, t)x, y \rangle = \langle A_N u_{s,x}, u_{t,y} \rangle = \overline{\langle A_N u_{t,y}, u_{s,x} \rangle} = \overline{\langle K(t, s)y, x \rangle},$$

whence it is clear that $K(s, t) \in \mathcal{L}(E; F)$ (and also that $K(s, t)^* = K(t, s)$). Finally, since every $u \in \mathcal{F}$ can be expressed as $u = \sum_{s \in Z} u_{s, u(s)}$ we have

$$\langle A_N u, v \rangle = \sum_{s,t \in Z} \langle A u_{s, u(s)}, u_{t, v(t)} \rangle = \sum_{s,t \in Z} \langle K(s, t)u(s), v(t) \rangle,$$

which means that A_N is induced by the kernel K . □

4. THE BANACH SPACE SETTING

In this section we are going to investigate the special case when the anti-duality is the evaluation on the pair of a fixed Banach space E and its conjugate topological dual E' . We will obtain a strengthening of the main result of [26], and we will show that Halmos' original result is indeed a corollary of our main theorem.

We remark that the Banach-Steinhaus theorem forces E' to be weakly sequentially complete, and hence everything has been proved in the preceding sections remains valid also for the anti-dual pair $\langle \bar{E}', E \rangle$.

We start our investigations with a suitable generalization of the Schwarz inequality (cf. Kadison [16, Theorem 1], and also Lemma 2.1 and Corollary 2.2 in [29]).

Proposition 4.1. *Let E be a Banach space and let $A_i : E \rightarrow \bar{E}'$ ($1 \leq i \leq n$) be positive operators. Then for every set of vectors $x_i \in E$ $1 \leq i \leq n$ the following inequality holds*

$$(4.1) \quad \left\| \sum_{j=1}^n A_j x_j \right\|^2 \leq \left\| \sum_{j=1}^n A_j \right\| \cdot \sum_{j=1}^n \langle A_j x_j, x_j \rangle.$$

Moreover, $C := \left\| \sum_{j=1}^n A_j \right\|$ is the smallest constant satisfying (4.1).

Proof. Consider the direct sum Hilbert-space $K := \bigoplus_{j=1}^n H_{A_j}$, and define a linear operator $V : E \rightarrow K$ by $Vx := (A_1 x, A_2 x, \dots, A_n x)$ for $x \in E$. It is easy to check that V is continuous by norm bound $\|V\|^2 \leq \left\| \sum_{j=1}^n A_j \right\|$. A straightforward calculation shows that its adjoint acts as $V^*(A_1 x_1, \dots, A_n x_n) := \sum_{j=1}^n A_j x_j$, for all $x_1, x_2, \dots, x_n \in E$. Since $\|V^*\| = \|V\|$, we infer that

$$\begin{aligned} \left\| \sum_{j=1}^n A_j x_j \right\|^2 &= \|V^*(A_1 x_1, A_2 x_2, \dots, A_n x_n)\|^2 \\ &\leq \|V^*\|^2 \cdot \|(A_1 x_1, A_2 x_2, \dots, A_n x_n)\|^2 \\ &\leq \left\| \sum_{j=1}^n A_j \right\| \cdot \sum_{j=1}^n \langle A_j x_j, x_j \rangle. \end{aligned}$$

In order to prove minimality, let us fix a constant $C \geq 0$ that satisfies (4.1) with $\left\| \sum_{j=1}^n A_j \right\|$ replaced by C . Then for $x \in E$ the following inequalities hold

$$\left\| \sum_{j=1}^n A_j x \right\|^2 \leq C \cdot \sum_{j=1}^n \langle A_j x, x \rangle \leq C \cdot \left\| \sum_{j=1}^n A_j \right\| \cdot \|x\|^2,$$

thus we have $\left\| \sum_{j=1}^n A_j \right\| \leq C$, indeed. \square

As an immediate consequence of the preceding proposition ($n = 1$) we have

$$(4.2) \quad \|Ax\|^2 \leq \|A\| \langle Ax, x \rangle, \quad x \in E.$$

Now we rephrase our main theorem in the Banach space setting, extending the results obtained in [26, Theorem 3.1] and also [2, Theorem 2.2].

Theorem 4.2. *Let E be a Banach space, and let $A : E \supseteq \text{dom } A \rightarrow \bar{E}'$ be a linear operator. Then the following statements are equivalent.*

- (i) A has a (continuous) positive extension $\tilde{A} \in \mathcal{L}(E; \bar{E}')$,
- (ii) There is a constant $M \geq 0$ such that

$$(4.3) \quad \|Ax\|^2 \leq M \cdot \langle Ax, x \rangle, \quad x \in \text{dom } A,$$

- (iii) For any $y \in H$ there exists $M_y \geq 0$ such that

$$|\langle Ax, y \rangle|^2 \leq M_y \cdot \langle Ax, x \rangle, \quad x \in \text{dom } A.$$

In any case, there exists the Krein-von Neumann extension A_N of A that is the smallest among the set of positive extensions of A . The norm of A_N satisfies

$$(4.4) \quad \|A_N\| = \inf\{M \geq 0 \mid \|Ax\|^2 \leq M \cdot \langle Ax, x \rangle, \quad x \in \text{dom } A\}.$$

If $B, C \in \mathcal{L}(E; E)$ are continuous operators leaving $\text{dom } A$ invariant such that $C^*A \subset AB$ and $B^*A \subset AC$ then the Krein-von Neumann extension of A satisfies

$$(4.5) \quad C^*A_N = A_N B, \quad \text{and} \quad B^*A_N = A_N C.$$

Proof. The equivalence between (i) and (iii) is clear from Theorem 3.1. That (i) implies (ii) follows from (4.2) with $M = \|A\|$. To see that (ii) implies (iii), fix $y \in E$ and observe that

$$|\langle Ax, y \rangle|^2 \leq \|Ax\|^2 \|y\|^2 \leq M \cdot \|y\|^2 \cdot \langle Ax, x \rangle, \quad x \in \text{dom } A.$$

Hence (iii) holds with $M_y = M \cdot \|y\|^2$.

To prove (4.4), let M_A denote the corresponding infimum. It follows from (4.2) that $M_A \leq \|A_N\|$. To see the converse, take $y \in E$, $\|y\| \leq 1$, then for every $x \in \text{dom } A$,

$$|\langle Ax, y \rangle|^2 \leq \|Ax\|^2 \leq M \cdot \langle Ax, x \rangle,$$

hence $\langle Ay, y \rangle \leq M$, according to formula (3.6). Consequently, $\|A_N\|^2 \leq M^2$ follows from

$$\sup_{\substack{y, z \in E, \\ \|y\| = \|z\| = 1}} |\langle A_N y, z \rangle|^2 \leq \sup_{\substack{y, z \in E, \\ \|y\| = \|z\| = 1}} \langle A_N y, y \rangle \langle A_N z, z \rangle \leq M^2.$$

The rest of the statement follows immediately from Theorem 3.3. \square

We remark here that implication (ii) \Rightarrow (iii) could be deduced also from (iii) \Rightarrow (iv) of Theorem 3.1 because inequality (4.3) expresses just that $W(A)$ is bounded in the functional norm, i.e., $W(A)$ is $\beta(\bar{E}', E)$ -bounded.

As an immediate consequence of Theorem 4.2 we recover [24, Theorem 1] concerning positive extendibility of suboperators on a Hilbert space.

Corollary 4.3. *Let H be a Hilbert space and let $A : H \supseteq \text{dom } A \rightarrow H$ be a linear operator. Then the following assertions are equivalent:*

- (i) A has a (continuous) positive extension $\tilde{A} \in \mathcal{B}(H)$,
- (ii) There is a constant $M \geq 0$ such that

$$\|Ax\|^2 \leq M \cdot (Ax | x), \quad x \in \text{dom } A,$$

- (iii) For any $y \in H$ there exists $M_y \geq 0$ such that

$$|(Ax | y)|^2 \leq M_y \cdot (Ax | x), \quad x \in \text{dom } A.$$

In any case, there exists is the Krein-von Neumann extension A_N of A whose norm satisfies

$$\|A_N\| = \inf\{M \geq 0 \mid \|Ax\|^2 \leq M \cdot (Ax | x), \quad x \in \text{dom } A\}$$

If $B, C \in \mathcal{B}(H)$ are bounded operators leaving $\text{dom } A$ invariant such that $C^*A \subset AB$ and $B^*A \subset AC$ then the Krein-von Neumann extension of A satisfies

$$C^*A_N = A_N B, \quad \text{and} \quad B^*A_N = A_N C.$$

Note that for a positive operator $A \in \mathcal{B}(H)$ one has $\|A\| \leq M$ if and only if $A \leq M \cdot I$, I being the identity operator of H . Combining this observation with Theorem 3.2 one immediately gets the following statement.

Corollary 4.4. *Let $A : \text{dom } A \rightarrow H$ be a positive operator satisfying the conditions of Corollary 4.3. Then, for every constant $M \geq \|A_N\|$ there is a positive extension A_{max}^M of A with $\|A_{max}^M\| \leq M$ such that for any positive extension \tilde{A} of A , $\|\tilde{A}\| \leq M$ one has $\tilde{A} \leq A_{max}^M$. In other words,*

$$A_{max}^M = \max\{\tilde{A} \in \mathcal{B}(H) \mid \tilde{A} \geq 0, A \subset \tilde{A}, \|\tilde{A}\| \leq M\}.$$

Furthermore, one has equality

$$[A_N, A_{max}^M] = \{\tilde{A} \in \mathcal{B}(H) \mid \tilde{A} \geq 0, A \subset \tilde{A}, \|\tilde{A}\| \leq M\}.$$

As a concluding result of this section we prove Halmos' result on positive completions of incomplete matrices (see [11, §5, Corollary 2]).

Corollary 4.5. *Let $A := \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & * \end{pmatrix}$ be an "incomplete" operator matrix in a Hilbert space $H = H_0 \oplus H_0^\perp$ with $\text{dom } A = H_0$. Assume that $A_{11} \geq 0$ and that $A_{21} = A_{21}^*$, then the following assertions are equivalent:*

- (i) *there exists $A_{22} \geq 0$ that makes A positive, i.e., $\begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \geq 0$,*
- (ii) *$A_{21}^* A_{21} \leq M \cdot A_{11}$, for some constant $M \geq 0$,*
- (iii) *$\text{ran } A_{21}^* \subseteq \text{ran } A_{11}^{1/2}$.*

Proof. Assume first (i) and prove (ii): Suppose that $B := \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \geq 0$. Since for every positive operator B one has $B^2 \leq \|B\| \cdot B$, we have for $x \in H_0$ that

$$\|A_{11}x\|^2 + \|A_{21}x\|^2 = \left\| B \begin{pmatrix} x \\ 0 \end{pmatrix} \right\|^2 \leq \|B\| \left(B \begin{pmatrix} x \\ 0 \end{pmatrix} \mid \begin{pmatrix} x \\ 0 \end{pmatrix} \right) = \|B\| (A_{11}x \mid x).$$

Since $A_{21}^2 \leq \|A_{11}\| \cdot A_{11}$, this obviously gives (ii).

The equivalence of (ii) and (iii) follows immediately from the Douglas factorization theorem [9]. Finally, assuming (ii) we clearly have

$$\left\| A \begin{pmatrix} x \\ 0 \end{pmatrix} \right\|^2 \leq \|A_{11}\| (A_{11}x \mid x) + M (A_{11}x \mid x), \quad x \in H_0,$$

hence A fulfills (i) of Corollary 4.3. Consequently, A has a bounded positive extension, or equivalently, A can be filled in to get a positive operator. \square

Finally we mention that Friedrichs extension of unbounded operators acting between reflexive Banach spaces and their duals has been investigated by Farkas and Matolcsi in [10].

5. FUNCTIONAL EXTENSIONS ON *-ALGEBRAS

The goal of this section is to show that how abstract operator extensions can be applied for positive functionals on a *-algebra.

Let \mathcal{A} be a not necessarily unital complex *-algebra. Recall that a linear functional f on \mathcal{A} is called *representable* if there exist a Hilbert space H , a *-representation (that is, a *-homomorphism) $\pi : \mathcal{A} \rightarrow \mathcal{B}(H)$ and a vector $\xi \in H$ such that

$$f(a) = (\pi(a)\xi \mid \xi), \quad a \in \mathcal{A}.$$

It is clear that every representable functional is positive, nevertheless the converse is not true in such generality.

Consider now a left ideal \mathcal{I} of \mathcal{A} and a linear functional $f : \mathcal{I} \rightarrow \mathbb{C}$. In this section we provide necessary and sufficient conditions under which f admits a representable extension to \mathcal{A} (cf. also [26] for the Banach- $*$ algebra setting). Recall that if $f : \mathcal{I} \rightarrow \mathbb{C}$ is a linear functional, then we can associate an operator $A : \mathcal{I} \rightarrow \mathcal{A}^*$ to f by setting

$$(5.1) \quad \langle Aa, x \rangle := f(x^*a), \quad x \in \mathcal{A}, a \in \mathcal{I}.$$

Clearly, A is positive if and only if f is positive, i.e., $f(a^*a) \geq 0$ holds for all $a \in \mathcal{I}$.

In what follows we are exclusively interested in representable extensions of functionals. Recall that f is said to be *Hilbert bounded* if there is constant $M \geq 0$ such that

$$(5.2) \quad |f(a)|^2 \leq Mf(a^*a) \quad \text{for all } a \in \mathcal{I},$$

and *admissible* if for any x in \mathcal{A} there exists $\lambda_x \geq 0$ such that

$$(5.3) \quad f(a^*x^*xa) \leq \lambda_x f(a^*a) \quad \text{for all } a \in \mathcal{I}.$$

Theorem 5.1. *Let \mathcal{A} be a $*$ -algebra, $\mathcal{I} \subseteq \mathcal{A}$ be a left ideal, and let $f : \mathcal{I} \rightarrow \mathbb{C}$ be a linear functional. The following assertions are equivalent:*

- (i) *there is a representable functional $\tilde{f} \in \mathcal{A}^*$ extending f ,*
- (ii) *f is admissible and Hilbert bounded.*

If there is any, then there is a minimal one (denoted by f_N) among the set of representable extensions.

Proof. Implication (i) \Rightarrow (ii) is trivial so we only have to prove that if a functional f on a left ideal \mathcal{I} of \mathcal{A} satisfies (5.2) and (5.3) then f admits a minimal representable extension. We are going to imitate the proof of [23, Theorem 1], combined with the construction of the proof of Theorem 3.1: suppose that f is admissible and Hilbert bounded and consider the positive operator $A : \mathcal{I} \rightarrow \mathcal{A}'$, defined by (5.1). Since the following chain of inequalities holds true for all $a \in \mathcal{I}$ and $x \in \mathcal{A}$

$$|\langle Aa, x \rangle|^2 = |f(x^*a)|^2 \leq Mf(a^*xx^*a) \leq \lambda_{x^*} Mf(a^*a) = \lambda_{x^*} M \langle Aa, a \rangle,$$

we conclude that A fulfills condition (iv) of Theorem 3.1, and thus the Krein-von Neumann extension $A_N = JJ^* : \mathcal{A} \rightarrow \mathcal{A}'^*$ exists. Consider the auxiliary Hilbert space H_A , and for every $x \in \mathcal{A}$ define the operator $\pi_f(x) \in \mathcal{B}(H_A)$ as the continuous extension of the densely defined bounded operator

$$\pi_f(x)(Aa) := A(xa), \quad a \in \mathcal{I}.$$

Note that boundedness of $\pi_f(x)$ is guaranteed by admissibility (5.3):

$$\|\pi_f(x)(Aa)\|_A^2 = \langle A(xa), xa \rangle = f(a^*x^*xa) \leq \lambda_x f(a^*a) = \lambda_x \|Aa\|_A^2.$$

We claim that $\pi_f : \mathcal{A} \rightarrow \mathcal{B}(H_A)$ is a $*$ -homomorphism. It is straightforward that π_f is linear and multiplicative. To see that π_f preserves involution, fix $x \in \mathcal{A}$ and $a, b \in \mathcal{I}$, then

$$\begin{aligned} (\pi_f(x)(Aa) | Ab)_A &= \langle A(xa), b \rangle = f(b^*xa) \\ &= f((x^*b)^*a) = \langle Aa, x^*b \rangle = (Aa | \pi_f(x^*)(Ab))_A, \end{aligned}$$

hence $\pi_f(x)^* = \pi_f(x^*)$, indeed. Our next goal is to show the existence of a vector $\zeta_f \in H_A$ such that $f(a) = (\pi_f(a)\zeta_f | \zeta_f)_A$, $a \in \mathcal{I}$. To this aim consider the densely defined linear functional

$$(5.4) \quad \varphi : H_A \supseteq \text{ran } A \rightarrow \mathbb{C}; \quad \varphi(Aa) = f(a), \quad a \in \mathcal{I},$$

which is well defined and continuous because of Hilbert boundedness (5.2):

$$|\varphi(Aa)|^2 \leq Mf(a^*a) = M\|Aa\|_A^2, \quad a \in \mathcal{I}.$$

By the Riesz representation theorem, φ is represented by a unique vector $\zeta_f \in H_A$:

$$(5.5) \quad f(a) = (Aa | \zeta_f)_A, \quad a \in \mathcal{I}.$$

We claim that

$$(5.6) \quad f_N(x) := \overline{\langle J\zeta_f, x \rangle}, \quad x \in \mathcal{A}$$

defines a representable functional such that $f \subset f_N$. To see this, observe first that the following useful identity holds true

$$(5.7) \quad \pi_f(x)\zeta_f = J^*x, \quad x \in \mathcal{A}.$$

Indeed, let $x \in \mathcal{A}$ and $a \in \mathcal{I}$ be fixed. Then by (5.5) we have

$$(Aa | \pi_f(x)\zeta_f)_A = (A(x^*a) | \zeta_f)_A = f(x^*a) = \langle Aa, x \rangle = \langle J(Aa), x \rangle = (Aa | J^*x)_A.$$

Consequently, $f(a) = (Aa | \zeta_f)_A = (J^*a | \zeta_f)_A = \overline{\langle J\zeta_f, a \rangle} = f_N(a)$ holds for $a \in \mathcal{I}$, thus $f \subset f_N$. On the other hand we have

$$(\pi_f(x)\zeta_f | \zeta_f)_A = (J^*x | \zeta_f)_A = \overline{\langle J\zeta_f, x \rangle} = f_N(x), \quad x \in \mathcal{A},$$

hence f_N is representable. All that remains is to prove the minimality of f_N . Let us calculate first the value of f_N on positive elements:

$$\begin{aligned} f_N(x^*x) &= \overline{\langle J\zeta_f, x^*x \rangle} = (J^*(x^*x) | \zeta_f)_A = (J^*x | J^*x)_A \\ &= \sup\{|(Aa | J^*x)_A|^2 | a \in \mathcal{I}, \|Aa\|_A^2 \leq 1\} \\ &= \sup\{|f(x^*a)|^2 | a \in \mathcal{I}, f(a^*a) \leq 1\}, \end{aligned}$$

for every $x \in \mathcal{A}$, where we used the density of $\text{ran } A$ in H_A . Let now \tilde{f} be any representable extension of f . Using identity $\tilde{f} = \tilde{f}_N$, we conclude that

$$\begin{aligned} \tilde{f}(x^*x) &= \sup\{|\tilde{f}(x^*a)|^2 | a \in \mathcal{A}, \tilde{f}(a^*a) \leq 1\} \\ &\geq \sup\{|f(x^*a)|^2 | a \in \mathcal{I}, f(a^*a) \leq 1\} = f_N(x^*x), \end{aligned}$$

hence $f_N \leq \tilde{f}$. The proof is complete. \square

Following the terminology of [26], we will call f_N the Krein–von Neumann extension of f . Note that in the above construction we gained an explicit formula for f_N and also for its values on positive elements.

Corollary 5.2. *Suppose that $f : \mathcal{I} \rightarrow \mathbb{C}$ is admissible and Hilbert-bounded, then the minimal extension f_N satisfies*

$$(5.8) \quad f_N(x) = \overline{\langle J\zeta_f, x \rangle}, \quad x \in \mathcal{A},$$

and

$$(5.9) \quad f_N(x^*x) = \sup\{|f(x^*a)|^2 | a \in \mathcal{I}, f(a^*a) \leq 1\},$$

In case when the algebra has a unit element the theorem can be more easily formulated. In fact, the following simple formula may suggest that extension theory of functionals and extension theory of operators fit nicely together, indeed.

Corollary 5.3. *Assume that \mathcal{A} is a unital $*$ -algebra with unit $1 \in \mathcal{A}$. Assume further that $f : \mathcal{S} \rightarrow \mathbb{C}$ is admissible. Then f is Hilbert bounded, and its Krein-von Neumann extension satisfies*

$$(5.10) \quad f_N(x) = \overline{\langle A_N 1, x \rangle}, \quad x \in \mathcal{A}.$$

Proof. Take $a \in \mathcal{S}$, then

$$|f(a)|^2 = |f(1^*a)|^2 \leq f(1)f(a^*a),$$

hence f is Hilbert bounded. On the other hand, for $a \in \mathcal{S}$,

$$(Aa | J^*1)_A = \langle Aa, 1 \rangle = f(a) = (Aa | \zeta_f)_A,$$

hence $J^*1 = \zeta_f$. This yields

$$\langle A_N 1, x \rangle = (J^*1 | J^*x)_A = (\zeta_f | J^*x)_A = \langle J\zeta_f, x \rangle, \quad x \in \mathcal{A},$$

and hence (5.10). \square

In the next theorem we provide the analogue of Theorem 3.2.

Theorem 5.4. *Let $f : \mathcal{S} \rightarrow \mathbb{C}$ be an admissible and Hilbert bounded functional and fix any representable functional $g \in \mathcal{A}^*$ such that $f_N \leq g$. Then there is a representable extension $f_{max}^g \in \mathcal{A}^*$ of f such that $f_{max}^g \leq g$, and for every representable extension \tilde{f} of f one has $\tilde{f} \leq f_{max}^g$. In other words,*

$$f_{max}^g = \max\{\tilde{f} \in \mathcal{A}^\# \mid \tilde{f} \leq g, f \subset \tilde{f}\}.$$

Furthermore, a representable functional $\tilde{f} \leq g$ is an extension of f if and only if $f_N \leq \tilde{f} \leq f_{max}^g$:

$$[f_N, f_{max}^g] = \{\tilde{f} \in \mathcal{A}^\# \mid \tilde{f} \leq g, f \subset \tilde{f}\}$$

Proof. By assumption, $0 \leq g - f_N \leq g$, hence $g - f_N$ is representable due to [20, Proposition 9.4.22]. Clearly, $g - f \subset g - f_N$, hence $(g - f)_N$ exists. Set

$$f_{max}^g := g - (g - f)_N,$$

which is again representable by [20, Proposition 9.4.22]. One can follow the argument of the proof Theorem 3.2 to show that f_{max}^g possesses every claimed properties.

Finally, let $\tilde{f} \in [f_N, f_{max}^g]$, then for $a \in \mathcal{S}$,

$$|(\tilde{f} - f)(a)|^2 = |(\tilde{f} - f_N)(a)|^2 \leq M'(\tilde{f} - f_N)(a^*a) \leq M'(f_{max}^g - f_N)(a^*a) = 0$$

for some $M' \geq 0$, because of Hilbert boundedness. Hence $f \subset \tilde{f}$, which completes the proof. \square

REFERENCES

- [1] D. Alpay, S. Gabrielyan, Positive definite functions and dual pairs of locally convex spaces, *Opuscula Math.* 38, no. 4 (2018), 463–482.
- [2] D. Alpay, O. Timoshenko, D. Volok, Carathodory functions in the Banach space setting, *Linear Algebra and its Applications*, 425 (23) 2007, 700–713.
- [3] T. Ando, W. Szymański, Order Structure and Lebesgue Decomposition of Positive Definite Operator Functions, *Indiana Univ. Math. J.*, 35 (1986), 157–173.
- [4] Gr. Arsene and A. Gheondea, Completing Matrix Contractions, *Journal of Operator Theory*, Vol. 7, No. 1 (1982), 179–189.
- [5] M. S. Ashbaugh, F. Gesztesy, A. Laptev, M. Mitrea, S. Sukhtaiev, A bound for the eigenvalue counting function for Krein-von Neumann and Friedrichs extensions, *Advances in Mathematics*, 304 (2017), 1108–1155.

- [6] M. S. Ashbaugh, F. Gesztesy, M. Mitrea, R. Shterenberg, G. Teschl, The Krein von Neumann extension and its connection to an abstract buckling problem, *Mathematische Nachrichten*, Vol. 283 (2) 2010, 165–179.
- [7] D. Baidiuk, S. Hassi, Completion, extension, factorization, and lifting of operators, *Math. Ann.* 364 (2016), no. 3-4, 1415–1450.
- [8] J. F. Brasche, H. Neidhardt, Some Remarks on Krein’s Extension Theory, *Mathematische Nachrichten*, Vol 165 (1994), 159–181.
- [9] R. G. Douglas, On majorization, factorization, and range inclusion of operators on Hilbert space, *Proc. Amer. Math. Soc.*, 17 (1966), 413–415.
- [10] B. Farkas, M. Matolcsi, Positive forms on Banach spaces, *Acta Mathematica Hungarica*, 2003 (99), 43–55.
- [11] P. R. Halmos, Subnormal operators and the subdiscrete topology, *Anniversary Volume on Approximation Theory and Functional Analysis*, Birkhuser, Basel, 49–65, 1984.
- [12] S. Hassi, M. Malamud, H. de Snoo, On Kren’s extension theory of nonnegative operators, *Mathematische Nachrichten*, Vol. 274-275 (2004), 40–73.
- [13] S. Hassi, A. Sandovici, H. de Snoo, H. Winkler, A general factorization approach to the extension theory of nonnegative operators and relations, *J. Operator Theory* 58 (2007), 351–386.
- [14] S. Hassi, Z. Sebestyén, H.S.V. de Snoo, On the nonnegativity of operator products, *Acta Math. Hungar.* 109 (2005), 1–14.
- [15] S. Hassi, Z. Sebestyén, H. de Snoo, Lebesgue type decompositions for nonnegative forms, *Journal of Functional Analysis*, 257 (2009), 3858–3894.
- [16] R. V. Kadison, A generalized Schwarz inequality and algebraic invariants for operator algebras, *Annals of Mathematics*, 56 (1952), 494–503.
- [17] M. G. Krein, The theory of self-adjoint extensions of semi-bounded Hermitian transformations and its applications. I, *Rec. Math. [Mat. Sbornik] N.S.* 20 (62) (1947), 431–495.
- [18] P. Masani, Dilations as Propagators of Hilbertian Varieties, *SIAM J. Math. Anal.*, 9(3), 414–456.
- [19] J. von Neumann, Allgemeine Eigenwerttheorie Hermitescher Funktionaloperatoren, *Math. Ann.* 102, 49–131.
- [20] T. W. Palmer, *Banach Algebras and the General Theory of *-Algebras II*, Cambridge University Press (2001).
- [21] H. H. Schaefer, Topological vector spaces, Third printing corrected. Graduate Texts in Mathematics, Vol. 3. *Springer-Verlag*, New York-Berlin, 1971.
- [22] C. F. Skau, Positive self-adjoint extensions of operators affiliated with a von Neumann algebra, *Mathematica Scandinavica*, 44 (1979), 171–195.
- [23] Z. Sebestyén, On representability of linear functionals on *-algebras, *Period. Math. Hungar.*, 15 (1984), 233–239.
- [24] Z. Sebestyén, Operator extensions on Hilbert space, *Acta Sci. Math. (Szeged)*, 57 (1993), 233–248.
- [25] Z. Sebestyén, Zs. Tarcsay, On the adjoint of Hilbert space operators, *Linear and Multilinear Algebra*, 2018, online first, DOI: 10.1080/03081087.2018.1430120.
- [26] Z. Sebestyén, Zs. Szűcs, Zs. Tarcsay, Extensions of positive operators and functionals, *Linear Algebra Appl.*, 472 (2015), 54–80.
- [27] W. Szymański, Positive forms and dilations, *Trans Amer Math.* 301(2) (1987), 761–780.
- [28] Zs. Tarcsay, Operator extensions with closed range, *Acta Math. Hungar.*, 135 (2012), 325–341.
- [29] Zs. Tarcsay, Closed range positive operators on Banach spaces, *Acta Math. Hungar.*, 142 (2014), 494–501.
- [30] A. Wilansky, Modern methods in topological vector spaces, *McGraw-Hill International Book Co.*, New York, 1978.

ZS. TARCSAY, DEPARTMENT OF APPLIED ANALYSIS AND COMPUTATIONAL MATHEMATICS,
EÖTVÖS LORÁND UNIVERSITY, PÁZMÁNY PÉTER SÉTÁNY 1/C., BUDAPEST H-1117, HUNGARY
E-mail address: `tarcsay@cs.elte.hu`

T. TITKOS, ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, HUNGARIAN ACADEMY OF SCIENCES,
REÁLTANODA UTCA 13-15., BUDAPEST H-1053, HUNGARY, AND BBS UNIVERSITY OF APPLIED
SCIENCES, ALKOTMÁNY U. 9., BUDAPEST H-1054, HUNGARY
E-mail address: `titkos.tamas@renyi.mta.hu`