



**PUSAT PENGAJIAN TEKNOLOGI INDUSTRI
UNIVERSITI SAINS MALAYSIA**

**BORANG PENYERAHAN DISERTASI MUTAKHIR
SATU (1) NASKAH**

Nama penyelia: Dr Maizura Murad

Bahagian: Teknologi Makanan

Saya telah menyemak semua pembetulan/pindaan yang dilaksanakan oleh Encik/Puan/Cik OON LI JUEN

mengenai disertasinya sebagaimana yang dipersetujui oleh Panel Pemeriksa di *Viva Voce*-nya.

2. Saya ingin mengesahkan bahawa saya berpuas hati dengan pembetulan/pindaan yang dilaksanakan oleh calon.

Sekian, terima kasih.

Maizura Murad, Ph.D
Lecturer
Food Technology Division
School of Industrial Technology
Universiti Sains Malaysia
11800 USM, Penang, Malaysia

(Tandatangan dan cop)

08 Ogos 2020

(Tarikh)



**DEVELOPMENT OF DRIED BLACK GRASS JELLY
(*MESONA CHINENSIS*) CONTAINING DIFFERENT
TAPIOCA AND SAGO STARCH RATIO**

by

OON LI JUEN

A dissertation submitted in partial fulfillment of the requirements for the degree of
Bachelor of Technology (B. Tech) in the field of Food Technology
School of Industrial Technology
Universiti Sains Malaysia

July 2020

DECLARATION BY AUTHOR

This dissertation is composed of my original work and contains no material previously published or written by another person except where due reference has been made in the text. The content of my dissertation is the result of work which I have carried out since the commencement of my research project and does not include a substantial part of work that has been submitted to qualify for the award of any other degree or diploma in any university or other tertiary institution.

A handwritten signature in black ink, appearing to read 'Oon Li Juén', is written above a horizontal line.

OON LI JUEN

JULY 2020

ACKNOWLEDGEMENTS

First and foremost, I would like to express my sincere gratitude and appreciation to my supervisor, Dr. Maizura Murad for allocating her precious time in providing valuable advice and guidance as well as knowledge throughout my final year research project despite her busy lecture and research schedules. Besides, I would also like to give special thanks to the School of Industrial Technology and Centre for Global Archaeological Research, Universiti Sains Malaysia (USM) for allowing me to use the equipment and facilities in the processing and research labs throughout my research project. Moreover, I would like to thank all the lab technicians, namely Mr. Maarof Salleh, Mr. Mohamad Firdaus bin Mohd Adnan and Mr. Abdul Rahim bin Md Sari for kindly providing their assistance aside from sharing their laboratory knowledge and technique in using both laboratory facilities and equipment. I was able to complete the analyses required for my study smoothly and on time due to their great assistance throughout my final year research project.

In addition, I would like to express my appreciation towards Miss Siti Rashima binti Romli, a Ph.D. student who is currently pursuing her studies in USM. Throughout my whole research project, she had guided me with patience and shared valuable knowledge on conducting the analyses required for my study. Last but not least, I would like to express my sincere thanks to my beloved family, coursemates and friends who have given me continuous encouragement, moral support and assistance without any hesitation upon completion of my final year research project.

OON LI JUEN

JULY 2020

TABLE OF CONTENTS

| | Page |
|---|-------------|
| Acknowledgements | iii |
| Table of Contents | iv |
| List of Tables | vii |
| List of Figures | viii |
| List of Abbreviations | ix |
| Abstrak | x |
| Abstract | xii |
| CHAPTER 1 INTRODUCTION | |
| 1.1 Research Background | 1 |
| 1.2 Problem Statement/Rationale of Study | 2 |
| 1.3 Hypothesis | 3 |
| 1.4 Objectives | 3 |
| CHAPTER 2 LITERATURE REVIEW | |
| 2.1 Black Grass Jelly | 4 |
| 2.1.1 Health Benefits of Black Grass Jelly | 4 |
| 2.1.2 Processing of Black Grass Jelly | 5 |
| 2.1.3 Textural Properties of Black Grass Jelly | 6 |
| 2.2 Starch | 7 |
| 2.2.1 Types of Starch | 8 |
| 2.2.1a Tapioca Starch | 8 |
| 2.2.1b Sago Starch | 9 |
| 2.2.2 Synergistic Interactive Effect of Starch Mixtures | 10 |
| 2.2.3 Retrogradation of Starch | 12 |

| | |
|---|----|
| 2.2.3a Effects of Mixed Starch System on The Rate of Retrogradation | 13 |
| 2.3 Effects of Drying on The Quality of Food Products | 15 |
| 2.3.1 Application of Different Drying Treatments on Food Products | 18 |
| 2.4 Effects of Rehydration on The Quality of Food Products | 20 |
| CHAPTER 3 MATERIALS AND METHODS | |
| 3.1 Materials | 23 |
| 3.2 Overall Experimental Design | 23 |
| 3.3 Phase 1 | 25 |
| 3.3.1 Preparation of Cincau Liquid Extract | 25 |
| 3.3.2 Preparation of Black Grass Jelly | 25 |
| 3.3.3 Physical Analysis of Fresh Black Grass Jelly | 26 |
| 3.3.3a Determination of Moisture Content | 26 |
| 3.3.3b Colour Measurement | 26 |
| 3.3.3c Texture Profile Analysis (TPA) | 27 |
| 3.3.3d Determination of Syneresis | 27 |
| 3.4 Phase 2 | 28 |
| 3.4.1 Sample Preparation | 28 |
| 3.4.1a Preparation of Dried Black Grass Jelly | 28 |
| 3.4.1b Preparation of Rehydrated Black Grass Jelly | 28 |
| 3.4.2 Physical Analysis of Dried Black Grass Jelly | 29 |
| 3.4.2a Determination of Moisture Content | 29 |
| 3.4.3 Physical Analysis of Rehydrated Black Grass Jelly | 29 |
| 3.4.3a Colour Measurement | 29 |
| 3.4.3b Rehydration Capacity | 30 |

| | |
|---|----|
| 3.5 Statistical Analysis | 30 |
| CHAPTER 4 RESULTS AND DISCUSSION | |
| 4.1 Moisture Content of Fresh Black Grass Jelly | 32 |
| 4.2 Colour Measurement of Fresh Black Grass Jelly | 35 |
| 4.3 Texture Profile Analysis (TPA) of Fresh Black Grass Jelly | 39 |
| 4.4 Syneresis of Fresh Black Grass Jelly | 44 |
| 4.5 Moisture Content of Black Grass Jelly After Drying | 46 |
| 4.6 Colour Measurement of Rehydrated Black Grass Jelly | 52 |
| 4.7 Rehydration Capacity of Black Grass Jelly | 57 |
| CHAPTER 5 CONCLUSIONS AND RECOMMENDATIONS | |
| 5.1 Conclusion | 62 |
| 5.2 Recommendations for Future Study | 63 |
| REFERENCES | 65 |
| APPENDICES | 75 |

LIST OF TABLES

| Table | Caption | Page |
|--------------|--|-------------|
| 3.1 | Formulations of black grass jelly | 25 |
| 3.2 | F values and P values (two-way ANOVA) of the interaction between two parameters (different tapioca and sago starch ratio and drying time) on the rate of syneresis for fresh black grass jelly and moisture content of dried black grass jelly | 31 |
| 4.1 | Moisture content of fresh black grass jelly | 32 |
| 4.2 | Colour parameters of fresh black grass jelly | 36 |
| 4.3 | Texture Profile Analysis (TPA) of fresh black grass jelly with different formulations | 40 |
| 4.4 | Syneresis of fresh black grass jelly with different formulations after storage at 4 °C for 24 and 48 hours | 45 |
| 4.5 | Moisture content of black grass jelly with different formulations after drying at 70 °C for 3, 6, 9, 12, 24 and 30 hours | 47 |
| 4.6 | Colour parameters of rehydrated black grass jelly | 52 |
| 4.7 | Rehydration ratio of dried black grass jelly | 57 |

LIST OF FIGURES

| Figure | Caption | Page |
|---------------|---|-------------|
| 3.1 | Overall flow chart of the experiment | 24 |
| 4.1 | Drying curve of black grass jelly | 50 |
| 4.2 | Colour difference (ΔE) of rehydrated black grass jelly samples with different formulations | 56 |
| 4.3 | Physical observation of dried black grass jelly before rehydration: (a) Control, sago and tapioca starch (0:100); (b) S2, sago and tapioca starch (50:50); (c) S4, sago and tapioca starch (100:0). | 59 |
| 4.4 | Physical observation of rehydrated black grass jelly: (a) Control, sago and tapioca starch (0:100); (b) S2, sago and tapioca starch (50:50); (c) S4, sago and tapioca starch (100:0). | 60 |

LIST OF ABBREVIATIONS

Abbreviations

M. chinensis

M_e

Wt.

Definitions

Mesona chinensis

Equilibrium moisture content

Weight

PENGHASILAN CINCAU HITAM KERING (*MESONA CHINENSIS*) YANG MENGANDUNGI NISBAH KANJI TAPIOKA DAN SAGU YANG BERBEZA

ABSTRAK

Cincau hitam ialah pencuci mulut yang dihasilkan dengan menggunakan ekstrak daun cincau kering daripada pokok *Mesona chinensis*. Cincau hitam komersial mempunyai jangka hayat pendek apabila disimpan pada suhu bilik. Ini adalah disebabkan oleh penggunaan kanji tapioka yang mengakibatkan kadar sineresis cincau menjadi lebih tinggi dengan tekstur yang tidak stabil. Oleh itu, objektif kajian ini adalah untuk menghasilkan cincau hitam kering dengan nisbah kanji tapioka dan sagu yang berbeza untuk meningkatkan jangka hayat cincau hitam. Cincau hitam dihasilkan dengan menggunakan nisbah kanji sagu dan tapioka yang berbeza (0:100, 25:75, 50:50, 75:25, 100:0) dan seterusnya dikeringkan pada masa pengeringan yang berbeza (0, 3, 6, 9, 12, 24, 30 jam) serta dihidrasikan semula pada 90 °C untuk 15 minit. Cincau hitam kering dengan menggunakan nisbah kanji sagu dan tapioka yang berbeza telah berjaya dihasilkan. Kesan nisbah kanji sagu dan tapioka yang berbeza serta masa pengeringan terhadap ciri-ciri fizikal cincau hitam telah diselidiki. Dengan ini, analisis-analisis seperti kandungan kelembapan, warna, tekstur, sineresis, kinetik pengeringan (kandungan kelembapan berlawanan masa pengeringan) dan keupayaan rehidrasi telah dijalankan. Kandungan kelembapan cincau hitam segar yang mengandungi kedua-dua jenis kanji berkurang secara signifikan ($p < 0.05$) dengan peningkatan nisbah kanji sagu. Nilai-nilai parameter warna serta tekstur kecuali keanjalan, cincau hitam segar bertambah secara signifikan ($p < 0.05$) berbanding dengan sampel kawalan apabila nisbah kanji sagu ditambah sehingga melebihi 75%. Penggunaan nisbah kanji sagu dan tapioka berbeza tidak mempunyai kesan signifikan ($p > 0.05$) terhadap sineresis cincau hitam segar selepas disimpan selama 24 jam. Sineresis cincau hitam segar bertambah

secara signifikan ($p < 0.05$) dengan penambahan masa penyimpanan. Hubungan tidak linear telah dikesan pada graf keluk pengeringan cincau hitam yang dianalisis. Cincau hitam dihidrasi semula yang dihasilkan dengan nisbah kanji sagu dan tapioka yang sama (50:50) mempunyai perbezaan warna yang paling tinggi ($p < 0.05$) serta nisbah rehidrasi yang paling rendah ($p < 0.05$) berbanding dengan sampel kawalan (100% kanji tapioka), dan ini adalah disebabkan oleh pemerhatian tahap pengecutan cincau hitam kering tersebut yang lebih tinggi. Kombinasi kanji sagu dan tapioka adalah tidak baik untuk sifat-sifat struktur cincau hitam selepas dikeringkan. Kesimpulannya, cincau hitam kering dihasilkan dengan 100% kanji tapioka atau kanji sagu mempunyai kualiti yang lebih baik berbanding dengan yang dihasilkan dengan nisbah kanji sagu dan tapioka yang sama (50:50).

DEVELOPMENT OF DRIED BLACK GRASS JELLY (*MESONA CHINENSIS*) CONTAINING DIFFERENT TAPIOCA AND SAGO STARCH RATIO

ABSTRACT

Black grass jelly is a herbal dessert made from dried leaves extracts of *Mesona chinensis* plant. Commercial black grass jelly exhibits a short shelf life when stored at room temperature due to the usage of tapioca starch which results in a higher syneresis rate and unstable texture. Hence, the objective of this research is to develop dried black grass jelly with different tapioca and sago starch ratio for increasing the shelf life of the black grass jelly. The black grass jelly was formulated with different sago and tapioca starch ratio (0:100, 25:75, 50:50, 75:25, 100:0), thereafter dried at different drying time (0, 3, 6, 9, 12, 24, 30 hr) and rehydrated at 90 °C for 15 min. The development of dried black grass jelly containing different tapioca and sago starch ratio was successful. The effects of different sago and tapioca starch ratio and drying time on the physical properties of black grass jelly were determined. Hence, analyses on the moisture content, colour, texture, syneresis, drying kinetics (moisture content against drying time) and rehydration capacity were conducted. The increase in sago starch ratio significantly decreased ($p < 0.05$) the moisture content of fresh black grass jelly consisting of both starches. The values of all the colour parameters and textural parameters except springiness, of the fresh black grass jelly increased significantly ($p < 0.05$) in comparison with control sample when the sago starch ratio was increased to more than 75%. Different sago and tapioca starch ratio did not significantly affect ($p > 0.05$) the syneresis of black grass jelly after storage for 24 hr, however the syneresis of fresh black grass jelly significantly ($p < 0.05$) increased with the increase in storage time. A non-linear relationship was observed on the drying curve of black grass jelly. Rehydrated black grass jelly with an equal sago to tapioca starch ratio (50:50) had the

greatest ($p < 0.05$) colour difference with a significantly lower ($p < 0.05$) rehydration ratio in comparison with the control sample (100% tapioca starch) due to the higher degree of shrinkage. The combination of sago and tapioca starches was not favourable for the black grass jelly structure properties after drying. Thus, it was deduced that the quality of dried black grass jelly containing either 100% tapioca or sago starch was better than that containing equal sago to tapioca starch ratio (50:50).