

Histopathological and cholinesterase changes in the gills of *Clarias gariepinus* as a result of cadmium exposure

ABSTRACT

Aim : The cholinesterase (ChE) based inhibition and histopathological studies from fish were investigated and represented in this study to develop as one of the great potential biomarkers for heavy metals monitoring. **Methodology :** In this study, the histopathological study of gills were observed a under microscope. The capability of ChE extracted from the gills of *Clarias gariepinus* was assessed for declining Cd. ChE was purified through affinity chromatography and continued with the optimisation and inhibition study (IC50) of cholinesterase. **Results :** Histopathological study of gills was carried out and several changes such as aneurysm, necrosis and lamella fusion were noted. Purification fold obtained from purified enzyme was 1.15 with 30% a yield specific activity 20.726. The optimum temperature for purified AChE was 35°C along with acetylthiocholine iodide (ATC) as a preferable substrate that had the highest Vmax value of 0.5452 U mg⁻¹ and the lowest Km value of 0.0311 mM. The optimum pH was observed to be 10 of Tris-HCl as a medium. Meanwhile, the IC50 of cadmium was 6.808 mg l⁻¹ with R2 value of 0.9532. **Interpretation :** The result of the study can be used as a tool for further developing a biomarker for the detection of heavy metals in aquatic ecosystems. In addition, the baseline data provided can also be used for designing a kit, which would give rapid and accurate result.