

Activation of Invariant NKT Cells Exacerbates Experimental Visceral Leishmaniasis

Amanda C. Stanley¹, Yonghong Zhou¹, Fiona H. Amante¹, Louise M. Randall¹, Ashraf Haque¹, Daniel G. Pellicci², Geoff R. Hill¹, Mark J. Smyth³, Dale I. Godfrey², Christian R. Engwerda^{1*}

1 Immunology and Infection Laboratory, Queensland Institute of Medical Research, Herston, Queensland, Australia, **2** Department of Microbiology and Immunology, University of Melbourne, Parkville, Victoria, Australia, **3** Cancer Immunology Program, Trescowthick Laboratories, Peter MacCallum Cancer Centre, East Melbourne, Victoria, Australia

Abstract

We report that natural killer T (NKT) cells play only a minor physiological role in protection from *Leishmania donovani* infection in C57BL/6 mice. Furthermore, attempts at therapeutic activation of invariant NKT (iNKT) cells with α -galactosylceramide (α -GalCer) during *L. donovani* infection exacerbated, rather than ameliorated, experimental visceral leishmaniasis. The inability of α -GalCer to promote anti-parasitic immunity did not result from inefficient antigen presentation caused by infection because α -GalCer-loaded bone marrow-derived dendritic cells were also unable to improve disease resolution. The immune-dampening affect of α -GalCer correlated with a bias towards increased IL-4 production by iNKT cells following α -GalCer stimulation in infected mice compared to naïve controls. However, studies in IL-4-deficient mice, and IL-4 neutralisation in cytokine-sufficient mice revealed that α -GalCer-induced IL-4 production during infection had only a minor role in impaired parasite control. Analysis of liver cell composition following α -GalCer stimulation during an established *L. donovani* infection revealed important differences, predominantly a decrease in IFN γ ⁺ CD8⁺ T cells, compared with control-treated mice. Our data clearly illustrate the double-edged sword of NKT cell-based therapy, showing that in some circumstances, such as when sub-clinical or chronic infections exist, iNKT cell activation can have adverse outcomes.

Citation: Stanley AC, Zhou Y, Amante FH, Randall LM, Haque A, et al. (2008) Activation of Invariant NKT Cells Exacerbates Experimental Visceral Leishmaniasis. PLoS Pathog 4(2): e1000028. doi:10.1371/journal.ppat.1000028

Editor: Ingrid Muller, Imperial College London, United Kingdom

Received: June 26, 2007; **Accepted:** January 7, 2008; **Published:** February 29, 2008

Copyright: © 2008 Stanley et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by grants from the Australian NHMRC and the UNICEF/UNDP/World Bank/WHO Special Programme for Research and Training in Tropical Diseases (TDR). CRE is an Australian NHMRC Career Development Fellow, DIG is an Australian NHMRC Principal Research Fellow, GRH is an Australian NHMRC Practitioner Fellow, and MJS is an Australian NHMRC Senior Principal Research Fellow.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: Christian.Engwerda@qimr.edu.au

Introduction

Natural killer T (NKT) cells are a unique subset of CD1d-restricted T cells that provide a link between innate and adaptive immune responses. In mice, invariant NKT (iNKT) cells express a semi-invariant TCR, consisting of a V α 14J α 18 TCR α -chain and a TCR β -chain biased towards V β 8.2, V β 2, and V β 7 expression (reviewed in [1]). Type II NKT cells are another cell subset in mice with more diverse TCR expression [1–3]. Upon stimulation, iNKT cells rapidly produce large quantities of pro- and anti-inflammatory cytokines, resulting in activation of other immune cells such as NK cells and conventional T cells [4–7] (also reviewed in [8]). NKT cells recognise and respond to glycolipid antigens presented on CD1d molecules. The most well-defined antigen for iNKT cells is α -galactosylceramide (α -GalCer), a marine sponge-derived glycolipid that specifically targets iNKT cells and no other lymphocyte populations directly [9]. The activation of iNKT cells by α -GalCer can enhance resistance in several infectious disease models, including viral, bacterial and parasitic infections (reviewed in [10–12]). Among parasitic infections studied, α -GalCer has been shown to enhance resistance to malaria [13], trypanosomiasis [14] and toxoplasmosis [15]. The ability of iNKT cells to produce IFN γ following stimulation

with α -GalCer is important for this therapeutic effect and host protection during infection, although the robust induction of TNF, IL-4 and IL-13 by iNKT cells also occurs (reviewed in [8,16,17]). However, the connection between therapeutically induced NKT cell responses and physiological NKT cell responses is not always clear. Nevertheless, there are parallels between physiological and therapeutic NKT cell responses in some disease models. For example, in experimental tumour models, the growth of methylcholanthrene (MCA)-induced sarcoma cell lines is restricted by physiological IFN γ produced by endogenous NKT cells [18], while many other experimental cell lines, including the B16-F10 melanoma are only controlled by NKT cells following therapeutic activation with α -GalCer, again in an IFN γ -dependent manner [7]. It is also possible that following pathogen challenge, NKT cells will be exposed to foreign glycolipids, or self glycolipids, that, in the inflammatory environment, can trigger NKT cell responses that are similar to those induced by α -GalCer [19–22] (reviewed in [23]).

Visceral leishmaniasis (VL) is a potentially fatal human disease caused by infection with *Leishmania donovani* or *L. infantum* (*chagasi*) [24,25]. Infection of genetically susceptible mice with *L. donovani* results in an acute but ultimately resolving infection in the liver associated with the development of granulomas around infected

Author Summary

Natural killer T (NKT) cells are a unique subset of T cells that can produce large quantities of inflammatory cytokines very rapidly upon stimulation. They are known to be strongly stimulated by a molecule called α -galactosylceramide (α -GalCer) that is derived from a marine sponge, and in this way α -GalCer is hoped to provide effective immunotherapy for a wide range of diseases. We attempted to stimulate NKT cells with α -GalCer in mice infected with *Leishmania donovani*, a protozoan parasite that causes a chronic disease known as visceral leishmaniasis in humans. *L. donovani* characteristically causes an acute resolving infection in the liver where NKT cells are abundant. Therefore, we hypothesised that by stimulating these cells with α -GalCer we would improve the rate of hepatic disease resolution. However, while α -GalCer administered prior to infection had no effect on hepatic parasite burden, α -GalCer administered during an established infection exacerbated hepatic disease, associated with a decrease in IFN γ -producing CD8⁺ T cells. These results are important as they demonstrate that therapies aimed at modulating NKT cell function are not always beneficial, and adverse consequences may occur in certain circumstances, such as in the presence of persistent and/or sub-clinical infections.

Kupffer cells [26]. In contrast, a chronic infection develops in the spleen, associated with severe immunopathology [27,28]. Parasite numbers increase more slowly in the spleen, and total splenic parasite burdens usually only reach 5–10% of maximum levels in the liver, often with greater variation between individual mice than in the livers of the same animals [28,29]. NKT cells have been shown to regulate CXCL10 expression in the livers of C57BL/6 mice in the early stages of *L. donovani* infection, an event thought to be important for hepatic granuloma development [30]. A subset of mouse NKT cells has also been reported to be activated by *L. donovani* lipophosphoglycan presented by host CD1d [31]. Studies in CD1d-deficient BALB/c mice have also suggested that NKT cells are important for the efficient control of *L. donovani* growth [31]. In addition, infection of immature human DC with *L. infantum* results in increased CD1d cell surface expression, and subsequent increased recognition and killing by IFN γ -producing iNKT cells [32]. CD4⁺ NKT cells have also been implicated in protection against the early stages of cutaneous leishmaniasis

caused by *L. major* infection in genetically resistant mouse strains [33]. However, a more recent study has shown that clinical recovery from *L. major* infection can occur in NKT cell-deficient mice in the same time period as wild type mice, despite this early NKT cell role in protection [34].

Here we investigated the role of NKT cells during experimental VL in C57BL/6 mice, and tested whether stimulation of iNKT cells with α -GalCer could enhance anti-parasitic activity. Our data indicate that NKT cells are neither required for the development of immunity in the liver, nor for the control of parasite burden in the spleen following *L. donovani* infection. Furthermore, iNKT cell activation by α -GalCer hinders disease resolution in the liver. These results have important implications for the modulation of iNKT cell function during established disease.

Results

The Role of NKT Cells in the Control of *L. donovani* Infection in C57BL/6 Mice

To determine the relative roles of iNKT cells and type II NKT cells during experimental VL, we infected C57BL/6 mice deficient in iNKT cells only (B6.J α 18^{-/-}) and all CD1d-restricted NKT cells (B6.CD1d^{-/-}) with *L. donovani* and compared their course of infection to C57BL/6 controls. Only minor differences in parasite burdens were found between NKT cell-deficient mice and C57BL/6 controls. B6.J α 18^{-/-} mice had significantly higher hepatic parasite burdens than C57BL/6 mice at day 56 post-infection (p.i.) ($p < 0.05$; Figure 1A), suggesting that iNKT cells may be required for optimal control of parasite growth at later stages of infection. However, C57BL/6 and B6.J α 18^{-/-} mice had resolved hepatic infection to a similar extent by day 90 and 180 p.i. (liver parasite burdens were 93 \pm 42 Leishman-Donovan units (LDU) versus 119 \pm 36 LDU in C57BL/6 versus B6.J α 18^{-/-} mice at day 90 p.i., and 0.4 \pm 0.5 LDU versus 6.2 \pm 3.0 LDU in C57BL/6 versus B6.J α 18^{-/-} mice at day 180 p.i.). A small but significant difference ($p < 0.05$) in hepatic parasite burdens was also observed between B6.CD1d^{-/-} and C57BL/6 mice at day 7 p.i. (Figure 1A) suggesting a minor role for type II NKT cells in the early control of *L. donovani* growth. Alternatively, the CD1d molecule itself may signal to antigen presenting cells independently of NKT cells [35], thereby contributing to anti-parasitic immunity. B6.CD1d^{-/-} mice had resolved hepatic *L. donovani* infection to the same extent as C57BL/6 control mice by day 56 p.i. (Figure 1A), and were not examined after this time point. There was no decrease in the total




Figure 1. NKT Cells Are Not Required for Control of *L. donovani* Infection. Female C57BL/6 (closed squares), B6.J α 18^{-/-} (open circles) and B6.CD1d^{-/-} (open triangles) mice were infected with *L. donovani*, and parasite burdens were monitored from day 7 to day 56 p.i. in the liver (A) and spleen (B). Data represent the mean \pm SEM of parasite burdens (LDU) from four mice per group for each time point. One representative experiment of two performed with similar outcome is shown. Significant differences of $p < 0.05$ between C57BL/6 and B6.J α 18^{-/-} mice (#) and C57BL/6 and B6.CD1d^{-/-} mice (*) are indicated.

doi:10.1371/journal.ppat.1000028.g001

numbers of IFN γ - or TNF-producing cells in the liver measured by ELISPOT at any time point in any group of mice (Figure S1). In addition, parasite burdens in the spleen were similar in all three groups of mice throughout the first 56 d of infection (Figure 1B) and until day 180 p.i. in C57BL/6 and B6.J α 18 $^{-/-}$ mice (splenic parasite burdens were 24 ± 17 LDU versus 18 ± 5 LDU in C57BL/6 versus B6.J α 18 $^{-/-}$ mice at day 90 p.i., and 5 ± 3 LDU versus 10 ± 3 LDU in C57BL/6 versus B6.J α 18 $^{-/-}$ mice at day 180 p.i.). Therefore, control of parasite growth in the liver was achieved in the absence of both iNKT and type II NKT cells, with only minor roles for these cells at different times following infection.

Activation of iNKT Cells with α -GalCer at the Time of *L. donovani* Infection Does Not Improve Disease Resolution

The links between physiological NKT cell responses and iNKT responses induced by ligands such as α -GalCer are not always apparent [8], and even though we were unable to detect a clear physiological role for NKT cells in protection against *L. donovani*, it was important to determine whether therapeutic iNKT cell activation with α -GalCer could improve disease resolution. Therefore, we next tested whether iNKT cells could be activated by α -GalCer at the time of infection to improve control of *L. donovani* growth. We focused these studies on the liver because NKT cells comprise a major cell population in this tissue site and parasite burdens are significantly higher and more consistent in the liver than in the spleen. The activity of α -GalCer was confirmed prior to all experiments by measuring IFN γ production by hepatic iNKT cells 2 h following α -GalCer administration (data not shown). A single injection of α -GalCer was administered so that the iNKT cell response was not tolerised or biased towards a Th2 response, as has previously been reported [12,36–38]. First, α -GalCer was administered the day prior to infection, and no differences in hepatic parasite burdens were observed in treated mice, compared with controls on day 14 p.i. (Figure 2A). The same result was observed when α -GalCer was administered to mice 2 h prior to infection (Figure 2B). Stimulation with α -GalCer-loaded bone marrow-derived dendritic cells (BMDC) results in more prolonged cytokine responses than with soluble α -GalCer in mice due to selective targeting of the glycolipid to DC and improved presentation of the antigen to iNKT cells [37]. *L. donovani* has also been shown to interfere with antigen processing pathways [39]. Although there is no direct evidence for CD1d antigen presentation being compromised during experimental VL, we sought to eliminate this possibility by delivering α -GalCer-loaded

BMDC to mice 2 h prior to infection. However, despite the stimulation of efficient iNKT cell activation by α -GalCer-loaded BMDC, demonstrated by increased serum cytokine levels (data not shown), and as previously reported [37] this mode of α -GalCer delivery also had no impact on hepatic parasite burdens relative to control mice that had received vehicle-loaded BMDC (Figure 2B).

α -GalCer Treatment at the Time of iNKT Cell Activation in Response to *L. donovani* Infection Enhances Parasite Growth

We next determined whether and when iNKT cells were first activated in response to *L. donovani* infection, so that we could test whether α -GalCer stimulation at this time was effective at activating these cells for anti-leishmanial effects. Although the percentage of iNKT cells in the liver was reduced at day 14 p.i. (Figure 3A), due to the hepatic recruitment of other leukocytes during granuloma formation (reviewed in [26]), total iNKT cell numbers doubled by day 7 p.i., and this increase was sustained at day 14 p.i. (Figure 3B) due to an approximate 10-fold increase in total liver mononuclear cell (MNC) number. There was also evidence of TCR down-regulation by iNKT cells at day 7 p.i. (data not shown) and day 14 p.i. (Figure 3A), but not earlier, suggesting TCR-dependent NKT cell activation, as seen following α -GalCer stimulation [40–42]. Furthermore, NK1.1 expression on iNKT cells was also reduced at these times, consistent with activation of these cells (Figure 3C). The expression of CD69 on iNKT cells also increased above normal intermediate levels following *L. donovani* infection from day 7 p.i. onwards (Figure 3C and 3D). Together, these data suggested that day 7 p.i. was an appropriate time to stimulate iNKT cells with α -GalCer to coincide with the time when hepatic iNKT cells were naturally activated following *L. donovani* infection. However, when C57BL/6 mice were given α -GalCer at this time, there was a significant increase in hepatic parasite burdens on day 14 p.i. ($p < 0.05$), relative to controls (Figure 3E). Thus, α -GalCer-stimulation of iNKT cells at the time when activation occurred naturally during infection suppressed, rather than improved, control of parasite growth in the liver.

Treatment with α -GalCer during an Established *L. donovani* Infection Impairs Disease Resolution in the Liver

To test whether the activation of iNKT cells during an established *L. donovani* infection had any therapeutic potential,




Figure 2. Stimulation of iNKT Cells with α -GalCer at the Time of Infection Does Not Enhance the Early Control of *L. donovani* Growth in the Liver. C57BL/6 mice were injected i.p. with either vehicle control (open bars) or 2 μ g α -GalCer (closed bars) the day prior to *L. donovani* infection. Parasite burdens were determined at day 14 p.i. and data represent the mean \pm SEM of parasite burdens (LDU) from four mice per group (A). C57BL/6 mice were injected i.p. with either vehicle control (open bars) or 2 μ g α -GalCer (closed bars), or i.v. with 5×10^5 vehicle-pulsed BMDC (open bars) or α -GalCer-pulsed BMDC (closed bars) 2 h prior to *L. donovani* infection, as indicated (B). Parasite burdens were determined in the liver at day 14 p.i. and data represent the mean \pm SEM of parasite burdens (LDU) from four mice per group. doi:10.1371/journal.ppat.1000028.g002




Figure 3. The Number of iNKT Cells and the Expression of CD69 by iNKT Cells Are Increased in the Liver by Day 7 after Infection, Yet Stimulation with α -GalCer at This Time Enhances Hepatic Parasite Growth. (A–D) C57BL/6 mice were infected with *L. donovani*, and killed at various time points p.i. for analysis of hepatic iNKT cells by flow cytometry. (A) iNKT cells were enumerated by labelling with CD1d/ α -GalCer tetramers and anti- α TCR, depicted for a naive liver, and at day 14 p.i., as indicated. The numbers in the top right hand corner represent the mean \pm SEM of gated cells. (B) The absolute number of hepatic TCR β^+ tetramer⁺ iNKT cells at each time point. (C) The surface expression of NK1.1 and CD69 (early activation marker) is shown on gated hepatic NKT cells from a naive animal (grey line) and at day 14 p.i. (solid black line). (D) Total number of hepatic NKT cells expressing enhanced levels of CD69, relative to naive mice. All data are from three mice per time point, and data in (B) and (D) represent the mean frequency \pm SEM for each time point. (E) C57BL/6 mice were infected with *L. donovani* and treated with either vehicle control (open bars) or 2 μ g α -GalCer (closed bars) on day 7 p.i. Parasite burdens were determined in the liver at day 14 p.i. and data represent the mean \pm SEM of parasite burdens (LDU) from four mice per group. Statistical differences of $p < 0.01$ (**) for vehicle versus α -GalCer treatment are indicated.

doi:10.1371/journal.ppat.1000028.g003

α -GalCer was administered on day 14 p.i., which is around the time of peak hepatic parasite burden in C57BL/6 mice (Figure 1A). Mice were sacrificed 1 wk later, and again, there were significant increases ($p < 0.05$) in hepatic parasite burdens in mice that received α -GalCer, compared to controls (Figure 4A). When α -GalCer-loaded BMDC were used instead of soluble α -GalCer,

there were no significant changes in hepatic parasite burdens, compared with controls that received vehicle-loaded BMDC (Figure 4A). Serum TNF levels were reduced in α -GalCer-treated mice on day 21 p.i. compared to controls ($p < 0.05$), but this did not occur in mice that received α -GalCer-loaded BMDC compared to controls that received vehicle-loaded BMDC (Figure 4B). A similar




Figure 4. Stimulation of iNKT Cells with α -GalCer during an Established Infection Exacerbates *L. donovani* Growth in the Liver. C57BL/6 mice were infected with *L. donovani* and treated with either vehicle control (open bars) or 2 μ g α -GalCer (closed bars) i.p., or injected i.v. with 5×10^5 vehicle-pulsed BMDC (open bars) or α -GalCer-pulsed BMDC (closed bars) on day 14 p.i.. Parasite burdens were determined in the liver at day 14 p.i. in untreated mice (grey bars; base-line parasite burden) and 1 wk later in treated groups, as indicated (A). Data represent the mean \pm SEM of parasite burdens (LDU) from four mice per group. Serum TNF (B) and IFN γ (C) levels at indicated time points are also shown. Statistical differences of $p < 0.05$ (*) for vehicle versus α -GalCer treatment are indicated. doi:10.1371/journal.ppat.1000028.g004

trend was observed for reductions in serum IFN γ levels in mice that received either soluble α -GalCer or α -GalCer-loaded BMDC compared to controls, but these differences were not statistically significant (Figure 4C). It should also be noted that vehicle-pulsed BMDC caused small reductions in serum TNF and IFN γ levels, relative to mice receiving vehicle alone (Figure 4B and 4C), suggesting that a DC-derived mediator might contribute to anti-protective effects seen following α -GalCer treatment. Thus, stimulation of iNKT cells with α -GalCer during an established infection, either in soluble form or loaded onto DC, did not improve disease control, and furthermore, soluble α -GalCer enhanced parasite growth in mice with an established infection.

To determine whether α -GalCer-treated mice recovered at later time points, α -GalCer was administered on day 14 p.i., and mice were sacrificed on both day 21 p.i. and day 70 p.i. (Figure S2). At day 70 p.i., hepatic parasite burdens had diminished significantly in mice receiving either vehicle or α -GalCer, and there was no significant difference between the groups. Therefore, the anti-protective effects of a single dose of α -GalCer during an established infection were transient, and anti-parasitic mechanisms were ultimately re-established in treated mice.

IL-10 Does Not Impede the Function of iNKT Cells During *L. donovani* Infection

L. donovani infection results in the generation of immune regulatory mechanisms that prevent efficient disease resolution. In particular, IL-10 suppresses pro-inflammatory responses and parasite clearance during experimental VL [27,43–45]. Previous work has reported an expansion of IL-10-producing CD4⁺ T cells

in mice following *L. donovani* infection [46], and more recently, high levels of IL-10 mRNA accumulation were found in T cells isolated from splenic aspirates taken from Indian VL patients [47]. Furthermore, activated NKT cells are capable of IL-10 production [48]. Therefore, mice were treated with either anti-IL-10 receptor (IL-10R) mAb or control rat IgG on day 12 p.i., 2 d prior to the administration of α -GalCer. Liver parasite burdens were measured 1 wk following α -GalCer treatment. Again, the administration of α -GalCer to control mice on day 14 p.i. resulted in increased hepatic ($p < 0.05$) parasite burdens compared to vehicle-treated controls (Figure 5A). IL-10R blockade was extremely potent in reducing parasite burdens (Figure 5A), as previously reported [43]. Stimulation of iNKT cells with α -GalCer in the absence of IL-10 signalling did not have an additive effect on the control of parasite growth, compared to vehicle-treated controls (Figure 5A), but instead resulted in a small, but significant ($p < 0.01$), increase in hepatic parasite burdens (Figure 5A). IL-10 blockade led to significantly elevated levels of serum TNF in both vehicle and α -GalCer treated mice ($p < 0.01$; Figure 5B), and a significant increase in serum IFN γ in mice that received α -GalCer ($p < 0.05$; Figure 5C). However, α -GalCer treatment resulted in increased hepatic parasite burdens despite these increased levels of inflammatory cytokines. α -GalCer treatment reduced the levels of serum TNF but not IFN γ relative to vehicle treatment in mice that received IL-10R blockade. Thus, stimulation of iNKT cells with α -GalCer in the absence of IL-10 signalling during an established *L. donovani* infection did not reduce parasite burdens in the liver, thereby indicating that this mechanism of immune regulation was not responsible for the failure of α -GalCer to improve control of parasite growth.




Figure 5. IL-10R Blockade Prior to α -GalCer Stimulation Does Not Improve Control of Parasite Growth During an Established Infection. (A) C57BL/6 mice were infected with *L. donovani* and treated with anti-IL-10R mAb or control rat IgG on day 12 p.i., as indicated. Two days later, mice were treated with vehicle control (open bars) or 2 μ g α -GalCer (closed bars) i.p., and parasite burdens were determined in the liver 1 wk later. Parasite burdens were also determined in untreated mice at day 14 p.i. (grey bars; base-line parasite burden), and at day 21 p.i. in mice that received no antibody treatment, as indicated. Data represent the mean \pm SEM of parasite burdens (LDU) from four mice per group. (B) Serum TNF and (C) IFN γ levels at indicated time points are also shown. Statistical differences of $p < 0.05$ (*) for vehicle versus α -GalCer treatment are indicated. doi:10.1371/journal.ppat.1000028.g005

L. donovani Infection Alters iNKT Cell Cytokine and Hepatic Regulatory Cytokine Production in Response to α -GalCer Stimulation

We next examined whether iNKT cells responded differently to α -GalCer stimulation during an established *L. donovani* infection. Naïve and infected mice (day 14 p.i.) were administered α -GalCer and sacrificed 2 h later. The down-regulation of TCR on iNKT cells following α -GalCer stimulation was noted in naïve mice (Figure 6A), as previously described [40], as was down-regulation of TCR on iNKT cells during experimental VL (Figure 6A). *L. donovani* infection did not alter the ratio of CD4⁺ to CD4⁻ iNKT cells in the liver, and therefore did not appear to selectively favour the expansion of a particular subset of iNKT cells (data not shown). IFN γ was readily detected in hepatic iNKT cells (Figure 6B), NK cells and conventional T cells (Figure S3) by day 14 p.i. Following α -GalCer stimulation, the frequency of hepatic iNKT cells producing IFN γ and TNF was significantly reduced in infected animals relative to naïve animals ($p < 0.05$), while the frequency of hepatic iNKT cells producing IL-4 was significantly

increased ($p < 0.05$; Figure 6B–6D). However, due to the increase in liver MNC number that had occurred by day 14 p.i. ($3.3 \times 10^6 \pm 1.8 \times 10^5$ versus $3.5 \times 10^7 \pm 7.9 \times 10^6$ in naïve versus day 14 p.i., respectively), there was an approximate 3-fold increase in the absolute number of IL-4-producing hepatic iNKT cells, but little difference in the absolute numbers of IFN γ -producing and TNF-producing iNKT cells following α -GalCer stimulation in infected mice relative to α -GalCer-treated naïve controls. These data suggest an iNKT cell bias towards Th2 cytokine production in the liver following α -GalCer stimulation during *L. donovani* infection. Importantly, there was also an increase in the number of hepatic NK cells producing IFN γ following α -GalCer stimulation in infected animals relative to α -GalCer-treated naïve animals ($p < 0.05$; $8.2 \times 10^4 \pm 5.3 \times 10^3$ versus $6.7 \times 10^5 \pm 1.7 \times 10^5$ in naïve versus day 14 p.i., respectively), indicating that *L. donovani* infection does not interfere with this important α -GalCer-mediated effector pathway.

To determine whether there was a general enhancement in Th2 and/or regulatory cytokine production in response to α -GalCer




Figure 6. Stimulation of iNKT Cells with α -GalCer during an Established *L. donovani* Infection Results in a Decreased Frequency of Hepatic iNKT Cells Producing IFN γ and TNF, and an Increased Frequency of Hepatic iNKT Cells Producing IL-4. C57BL/6 mice were infected with *L. donovani* and on day 14 p.i., infected mice and uninfected controls were treated with vehicle control or 2 μ g α -GalCer i.p. and killed 2 h later for FACS analysis of iNKT cell intracellular cytokine production. Hepatic lymphocytes were labelled with CD1d/ α -GalCer tetramers and anti- α TCR, and iNKT cells were electronically gated as shown (A). The mean \pm SEM of the percentage of gated cells for each group is shown in the top right-hand corner. From left to right data represent: naïve mice that received vehicle control, naïve mice that received α -GalCer, infected mice that received vehicle control on day 14 p.i., and infected mice that received α -GalCer on day 14 p.i.. Hepatic iNKT cells were examined for expression of NK1.1 and intracellular IFN γ (B), TNF (C), IL-4 (D), or rat IgG1 (E) (isotype control). Regions were set based on background levels of the isotype control as shown, and the numbers expressed in the bottom right hand corner are the mean \pm SEM in this region for each group (4 mice per group). doi:10.1371/journal.ppat.1000028.g006

treatment in *L. donovani*-infected mice, we next measured hepatic IL-10, TGF β , IL-13 and IL-4 mRNA accumulation at day 21 p.i. in mice that had received either vehicle or α -GalCer at day 14 p.i. (Figure 7A–7D). We found no significant differences in hepatic IL-10 (Figure 7A) or TGF β (Figure 7B) mRNA accumulation between vehicle- and α -GalCer-treated mice. Similar to the increased iNKT cell IL-4 production observed 2 h following α -GalCer administration shown above, we found a significant ($p < 0.05$) increase in hepatic IL-4 mRNA accumulation (Figure 7C) 1 wk following α -GalCer administration, relative to vehicle-treated controls. Interestingly, we also found a similar increase ($p < 0.01$) in hepatic IL-13 mRNA accumulation (Figure 7D) at the same time. Therefore, α -GalCer treatment of *L. donovani*-infected mice did not cause a general increase in Th2 and/or regulatory cytokine production, but instead, resulted in a selective increase in IL-4 and IL-13 production.

We next tested whether increased IL-4 production following α -GalCer stimulation during an established *L. donovani* infection was responsible for impaired disease resolution. C57BL/6 and B6.IL-4^{-/-} mice were infected and administered α -GalCer or vehicle control 14 days later. At this time, B6.IL-4^{-/-} mice had similar hepatic parasite burdens to C57BL/6 mice (Figure 8A). α -GalCer treatment resulted in increased hepatic parasite burdens ($p < 0.05$) in both C57BL/6 and B6.IL-4^{-/-} mice, compared to vehicle-treated controls (Figure 8A). To ensure that redundancies in IL-4

deficient mice were not influencing the outcome of these experiments, we also neutralised IL-4 activity in C57BL/6 mice using a mAb at a dose previously shown to improve control of *L. donovani* growth in a Th2 environment [49] (Figure 8B). Again, we found that *L. donovani*-infected mice treated with α -GalCer had significantly higher ($p < 0.01$) hepatic parasite burdens, compared with vehicle controls, regardless of whether they received anti-IL-4 mAb or control rat IgG. Together, these data suggest that increased IL-4 production in response to α -GalCer treatment in *L. donovani*-infected mice, either by iNKT cells or another cell type following iNKT cell activation, could not alone explain the impaired clearance of parasites.

Treatment with α -GalCer during an Established *L. donovani* Infection Alters Liver Cell Composition and Cytokine Production

To test whether the activation of iNKT cells during an established *L. donovani* infection had any impact on the cellular composition of the liver that could explain the impaired parasite clearance, α -GalCer was administered on day 14 p.i., and the numbers of different hepatic mononuclear cell populations were assessed 7 d later. We found no significant differences in hepatic B cell, NKT cell, neutrophil, monocyte/macrophage or DC numbers in mice that had received α -GalCer, relative to vehicle-treated control animals (Figure S4). However, we did observe a




Figure 7. Th2 and Regulatory Cytokine mRNA Accumulation in the Liver Following α -GalCer Treatment during an Established *L. donovani* Infection. C57BL/6 mice were infected with *L. donovani* and treated with vehicle control (open bars) or 2 μ g α -GalCer (closed bars) i.p. on day 14 p.i.. IL-10 (A), TGF β (B), IL-4 (C), and IL-13 (D) mRNA accumulation were determined in the liver 1 wk later, as indicated. Cytokine mRNA accumulation was also determined in naïve mice (hatched bars) and untreated mice at day 14 p.i. (grey bars). Data represent the mean \pm SEM of cytokine mRNA accumulation from four mice per group. Statistical differences of $p < 0.05$ (*) and $p < 0.01$ (**) for vehicle versus α -GalCer treatment are indicated.

doi:10.1371/journal.ppat.1000028.g007




Figure 8. IL-4 Production following α -GalCer Stimulation during an Established *L. donovani* Infection Has Only a Minor Role in the Impaired Control of Parasite Growth. (A) C57BL/6 mice and B6.IL-4^{-/-} mice were infected with *L. donovani* and treated with vehicle control (open bars) or 2 μ g α -GalCer (closed bars) i.p. on day 14 p.i., and parasite burdens were determined in the liver 1 wk later. Parasite burdens were also determined in untreated mice at day 14 p.i. (grey bars; baseline parasite burden). Data represent the mean \pm SEM of parasite burdens (LDU) from four mice per group. (B) C57BL/6 mice were also infected with *L. donovani* and treated with anti-IL-4 mAb or control rat IgG on days 14, 16 and 18 p.i., as indicated. Following mAb injection on day 14 p.i. mice were treated with vehicle control (open bars) or 2 μ g α -GalCer (closed bars) i.p., and parasite burdens were determined in the liver 1 wk later. Parasite burdens were also determined in untreated mice at day 14 p.i. (grey bars; baseline parasite burden). Data represent the mean \pm SEM of parasite burdens (LDU) from four mice per group. Statistical differences of $p < 0.01$ (**) for vehicle versus α -GalCer treatment are indicated. doi:10.1371/journal.ppat.1000028.g008

small decrease in the number of hepatic CD4⁺ T cells (Figure 9A) and a larger, significant ($p < 0.05$) decrease in the number of hepatic CD8⁺ T cells 7 d after α -GalCer-treatment (Figure 9B)

relative to controls. We also found a small, but significant ($p < 0.05$) increase in hepatic NK cell numbers at the same time (Figure 9C).

We next examined whether the activation status of various cell populations had been altered by α -GalCer administration at day 14 p.i.. We first focused on both hepatic DC and monocytes/macrophages and found no alterations in the expression of MHC-II, CD40, CD80 or CD86 on either population at 1, 2, 3, or 7 d following α -GalCer treatment, compared with vehicle-treated controls (data not shown). In addition, TNF production by monocytes/macrophages was not different between treatment groups at these times after infection, and no TNF was detected in DC (data not shown). Therefore, the activation status of the major hepatic antigen-presenting cell populations did not appear to be modulated by α -GalCer. To investigate whether lymphocyte activation had been altered, we examined IFN γ production following α -GalCer or vehicle treatment and found no difference in the number of hepatic IFN γ ⁺ NK cells (Figure 9C) or IFN γ ⁺ NKT cells (data not shown) by 7 d post-treatment. However, we did observe a small decrease in the number of IFN γ ⁺ CD4⁺ T cells (Figure 9A) and a larger, significant ($p < 0.05$) decrease in the number of IFN γ ⁺ CD8⁺ T cells 7 d after α -GalCer-treatment (Figure 9B). Given the importance of IFN γ [50] and CD8⁺ T cells [51] for the effective resolution of hepatic *L. donovani* infection, this loss of IFN γ ⁺ CD8⁺ T cells after α -GalCer treatment is a likely explanation for the enhanced parasite growth observed in our studies.

Discussion

We report that CD1d-restricted NKT cells have only a minor physiological role in experimental VL in C57BL/6 mice. More importantly, our results indicate that intentional stimulation of iNKT cells with the glycolipid α -GalCer during *L. donovani* infection can exacerbate disease. This suggests that the therapeutic use of NKT cell activators, such as α -GalCer, may not always provide beneficial effects, particularly in situations where sub-clinical or chronic infection may be present.

Previous studies in CD1d-deficient BALB/c mice suggested that NKT cells were required for efficient control of hepatic *L. donovani* infection [31]. The most likely explanation for this discrepancy with our data is a different requirement for NKT cells in C57BL/6 and BALB/c mice, possibly reflecting different kinetics of parasite control between these mouse strains [28]. Alternatively, NKT cell responses may be influenced by host genetic background, as previously reported in experimental cerebral malaria [52]. Another study in B6.J α 18^{-/-} mice suggested that NKT cells were required for the efficient induction of CXCL10 production during the early stages of *L. donovani* infection in the liver [30]. However, given the limited differences observed in hepatic parasite burdens between B6.J α 18^{-/-} mice and control mice at early time points (Figure 1A), our data suggest that NKT-independent chemokine signals can compensate for this lack of early CXCL10 production.

Although only a limited physiological role was found for CD1d-restricted NKT cells during experimental VL, the activation of these cells has been shown to improve disease outcome in many infections, often via the increased production of IFN γ [11–15,53]. This can occur even when there appears to be little requirement for NKT cells during natural infection, and is often due to downstream activation of NK cells. For example, although iNKT cells have only a minor role in the clearance of murine cytomegalovirus, the activation of iNKT cells by α -GalCer during infection resulted in reduced viral replication in visceral organs [53]. However, despite iNKT cell IFN γ and TNF production, and




Figure 9. Changes in Liver Cell Composition and IFN γ ⁺ Cell Numbers following α -GalCer Treatment during an Established *L. donovani* Infection. C57BL/6 mice were infected with *L. donovani* and treated with vehicle control (open bars) or 2 μ g of α -GalCer (closed bars) i.p. on day 14 p.i. The number of hepatic CD4⁺ T cells (A), CD8⁺ T cells (B), and NK cells (C) were determined 7 d later, as was the number of these cells producing IFN γ , as indicated. Cell composition was also determined in naïve mice (hatched bars) and untreated mice at day 14 p.i. (grey bars). Data represent the mean \pm SEM of cell numbers from four mice per group. Statistical differences of $p < 0.05$ (*) and $p < 0.01$ (**) for vehicle versus α -GalCer treatment are indicated.

doi:10.1371/journal.ppat.1000028.g009

enhanced NK cell IFN γ production in the liver following α -GalCer administration during infection, the control of *L. donovani* was not improved. In fact, α -GalCer treatment during an established infection suppresses parasite clearance from the liver. In contrast, the administration of another strong inducer of a Th1

response, recombinant IL-12, does enhance disease resolution in experimental VL [54], indicating that α -GalCer has a different qualitative effect despite its ability to induce a strong Th1 response in *L. donovani*-infected mice. This may be related to the ability of α -GalCer to stimulate both Th1 and Th2 cytokine production.

The reason for the different effects of α -GalCer stimulation on parasite growth before and after infection is unclear, but it seems likely that changes in local tissue environments during infection may alter the outcome of iNKT cell activation with an impact on anti-leishmanial immunity. This explanation is supported by the finding that the timing of α -GalCer administration was also critical to the effect on murine experimental conjunctivitis [55]. The effect of other regimes of α -GalCer therapy, other sites of administration and/or other analogues such as α -C-GalCer [56,57] have not yet been tested in experimental VL, and it remains possible that these other strategies might represent a therapeutic regime by which iNKT cell activation may improve disease resolution in this model. However, our data clearly show that α -GalCer therapy can exacerbate disease in *L. donovani*-infected mice.

A major functional change following α -GalCer-stimulation during *L. donovani* infection in the current study was rapidly increased numbers of IL-4 producing hepatic iNKT cells and a sustained increase in hepatic IL-4 mRNA accumulation 1 wk later. However, when *L. donovani*-infected B6.IL-4^{-/-} mice were treated with α -GalCer, hepatic parasite burdens were still increased compared to vehicle-treated B6.IL-4^{-/-} mice. A similar failure to reverse the immune-dampening effect of α -GalCer was also found when mice were treated with anti-IL-4 mAb before and after α -GalCer treatment. Therefore, the increased α -GalCer induced IL-4 production by iNKT cells or another hepatic cell population during *L. donovani* infection could not alone explain the enhanced hepatic parasite growth. It is possible that the induction of other Th2 and/or regulatory cytokines by α -GalCer might contribute to the increased parasite burden observed. Although no α -GalCer-related increase in hepatic IL-10 or TGF β mRNA accumulation was found, we did observe a significant increase in IL-13 mRNA accumulation. Studies in IL-13-deficient mice indicate that IL-13 itself does not influence resistance to *L. donovani* infection [58]. However, the enhanced production of both IL-4 and IL-13 may work together to exacerbate parasite growth. This possibility is currently being examined.

Other major changes observed in α -GalCer-treated infected mice were alterations to liver cell composition. Total hepatic NK cell number increased in mice that received α -GalCer, yet the number of IFN γ ⁺ NK cells remained similar to vehicle-treated mice. NK cells can produce regulatory cytokines and modulate antigen presenting cell function (reviewed in [59]) and in this way NK cells may have contributed to the enhanced parasite growth seen following α -GalCer treatment. Studies to further elucidate the impact of NK cells on the immune dampening effect of α -GalCer in *L. donovani*-infected mice are underway. The other significant change in liver cell composition that occurred in α -GalCer-treated mice was a decrease in the number of CD8⁺ T cells, and importantly, a decrease in the number of IFN γ ⁺ CD8⁺ T cells. Previous studies by Murray and colleagues have demonstrated a critical role for both CD8⁺ T cells [51] and IFN γ [50] for the effective resolution of hepatic *L. donovani* infection. Thus, it is likely that such a change to CD8⁺ T cell effector numbers will have a major impact on the ability of an animal to control parasite growth. The reason for this reduction in CD8⁺ T cell number is currently unknown, but could involve enhanced activation-induced cell death (AICD) and/or failure of expansion in this cell population. These possibilities are being investigated.

In summary, our data indicate that although NKT cells have a relatively minor physiological role in experimental VL, intentional stimulation of iNKT cells with α -GalCer can exacerbate infection. This finding is significant because it highlights the fact that therapies aimed at modulating NKT cell function are not always beneficial to the host, and it is important to take into account the

possibility of adverse consequences, such as may occur in the presence of pre-existing infections.

Materials and Methods

Mice

Female C57BL/6 and BALB/c mice were purchased from the Australian Resource Centre (Canning Vale, Western Australia), and maintained under conventional conditions. B6.RAG1/J^{-/-} mice were bred at the Queensland Institute of Medical Research. B6.J α 18^{-/-} mice [60] were bred at the Peter MacCallum Cancer Centre and the Queensland Institute of Medical Research. B6.CD1d^{-/-} mice [61] and B6.IL-4^{-/-} mice (B6.IL4^{tm1Nnt}, The Jackson Laboratory, Bar Harbor, ME) [62] were bred at the Peter MacCallum Cancer Centre. These mouse strains were originally generated on either the C57BL/6 background or the 129sv background and backcrossed 10 or more generations to C57BL/6 mice before use. Mice used were sex- and age-matched (6–10 wk), and all animal procedures were approved by the Queensland Institute of Medical Research Animal Ethics Committee.

Parasites and infection of mice

L. donovani (LV9) was maintained by passage in BALB/c or B6.RAG1/J^{-/-} mice, and amastigotes were isolated from the spleens of chronically infected mice. Mice were infected by injecting 2 \times 10⁷ amastigotes i.v. via the lateral tail vein, killed at the times indicated in the text by CO₂ asphyxiation and bled via cardiac puncture. Spleens and perfused livers were removed, and parasite burdens were determined from Diff-Quick-stained impression smears (Lab Aids, Narrabeen, Australia) and expressed in Leishman-Donovan units (LDU) (the number of amastigotes per 1,000 host nuclei multiplied by the organ weight in grams) [63]. Hepatic mononuclear cells (MNC) and splenocytes were isolated immediately following death as previously described [64,65].

Antibodies

Allophycocyanin (APC)-conjugated anti-TCR β chain, phycoerythrin (PE)-conjugated anti-CD8 α , PE-Cy5-conjugated anti-CD4, PE-conjugated anti-Ly6G, fluorescein isothiocyanate (FITC)-conjugated anti-Ly6C, APC-conjugated anti-B220, FITC-conjugated anti-CD19, APC-conjugated anti-CD11c, FITC-conjugated anti-I-A/I-E (MHC-II), PE-Cy5-conjugated anti-CD11b, PE-conjugated anti-CD69, PE-conjugated anti-IFN γ , PE-conjugated anti-TNF, PE-conjugated anti-IL-4, PE-conjugated rat IgG1, PE-conjugated rat IgG2a, and biotinylated anti-NK1.1 mAbs were purchased from Biolegend (San Diego, CA) or BD Biosciences (Franklin Lakes, NJ). PE-Cy5-conjugated α -GalCer-loaded or vehicle-loaded mouse CD1d tetramers were produced as previously described [66]. Biotinylated antibodies were detected using Alexa Fluor 488-conjugated streptavidin (Invitrogen Life Technologies, Eugene, OR). Anti-IL-10R (1B1.3a; rat IgG1) was purchased from BD Biosciences. Purified control rat IgG was purchased from Sigma-Aldrich (Castle Hill, Australia).

Flow cytometry

The labelling of hepatic MNC for cell surface antigens and intracellular cytokines for FACS analysis was performed as previously described [40,65]. Flow cytometric analysis was performed on a FACScalibur flow cytometer and analysed using Cell Quest Pro (BD Biosciences) and FlowJo Software (Tree Star, Ashland, OR). Cell populations in the liver were defined as follows: CD4⁺ T cells (CD4⁺ TCR β ⁺), CD8⁺ T cells (CD8 α ⁺ TCR β ⁺), B cells (B220⁺ CD19⁺), neutrophils (CD11b⁺ Ly6G⁺), macrophages/monocytes (CD11b⁺ Ly6C⁺), dendritic cells (DC;

CD11c^{hi} MHCII^{hi}), NK cells (NK1.1⁺ TCRβ⁻), NKT cells (CD1d α-GalCer tetramer⁺ TCRβ⁺ or NK1.1⁺ TCRβ⁺, as indicated in the text).

In vivo α-GalCer treatment

α-GalCer (KRN7000) was provided by the Kirin Pharmaceutical Research laboratories (Kirin Brewery, Gumna, Japan) and was prepared in saline/0.5% (w/v) polysorbate-20. No endotoxin contamination was detected in the supplied material (Tomoaki Kuwaki, Kirin Brewery, personal communication). Mice received a single i.p. injection of 2 μg α-GalCer or vehicle control (saline/0.5% (w/v) polysorbate-20) at the times indicated in the text.

Delivery of α-GalCer in bone marrow-derived dendritic cells (BMDC)

Bone marrow cells were isolated from the femurs of C57BL/6 mice and cultured in 20 ng/ml recombinant mouse GM-CSF and 20 ng/ml recombinant mouse IL-4 (R&D Systems, Minneapolis, MN), as previously described [37]. BMDC were supplemented with fresh media and 100 ng/ml α-GalCer or vehicle control on days 6–8 of culture. BMDC (5 × 10⁵) were injected i.v. at the times indicated in the text.

IL-10 and IL-4 neutralisation

Mice were injected i.p. with 1 mg of anti-IL-10R mAb (1B1.3a) or purified control rat IgG (Sigma-Aldrich) at day 12 p.i.. Mice were injected i.p. with 1 mg of anti-IL-4 mAb (11B11) or purified control rat IgG on days 14, 16 and 18 p.i..

Serum cytokine analysis

Serum samples were analysed for IFNγ and TNF using a cytometric bead array (mouse inflammation kit, BD Biosciences) on a FACScan cytometer equipped with Cell Quest Pro and CBA Software (BD Biosciences).

ELISPOT analysis

The frequency of liver MNC producing IFNγ or TNF was determined by direct *ex vivo* ELISPOT assay, as previously described [67] using mAbs R46A2 and TN3-19.12, respectively, as capture antibodies and polyclonal anti-murine IFNγ or TNF as detecting antibodies (BD Biosciences).

Real-time reverse transcriptase–polymerase chain reaction

Total RNA was extracted from liver tissue using TRIzol reagent (Invitrogen Life Technologies), and an RNeasy Mini Kit with on-column DNase digestion (Qiagen, Valencia, CA). RNA samples were reverse transcribed into cDNA using the High Capacity cDNA Reverse Transcription Kit (Applied Biosystems, Foster City, CA) according to the manufacturer's instructions. The number of IL-10, TGFβ, IL-4, IL-13 and HPRT (house-keeping gene) cDNA molecules in each sample were calculated using Taqman Gene Expression Assays (Applied Biosystems). All real-time RT-PCR were performed on a Corbett Research RG-3000 Rotor Gene (Corbett Life Sciences, Sydney, Australia). Standard curves were generated with known amounts of cDNA for each gene, and the number of cytokine molecules per 1,000 HPRT molecules in each sample was calculated.

Statistical analysis

Statistical significance of differences in parasite burdens over a time course of infection were determined using a one way analysis of

variance (ANOVA) followed by a Tukey's Post-Test. Statistical significance of other parameters were determined using a Mann-Whitney U test or Student's t-test. All analysis was conducted using GraphPad Prism version 4.03 for Windows (GraphPad Software, San Diego, CA) and *p* < 0.05 was considered statistically significant.

Supporting Information

Figure S1 The Number of IFNγ- or TNF-Producing Cells Is Similar in *L. donovani*-Infected C57BL/6 Mice and Mice Lacking NKT Cells. Female C57BL/6 (closed squares), B6.Jα18^{-/-} (open circles) and B6.CD1d^{-/-} (open triangles) mice were infected with *L. donovani*, and the total numbers of IFNγ- (A) or TNF-producing cells (B) in the liver were measured by ELISPOT from day 7 to day 56 p.i.. Data represent the mean ± SEM of cytokine-producing cells from four mice per group for each time point.

Found at: doi:10.1371/journal.ppat.1000028.s001 (1.24 MB TIF)

Figure S2 α-GalCer-Treated Mice Recover at Later Time Points. C57BL/6 mice were infected with *L. donovani* and treated with either vehicle control (open bars) or 2 μg α-GalCer (closed bars) i.p. on day 14 p.i.. Parasite burdens were determined in the liver at day 14 p.i. in untreated mice (grey bars; baseline parasite burden), 1 wk later in treated groups or 56 d later in treated groups, as indicated. Statistical differences of *p* < 0.001 (***) for vehicle versus α-GalCer treatment are indicated (*n* = 4 mice per group).

Found at: doi:10.1371/journal.ppat.1000028.s002 (0.90 MB TIF)

Figure S3 IFNγ Is Readily Detected in Hepatic NK Cells and Conventional T Cells 14 Days After *L. donovani* Infection. Naïve C57BL/6 mice or mice infected with *L. donovani* were treated with vehicle control on day 14 p.i. and killed 2 h later for FACS analysis of intracellular IFNγ production, as indicated. Hepatic lymphocytes were labelled with CD1d/α-GalCer tetramers, anti-αβTCR, anti-NK1.1, and anti-IFNγ. T cells (A) and NK cells (B) were electronically gated as shown, and examined for expression of IFNγ or isotype control antibody. The percentage of gated cells is shown in the top left-hand corner. One representative animal of four examined is shown.

Found at: doi:10.1371/journal.ppat.1000028.s003 (5.38 MB TIF)

Figure S4 Liver Cell Composition Following α-GalCer Treatment. C57BL/6 mice were infected with *L. donovani* and treated with either vehicle control (open bars) or 2 μg α-GalCer (closed bars) i.p. on day 14 p.i.. Liver cell numbers were determined by FACS in naïve mice (hatched bars), at day 14 p.i. in untreated mice (grey bars), and 1 wk later in treated groups, as indicated (*n* = 4 mice per group).

Found at: doi:10.1371/journal.ppat.1000028.s004 (4.14 MB TIF)

Acknowledgments

We thank Grace Chojnowski and Paula Hall for expert assistance with flow cytometry and cell sorting, and Kirin Brewery Pharmaceutical Research laboratories (Gumna, Japan) for providing α-GalCer (KRN7000).

Author Contributions

Conceived and designed the experiments: AS GH MS DG CE. Performed the experiments: AS YZ FA LR AH CE. Analyzed the data: AS MS DG CE. Contributed reagents/materials/analysis tools: GH. Wrote the paper: AS CE. Produced the PE-Cy5-conjugated α-GalCer-loaded CD1d tetramers: DP.

References

- Godfrey DI, MacDonald HR, Kronenberg M, Smyth MJ, Van Kaer L (2004) NKT cells: What's in a name? *Nat Rev Immunol* 4: 231–237.
- Behar SM, Podrebarac TA, Roy CJ, Wang CR, Brenner MB (1999) Diverse TCRs recognize murine CD1. *J Immunol* 162: 161–167.
- Cardell S, Tangri S, Chan S, Kronenberg M, Benoist C, et al. (1995) CD1-restricted CD4+ T cells in major histocompatibility complex class II-deficient mice. *J Exp Med* 182: 993–1004.
- Carnaud C, Lee D, Donnars O, Park SH, Beavis A, et al. (1999) Cutting edge: Cross-talk between cells of the innate immune system: NKT cells rapidly activate NK cells. *J Immunol* 163: 4647–4650.
- Hayakawa Y, Takeda K, Yagita H, Kakuta S, Iwakura Y, et al. (2001) Critical contribution of IFN-gamma and NK cells, but not perforin-mediated cytotoxicity, to anti-metastatic effect of alpha-galactosylceramide. *Eur J Immunol* 31: 1720–1727.
- Nishimura T, Kitamura H, Iwakabe K, Yahata T, Ohta A, et al. (2000) The interface between innate and acquired immunity: Glycolipid antigen presentation by CD1d-expressing dendritic cells to NKT cells induces the differentiation of antigen-specific cytotoxic T lymphocytes. *Int Immunol* 12: 987–994.
- Smyth MJ, Crowe NY, Pellicci DG, Kyparissoudis K, Kelly JM, et al. (2002) Sequential production of interferon-gamma by NK1.1(+) T cells and natural killer cells is essential for the antimetastatic effect of alpha-galactosylceramide. *Blood* 99: 1259–1266.
- Godfrey DI, Kronenberg M (2004) Going both ways: Immune regulation via CD1d-dependent NKT cells. *J Clin Invest* 114: 1379–1388.
- Kawano T, Cui J, Koezuka Y, Toura I, Kaneko Y, et al. (1997) CD1d-restricted and TCR-mediated activation of valpha14 NKT cells by glycosylceramides. *Science* 278: 1626–1629.
- Hayakawa Y, Godfrey DI, Smyth MJ (2004) Alpha-galactosylceramide: Potential immunomodulatory activity and future application. *Curr Med Chem* 11: 241–252.
- Parekh VV, Wilson MT, Van Kaer L (2005) iNKT-cell responses to glycolipids. *Crit Rev Immunol* 25: 183–213.
- Taniguchi M, Harada M, Kojo S, Nakayama T, Wakao H (2003) The regulatory role of Valpha14 NKT cells in innate and acquired immune response. *Annu Rev Immunol* 21: 483–513.
- Gonzalez-Aseguinolaza G, de Oliveira C, Tomaska M, Hong S, Bruna-Romero O, et al. (2000) alpha-galactosylceramide-activated Valpha 14 natural killer T cells mediate protection against murine malaria. *Proc Natl Acad Sci U S A* 97: 8461–8466.
- Duthie MS, Kahn SJ (2002) Treatment with alpha-galactosylceramide before *Trypanosoma cruzi* infection provides protection or induces failure to thrive. *J Immunol* 168: 5778–5785.
- Ronet C, Darche S, Leite de Moraes M, Miyake S, Yamamura T, et al. (2005) NKT cells are critical for the initiation of an inflammatory bowel response against *Toxoplasma gondii*. *J Immunol* 175: 899–908.
- Smyth MJ, Godfrey DI (2000) NKT cells and tumor immunity—a double-edged sword. *Nat Immunol* 1: 459–460.
- Wilson SB, Delovitch TL (2003) Janus-like role of regulatory iNKT cells in autoimmune disease and tumour immunity. *Nat Rev Immunol* 3: 211–222.
- Crowe NY, Smyth MJ, Godfrey DI (2002) A critical role for natural killer T cells in immunosurveillance of methylcholanthrene-induced sarcomas. *J Exp Med* 196: 119–127.
- Brigl M, Bry L, Kent SC, Gumperz JE, Brenner MB (2003) Mechanism of CD1d-restricted natural killer T cell activation during microbial infection. *Nat Immunol* 4: 1230–1237.
- Kinjo Y, Wu D, Kim G, Xing GW, Poles MA, et al. (2005) Recognition of bacterial glycosphingolipids by natural killer T cells. *Nature* 434: 520–525.
- Mattner J, Debord KL, Ismail N, Goff RD, Cantu C 3rd, et al. (2005) Exogenous and endogenous glycolipid antigens activate NKT cells during microbial infections. *Nature* 434: 525–529.
- Kinjo Y, Tupin E, Wu D, Fujio M, Garcia-Navarro R, et al. (2006) Natural killer T cells recognize diacylglycerol antigens from pathogenic bacteria. *Nat Immunol* 7: 978–986.
- Bendelac A, Savage PB, Teyton L (2007) The biology of NKT cells. *Annu Rev Immunol* 25: 297–336.
- Gardener PJ (1977) Taxonomy of the genus *Leishmania*: A review of nomenclature and classification. *Trop Dis Bull* 74: 1069–1088.
- Grimaldi G Jr, Tesh RB, McMahon-Pratt D (1989) A review of the geographic distribution and epidemiology of leishmaniasis in the New World. *Am J Trop Med Hyg* 41: 687–725.
- Murray HW (2001) Tissue granuloma structure-function in experimental visceral leishmaniasis. *Int J Exp Pathol* 82: 249–267.
- Ato M, Stager S, Engwerda CR, Kaye PM (2002) Defective CCR7 expression on dendritic cells contributes to the development of visceral leishmaniasis. *Nat Immunol* 3: 1185–1191.
- Engwerda CR, Ato M, Cotterell SE, Mynott TL, Tschannerl A, et al. (2002) A role for tumor necrosis factor-alpha in remodeling the splenic marginal zone during *Leishmania donovani* infection. *Am J Pathol* 161: 429–437.
- Engwerda CR, Kaye PM (2000) Organ-specific immune responses associated with infectious disease. *Immunol Today* 21: 73–78.
- Svensson M, Zubairi S, Maroof A, Kazi F, Taniguchi M, et al. (2005) Invariant NKT cells are essential for the regulation of hepatic CXCL10 gene expression during *Leishmania donovani* infection. *Infect Immun* 73: 7541–7547.
- Amprey JL, Im JS, Turco SJ, Murray HW, Illarionov PA, et al. (2004) A subset of liver NK T cells is activated during *Leishmania donovani* infection by CD1d-bound lipophosphoglycan. *J Exp Med* 200: 895–904.
- Campos-Martin Y, Colmenares M, Gozalbo-Lopez B, Lopez-Nunez M, Savage PB, et al. (2006) Immature human dendritic cells infected with *Leishmania infantum* are resistant to NK-mediated cytotoxicity but are efficiently recognized by NKT cells. *J Immunol* 176: 6172–6179.
- Ishikawa H, Hisaeda H, Taniguchi M, Nakayama T, Sakai T, et al. (2000) CD4(+) v(alpha)14 NKT cells play a crucial role in an early stage of protective immunity against infection with *Leishmania major*. *Int Immunol* 12: 1267–1274.
- Mattner J, Donhauser N, Werner-Felmayer G, Bogdan C (2006) NKT cells mediate organ-specific resistance against *Leishmania major* infection. *Microbes Infect* 8: 354–362.
- Yue SC, Shaulov A, Wang R, Balk SP, Exley MA (2005) CD1d ligation on human monocytes directly signals rapid NF-kappaB activation and production of bioactive IL-12. *Proc Natl Acad Sci U S A* 102: 11811–11816.
- Parekh VV, Wilson MT, Olivares-Villagomez D, Singh AK, Wu L, et al. (2005) Glycolipid antigen induces long-term natural killer T cell anergy in mice. *J Clin Invest* 115: 2572–2583.
- Fujii S, Shimizu K, Kronenberg M, Steinman RM (2002) Prolonged IFN-gamma-producing NKT response induced with alpha-galactosylceramide-loaded DCs. *Nat Immunol* 3: 867–874.
- Uldrich AP, Crowe NY, Kyparissoudis K, Pellicci DG, Zhan Y, et al. (2005) NKT cell stimulation with glycolipid antigen in vivo: Costimulation-dependent expansion, Bim-dependent contraction, and hyporesponsiveness to further antigenic challenge. *J Immunol* 175: 3092–3101.
- Reiner NE, Ng W, McMaster WR (1987) Parasite-accessory cell interactions in murine leishmaniasis. II. *Leishmania donovani* suppresses macrophage expression of class I and class II major histocompatibility complex gene products. *J Immunol* 138: 1926–1932.
- Crowe NY, Uldrich AP, Kyparissoudis K, Hammond KJ, Hayakawa Y, et al. (2003) Glycolipid antigen drives rapid expansion and sustained cytokine production by NK T cells. *J Immunol* 171: 4020–4027.
- Wilson MT, Johansson C, Olivares-Villagomez D, Singh AK, Stanic AK, et al. (2003) The response of natural killer T cells to glycolipid antigens is characterized by surface receptor down-modulation and expansion. *Proc Natl Acad Sci U S A* 100: 10913–10918.
- Harada M, Seino K, Wakao H, Sakata S, Ishizuka Y, et al. (2004) Down-regulation of the invariant Valpha14 antigen receptor in NKT cells upon activation. *Int Immunol* 16: 241–247.
- Murray HW, Lu CM, Mauze S, Moreira AL, et al. (2002) Interleukin-10 (IL-10) in experimental visceral leishmaniasis and IL-10 receptor blockade as immunotherapy. *Infect Immun* 70: 6284–6293.
- Murphy ML, Wille U, Villegas EN, Hunter CA, Farrell JP (2001) IL-10 mediates susceptibility to *Leishmania donovani* infection. *Eur J Immunol* 31: 2848–2856.
- Nylen S, Sacks D (2007) Interleukin-10 and the pathogenesis of human visceral leishmaniasis. *Trends Immunol* 28: 378–384.
- Stager S, Maroof A, Zubairi S, Sanos SL, Kopf M, et al. (2006) Distinct roles for IL-6 and IL-12p40 in mediating protection against *Leishmania donovani* and the expansion of IL-10+ CD4+ T cells. *Eur J Immunol* 36: 1764–1771.
- Nylen S, Maurya R, Eidsmo L, Manandhar KD, Sundar S, et al. (2007) Splenic accumulation of IL-10 mRNA in T cells distinct from CD4+CD25+ (Foxp3) regulatory T cells in human visceral leishmaniasis. *J Exp Med* 204: 805–817.
- Arase H, Arase N, Nakagawa K, Good RA, Onoe K (1993) NK1.1+ CD4+ CD8- thymocytes with specific lymphokine secretion. *Eur J Immunol* 23: 307–310.
- Murray HW, Hariprasad J, Coffman RL (1997) Behavior of visceral *Leishmania donovani* in an experimentally induced T helper cell 2 (Th2)-associated response model. *J Exp Med* 185: 867–874.
- Squires KE, Schreiber RD, McElrath MJ, Rubin BY, Anderson SL, et al. (1989) Experimental visceral leishmaniasis: Role of endogenous IFN-gamma in host defense and tissue granulomatous response. *J Immunol* 143: 4244–4249.
- Stern JJ, Oca MJ, Rubin BY, Anderson SL, Murray HW (1988) Role of L3T4+ and LyT-2+ cells in experimental visceral leishmaniasis. *J Immunol* 140: 3971–3977.
- Hansen DS, Siomos MA, Buckingham L, Scalzo AA, Schofield L (2003) Regulation of murine cerebral malaria pathogenesis by CD1d-restricted NKT cells and the natural killer complex. *Immunity* 18: 391–402.
- van Dommelen SL, Tabarias HA, Smyth MJ, Degli-Esposti MA (2003) Activation of natural killer (NK) T cells during murine cytomegalovirus infection enhances the antiviral response mediated by NK cells. *J Virol* 77: 1877–1884.
- Murray HW, Hariprasad J (1995) Interleukin 12 is effective treatment for an established systemic intracellular infection: Experimental visceral leishmaniasis. *J Exp Med* 181: 387–391.

55. Fukushima A, Sumi T, Fukuda K, Yamaguchi T, Kumagai N, et al. (2006) Modulation of murine experimental allergic conjunctivitis by treatment with alpha-galactosylceramide. *Immunol Lett* 107: 32–40.
56. Schmieg J, Yang G, Franck RW, Tsuji M (2003) Superior protection against malaria and melanoma metastases by a C-glycoside analogue of the natural killer T cell ligand alpha-Galactosylceramide. *J Exp Med* 198: 1631–1641.
57. Fujii S, Shimizu K, Hemmi H, Fukui M, Bonito AJ, et al. (2006) Glycolipid alpha-Galactosylceramide is a distinct inducer of dendritic cell function during innate and adaptive immune responses of mice. *Proc Natl Acad Sci U S A* 103: 11252–11257.
58. Murray HW, Tsai CW, Liu J, Ma X (2006) Visceral *Leishmania donovani* infection in interleukin-13^{-/-} mice. *Infect Immun* 74: 2487–2490.
59. Shi FD, Van Kaer L (2006) Reciprocal regulation between natural killer cells and autoreactive T cells. *Nat Rev Immunol* 6: 751–760.
60. Cui J, Shin T, Kawano T, Sato H, Kondo E, et al. (1997) Requirement for Valpha14 NKT cells in IL-12-mediated rejection of tumors. *Science* 278: 1623–1626.
61. Mendiratta SK, Martin WD, Hong S, Boesteanu A, Joyce S, et al. (1997) CD1d1 mutant mice are deficient in natural T cells that promptly produce IL-4. *Immunity* 6: 469–477.
62. Metwali A, Elliott D, Blum AM, Li J, Sandor M, et al. (1996) The granulomatous response in murine *Schistosomiasis mansoni* does not switch to Th1 in IL-4-deficient C57BL/6 mice. *J Immunol* 157: 4546–4553.
63. Smelt SC, Engwerda CR, McCrossen M, Kaye PM (1997) Destruction of follicular dendritic cells during chronic visceral leishmaniasis. *J Immunol* 158: 3813–3821.
64. Alexander CE, Kaye PM, Engwerda CR (2001) CD95 is required for the early control of parasite burden in the liver of *Leishmania donovani*-infected mice. *Eur J Immunol* 31: 1199–1210.
65. Engwerda CR, Ato M, Stager S, Alexander CE, Stanley AC, et al. (2004) Distinct roles for lymphotoxin-alpha and tumor necrosis factor in the control of *Leishmania donovani* infection. *Am J Pathol* 165: 2123–2133.
66. Matsuda JL, Naidenko OV, Gapin L, Nakayama T, Taniguchi M, et al. (2000) Tracking the response of natural killer T cells to a glycolipid antigen using CD1d tetramers. *J Exp Med* 192: 741–754.
67. Engwerda CR, Murphy ML, Cotterell SE, Smelt SC, Kaye PM (1998) Neutralization of IL-12 demonstrates the existence of discrete organ-specific phases in the control of *Leishmania donovani*. *Eur J Immunol* 28: 669–680.



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Stanley, AC; Zhou, Y; Amante, FH; Randall, LM; Haque, A; Pellicci, DG; Hill, GR; Smyth, MJ; Godfrey, DI; Engwerda, CR

Title:

Activation of invariant NKT cells exacerbates experimental visceral leishmaniasis

Date:

2008-02-01

Citation:

Stanley, A. C., Zhou, Y., Amante, F. H., Randall, L. M., Haque, A., Pellicci, D. G., Hill, G. R., Smyth, M. J., Godfrey, D. I. & Engwerda, C. R. (2008). Activation of invariant NKT cells exacerbates experimental visceral leishmaniasis. PLOS PATHOGENS, 4 (2), <https://doi.org/10.1371/journal.ppat.1000028>.

Persistent Link:

<http://hdl.handle.net/11343/242616>

File Description:

published version

License:

CC BY