



Article (refereed) - postprint

Harmens, Harry; Schnyder, Elvira; Thöni, Lotti; Cooper, David M.; Mills, Gina; Leblond, Sebastien; Mohr, Karsten; Poikolainen, Jarmo; Santamaria, Jesus; Skudnik, Mitja; Zechmeister, Harald G.; Lindroos, Antti-Jussi; Hanus-Ilmar, Andrea. 2014. **Relationship between site-specific nitrogen concentrations in mosses and measured wet bulk atmospheric nitrogen deposition across Europe.**

Copyright © 2014 Elsevier B.V.

This version available <http://nora.nerc.ac.uk/508021/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at

<http://nora.nerc.ac.uk/policies.html#access>

NOTICE: this is the author's version of a work that was accepted for publication in *Atmospheric Environment*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Atmospheric Environment*, 89. 35-42. [10.1016/j.atmosenv.2014.02.012](https://doi.org/10.1016/j.atmosenv.2014.02.012)

www.elsevier.com/

Contact CEH NORA team at
noraceh@ceh.ac.uk

1 **Relationship between site-specific nitrogen concentrations in mosses and measured wet**
2 **bulk atmospheric nitrogen deposition across Europe**

3

4 Harry Harmens^{a*}, Elvira Schnyder^b, Lotti Thöni^b, David M. Cooper^a, Gina Mills^a, Sébastien
5 Leblond^c, Karsten Mohr^d, Jarmo Poikolainen^e, Jesus Santamaria^f, Mitja Skudnik^g, Harald G.
6 Zechmeister^h, Antti-Jussi Lindroosⁱ, Andrea Hanus-Illnar^j

7

8 ^a Centre for Ecology and Hydrology, Environment Centre Wales, Deiniol Road, Bangor,
9 Gwynedd LL57 2UW, UK. hh@ceh.ac.uk; cooper@ceh.ac.uk; gmi@ceh.ac.uk

10 ^b FUB-Research Group for Environmental Monitoring, Alte Jonastrasse 83, CH-8640
11 Rapperswil, Switzerland. fub@fub-ag.ch

12 ^c Muséum National d'Histoire Naturelle, 57 rue Cuvier, Case 39, 75005 Paris, France.
13 sleblond@mnhn.fr

14 ^d Landwirtschaftskammer Niedersachsen, Mars-la-Tour Str. 1-13, 26121 Oldenburg,
15 Germany, Karsten.Mohr@LWK-Niedersachsen.de

16 ^e Finnish Forest Research Institute, P.O. Box 413, FI-90014 University of Oulu, Finland.
17 jarmo.poikolainen@metla.fi

18 ^f University of Navarra, Irunlarrea No 1, 31008 Pamplona, Spain. chusmi@unav.es

19 ^g Slovenian Forestry Institute, Vecna pot 2, 1000 Ljubljana, Slovenia.
20 mitja.skudnik@gozdis.si.

21 ^h University of Vienna, CVL, Rennweg 14, 1020 Vienna, Austria.
22 Harald.Zechmeister@univie.ac.at

23 ⁱ Finnish Forest Research Institute, P.O. Box 18, FI-01301 Vantaa, Finland.
24 antti.lindroos@metla.fi

25 ^j Umweltbundesamt, Spittelauer Lände 5, 1090 Vienna.
26 andrea.hanus-illnar@umweltbundesamt.at

27

28 Corresponding Author: Harry Harmens

29 E-mail address: hh@ceh.ac.uk

30 Tel.: +44-1248-374512

31 Fax: +44-1248-362133

32

33

34 **Abstract**

35

36 To assess the relationship between nitrogen concentrations in mosses and wet bulk nitrogen
37 deposition or concentrations in precipitation, moss tissue and deposition were sampled within
38 a distance of 1 km of each other in seven European countries. Relationships for various forms
39 of nitrogen appeared to be asymptotic, with data for different countries being positioned at
40 different locations along the asymptotic relationship and saturation occurring at a wet bulk
41 nitrogen deposition of ca. 20 kg N ha⁻¹ yr⁻¹. The asymptotic behaviour was more pronounced
42 for ammonium-N than nitrate-N, with high ammonium deposition at German sites being most
43 influential in providing evidence of the asymptotic behaviour. Within countries, relationships
44 were only significant for Finland and Switzerland and were more or less linear. The results
45 confirm previous relationships described for modelled total deposition. Nitrogen
46 concentration in mosses can be applied to identify areas at risk of high nitrogen deposition at
47 European scale.

48

49 **Capsule:** Nitrogen concentration in mosses shows saturation occurring at a measured wet
50 bulk nitrogen deposition of ca. 20 kg N ha⁻¹ yr⁻¹.

51

52 **Keywords:** biomonitoring; moss survey; bulk nitrogen deposition; ammonium; nitrate.

53

54 **1. Introduction**

55 For ectohydric moss species, the lack of a well-developed root system, vascular
56 system and protective cuticle means that they receive and take up water, nutrients and
57 contaminants mainly from atmospheric deposition (dry, wet and occult). Hence, such mosses
58 have shown to be suitable indicators of atmospheric deposition of, for example, nitrogen
59 (Harmens et al., 2011; Pitcairn et al., 2006; Salemaa et al., 2008; Solga et al., 2005;
60 Zechmeister et al., 2008), heavy metals (Harmens et al., 2010; Harmens et al., 2012; Schröder
61 et al., 2010b) and selected persistent organic pollutants (Foan et al., 2010, 2014; Harmens et
62 al., 2013a). The moss monitoring technique provides a complementary, time-integrated
63 measure of element deposition from the atmosphere to terrestrial systems. As it is easier and
64 cheaper than conventional deposition analysis, a much higher sampling density can be
65 achieved than with conventional deposition analysis. Hence, passive biomonitoring of
66 atmospheric nitrogen deposition using mosses would allow the determination of the variation
67 in atmospheric nitrogen deposition at a high spatial resolution, including in countries or areas
68 where nitrogen deposition monitoring networks are absent.

69 For nitrogen, sometimes the relationship between atmospheric deposition rates and the
70 concentration in mosses is weak (Stevens et al., 2011) or shown to be species-specific
71 (Arroniz-Crespo et al., 2008; Salemaa et al., 2008). One possible explanation for the weak
72 relationship between the deposition and accumulation of nitrogen is the regulation of tissue
73 loads in mosses because nitrogen is known to play an important role in the metabolism of
74 organisms (e.g., Koranda et al. 2007; Arróniz-Crespo et al. 2008), in contrast to for example
75 non-essential heavy metals such as cadmium and lead. Such regulation may distort the
76 patterns of nitrogen deposition identified by biomonitoring with terrestrial mosses. Schröder
77 et al. (2010a) have shown that atmospheric nitrogen deposition, as modelled by the European
78 Monitoring and Evaluation Programme (EMEP), is the primary factor determining total
79 nitrogen concentrations in mosses. Harmens et al. (2011) observed an asymptotic relationship

80 between the total nitrogen concentrations in mosses and EMEP modelled total nitrogen
81 deposition (averaged per 50 km x 50 km grid) across Europe, with saturation (i.e. no further
82 increasing nitrogen concentration in moss tissues with increasing nitrogen deposition)
83 occurring at a total deposition rate of ca. 15 kg N ha⁻¹ yr⁻¹. Whether such a relationship also
84 holds when both the nitrogen concentration in moss and atmospheric wet nitrogen deposition
85 are measured at nearby sites across Europe, is unknown.

86 Only a few studies have examined the relationship between the nitrogen concentration
87 in mosses and measured (as opposed to modelled) nitrogen deposition in the immediate
88 vicinity of the moss sampling sites (Skudnik et al., 2014; Solga et al., 2005; Thöni et al., 2008;
89 Zechmeister et al., 2008), in monitoring studies not conducted in the immediate vicinity of
90 local sources (e.g. Pitcairn et al., 2006). These studies were all conducted at the (sub-)national
91 scale and such data is not available at the European scale. The strength and shape of the
92 relationship observed in these (sub-)national studies varies between countries. For example, in
93 Switzerland, a strong, significant ($r^2 = 0.91$) linear relationship was found between the total
94 nitrogen concentration in mosses and measured site-specific wet bulk nitrogen deposition
95 (Harmens et al., 2011; Thöni et al., 2008). Less strong but still significant linear relationships
96 were also reported for North Rhine-Westphalia in Germany (Solga et al., 2005) and Austria
97 (Zechmeister et al., 2008). Skudnik et al. (2014) showed a weak but significant linear-
98 logarithmic relationship between the nitrogen concentration in mosses and atmospheric bulk
99 nitrogen deposition. To investigate the strength and shape of the relationship at the European
100 scale, data on nitrogen concentrations in mosses and measured wet bulk nitrogen deposition
101 were collected in seven European countries. Only monitoring sites where the distance
102 between the moss sampling site and the atmospheric deposition was less than 1 km were
103 considered.

104 As different moss species were used in the current study, we also investigated whether
105 moss species differ in their nitrogen concentration when sampled at the same sites, as this

106 might confound the relationship between atmospheric nitrogen deposition and the nitrogen
107 concentration in mosses (Arroniz-Crespo et al., 2008; Salemaa et al., 2008). Although there
108 are other factors potentially confounding the relationship between atmospheric nitrogen
109 deposition and its concentration in mosses, these were not investigated here but have been
110 discussed previously in more detail (Harmens et al., 2011, Schröder et al., 2010a) and some
111 are further discussed in the results and discussion section.

112 Despite the sometimes reported linear relationship between the nitrogen concentration
113 in mosses and measured wet bulk nitrogen deposition at the (sub-) national scale (Harmens et
114 al., 2011; Solga et al., 2005; Thöni et al., 2008), we hypothesise that the relationship will
115 show an asymptotic behaviour at the European scale (conform Harmens et al., 2011, using
116 modelled nitrogen deposition) when higher deposition rates are included. However, we expect
117 less scatter in the underlying data than for modelled deposition (Harmens et al., 2011). We
118 also tested whether the relationship is affected by nitrogen speciation in deposition and
119 whether the strength of the relationship differs for nitrogen deposition or nitrogen
120 concentration in precipitation.

121

122 **2. Materials and methods**

123 *Sites*

124 Mosses were collected between 1998 and 2012 at selected sites in seven European countries
125 (Figure 1): Austria (AT), Switzerland (CH), the German Bundesland Niedersachsen (DE-NI),
126 Spain (ES), Finland (FI), France (FR), and Slovenia (SI, although some of sites were in
127 Austria and Italy close to the Slovenian border). For this study, moss data were only included
128 from sites (97 in total) where the distance to the deposition monitoring site was less than 1 km
129 (the maximum distance recorded was 900 m). At some sites (s) sampling was repeated in
130 time, leading to 160 data points (p) for comparison (AT 26s, 26p; CH 18s, 33p; DE-
131 NI 6s, 33p; FR 24s, 36p; SI 11s, 11p; FI 11s, 19p; ES 1s, 2p). At some forested sites the

132 deposition was characterised as throughfall below the canopy of trees rather than bulk
133 deposition only. This was the case for the majority of data points in Germany, all sites in
134 France and the one site in Spain. Including throughfall for forested sites in Germany allowed
135 the inclusion of high deposition data beyond the level that was included in the study described
136 previously by Harmens et al. (2011).

137

138 *Moss species and sample preparation*

139 The main moss species sampled were *Pleurozium schreberi* (Willd. ex Brid.) Mitt. (*Ps*, at
140 44.4% of the sites) and *Hypnum cupressiforme* Hedw. (*Hc*, 36.3 %). Where neither of these
141 could be found, other species were collected (19.4 %): *Hylocomium splendens* (Hedw.)
142 Schimp. (*Hs*; 6.3%), *Pseudoscleropodium purum* (Hedw.) M.Fleisch. (*Pp*; 6.3%), *Thuidium*
143 *tamariscinum* (Hedw.) Schimp. (*Tt*; 5.6%) or *Abietinella abietina* (Hedw.) M.Fleisch. (*Aa*;
144 1.3%; Figure 1). Moss sampling and preparation were conducted according to guidelines
145 described in the ICP Vegetation moss monitoring manual (ICP Vegetation, 2010). Moss
146 samples were either collected below the canopy of trees but not from stems (hence, exposed
147 to throughfall deposition), or in open areas or forest clearings at least 3 m away from tree
148 crowns (see Table 1 for details). Litter and other debris was removed from the mosses and
149 green and brownish parts were separated for analysis (estimated 2 to 3 years' growth). After
150 drying the mosses were ground to a powder for the determination of nitrogen.

151

152 *Deposition sampling*

153 Most countries collected precipitation using bulk samplers with open funnels, although France
154 collected precipitation in gutters beneath the canopy of trees; Finland and Slovenia also used
155 snow collectors during winter, i.e. bulk samplers designed for winter conditions (Table 1).
156 Often, deposition was sampled according the manuals of the ICP Forests (see Table 1 for
157 details). Precipitation was collected in two or four week intervals. Wet bulk nitrogen

158 deposition (open field or throughfall) was determined from nitrogen concentration in the
159 samples and the amount of precipitation. Where possible, the averages of three years of
160 deposition data (year of moss sampling and the previous two years) were calculated to
161 correspond with the estimated two to three years of moss growth and to allow for the variation
162 in deposition between years. For Germany, 10 data points have deposition data from only one
163 year and 11 data points have only averages of two years.

164

165 *Nitrogen analysis*

166 The nitrogen concentration in mosses was determined using the Kjeldahl method (Kjeldahl,
167 1883), a modified micro-Kjeldahl method (Kubin and Siira, 1980), or by elemental analysis
168 following the Dumas method (Dumas, 1831; Table 1). Various methods were applied to
169 determine the nitrogen concentration in precipitation and throughfall (see Table 1 for details).
170 Nitrogen deposition in precipitation or throughfall was also calculated as the sum of N-NH_4^+
171 and N-NO_3^- as collected by the samplers and we will refer to this as ‘bulk nitrogen’
172 deposition. In addition, some countries (Finland and Germany) measured dissolved organic
173 nitrogen (DON) or the total nitrogen concentration (France and Slovenia) in precipitation (96
174 data points for comparison). We will refer to this as ‘total bulk nitrogen’ deposition, either
175 measured (France and Slovenia) or calculated from ‘bulk nitrogen’ plus organic nitrogen
176 deposition (other countries). One should bear in mind that this is not total nitrogen deposition
177 as the total dry deposition of nitrogen from aerosols and gas was not determined. In contrast
178 to wet-only collectors, bulk samplers often contain a fraction of total dry deposition, so open
179 bulk samplers do not only collect wet deposition (Thimonier, 1998, and reference therein).

180

181 *Quality assurance*

182 Participating laboratories, except for Germany, determined the nitrogen concentration in moss
183 reference material M2 and M3 (Steinnes et al., 1997) for quality assurance purposes (Table

184 2). Generally, the results from participating laboratories agreed well with the recommended
185 values (Harmens et al., 2010) for the nitrogen concentration in M2 and M3. In France, the
186 laboratory practise differed between 2006 and 2011, resulting in higher nitrogen
187 concentrations in the reference material M2 and M3 (Table 2). Hence, the 2011 data for
188 France were adjusted to reduce variability in the French data due to inter-laboratory
189 difference. In addition, some laboratories used other certified reference material to assure
190 good quality data, whereas the German laboratory was accredited according to standards
191 developed by the International Organization for Standardization (DIN EN ISO 17025). In
192 many countries the deposition sampling was conducted according to protocols and procedure
193 developed by International Cooperative Programmes and the determination of the
194 concentration of different nitrogen forms in deposition was subject to ring tests, inter-
195 laboratory calibration exercises and standards developed by the International Organization for
196 Standardization (see table 1 for details).

197

198 *Statistical analysis*

199 Statistical analysis was conducted using the R statistical package (www.r-project.org). A test
200 for differences between moss species was carried out by fitting a linear mixed model to moss
201 nitrogen concentrations, taking species as a factor and site as a random effect. The routine lme
202 of the R statistical package was used for this purpose. When the nitrogen concentration in
203 mosses was plotted against the various forms of measured nitrogen deposition or
204 concentration in precipitation, the moss concentrations were adjusted to allow for the
205 variability between moss species. An asymptotic relationship has been fitted to the data using
206 the R package non-linear least squares package gnls. The asymptotic relationship fitted to the
207 data can be described by the following equation:

$$208 \quad y = c + A \times (1 - \exp(-bx))$$

209 y = nitrogen concentration in mosses;

210 c = intercept on the y-axis;
211 x = deposition or concentration in precipitation of various nitrogen forms;
212 $A + c$ = the asymptote;
213 $\exp(-bx)$ represents the rate at which the asymptote is approached.

214 A non-linear mixed model was fitted to the data with parameter b being allowed to vary with
215 site as a random effect. Clear statistical outliers in the data were omitted from the analysis.

216

217 **3. Results and discussion**

218 *Interspecies variation in nitrogen concentration*

219 Previous studies have shown that the relationship between atmospheric nitrogen deposition
220 rates and the nitrogen concentration in mosses can be species-specific (Arroniz-Crespo et al.,
221 2008; Salemaa et al., 2008). Hence, the sampling of different moss species in the current
222 study might be a confounding factor and introduce ambiguity into the interpretation of the
223 possible causes of variability in the nitrogen concentration in mosses. Although atmospheric
224 nitrogen deposition was identified as the primary factor determining the total nitrogen
225 concentration in mosses, the use of different mosses species in biomonitoring programmes
226 across Europe also contributes to the spatial variation of nitrogen concentrations in mosses
227 (Schröder et al., 2010a; Harmens et al., 2011).

228 In the current study, the nitrogen concentration in mosses was determined for different
229 mosses species at a selection of sites (Figure 2). The analysis indicates significant differences
230 ($F=76.6$; 4 and 125 df) between moss species. At the extremes of the range are *H.*
231 *cupressiforme* (lowest) and *T. tamariscinum* (highest), showing a significant difference ($p <$
232 0.0001) of 2.15 mg N g^{-1} dry wt. Other species fall between *H. cupressiforme* and *T.*
233 *tamariscinum*, with overlapping confidence intervals. For a single set of paired values, the
234 analysis is equivalent to a paired t-test, showing significant differences ($p < 0.05$) for some
235 paired species: *H. cupressiforme* contained less nitrogen than *T. tamariscinum* and *P. purum*,

236 *P. schreberi* contained less nitrogen than *H. splendens* (Figure 2). For further analysis of the
237 data (see below), *H. cupressiforme* was taken as a baseline species and responses of other
238 species were linearly adjusted for bias with respect to *H. cupressiforme*. The maximum
239 adjustment was -2.14 mg g^{-1} for *T. tamariscinum*. Plots of the nitrogen concentration in
240 mosses by paired species (Figure 2) suggested that a simple bias adjustment was sufficient.

241

242 *Relationship between nitrogen concentration in mosses and various forms of wet nitrogen*
243 *deposition or concentrations in precipitation*

244 Figure 3 and 4 show the relationship between the nitrogen concentration in mosses and
245 the various forms of wet nitrogen deposition (NO_3^- , NH_4^+ , sum of NO_3^- and NH_4^+ ('bulk
246 nitrogen') and sum of NO_3^- , NH_4^+ and organic N ('total bulk nitrogen')) or concentrations in
247 precipitation respectively. Following inspection of the data and preliminary model fitting, the
248 parameter c (intercept on the y-axis) was set at 2 mg N g^{-1} dry weight, ensuring that the
249 modelled data also showed a good fit at the lower range, representing the Finnish data.
250 Parameter c is an approximation of the apparent nitrogen concentration in mosses in the
251 absence of any nitrogen deposition. While there is the appearance of an asymptotic
252 relationship, there is considerable scatter, with differing variability between countries, and
253 data for different countries positioned at different locations along the asymptotic relationship.
254 The model is therefore a first attempt to show the relationship between the nitrogen
255 concentration in mosses and the various deposition and concentration in precipitation
256 variables across Europe. It does not take full account of the correlations between some data
257 points.

258 The lowest wet bulk nitrogen deposition rates were found in Finland (Figure 3),
259 resulting in the lowest nitrogen concentrations in mosses (Poikolainen et al., 2009). The
260 Finnish data are at the lower end of the relationship, more or less within the initial linear part.
261 In Finland, the nitrogen concentration in mosses is strongly correlated ($p < 0.05$; F-test 1, 16

262 df) with all forms of nitrogen deposition and concentration in precipitation. The same is true
263 for Switzerland ($p < 0.05$; F-test 1, 30 df), where the relationship between nitrogen
264 concentration in mosses and wet bulk nitrogen deposition is more or less linear (Harmens et
265 al., 2011, 2013b). Although the moss and deposition data for Austria, France and Slovenia are
266 in a similar range as those for Switzerland, representing the middle range of the data across all
267 countries, a nationwide analysis of the data shows a lot of scatter with no significant
268 relationship ($p > 0.05$; F-test on 1 and appropriate df by country) between the nitrogen
269 concentration in mosses and all forms of wet nitrogen deposition or concentration in
270 precipitation. Especially the data for France are well-scattered regionally and not consistent
271 with the overall asymptotic behaviour shown in the Europe-wide data.

272 The German data were restricted to a few sites in Niedersachsen (North-West
273 Germany), which were sampled in various years (Mohr, 1999; Mohr et al., 2009). The
274 German throughfall data, associated with high nitrogen deposition and concentration in
275 precipitation, are the only data that lie along the asymptotic part of the relationships shown in
276 Figure 3 and 4. A few data points were available for Germany from non-throughfall sites and
277 these points fall within the mid-range of the asymptotic curves. The inclusion of the German
278 throughfall data allowed us to verify whether the asymptotic relationship observed in an
279 earlier Europe-wide study with modelled total deposition data (Harmens et al., 2011) would
280 also hold when using measured wet bulk deposition data, including bulk nitrogen deposition
281 data above $20 \text{ kg N ha}^{-1} \text{ yr}^{-1}$. Kluge et al. (2013) and Skudnik et al. (2014) found significantly
282 higher nitrogen concentrations in mosses when exposed to throughfall in forests compared to
283 exposure to atmospheric nitrogen deposition in open fields.

284 A priori, there was no reason to assume that the inclusion of throughfall nitrogen
285 might make a qualitative difference to the relationship between nitrogen deposition and
286 nitrogen concentration in mosses. However, nitrogen speciation in throughfall might differ
287 from that in wet deposition due to canopy exchange processes, possibly affecting the

288 ammonium-N to nitrate-N ratio (Draaijers et al., 1997; Adriaenssens et al., 2012) and the
289 contribution of dissolved organic nitrogen (DON; Drápelová, 2012), potentially affecting the
290 uptake of nitrogen in mosses (see below). Although mosses have a preference for ammonium
291 uptake (see below), which might suppress the utilization of DON, the contribution of
292 atmospheric DON to the nitrogen concentration in mosses could be significant (Liu et al.,
293 2013). In addition, the microclimate in forest undergrowth is likely to differ from more
294 exposed locations and such microclimate differences might affect the relationships studied
295 here (Harmens et al., 2011). The data from the current study do not allow direct assessment of
296 the impact of throughfall as at none of the sites a comparison was made between nitrogen
297 concentrations in mosses sampled under the influence of tree canopies and mosses sampled in
298 the open field.

299 Ammonium and nitrate deposition are generally of the same order, with the exception
300 of the throughfall sites in Germany, where ammonium deposition exceeded nitrate by a factor
301 of two to three. The German sites with high ammonium deposition rates are the most
302 influential in providing evidence of asymptotic behaviour in the nitrogen concentration in
303 mosses. That is to say, the nitrogen concentration in mosses does not appear to respond to
304 increasing ammonium-N deposition of over $12 \text{ kg ha}^{-1} \text{ yr}^{-1}$, and wet bulk N deposition of over
305 $20 \text{ kg ha}^{-1} \text{ yr}^{-1}$ (Figure 3), or at ammonium-N concentration in precipitation of over 2 mg l^{-1}
306 (Figure 4). In contrast, the asymptotic behaviour is very weak with respect to nitrate-N
307 deposition or concentration in precipitation. The asymptotic behaviour with respect to
308 ammonium-N is even more pronounced when precipitation concentrations are considered,
309 because rainfall at the throughfall sites in Germany is relatively low (Figure 4). Saturation of
310 nitrogen concentration in mosses at high ammonium deposition or concentration in
311 precipitation might reflect a lower uptake efficiency at higher nitrogen exposure (Pitcairn et
312 al., 2006; Wiedermann et al., 2009). Previous studies have reported a higher uptake of
313 ammonium than nitrate in mosses (Forsum et al., 2006; Jauhiainen et al., 1998; Liu et al.,

314 2013; Pearce et al., 2003; Soares and Pearson, 1997; Wiedermann et al., 2009), which is
315 probably due to the high cation-exchange capacity common for mosses (Bates, 1992).
316 Utilising NH_4^+ as a nitrogen source as opposed to NO_3^- is commonly regarded as being more
317 energy efficient, achieving greater specific growth rates. NO_3^- assimilation in mosses was
318 found to be negligible when the supply rate of reduced dissolved nitrogen (NH_4^+ plus DON)
319 was significantly higher than that of NO_3^- (Liu et al., 2012). However, in the current study,
320 the supply rate of NH_4^+ and NO_3^- was similar at most sites except in Germany, so it is
321 unknown whether NO_3^- assimilation was low. If NO_3^- assimilation was low, the effect of
322 NO_3^- deposition on the nitrogen concentration in mosses is likely to be overestimated.

323 The Akaike Information Criterion (AIC), an indicator of model fit, suggests that the
324 best fit is obtained by the combined concentration of ammonium and nitrate in rainfall for
325 data including all countries (Table 3). Analysis for Finland, France, Germany and Slovenia
326 only indicates that there is no further improvement in fit using total nitrogen concentration in
327 precipitation. In Germany and France, the average contribution of DON to the total wet bulk
328 deposition ranged from 6 to 28% respectively, which is similar to the range reported for the
329 Czech Republic (Drápelová, 2012).

330

331 *Uncertainty in the contribution of other sources to the nitrogen concentration in mosses*

332 The lack of data on other nitrogen sources potentially contributing to the nitrogen
333 concentration in mosses is likely to contribute to the scatter in the data and the uncertainty of
334 the relationships shown in Figure 3 and 4. In the current study, we only included nitrogen
335 from wet bulk deposition as data on dry deposition was lacking for most sites (although some
336 dry deposition will be included in wet bulk deposition samplers; Thimonier, 1998). Pitcairn et
337 al. (2006) have shown that nitrogen concentration in mosses respond differently to wet and
338 dry deposited nitrogen. For a $1 \text{ kg ha}^{-1} \text{ yr}^{-1}$ increase in nitrogen deposition, tissue nitrogen
339 increased by 0.1 mg g^{-1} at wet deposition sites but by 0.3 mg g^{-1} at sites dominated by dry

340 deposited ammonia downwind of livestock (poultry and pig) farms in Scotland. Larger
341 concentrations of nitrogen (up to 40 mg g⁻¹) occurred in mosses at sites where nitrogen
342 deposition was dominated by dry deposited ammonia and where rainfall (and therefore
343 leaching losses) was small, compared with sites where deposition was dominated by wet
344 deposition (up to 16 mg g⁻¹). In the current study, the maximum nitrogen concentration in
345 mosses was 25 mg g⁻¹ at throughfall sites in the agriculturally intensive region of Germany,
346 where dry nitrogen deposition is high. Thus, the critical nitrogen concentration of 20 mg g⁻¹,
347 specified by Pitcairn et al. (1998) was exceeded considerably in Germany.

348 In addition to inorganic nitrogen, mosses also take up DON, hence analysis of DON
349 should be included to account fully for nitrogen input to mosses from precipitation (Forsum et
350 al., 2006). Several studies have reported the preferred uptake of ammonia, followed by DON
351 or amino acids, over nitrate (Forsum et al., 2006; Hill et al., 2011; Liu et al., 2013; Wanek and
352 Portl, 2008; Wiedermann et al., 2009) and in some cases amino acids may be the preferred
353 source of nitrogen for certain moss species (Kielland, 1997; McKane et al., 1993). In the
354 current study, the relationships shown in Figure 3 and 4 did not change much when DON
355 (total bulk nitrogen deposition or total nitrogen precipitation) was included in addition to
356 ammonia and nitrate.

357 Scatter in the data might also be caused by uptake of nitrogen from the soil (Ayres et
358 al., 2006). Although Liu et al. (2013) reported that the uptake of nitrogen from the soil might
359 contribute significantly (ca. 37%) to the nitrogen concentration in terricolous mosses, this is
360 in contrast to other studies stating that mosses receive most of the nitrogen from deposition,
361 leaching and throughfall (Kotanen, 2002; Li and Vitt, 1997; Rousk et al., 2013a; Turetsky,
362 2003). In the current study, mosses in forested areas were sampled from tree stumps where
363 possible, where uptake of soil nitrogen is unlikely to play a significant role.

364 At some lower nitrogen deposition sites with relatively high nitrogen concentration in
365 mosses, cyanobacteria living in association with mosses could potentially be responsible for

366 the high nitrogen concentration in mosses (Rousk et al., 2013b). However, the number of
367 cyanobacteria cells was shown to decline significantly at nitrogen deposition rates of 5 kg ha^{-1}
368 yr^{-1} or more compared to the background deposition rate of $2 \text{ kg ha}^{-1} \text{ yr}^{-1}$ (Gundale et al.,
369 2011). In the current study, sites with relatively high nitrogen concentrations in mosses at low
370 nitrogen deposition were found in Austria, where cyanobacterial associations were not
371 observed, and in France, where the drier climate is not conducive to high cyanobacterial
372 activity (Rousk et al., 2013b). A relatively high nitrogen concentration in mosses was also
373 observed at one Finnish site, however, this is unlikely to be due to cyanobacterial fixation of
374 nitrogen as at many other Finnish sites with lower nitrogen deposition rates the nitrogen
375 concentration in mosses was much lower. Leppänen et al. (2013) showed that nitrogen
376 fixation associated with mosses increased towards the north and was hardly observed in the
377 south of Finland, where nitrogen deposition rates are higher.

378 Other factors that are likely to contribute to the scatter in the data (e.g. effects of
379 nitrogen and microclimate on moss growth, surrounding vegetation type and land use) have
380 been discussed in more detail elsewhere (Harmens et al., 2011; Schröder et al., 2010a). In the
381 current study, the distance between the moss sampling sites and the deposition measurement
382 sites varied between 1 – 900 m. In general, there is a high spatial and temporal variability in
383 throughfall (Thimonier, 1998) and in wet deposition of nitrogen (Harmens et al., 2011 and
384 references therein), especially in mountainous regions. Hence, a distance of up to 900 m
385 between moss sampling and deposition measurement site could also contribute to the scatter
386 in the data.

387

388 **Conclusions**

389 As previously described for modelled nitrogen deposition, the relationship between nitrogen
390 concentration in mosses and measured (total) wet bulk deposition or concentration in
391 precipitation across Europe is best described by an asymptotic relationship. The asymptotic

392 relationship is much stronger for ammonia-N than for nitrate-N in bulk deposition or
393 precipitation. Saturation appears to occur at wet bulk nitrogen deposition rates of ca. 20 kg ha⁻¹
394 yr⁻¹. Up to such deposition rates, linear relationships have been observed in some countries
395 (Finland and Switzerland) but not in others. Considerable scatter was observed in the
396 relationship at the European level, although less than previously found with modelled total
397 deposition (Harmens et al., 2011). The scatter in the data might potentially be reduced by
398 repeating this study with:

- 399 • Both mosses and precipitation sampled at the same site, rather than up to 1 km apart,
400 to minimise the influence of spatial variation in nitrogen deposition;
- 401 • Including analysis of ammonia and nitrogen dioxide measured with passive samplers
402 as an indication of dry deposition and measurements of DON to calculate total
403 nitrogen deposition;
- 404 • Further harmonising and improving the methodology of moss and deposition
405 sampling, and chemical analysis, and minimise the potential uptake of nitrogen from
406 soil;
- 407 • Measuring nitrogen concentration in mosses at more sites with high nitrogen
408 deposition or concentration in precipitation.

409 The moss technique remains a valuable tool to identify areas at risk of high nitrogen
410 deposition at a high spatial resolution in a cost-effective manner and appears to be a
411 complementary tool for estimating wet bulk nitrogen deposition in low to medium nitrogen
412 deposition areas. In addition, data for various years will allow analysis of temporal trends in
413 atmospheric nitrogen deposition (Harmens et al., 2013b).

414

415 **Acknowledgements**

416 We thank the United Kingdom Department for Environment, Food and Rural Affairs (Defra;
417 contract AQ0816), the UNECE (Trust Fund) and the Natural Environment Research Council

418 (NERC) for funding the ICP Vegetation Programme Coordination Centre at CEH Bangor,
419 UK. We thank David Norris (CEH Bangor) for producing figure 1. We thank the following
420 persons for supplying bulk deposition and concentration data from national monitoring
421 networks:

- 422 - Sabine Braun, Institute for Applied Plant Biology, Schönenbuch, Switzerland;
- 423 - Maria Schmitt, Anne Thimonier, Peter Waldner, Long Term Ecosystem Research,
424 Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Switzerland;
- 425 - Manuel Nicolas, Office National des Forêts, French Permanent Plot Network for the
426 Monitoring of Forest Ecosystems (RENECOFOR), Fontainebleau, France;
- 427 - Daniel Žlindra, Primož Simončič, Intensive monitoring of forest ecosystems,
428 Slovenian Forestry Institute, Slovenia;
- 429 - Ferdinand Kristöfel, Markus Neumann, Federal Research Centre for Forest, Austria;
- 430 - Enrico Pompei, National Integrated Programme for Forest Ecosystems Monitoring
431 (CONECOFOR), Corpo forestale dello Stato, Italy.

432

433 **References**

- 434 Adriaenssens, S., Hansen, K., Staelens, J., Wuyts, K., De Schrijver, A., Baeten, L., Boeckx,
435 P., Samson, R., Verheyen, K., 2012. Throughfall deposition and canopy exchange
436 processes along a vertical gradient within the canopy of beech (*Fagus sylvatica* L.) and
437 Norway spruce (*Picea abies* (L.) Karst). *Science of the Total Environment* 420: 168–
438 182.
- 439 Arroniz-Crespo, M., Leake, J.R., Horton, P., Phoenix, G.K., 2008. Bryophyte physiological
440 responses to, and recovery from, long-term nitrogen deposition and phosphorus
441 fertilisation in acidic grassland. *New Phytologist* 180: 864-874.
- 442 Ayres, E., Van der Wal, R., Sommerkorn, M., Bardgett, R.D., 2006. Direct uptake of soil
443 nitrogen by mosses. *Biology Letters* 2: 286-288.

444 Bates, J.W., 1992. Mineral nutrient acquisition and retention by bryophytes. *Journal of*
445 *Bryology* 17: 223-240.

446 Clarke, N., Žlindra, D., Ulrich, E., Mosello, R., Derome, J., Derome, K., König, N., Lövblad,
447 G., Draaijers, G.P.J., Hansen, K., Thimonier, A., Waldner, P., 2010. Sampling and
448 Analysis of Deposition - Part XIV. In: UNECE, I.F. (Ed.), *Manual on methods and*
449 *criteria for harmonized sampling, assessment, monitoring and analysis of the effects of*
450 *air pollution on forests*. vTI - Institute for World Forestry, Hamburg, pp. 66.

451 Delgado, V., Alicia Ederra, A., Santamaría, J.M., 2013. Nitrogen and carbon contents and
452 $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ signatures in six bryophyte species: assessment of long-term deposition
453 changes (1980-2010) in Spanish beech forests. *Global Change Biology* 19: 2221-2228.

454 Draaijers, G. P. J., Erisman, J. W., Van Leeuwen, N. F. M., Romer, F. G., Te Winkel, B. H.,
455 Veltkamp, A. C., Vermeulen, A.T., Wyers, G. P., 1997. Impact of canopy exchange on
456 differences observed between the atmospheric deposition and throughfall fluxes.
457 *Atmospheric Environment* 31: 387-397.

458 Drápelová, 2012. Organic and inorganic nitrogen in precipitation and in forest throughfall at
459 the Bílý Kříž site (Beskydy Mts., Czech Republic). *Journal of Forest Science* 58: 88–
460 100.

461 Dumas, J.B.A., 1831. *Procedes de l'Analyse Organique*. *Annales de Chimie et de Physique*
462 247, 198-213.

463 Foan, L., Sablayrolles, C., Elustondo, D., Laceras, E., González, L., Hederá, A., Simon, V.,
464 Santamaría, J.M., 2010. Reconstructing historical trends of polycyclic aromatic
465 hydrocarbon deposition in a remote area of Spain using herbarium moss material.
466 *Atmospheric Environment* 44: 3207-3214.

467 Foan, L., Leblond, S., Thöni, L., Raynaud, C., Santamaria, J.M, Sebiló, M., Simon, V., 2014.
468 Spatial distribution of PAH concentrations and stable isotopic signatures ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$) in

469 mosses from three European areas - Characterization by multivariate analysis.
470 Environmental Pollution 184: 113-122.

471 Forsum, Å., Dahlman, L., Näsholm, T., Nordin, A., 2006. Nitrogen utilization by
472 *Hylocomium splendens* in a boreal forest fertilization experiment. Functional Ecology
473 20: 421–426.

474 Gundale, M.J., Deluca, T.H., Nordin, A., 2011. Bryophytes attenuate anthropogenic nitrogen
475 inputs in boreal forests. Global Change Biology 17: 2743-2753.

476 Hansen, K., Thimonier, A., Clarke, N., Staelens, J., Žlindra, D., Waldner, P., Marchetto, A.,
477 2013. Chapter 18 - Atmospheric Deposition to Forest Ecosystems. In: Ferretti, M.,
478 Fisher, R. (Eds.), Developments in Environmental Science. Elsevier, pp. 337-374.

479 Harmens, H., Ilyin, I., Mills, G., Aboal, J.R., Alber, R., Blum, O., Coşkun, M., De
480 Temmerman, L., Fernández, J.A., Figueira, R., Frontasyeva, M., Godzik, B., Goltsova,
481 N., Jeran, Z., Korzekwa, S., Kubin, E., Kvietkus, K., Leblond, S., Liiv, S., Magnússon,
482 S.H., Maňková, B., Nikodemus, O., Pesch, R., Poikolainen, J., Radnović, D., Rühling,
483 Å., Santamaria, J.M., Schröder, W., Spiric, Z., Stafilov, T., Steinnes, E., Suchara, I.,
484 Tabor, G., Thöni, L., Turcsányi, G., Yurukova, L., Zechmeister, H.G., 2012. Country-
485 specific correlations across Europe between modelled atmospheric cadmium and lead
486 deposition and concentrations in mosses. Environmental Pollution 166: 1-9.

487 Harmens, H., Norris, D. A., Cooper, D.M., Mills, G., Steinnes E., Kubin, E., Thöni, L., Aboal,
488 J.R., Alber, R., Carballeira, A., Coşkun, M., De Temmerman, L., Frolova, M.,
489 Frontasyeva, M., Gonzáles-Miqueo, L., Jeran, Z., Leblond S., Liiv, S., Maňková, B.,
490 Pesch, R., Poikolainen, J., Rühling, Å., Santamaria, J. M., Simonè, P., Schröder, W.,
491 Suchara, I., Yurukova, L., Zechmeister, H. G., 2011. Nitrogen concentrations in
492 mosses indicate the spatial distribution of atmospheric nitrogen deposition in Europe.
493 Environmental Pollution 159: 2852-2860.

494 Harmens, H., Foan, L., Simon, V., Mills, G., 2013a. Terrestrial mosses as biomonitors of
495 atmospheric POPs pollution: A review. *Environmental Pollution* 173: 245-254.

496 Harmens, H., Norris, D., Mills, G., and the participants of the moss survey, 2013b. Heavy
497 metals and nitrogen in mosses: spatial patterns in 2010/2011 and long-term temporal
498 trends in Europe. ICP Vegetation Programme Coordination Centre, CEH Bangor, UK.
499 ISBN: 978-1-906698-38-6. <http://icpvegetation.ceh.ac.uk>

500 Harmens, H., Norris, D.A., Steinnes, E., Kubin, E., Piispanen, J., Alber, R., Aleksiyenak,
501 Y., Blum, O., Coşkun, M., Dam, M., De Temmerman, L., Fernández, J.A., Frolova, M.,
502 Frontasyeva, M., González-Miqueo, L., Grodzińska, K., Jeran, Z., Korzekwa, S.,
503 Krmar, M., Kvietkus, K., Leblond, S., Liiv, S., Magnússon, S.H., Maňkiovská, B.,
504 Pesch, R., Rühling, Å., Santamaria, J.M., Schröder, W., Spiric, Z., Suchara, I., Thöni,
505 L., Urumov, V., Yurukova, L., Zechmeister, H.G., 2010. Mosses as biomonitors of
506 atmospheric heavy metal deposition: spatial and temporal trends in Europe.
507 *Environmental Pollution* 158, 3144-3156.

508 Hill, P.W., Farrar, J., Roberts, P., Farrell, M., Grant, H., Newsham, K., Hopkins, D.W.,
509 Bardgett, R.D., Jones, D.L., 2011. Vascular plant success in a warming Antartic may be
510 due to efficient nitrogen acquisition. *Nature Climate Change* 1: 50 – 53.

511 ICP Vegetation, 2010. Heavy metals in European mosses: 2010 survey. Monitoring manual.
512 UNECE ICP Vegetation Coordination Centre, CEH Bangor, UK.
513 <http://icpvegetation.ceh.ac.uk>

514 Jauhiainen, J., Wallén, B., Malmer, N., 1998. Potential NH_4^+ and NO_3^- uptake in seven
515 *Sphagnum* species. *New Phytologist* 138: 287–293.

516 Kielland, K., 1997. Role of free amino acids in the nitrogen economy of arctic cryptogams.
517 *Ecoscience* 4: 75–79.

518 Kjeldahl, J., 1883. A new method for the determination of nitrogen in organic matter.
519 *Zeitschrift für Analytische Chemie* 22, 366.

520 Kluge, M., Pesch, R., Schröder, W., Hoffmann, A., 2013. Accounting for canopy drip effects
521 of spatiotemporal trends of concentrations of N in mosses, atmospheric N depositions
522 and critical load exceedances: a case study from North-Western Germany.
523 Environmental Sciences Europe 25: 26.

524 König, N., Cools, N., Derome, K., Kowalska, A., De Vos, B., Fürst, A., Marchetto, A., O'Dea,
525 P., Tartari, G.A., 2013. Chapter 22 - Data Quality in Laboratories: Methods and Results
526 for Soil, Foliar, and Water Chemical Analyses. In: Ferretti, M., Fisher, R. (Eds.),
527 Developments in Environmental Science. Elsevier, pp. 415-453.

528 Koranda, M., Kerschbaum, S., Wanek, W., Zechmeister, H.A., Richter, A., 2007.
529 Physiological responses of bryophytes *Thuidium tamariscinum* and *Hylocomium*
530 *splendens* to increased nitrogen deposition. Annals of Botany 99:161–169

531 Kotanen, P.M., 2002. Fates of added nitrogen in freshwater arctic wetlands grazed by snow
532 geese: the role of mosses. Arctic Antarctic Alpine Research 34: 219-225.

533 Kubin, E., Siira, J., 1980. On the suitability of the phenyl hypochlorite procedure for the
534 determination of total nitrogen from plant and soil samples. Aquilo Ser. Botanica 17,
535 11-17.

536 Leder, K.; Puxbaum, H.; Heimbürger, G.; Reinisch, R. Nasse, 2005. Deposition im Land
537 Kärnten, Oktober 2004 bis September 2005. Institut für Chemische Technologien und
538 Analytik, 06/05, TU-Wien, Österreich.

539 Leonardi S., Flückiger W., 1987. Short-term canopy interactions of beech trees: mineral ion
540 leaching and absorption during rainfall. Tree Physiology 3: 137-145.

541 Leppänen, S.M., Salemaa, M., Smolander, A., Mäkipää, R., Tirola, M., 2013. Nitrogen
542 fixation and methanotrophy in forest mosses along a N deposition gradient.
543 Environmental and Experimental Botany 90: 62-69.

544 Li, Y., Vitt, D.H., 1997. Patterns of retention and utilization of aerially deposited nitrogen in
545 boreal peatlands. Ecoscience 4: 106-116.

546 Liu, X.-Y., Koba, K., Liu, C.-Q., Li, X.-D., Yoh, M., 2012. Pitfalls and new mechanisms in
547 moss isotopic bio-monitoring of atmospheric nitrogen deposition. *Environmental*
548 *Science and Technology* 46: 12557-12566.

549 Liu, X.-Y., Koba, K., Makabe, A., Li, X.-D., Yoh, M., Liu, C.-Q., 2013. Ammonium first:
550 natural mosses prefer atmospheric ammonium but vary utilization of dissolved organic
551 nitrogen depending on habitat and nitrogen deposition. *New Phytologist* 199: 407-419.

552 Marchetto, A., Mosello, R., Tartari, G., Derome, J., Derome, K., König, N., Clarke, N.,
553 Kowalska, A., 2009a. Atmospheric deposition and soil working ring test 2009. CNR-
554 ICE Report. 56 p.

555 Marchetto, A., Mosello, R., Tartari, G., Tornimbeni, O., Derome, J., Derome, K., Sorsa, P.,
556 König, N., Clarke, N., Ulrich, E., Kowalska, A., 2009b. Influence of QA/QC
557 procedures on non-sampling error in deposition monitoring in forests. *Journal of*
558 *Environmental Monitoring* 11: 745-750.

559 McKane, R., Johnson, L., Shaver, G., Nadelhoffer, K., Fry, B., Rastetter, E., Giblin, A.,
560 Laundre, J., 1993. Differentiation in uptake of ^{15}N by depth, season, and chemical form
561 in an arctic tussock tundra plant community. 78th Ann. ESA Meeting, 31 July - 4 August
562 1993. *Bull. Ecol. Soc. Amer. Program and Abstracts, Suppl. vol 74(2): 354.*

563 Mohr, K., 1999. Passive Monitoring von Stickstoffeinträgen in Kiefernforsten mit dem
564 Rotstengelmoos (*Pleurozium schreberi* (Brid.) Mitt.). *Umweltwissenschaften und*
565 *Schadstoff-Forschung - Zeitschrift für Umweltchemie und Ökotoxikologie* 11: 721 –
566 735.

567 Mohr K., Holy M., Pesch R., Schröder W., 2009. Bioakkumulation von Metallen und
568 Stickstoff zwischen 1990 und 2005 in Niedersachsen. *Umweltwissenschaften und*
569 *Schadstoff-Forschung - Zeitschrift für Umweltchemie und Ökotoxikologie* 21: 454-469.

570 Mosello, R., Brizzio, M.C., Kotzias, D., Marchetto, A., Rembges, D., Tartari, G., 2002. The
571 chemistry of atmospheric deposition in Italy in the framework of the National
572 Programme for Forest Ecosystems Control (CONECOFOR).

573 Pearce, I.S.K., Woodin, S.J., Van der Wal, R., 2003. Physiological and growth responses of
574 the montane bryophyte *Racomitrium lanuginosum* to atmospheric nitrogen deposition.
575 *New Phytologist* 160: 145-155.

576 Pitcairn, C., Fowler, D., Leith, I., Sheppard, L., Tang, S., Sutton, M., Famulari, D., 2006.
577 Diagnostic indicators of elevated nitrogen deposition. *Environmental Pollution* 144:
578 941-950.

579 Pitcairn, C., Leith, I.D., Sheppard, L.J., Sutton, M.A. Fowler, D. Munro, R.C. Tang, S.,
580 Wilson, D., 1998. The relationship between nitrogen deposition, species composition
581 and foliar nitrogen concentration in woodland flora in the vicinity of livestock farms.
582 *Environmental Pollution* 102: 41-48.

583 Poikolainen, J., Piispanen, J., Karhu, J., Kubin, E., 2009. Long-term changes in nitrogen
584 deposition in Finland (1990-2006) monitored using the moss *Hylocomium splendens*.
585 *Environmental Pollution* 157:, 3091-3097.

586 Rousk, K., Rousk, J., Jones, D.L., Zackrisson, O., DeLuca, T.H., 2013a. Feather moss
587 nitrogen acquisitions across natural fertility gradients in boreal forests. *Soil Biology and*
588 *Biochemistry* 61: 86-95.

589 Rousk, K., Jones, D.L., DeLuca, T.H., 2013b. Moss-cyanobacteria associations as biogenic
590 sources of nitrogen in boreal forest ecosystems. *Frontiers in Microbiology* 4 (150): 1-
591 10.

592 Salemaa, M., Mäkipää, R., Oksanen, J., 2008. Differences in the growth response of three
593 bryophyte species to nitrogen. *Environmental Pollution* 152: 82-91.

594 Schröder, W., Holy, M., Pesch, R., Harmens, H., Fagerli, H., Alber, R., Coşkun, M., De
595 Temmerman, L., Frolova, M., González-Miqueo, L., Jeran, Z., Kubin, E., Leblond, S.,

596 Liiv, S., Maňkovská, B., Piispanen, J., Santamaría, J.M., Simonèè, P., Suchara, I.,
597 Yurukova, L., Thöni, L., Zechmeister, H.G., 2010a. First Europe-wide correlation
598 analysis identifying factors best explaining the total nitrogen concentration in mosses.
599 Atmospheric Environment 44: 3485-3491.

600 Schröder, W., Holy, M., Pesch, R., Harmens, H., Ilyin, I., Steinnes, E., Alber, R.,
601 Aleksiyenak, Y., Blum, O., Coşkun, M., Dam, M., De Temmerman, L., Frolova, M.,
602 Frontasyeva, M., González-Miqueo, L., Grodzińska, K., Jeran, Z., Korzekwa, S.,
603 Krmar, M., Kubin, E., Kvietkus, K., Leblond, S., Liiv, S., Magnússon, S., Maňkovská,
604 B., Piispanen, J., Rühling, Å., Santamaria, J., Spiric, Z., Suchara, I., Thöni, L., Urumov,
605 V., Yurukova, L., Zechmeister, H.G., 2010b. Are cadmium, lead and mercury
606 concentrations in mosses across Europe primarily determined by atmospheric deposition
607 of these metals? Journal of Soils and Sediments 10: 1572-1584.

608 Skudnik, M., Jeran, Z., Batic, F., Simončič, P., Lojen, S., Kastelec, D., 2014. Influence of
609 canopy drip on the indicative N, S and $\delta^{15}\text{N}$ content in moss *Hypnum cupressiforme*.
610 Environmental Pollution 190: 27-35.

611 Smidt, S., 2007. 10 Jahre Depositionsmessungen im Rahmen des Europäischen
612 Waldschadensmonitorings. Ergebnisse 1996-2005. BFW-Berichte 138: 1-79.

613 Soares, A., Pearson, J., 1997. Short-term physiological responses of mosses to atmospheric
614 ammonium and nitrate. Water, Air and Soil Pollution 93: 225-242.

615 Solga, A., Burkhardt, J., Zechmeister, H.G., Frahm, J.-P., 2005. Nitrogen content, ^{15}N natural
616 abundance and biomass of the two pleurocarpous mosses *Pleurozium schreberi* (Brid.)
617 Mitt., and *Scleropodium purum* (Hedw.) Limpr. in relation to atmospheric nitrogen
618 deposition. Environmental Pollution 134 : 465-473.

619 Steinnes, E., Rühling, Å., Lippo, H., Mäkinen, A., 1997. Reference material for large-scale
620 metal deposition surveys. Accreditation and Quality Assurance 2: 243-249.

621 Stevens, C.J., Duprè, C., Dorland, E., Gaudnik, C., Gowing, D.J.G., Bleeker, A., Diekmann,
622 M., Alard, D., Bobbink, R., Fowler, D., Corcket, E., Mountford, J.O., Vandvik, V.,
623 Aarrestad, P.A., Muller, S., Dise, N.B., 2011. The impact of nitrogen deposition on acid
624 grasslands in the Atlantic region of Europe. *Environmental Pollution* 159: 2243-2250.

625 Thimonier, A., 1998. Measurement of atmospheric deposition under forest canopies: Some
626 recommendations for equipment and sampling design. *Environmental Monitoring and*
627 *Assessment* 53: 353–387.

628 Thimonier, A, Schmitt, M., Waldner, P., Rihm, B., 2005. Atmospheric deposition on Swiss
629 long-term forest ecosystem research (LWF) plots. *Environmental Monitoring and*
630 *Assessment* 104:, 81-118.

631 Thöni, L., Matthaei, D., Seitler, E., Bergamini, A., 2008. Deposition von Luftschadstoffen in
632 der Schweiz. Moosanalysen 1990 - 2005. Umwelt Zustand Nr. 0827, Bundesamt für
633 Umwelt, Bern, Switzerland. www.umwelt-schweiz.ch/uz-0827-d

634 Thöni, L., Seitler, E., 2010. Stickstoff- & Elementdeposition OSTLUFT, 2000 bis 2008.
635 http://www.ostluft.ch/fileadmin/intern/LZ_Information/Publikationen/Fachberichte/BE_
636 [StickstoffElementdeposition2000-2008_PT_201002.pdf](http://www.ostluft.ch/fileadmin/intern/LZ_Information/Publikationen/Fachberichte/BE_StickstoffElementdeposition2000-2008_PT_201002.pdf)

637 Turetsky, M.R., 2003. The role of bryophytes in carbon and nitrogen cycling. *Bryologist* 106:
638 395-409.

639 Wanek, W., Portl, K., 2008. Short-term ¹⁵N uptake kinetics and nitrogen nutrition of
640 bryophytes in a lowland rainforest, Costa Rica. *Functional Plant Biology* 35: 51-62.

641 Wiedermann, M.M., Gunnarsson, U., Ericson, L., Nordin, A., 2009. Ecophysiological
642 adjustment of two *Sphagnum* species in response to anthropogenic nitrogen deposition.
643 *New Phytologist* 181: 208-217.

644 Zechmeister, H.G., Richter, A., Smidt, S., Hohenwallner, D., Roder, I., Maringer, S., Wanek,
645 W., 2008. Total nitrogen content and $\delta^{15}\text{N}$ signatures in moss tissue: indicative value for

646 nitrogen deposition patterns and source allocation on a nationwide scale. Environmental
647 Science and Technology 42: 8661-8667.

648 Žlindra, D., Skudnik, M., Rupel, M., Simončič, P., 2011. Measuring of precipitation quality in
649 the open and in a stand on the plots for intensive monitoring of forest ecosystems.
650 Gozdarski Vestnik 69: 279-288.

651 **Table 1.** Overview of methods applied in selected European countries to determine nitrogen concentrations in mosses and bulk precipitation.

Country	Institute	Deposition or throughfall	Moss species*	Sample years	Analysis N in moss	Moss QA method	Sampler type	Analysis of N deposition	QA method deposition	Monitoring network	Reference
Austria (AT)	Umweltbundesamt Wien	Deposition	Aa, Hc, Hs, Ps	2005	Elemental analysis	Standards M2 & M3; ÖNORM CEN/TS 15407	Bulk sampler	Chemoluminescens	Multiple sampling	ICP Forests (Smidt, 2007) National network (Leder et al., 2005)	Leder et al., 2005 Smidt, 2007 Zechmeister et al., 2008
Finland (FI)	Finnish Forest Research Institute (Metla)	Deposition	Ps	2009 & 2011	Modified Kjeldahl (Kubin and Siira, 1980)	Moss standards M2 & M3	Bulk sampler, incl. snow collector	NO ₃ -N: Ion chromatography (IC); NH ₄ -N, N _{tot} : Flow injection analysis	ICP Forests Manual (Clarke et al., 2010)	ICP Forests	Clarke et al., 2010 Kubin and Siira, 1980
France (FR)	Muséum national d'Histoire naturelle, Office National des Forêts	Throughfall	Hc, Hs, Pp, Tt	2006 & 2011	Elemental analysis	Moss standards M2 & M3, repeated sampling	Gutters beneath canopy	NO ₃ ⁻ , NH ₄ ⁺ : IC N total: chemoluminescence	ICP Forests Manual (Clarke et al., 2010) Ring test (Marchetto et al., 2009b)	RENECOFOR network (ICP Forests) BRAMM network (ICP Vegetation)	Clarke et al., 2010
Germany - Niedersachsen (DE-NI)	Landwirtschaftskammer Niedersachsen	Deposition and throughfall	Hc, Ps	1998-2010	Kjeldahl	Accredited DIN EN ISO 17025	Bulk sampler	Continous flow analyzer			
Slovenia** (SI)	Slovenian Forestry Institute	Deposition	Hc	2010	Elemental analysis	Standards M2 & M3	Bulk samplers, incl. snow collector	NO ₃ ⁻ , NH ₄ ⁺ : IC N total: UV-Vis Spectrophotometer	QC standards: Use of reference materials and ring tests (König et al., 2013)	ICP Forests intensive monitoring plots	Clarke et al., 2010 Hansen et al., 2013 König et al., 2013 Mosello et al., 2002 Smidt, 2007 Žlindra et al., 2011
Spain (ES)	University of Navarra	Throughfall	Hc	2010 & 2012	Elemental analysis	Standards M2 & M3	Bulk sampler	NO ₃ ⁻ , NH ₄ ⁺ : IC	Intercalibration ICP Waters; certified material	ICP Integrated Monitoring	Delgado et al., 2013
Switzerland (CH)	FUB - Research Group for Environmental Monitoring	Deposition	Hc, Ps	2005 & 2010	Kjeldahl	Standards M2 & M3, NIST-SRM 1515, repeated sampling	Bulk sampler	NO ₃ ⁻ : IC NH ₄ ⁺ : Flow injection analysis & Indophenolmethod (Spectrophotometer)	Reference material simulated rain: CRM 408 CEC bcr 1993; Ring test (Marchetto et al., 2009a)	ICP Forest (Thimonier et al., 2005) Swiss intercantonal research project	Leonardi and Flückiger, 1987 Marchetto et al., 2009a Thimonier et al., 2005 Thöni and Seitler, 2010

652

653

654

655

656

* Hs: *Hylocomium splendens*, Hc: *Hypnum cupressiforme*, Ps: *Pleurozium shreberi*, Pp: *Pseudoscleropodium purum*, Tt: *Thuidium tamariscinum*. Aa: *Abietinella abietina*
 ** A few moss samples were collected in Italy (1 site) and Austria (3 sites) near the Slovenian border, where deposition was sampled by the Ministry for Agriculture and Forestry Policies, CONECOFOR Service, National Forest Service and the Institut für Waldwachstum und Waldbau Waldschadenserfassung respectively.

657 **Table 2.** Nitrogen concentration (mg N g^{-1} dry weight; mean \pm one standard deviation) in the moss standards M2 and M3 (Harmens et al., 2010). N
 658 is the number of repeated analyses of the standard; the value in parenthesis indicates the year of analysis for those countries who repeated the
 659 sampling with time.
 660

Moss standard	Recommended value	Austria	Switzerland (2010)	France	Finland	Slovenia	Spain
M2 (mg N g^{-1} dry wt)	8.36 ± 0.62 (N = 10)	6.95 ± 0.28 (N=2)	7.81 ± 0.62 (N = 6)	8.32 ± 0.11 (N = 5) (2006) 9.05 ± 0.31 (N = 17) (2011)		8.27 ± 0.23 (N = 6)	8.80 ± 0.13 (N = 6)
M3 (mg N g^{-1} dry wt)	6.81 ± 0.52 (N = 8)	6.06 ± 0.29 (N=2)	6.93 ± 0.26 (N = 4)	6.57 ± 0.13 (N = 10) (2006) 7.48 ± 0.28 (N = 17) (2011)	6.82 ± 0.29 (N = 6) (2009) 6.66 ± 0.13 (N = 5) (2011)	6.72 ± 0.26 (N = 6)	7.30 ± 0.11 (N = 6)

661 **Table 3.** Parameters of the asymptotic relationship between nitrogen concentration in mosses
 662 and wet bulk deposition or concentration in precipitation for different nitrogen forms. The
 663 asymptotic relationship is described as $y = c + A \times (1 - \exp(-bx))$; AIC = Akaike Information
 664 Criterion.

665

Bulk deposition/concentration variable	A	b	AIC*
NH ₄ -N deposition	20.5	0.1911	542.3
NO ₃ -N deposition	22.5	0.1843	635.9
NO ₃ -N + NH ₄ -N deposition	21.4	0.0919	517.3
Total N deposition**	21.3	0.0781	329.2
NH ₄ -N concentration	20.0	0.0017	487.2
NO ₃ -N concentration	22.0	0.0016	544.5
NO ₃ -N + NH ₄ -N concentration	20.7	0.0008	454.5
Total N concentration**	20.6	0.0006	288.1

666 * AIC for total N deposition and precipitation cannot be compared to other AIC due to different number of data
 667 involved.

668 ** Finland, France, Germany and Slovenia only.

669

670 **Figure legends**

671

672 **Figure 1.** Sites where mosses and bulk precipitation were sampled for nitrogen analysis.

673


674 **Figure 2.** Deviation of the relationship between nitrogen concentration in paired moss species
675 from the 1:1 relationship (solid line). Paired moss species where sampled at the same sites in
676 one or more countries; n.s. = no significant difference between species.

677


678 **Figure 3.** Relationship between the deposition of different nitrogen forms in wet bulk
679 deposition (mean of 3 years of deposition) and the nitrogen concentration in mosses. Moss
680 and precipitation samples were collected less than 1 km apart in Austria (AT), Switzerland
681 (CH), Germany – Niedersachsen (DE-NI), Spain (ES), Finland (FI), France (FR) and Slovenia
682 (SI). Total wet bulk nitrogen deposition (i.e. including dissolved organic nitrogen) was only
683 determined in four countries (DE-NI, FI, FR, SI).

684

685 **Figure 4.** Relationship between the concentration of different nitrogen forms in precipitation
686 (mean of 3 years of deposition) and the nitrogen concentration in mosses. Moss and
687 precipitation samples were collected less than 1 km apart in Austria (AT), Switzerland (CH),
688 Germany – Niedersachsen (DE-NI), Spain (ES), Finland (FI), France (FR) and Slovenia (SI).
689 Total bulk nitrogen concentration (i.e. including dissolved organic nitrogen) was only
690 determined in four countries (DE-NI, FI, FR, SI).




691
 692 **Figure 1.**
 693




694
695
696

Figure 2.



697
698 **Figure 3.**



699
700 **Figure 4.**