

1 Lichen photobiont diversity and selectivity at the southern limit of the maritime Antarctic  
2 region (Coal Nunatak, Alexander Island)

3

4 Andreas Engelen<sup>1</sup>, Peter Convey<sup>2</sup>, Ovidiu Popa<sup>3</sup>, Sieglinde Ott<sup>1</sup>

5

6 <sup>1</sup> Institute of Botany, Heinrich-Heine-University, Universitaetsstr. 1, 40225 Duesseldorf,  
7 Germany

8 <sup>2</sup> British Antarctic Survey, Natural Environment Research Council, High Cross, Madingley  
9 Road, Cambridge CB3 0ET, England

10 <sup>3</sup> Genomic Microbiology Group, Institute of Microbiology, Christian-Albrechts-University  
11 Kiel, ZMB, Am Botanischen Garten 11, 24118 Kiel, Germany

12

13 Corresponding author: Sieglinde Ott

14 telephone: 0049 211 8113537

15 fax: 0049 211 8112881

16 e-mail: [otts@uni-duesseldorf.de](mailto:otts@uni-duesseldorf.de)

17

18 Running title:

19 Lichen photobiont diversity at Coal Nunatak, Antarctica

20

21

22 number of figures: 2

23 number of tables: 1

24

25 Abstract

26 Antarctic ice-free inland sites provide an unique perspective on the strategies coevolving  
27 organisms have developed for survival at the limits of life. Here, we provide the first combined  
28 description of the ecological and genetic diversity of lichen photobionts colonising an isolated  
29 Antarctic inland site, Coal Nunatak, on south-east Alexander Island (Antarctic Peninsula).  
30 Photobionts of 14 lichen species (42 samples) representing the entire lichen community of Coal  
31 Nunatak were investigated using the internal transcribed spacer region (ITS) of the nuclear  
32 ribosomal DNA. The study attempted to address the hypothesis that mycobiont selectivity for  
33 the photobiont partner is lower in more extreme environments. This hypothesis did not appear  
34 to hold true for the entire lichen community except one species. Another aspect focuses on the  
35 relevance of the reproduction modus concerning the distribution of photobiont haplotypes in  
36 the lichen community. Dispersal of generative mycobiont diaspores depends on lichenisation  
37 processes while by dispersal of vegetative diaspores both symbiotic bionts get dispersed.

38

39 Keywords: inland site, extreme environment, symbiotic association, community, genetic  
40 diversity, photobiont haplotypes

41

## 42 Introduction

43 Antarctica is the windiest, coldest and highest continent on Earth. Less than 0.5% of the  
44 Antarctic continent is permanently or seasonally free of ice cover (British Antarctic Survey,  
45 2004). Many lichens are successful colonisers of extreme environments, and can be found  
46 worldwide in deserts, high mountain ranges, tropical and polar regions. Lichens colonising  
47 Antarctic habitats are exposed to some of the most extreme environmental conditions faced in  
48 terrestrial environments on Earth (Peck *et al.*, 2006), including high levels of UV radiation,  
49 both extremely low and very variable temperatures, lack of liquid water and desiccation stress,  
50 and high wind speeds. With two flowering plants and approximately 50 liverworts and 104  
51 bryophytes known, the roughly 427 recorded lichen species form the dominant element in the  
52 diversity of the Antarctic flora (Ochyra, 1998; Bednarek-Ochyra *et al.*, 2000; Øvstedal &  
53 Smith, 2001; Convey, 2013).

54

55 The success of lichens under extreme environmental conditions is based on a remarkable  
56 symbiotic relation between, at least, two bionts. Approximately 21% of all fungi are known to  
57 form lichens (Hawksworth, 1988). In this obligate symbiosis the fungus (mycobiont) is  
58 associated with one or more photosynthesizing symbionts, the photobionts, which either can be  
59 eukaryotic green algae or prokaryotic cyanobacteria (Hawksworth, 1988). The green algae  
60 most commonly found as photobionts in lichens belong to the genus *Trebouxia* (Peveling,  
61 1988). Although between 14 000 and 20 000 lichen-forming mycobionts, mainly ascomycota,  
62 are estimated to exist (Feurerer & Hawksworth, 2007) they are associated with only a few  
63 different photobiont species (Tschermak-Woess, 1988).

64

65 To date, studies of the diversity of lichen photobionts in Antarctica have concentrated on  
66 coastal regions along the Antarctic Peninsula (Romeike *et al.*, 2002; Brinkmann, 2002;

67 Langohr, 2004; Siegesmund, 2005). Inland sites, where environmental conditions are generally  
68 more extreme and terrestrial diversity and levels of community development much lower  
69 (Convey & Smith, 1997) have received little attention (Neuburg, 2007; Pérez-Ortega *et al.*,  
70 2012). Until recently most studies have been carried out based on traditional taxonomic  
71 approaches. However, few studies have addressed the degree of selectivity between bionts and  
72 any correlation this may have with ecological factors in more extreme environments as typified  
73 by Antarctic inland sites (Pérez-Ortega *et al.*, 2012).

74

75 In this study diversity data are reported from Coal Nunatak (south-eastern Alexander Island,  
76 70°03'S 068°31'W), an inland nunatak ecosystem at the extreme southern limit of the maritime  
77 Antarctic. A very limited lichen and bryophyte flora is present with only 14 lichen species  
78 recorded. These 14 species were collected on Coal Nunatak, in order to investigate the  
79 hypothesis that mycobiont selectivity for the photobiont partner will be lower in more extreme  
80 environments (Romeike *et al.* 2002). The slow rates of community development at locations  
81 such as Coal Nunatak, which is characterized by very harsh environmental conditions, provide  
82 an opportunity to study lichen photobiont diversity. A second hypothesis relates to the  
83 reproductive tactics of the mycobionts, where it is postulated that the distribution of photobiont  
84 haplotypes is dependent on either the asexual or sexual reproduction of the mycobiont.

85

86 Coal Nunatak is located on south-eastern Alexander Island off the west coast of the Antarctic  
87 Peninsula. It is protected from the direct influence of the open sea, over 200 km due west in  
88 summer, by the high landmass of Alexander Island and the ice shelves that fringe its west coast,  
89 and by the permanent ice shelf that occupies George VI Sound to the east (6 km from the study  
90 site) and south (20 km from the study site). Located at the extreme southern limit of the  
91 maritime Antarctic, this region's climate is considered to be intermediate between that of the

92 more moist maritime region and the colder and drier continental zone (Smith, 1988; Convey &  
93 Smith, 1997). Coal Nunatak is snow-free during the Antarctic summer for approximately three  
94 months. The ecosystem at this site is characterised by its low developmental level (Brinkmann  
95 *et al.*, 2007; Engelen *et al.*, 2008). Small and often barely visible populations of different lichen  
96 species, occasionally associated with the few recorded bryophyte species, can be found in  
97 microniches restricted to rock surfaces and crevices and to the margins of soil polygons.

98

99 Reproductive tactics:

100 Lichens disperse over long distances by utilising two fundamentally different mechanisms.

101 There may be joint asexual dispersal of both symbionts in specific structures, either by means

102 of a vegetative thallus fragment or by specialized dispersal organs as soredia, which are small

103 (100 - 150  $\mu\text{m}$  in diameter) dispersal units that are produced in specialized cup-like structures

104 called soralia composed of both myco- and photobiont cells. These diaspores can easily be

105 distributed over long distances by wind at high altitudes. The sexual mechanism of lichen

106 dispersal involves the independent dispersal of the mycobiont (as ascospores) and the

107 photobiont (as vegetative cells). Both can grow individually in a new habitat, before coming

108 into contact through a recognition process and forming a new lichen thallus at that location *de*

109 *novo* by lichenisation (Ott, 1987). Dispersal of the bionts separately clearly requires the process

110 of relichenisation. Environmental conditions and physiological factors influence the success of

111 the recognition process (Meeßen & Ott, 2013; Meeßen *et al.*, 2013).

112

113 Selectivity and specificity:

114 The species diversity of lichen-forming fungi is much greater than that of the photobionts,

115 especially if only green-algal partners, that constitute the photobionts in the majority of lichens,

116 are considered. Algal lineages are widely shared among taxonomic mycobiont groups.

117  
118 Previous studies have demonstrated that mycobionts and photobionts cannot simply be  
119 combined randomly (Ahmadjian & Jacobs, 1981, 1982, 1983), indicating a degree of selectivity  
120 between the two bionts. Successful and complete lichenisation can only take place when both  
121 symbionts possess the appropriate adaptations (Schaper & Ott, 2003). The degree of specificity  
122 and selectivity of the mycobiont partner for particular photobionts varies between species.  
123 Rambold *et al.* (1998) defined specificity as the taxonomic range of photobionts associated  
124 with a mycobiont and selectivity as the exclusiveness with which specific photobionts are  
125 selected as partners. Galun & Bubrick (1984) defined ‘selectivity’ as the preferred interaction  
126 between two bionts, and ‘specificity’ as the exclusive interaction between photo- and  
127 mycobiont. Some fungi are only able to lichenise if a specific algal species is available (Galun,  
128 1988) while, in contrast, other species of fungi are able to form a lichen thallus with several  
129 members of the same genus of photobiont, and sometimes with partners related at an even  
130 higher systematic level (Piercey-Normore & DePriest, 2001; Helms *et al.*, 2001; Beck *et al.*,  
131 2002; Brinkmann, 2002; Romeike *et al.*, 2002). Symbiont selectivity and specificity are not  
132 only species-specific, but also can vary during the life-cycle of the partners and due to partner  
133 availability and environmental conditions. Graduated selectivity is expressed in the form of  
134 symbiotic contact achieved between myco- and photobiont. All stages, ranging from the  
135 intimate mutualistic contact of both symbionts in a well-developed lichen thallus to a loose-  
136 fitting parasitic contact, where the fungus penetrates the algal cells using haustoria and  
137 subsequently even kills the algae, are possible (Schaper & Ott, 2003).

138  
139 High levels of selectivity shown by a mycobiont are linked with a low diversity of suitable  
140 photobionts being present in a lichen genus as, for example, found in the family Cladoniaceae  
141 (Piercey-Normore & DePriest, 2001), the genus *Physcia* (Helms *et al.*, 2001) and the genus

142 *Letharia* (Kroken & Taylor, 2000). In contrast, a lower level of selectivity for photobionts has  
143 been reported in the Antarctic species *Umbilicaria antarctica* (Romeike *et al.*, 2002). The  
144 lower selectivity was interpreted as a form of flexibility and/or plasticity that acts as an  
145 adaptation to extreme environmental conditions. Yahr *et al.* (2004, 2006) concluded that, in  
146 particular *Cladonia* species, interactions were highly specific and that locally realized  
147 associations were probably influenced by environmental conditions. The more extreme lichen-  
148 dominated habitats become, the more it can be expected that environmental factors and  
149 community structure and composition will influence individual symbiotic interactions.

150

151 In this study photobiont diversity was assessed through sequencing of the ITS1, 5.8S and ITS2  
152 region of the ribosomal DNA (Friedl & Rokitta, 1997; Rambold *et al.*, 1998; Beck, 1999;  
153 Helms *et al.*, 2001; Kroken & Taylor, 2000; Piercey-Normore & DePriest, 2001; Romeike *et*  
154 *al.*, 2002; Fernandez-Mendoza *et al.*, 2011).

155

## 156 Material and Methods

### 157 Study site:

158 Coal Nunatak is located on south-east Alexander Island off the south-west coast of the  
159 Antarctic peninsula (70°03'S 068°31'W) (Fig. 1). The mountain ridge of the nunatak is about 4  
160 km, in a north-east to south-west orientation. Its ice free summit rises 380-424m above sea  
161 level. The research site was situated at the north-eastern end of the nunatak, covering  
162 approximately 3500 m<sup>2</sup>.

163

164 Coal Nunatak belongs to the Le May Group and is composed mainly of greywacke, a coarse  
165 grained sedimentary rock type (Burn, 1983). Surface geomorphology is characterised by  
166 extensive development of patterned ground and other typical periglacial features (e.g. frost-

167 sorted soil polygons, stone stripes) and bare rocks (Brinkmann *et al.*, 2007; Engelen *et al.*,  
168 2008).

169  
170 Coal Nunatak experiences a continental rather than a maritime climate. From March until mid-  
171 December the study site is covered by snow, becoming mostly snow-free during the short  
172 summer period from mid-December to early March. Terrestrial ecosystems at this site are at a  
173 very low or early stage of development. Much of the ground is barren to the naked eye, with  
174 colonisation by macroscopic vegetation restricted to small and generally sheltered micro-niches  
175 on rocks, in crevices, and on soil sheltered by rocks or associated with longer-lying snow  
176 patches. Investigations of the vegetation on the nunatak revealed a total of 14 lichen species  
177 and a small number of mosses are known from the nunatak (Brinkmann *et al.*, 2007; Engelen *et*  
178 *al.*, 2008).

179  
180 Lichen material:

181 Most lichen species were collected from the north-eastern part of the study area on the north-  
182 east of Coal Nunatak. *Xanthoria elegans* was obtained from the west exposed part of the study  
183 area. Three independent samples were obtained for each lichen species, with the quantities  
184 sampled being limited by the requirement not to damage the lichen community. After short-  
185 term storage at ambient conditions at the field site, lichen samples were transported to the  
186 British Antarctic Survey's Rothera Research Station (Adelaide Island). There the samples were  
187 stored at -20°C and returned frozen to the laboratory in Düsseldorf. Determination of the lichen  
188 species was carried out by taxonomic experts (H. Hertel, Munich; D. Øvstedal, Bergen; N.  
189 Wirtz, Frankfurt).

190  
191 Mycobionts included in the study:



192 The lichen species examined in this study are listed in Table 1. Seven species of the 14  
193 obtained from Coal Nunatak were epilithic (crustose lichens: *Tephromela disciformis*,  
194 *Tephromela atra*, *Caloplaca johnstonii*, *Lecidella pataviana*; macro lichens: *Usnea lambii*,  
195 *Pseudephebe minuscula*, *Xanthoria elegans*) and seven colonised soil-surface habitats (crustose  
196 lichens: *Buellia papillata*, *Candelariella flava*, *Caloplaca lewis-smithii*, *Lepraria cacuminum*,  
197 *Lepraria borealis*, *Ochrolechia frigida* and *Psoroma cf. tenue*). *Psoroma cf. tenue* was the only  
198 lichen colonising soil sites that has a well differentiated thallus.

199

200 Laboratory procedures:

201 Identification of the unicellular green algal photobionts using morphological and anatomical  
202 characters is known to be challenging. Therefore, we used a molecular approach to assess  
203 photobiont diversity. The nuclear internal transcribed spacer (ITS) region of the rDNA was  
204 analysed, including ITS1, ITS2 and the gene coding for the 5.8S ribosomal subunit. The region  
205 is located between the genes coding for the 18S and 26S ribosomal units in the ribosomal DNA  
206 tandem repeats and has been used routinely in molecular studies of green algal photobionts  
207 (Friedl & Rokitta, 1997; Rambold *et al.*, 1998; Beck, 1999; Helms *et al.*, 2001; Kroken &  
208 Taylor, 2000; Piercey-Normore & DePriest, 2001; Romeike *et al.*, 2002; Schaper & Ott, 2003;  
209 Yahr *et al.*, 2004; Yahr *et al.*, 2006).

210 To obtain photobiont DNA, conglomerates of photobiont cells were first carefully removed  
211 from the lichen thalli. This avoided the disruption of molecular procedures by secondary lichen  
212 metabolites such as phenolic substances. The clusters of photobiont cells were fragmented  
213 using liquid nitrogen and quartz sand. For DNA extraction the DNeasy Plant Mini Kit (Qiagen,  
214 Hilden, Germany) was used. After extraction the isolated DNA was stored at -20°C.

215

216 For a 25 µl PCR reaction, 2.5 µl template, 9 µl sterilized water, 12.5 µl HotStartTaq™ Master  
217 Mix (Qiagen) and 0.5 µl of each primer were used. The green alga specific primer with 5′-  
218 3′orientation is Al 1700f (Helms *et al.*, 2001). The primer used with 3′-5′orientation (LR3,  
219 <http://www.biology.duke.edu/fungi/mycolab/primers.htm>) is not specific for green algae (Freidl  
220 & Rokitta, 1997). For the amplification of the photobiont ITS-region a thermocycler (Biometra,  
221 Goettingen, Germany) was used as follows. The taq-polymerase was activated for one minute  
222 95°C. The DNA was denatured for one minute at 94°C. The annealing temperature of the  
223 primers was set to 53°C for one minute. The elongation of the annealed primers by taq-  
224 polymerase took place for 1.5 minutes at 72°C. The denaturation, annealing and elongation  
225 steps were repeated 35 times, after which the final extension of partially elongated products  
226 took 10 minutes at a temperature of 72°C. After final extension the PCR product was cooled at  
227 4°C. The amplified PCR products were purified using the QIAquick PCR Purification Kit  
228 (Qiagen, Hilden, Germany).

229

230 DNA sequencing was carried out by GATC-Biotech (Konstanz, Germany) using an ABI 3730  
231 XL Sequencer. Non algal specific primers used for sequencing were 1800f (5′-3′orientation)  
232 (Friedl, 1996) and ITS4 (3′-5′orientation) (White *et al.*, 1990). The resulting ITS rDNA  
233 sequences were edited using the application 'Bioedit for Windows'  
234 (<http://www.mbio.ncsu.edu/bioedit/bioedit.html>). NCBI-BLAST searches of GenBank records  
235 were performed to confirm that the amplified and sequenced DNA fragments originated from  
236 the photobiont and to identify the taxonomic classification of the closest hit.

237 The alignment of all sequences was carried out using the online application MAFFT version 7  
238 (<http://mafft.cbrc.jp/alignment/server/>) based on the HKY substitution model (Hasegawa *et al.*,  
239 1985). The calculation of a phylogenetic maximum likelihood tree using PhyML 3.0 (Guindon  
240 *et al.*, 2010) was supported by 1000 bootstrap steps. All sequences obtained from samples of

241 lichen photobionts from Coal Nunatak have been added to the database of the National Center  
242 for Biotechnology Information (accession numbers: FJ426284 - FJ426299).

243

## 244 Results

245 In 14 lichen species of 11 genera from Coal Nunatak we found seven different haplotypes of  
246 the genus *Trebouxia* and one haplotype of the genus *Asterochloris*. Sexual reproduction was  
247 noted only in crustose species, with six lichens producing fruiting bodies (Table 1). The  
248 reproduction of these lichen species using ascospores in Antarctic habitats has previously been  
249 noted by Øvstedal & Smith (2001). In most of the lichens included in this study the mycobiont  
250 was associated with green algal photobionts, with the exception of *Psoroma cf. tenue*. This  
251 lichen species also forms thallus structures (cephalodia) with cyanobacteria of the genus  
252 *Nostoc*. The mycobiont of this lichen is, therefore, associated with both a green algal species  
253 and a cyanobacterial species.

254

255 In the lichen species examined, haplotype 8 was found as photobiont of three lichen species:  
256 *Lepraria borealis*, *Usnea lambii* and *Pseudephebe minuscula*. *Trebouxia* haplotype 7 was the  
257 dominant haplotype in the samples investigated at this locality. Six species (*Lecidella*  
258 *pataviana*, *Lepraria borealis*, *Lepraria cacuminum*, *Tephromela atra*, *Tephromela disciformis*  
259 and *Xanthoria elegans*) were associated with this photobiont. Several haplotypes were  
260 restricted to a single lichen species, including haplotypes 2 (*Buellia papillata*), 3 (*Candelariella*  
261 *flava*), 4 (*Psoroma cf. tenue*), 5 (*Caloplaca lewis-smithii*) and 6 (*Caloplaca johnstonii*). The  
262 algal genus *Asterochloris* (haplotype 1) was found as photobiont in the two lichen species  
263 *Lepraria borealis* and *Ochrolechia frigida* (Table 1).

264

265 Thirteen of the 14 mycobiont species from Coal Nunatak were associated with only a single  
266 algal haplotype, the exception being *L. borealis*. This species contained three different  
267 photobiont haplotypes 1, 7 and 8 (Table 1). The ITS rDNA sequences of the photobionts  
268 detected in *L. borealis* were identical to the photobionts found in lichen species that were  
269 colonized by thalli of *L. borealis* (Engelen *et al.*, 2010). Respectively, *L. borealis* when  
270 growing in close association with *O. frigida* contained haplotype 1, in association with *T.*  
271 *disciformis* haplotype 7 and in association with *U. lambii* haplotype 8 (Engelen *et al.*, 2010).  
272

273 The phylogenetic tree showed three highly supported basal clades (Fig. 2). The first clade as  
274 outgroup only consisted of two sequences of haplotype 1. These sequences closest similarities  
275 in a BLAST search were to the genus *Asterochloris*. The sequences of the second clade  
276 belonging to haplotype 8 showed highest BLAST hits with taxa identified as *Trebouxia*  
277 *jamesii*. The third clade consisted of two subclades. One included haplotypes 6 and 7, and had  
278 highest BLAST similarities with taxa also identified as *T. jamesii*, and the other consisted of  
279 haplotypes 2 to 5, which were most similar to *Trebouxia impressa*. Most substitutions were  
280 found in the subclade consisting of haplotypes 2 to 5 (*T. impressa*). The other clades, involving  
281 the two different groups of *Trebouxia jamesii* and one group of *Asterochloris* sp., were  
282 composed of almost identical sequences within each clade.  
283

284 With the exception of *Lepraria borealis*, the photobiont clades were correlated to groups of  
285 lichens characterised by sharing particular morphological and ecological features. The clade  
286 consisting of the photobiont haplotype 8 (*T. jamesii*) was associated with fruticose lichens  
287 growing on rocks, whereas photobionts of the clade consisting of haplotypes 6 and 7 (*T.*  
288 *jamesii*) were found in crustose lichens on rocks. Crustose lichens growing on soil and/or  
289 mosses were either associated with photobionts of the clade consisting of haplotypes 2-5 (*T.*

290 *impressa*) or with the photobiont clade of haplotype 1 (*Asterochloris*). Photobionts of *L.*  
291 *borealis* were present in all clades with the exception of the *T. impressa* group.

292

## 293 Discussion

294 This study is the first to document lichen photobiont diversity in the southern region of the  
295 Antarctic Peninsula. Molecular studies on photobionts have shown that identical haplotypes are  
296 widespread and can be found across geographic regions (Kroken & Taylor, 2000; Yahr *et al.*,  
297 2004; Yahr *et al.*, 2006), continents (Piercey-Normore & DePriest, 2001) and even  
298 hemispheres. *Trebouxia jamesii* identified here at Coal Nunatak, has been described from a  
299 range of other localities in the maritime and continental Antarctic (Romeike *et al.*, 2002; Pérez-  
300 Ortega *et al.*, 2012) as well as from Europe (Beck, 1999), supporting effective dispersal across  
301 distances up to intercontinental and global scales.

302

303 Patterns of haplotype distribution being unique to specific habitats suggest a process of local  
304 adaptation or might be pre-adapted to local environmental conditions and communities (Pérez-  
305 Ortega *et al.*, 2012). Such an interpretation is in line with the conclusions drawn by Yahr *et al.*  
306 (2004) in an extensive community study, who found a homogeneous photobiont pool across  
307 geographic distances and proposed that local adaptation of the photobiont was of importance at  
308 some sites.

309

310 All localities analyzed by Yahr *et al.* (2004) shared the same habitat and vegetation type,  
311 reflected by the occurrence of similar *Cladonia sp.* communities. Such a sampling design  
312 enhances the detection of environmental selection acting in photobiont lineages that might  
313 cause ecological specialization.

314

315 The ITS sequence of photobiont haplotype 8 found on Coal Nunatak in this study was also  
316 found in lichens from the maritime Antarctic sites Lagoon Island, Rothera Point and Charcot  
317 Island by Romeike *et al.* (2002). It is also identical to the sequence of *T. jamesii* cultured from  
318 *Lecidea silacea*, collected from siliceous and heavy-metal containing rocks at localities in  
319 Austria (Beck, 1999). It seems that haplotype 8 (*T. jamesii*) is a generalist with a bi-  
320 hemispherical distribution that occurs ranging from extreme habitats at the limits of vegetation  
321 to more moderate maritime and polar habitats to comparably benign temperate localities.

322

323 Romeike *et al.* (2002) noted that this haplotype has only been described to date from iron-rich  
324 sites (Beck, 1999). However, the study on Coal Nunatak does not have atypically high iron  
325 concentration. Haplotype 8 was the second most abundant photobiont at Coal Nunatak.

326

327 At Coal Nunatak photobionts are associated with a relatively low number of mycobionts.

328 Haplotype 7 was the most abundant photobiont amongst 14 lichen species recorded on Coal

329 Nunatak, being present in 6 different species while haplotypes 1-6 and 8 were distributed

330 amongst 9 lichen species (Table 1). This is suggestive of selectivity and specificity of the

331 symbionts of the respective lichen species. Eight different photobiont haplotypes were found in

332 the 14 lichen species (Table 1). With one exception *Lepraria borealis* (Engelen *et al.*, 2010) all

333 mycobionts were associated with a single photobiont haplotype.

334 This might be due to the harsh environmental conditions and the overall short growing season

335 at the inland site that limits photosynthetic activity and thus photobiont productivity. In such a

336 life-averse habitat which effects a limited primary production (Sadowsky & Ott, 2012)

337 symbiont interactions can be expected to be fine-tuned for the holobiont to survive and to

338 successfully colonise and keep the habitat.

339

340 To form an own thallus *L. borealis* takes over the photobiont haplotypes 1, 7 and 8. Our data  
341 suggest that the species may be able to obtain these photobionts from physically adjacent thalli  
342 of other lichen species, such as *Ochrolechia frigida* (Pertusiales), *Tephromela disciformis* and  
343 *Usnea lambii* (Lecanorales s. str.) (Table 1). When growing close to *T. disciformis* or *U. lambii*,  
344 *L. borealis* incorporated the identical *Trebouxia jamesii* haplotypes 7 and 8 as the photobionts  
345 of these immediately adjacent lichens. Similarly, when growing in association with  
346 *Ochrolechia frigida* the same *Asterochloris* haplotype 1 was present in both lichens (Engelen *et*  
347 *al.*, 2010). Thus, only the *L. borealis* mycobiont shows low selectivity towards potential  
348 photobionts consistent with the prediction of Romeike *et al.* (2002).

349

350 The diversity of the photobiont haplotypes within the lichen community may be influenced by  
351 the mechanism of reproduction. Sexual reproduction requires a lichenisation process during  
352 which the mycobiont must encounter a suitable algal partner amongst those available in its  
353 immediate vicinity. The data obtained in the current study give no indication of any difference  
354 in diversity of photobiont haplotypes between the mycobionts reproducing asexually or  
355 sexually (Table. 1).

356

357 For the degree of selectivity of the mycobiont to the photobiont partner characteristic features  
358 of the respective lichen symbiosis may be primarily responsible at extreme habitats.

359 Environmental conditions might also effect the degree of selectivity (Romeike *et al.* 2002).

360 However, based on the results presented the potential of the symbiotic state of lichens seems to  
361 be the more relevant factor considering the success of colonisation processes particularly at  
362 extreme environments.

363

364

365 Acknowledgements

366 We thank the British Antarctic Survey for logistic support allowing access to the study sites on  
367 Alexander Island and its staff at Rothera Research Station for their support. We are especially  
368 thankful to the BAS field assistants Neil Stevenson and Robin Jarvis for their kind and  
369 invaluable technical support in the field. Thanks are due to Nora Wirtz, Dag Øvstedal and  
370 Hannes Hertel for determination of the lichen species. This project was funded by a grant of the  
371 Deutsche Forschungsgemeinschaft (DFG) to SO (Ot96/10-1/2) as part of the priority program  
372 SPP 1158 and the Düsseldorf Entrepreneurs Foundation. PC is supported by NERC core  
373 funding to the BAS Ecosystems programme. This paper also forms an output of the SCAR  
374 AntEco and AnT-ERA scientific programmes.

375

376

377 References

378

379 Ahmadjian V & Jacobs JB (1981) Relationship between fungus and alga in the lichen *Cladonia*  
380 *crustatella* Tuck.. *Nature* 289: 169-172.

381

382 Ahmadjian V & Jacobs JB (1982) Artificial reestablishment of lichens. III. Synthetic  
383 development of *Usnea strigosa*. *Journal of Hattori Botanical Laboratory* 52: 393-399.

384

385 Ahmadjian V & Jacobs JB (1983) Algal-fungal relationships in lichens: recognition, synthesis,  
386 and development. *Algal Symbiosis* (Goff LJ, ed), pp. 147-172. Cambridge University Press,  
387 Cambridge, UK.

388



389 Beck A (1999) Photobiont inventory of a lichen community growing on heavy-metal-rich rock..  
390 *Lichenologist* 31: 501-510.  
391

392 Beck A, Kasalicky G & Rambold G (2002) Myco-photobiontal selection in a Mediterranean  
393 cryptogam community with *Fulgensia fulgida*. *New Phytol* 153: 317-326.  
394

395 Bednark-Ochyra H, Vána J, Ochyra L & Smith RIL (2000) *The liverwort flora of Antarctica*.  
396 Polish Academy of Sciences, Cracow.  
397

398 Brinkmann M (2002) Genetische Diversität von Bionten in der Flechtensymbiose. Diploma  
399 thesis, University of Düsseldorf, Germany.  
400

401 Brinkmann M, Pearce DA, Convey P & Ott S (2007) The cyanobacterial community of  
402 polygon soils at an inland Antarctic nunatak. *Polar Biol* 30: 1505-1511.  
403

404 British Antarctic Survey (2004) *Antarctica, 1:10 000 000*  
405 *scale map. BAS (Misc) 11*. British Antarctic Survey, Cambridge.  
406

407 Burn RW (1983) Geology of the Le May Group, Alexander Island. *British Antarctic Survey*  
408 *Scientific Report* 109: 1-64.  
409

410 Convey P & Smith RIL (1997) The terrestrial arthropod fauna and its habitats in northern  
411 Marguerite Bay and Alexander Island, maritime Antarctic. *Antarct sci* 9: 12-26.  
412

413 Engelen A, Convey P, Hodgson DA, Worland MR & Ott S (2008) Soil properties of an  
414 Antarctic inland site: implications for ecosystem development. *Polar Biol* 31: 1453-1460.  
415

416 Engelen A, Convey P & Ott S (2010) Life history strategy of *Lepraria borealis* at an Antarctic  
417 inland site, Coal Nunatak. *Lichenologist* 42(3): 339-346.  
418

419 Fernández-Mendoza F, Domaschke S, García MÁ, Jordan P, Martín M & Printzen C (2011)  
420 Population structure of mycobionts and photobionts of the widespread lichen *Cetraria aculeata*.  
421 *Mol Ecol* 20: 1208-1232.  
422

423 Feuerer T & Hawksworth DL (2007) Biodiversity of lichens, including a world-wide analysis  
424 of checklist data based on Takhtajan's floristic regions. *Biodivers Conserv* 16: 85-98.  
425

426 Friedl T & Rokitta C (1997) Species relationships in the lichen genus *Trebouxia* (Chlorophyta,  
427 Trebouxiphyceae): molecular phylogenetic analyses of nuclear-encoded large subunit rRNA  
428 gene sequences. *Symbiosis* 23: 125-148.  
429

430 Friedl T (1996) Evolution of the polyphyletic genus *Pleurastrum* (Chlorophyta): inferences  
431 from nuclear-encoded DNA Sequences and motile cell ultrastructure. *Phycologia* 35: 456-469.  
432

433 Galun M (1988) Lichenization. *Handbook of Lichenology II* (Galun M, ed.), pp. 153-169. CRC  
434 Press, Boca Raton, Florida.  
435

436 Galun M & Bubrick P (1984) Physiological interactions between the partners of the lichen  
437 symbiosis. *Cellular Interactions* (Linskens HF, Heslop-Harrison J, eds), pp. 362-401.  
438 Encyclopedia of Plant Physiology, Berlin, Germany.  
439

440 Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W & Gascuel O (2010) New  
441 algorithms and methods to estimate maximum-likelihood phylogenies: assessing the  
442 performance of PhyML 3.0. *Syst Biol* 59: 307-321.  
443

444 Hasegawa M, Kishino H & Yano TA (1985) Dating of the human-ape splitting by a molecular  
445 clock of mitochondrial DNA. *J Mol Evol* 22: 160-174.  
446

447 Hawksworth DL (1988) The variety of fungal-algal symbioses, their evolutionary significance,  
448 and the nature of lichens. *Bot J Linn Soc* 96: 3-20.  
449

450 Helms G, Friedl T, Rambold G & Mayrhofer H (2001) Identification of photobionts from  
451 lichen family Physciaceae using algal-specific ITS rDNA sequencing. *Lichenologist* 33: 73-86.  
452

453 Kroken S & Taylor JW (2000) Phylogenetic Species, reproductive Mode, and Specificity of the  
454 Green Alga *Trebouxia* forming Lichens with the Fungal Genus *Letharia*. *Bryologist* 103: 645-  
455 650.  
456

457 Langohr J (2004) Genetische Diversität der Photobionten der Gattung *Usnea* subgenus  
458 *Neuropogon* in der Antarktis. Diploma thesis, University of Düsseldorf, Germany.  
459

460 Meeßen & Ott S (2013) Recognition mechanisms during the pre-contact state of lichens: I.  
461 Mycobiont-photobiont interactions of the mycobiont of *Fulgensia bracteata*. *Symbiosis* 59(3):  
462 121-130.  
463

464 Meeßen J, Eppenstein S & Ott S (2013) Recognition mechanisms during the pre-contact state  
465 of lichens: II. Influence of algal exudates and ribitol on the response of the mycobiont of  
466 *Fulgensia bracteata*. *Symbiosis* 59(3): 131-143  
467 .

468 Neuburg M (2007) Genetische Diversität von Photobionten aus Flechten der kontinentalen  
469 Antarktis. Diploma thesis, University of Duesseldorf, Germany.  
470

471 Ochyra R (1998) *The moss flora of King George Island, Antarctica*. Polish Academy of  
472 Sciences, Cracow.  
473

474 Ott S (1987) Reproductive strategies in lichens. *Bibliotheca Lichenologica* 25: 81-93.  
475

476 Øvstedal DO & Smith RIL (2001) *Lichens of Antarctica and South Georgia. A guide to their*  
477 *identification and ecology*. Cambridge University Press, Cambridge.  
478

479 Peck LS, Convey P & Barnes DKA (2006) Environmental constraints on life histories in  
480 Antarctic ecosystems: tempos, timings and predictability. *Biol Rev* 81: 75-109.  
481

482 Pérez-Ortega S, Ortiz-Álvarez R, Allan Green TG & de Los Ríos A (2012) Lichen myco- and  
483 photobiont diversity and their relationships at the edge of life (McMurdo Dry Valleys,  
484 Antarctica). *FEMS Microbiol Ecol* 82(2): 429-48.

485

486 Peveling E (1988) Beziehungen zwischen den Symbiosepartnern in Flechten.  
487 *Naturwissenschaften* 75: 77-86.

488

489 Piercey-Normore MD & DePriest PT (2001) Algal switching among lichen symbioses. *Am J*  
490 *Bot* 8: 1490-1498.

491

492 Rambold G, Friedl T & Beck A (1998) Photobionts in lichens: possible indicators of  
493 phylogenetic relationships? *Bryologist* 101: 392-397.

494

495 Romeike J, Friedl T, Helms G & Ott S (2002) Genetic diversity of algal and fungal partners in  
496 four species of *Umbilicaria* (Lichenized Ascomycetes) along a transect of the Antarctic  
497 Peninsula. *Mol Biol Evol* 19: 1209-1217.

498

499 Sadowsky A & Ott S (2012) Photosynthetic Symbionts in Antarctic Terrestrial Ecosystems:  
500 The Physiological Response of Lichen Photobionts to Drought and Cold. *Symbiosis* 58: 81-90.

501

502 Schaper T & Ott S (2003) Photobiont Selectivity and Interspecific Interactions in Lichen  
503 Communities. I. Culture Experiments with the mycobiont *Fulgensia bracteata*. *Plant Biol* 5:  
504 441-450.

505

506 Siegesmund M (2005) *Phylogenetische Analyse der evolutionären Beziehungen von*  
507 *Flechtensymbionten in der Antarktis*. Diploma thesis, University of Düsseldorf, Germany.

508

509 Smith RIL (1988) Bryophyte oases in ablation valleys on Alexander Island, Antarctica.  
510 *Bryologist* 91: 45-50.  
511  
512 Tschermak-Woess E (1988) The algal partner. *CRC Handbook of Lichenology I* (Galun M,  
513 ed.), pp. 39-92 . CRC Press, Boca Ration, Florida.  
514  
515 White TJ, Burns T, Lee S & Taylor J (1990) Amplification and direct sequencing of fungal  
516 ribosomal RNA genes for phylogenetics. *PCR Protocols. A Guide to Methods and Applications*  
517 (Innis M, Gelfand D, Sninsky J, White T, Orlando FL, eds.), pp. 315-322. Academic Press,  
518 London, UK.  
519  
520 Yahr R, Vilgalys R & DePriest PT (2004) Strong fungal specificity and selectivity for algal  
521 symbionts in Florida scrub *Cladonia* lichens. *Mol Ecol* 13: 3367-3378.  
522  
523 Yahr R, Vilgalys R & DePriest PT (2006) Geographic variation in algal partners of *Cladonia*  
524 *subtenuis* (Cladoniaceae) highlights the dynamic nature of a lichen symbiosis. *New Phytol* 171:  
525 847-860.  
526

527 Tab. 1: Reproductive mode and distribution of photobiont haplotypes in lichen species found  
 528 on Coal Nunatak.

529 \* haplotype 1: genus *Asterochloris*; haplotypes 2-8: genus *Trebouxia*

530

531

532

533

534

lichen species	reproductive mode		haplotypes*							
	sexual	asexual	1	2	3	4	5	6	7	8
<i>Buellia papillata</i>	X			X						
<i>Caloplaca johnstonii</i>	X								X	
<i>Caloplaca lewis-smithii</i>	X							X		
<i>Candelariella flava</i>		X			X					
<i>Lecidella pataviana</i>	X									X
<i>Lepraria borealis</i>		X	X						X	X
<i>Lepraria cacuminum</i>		X							X	
<i>Ochrolechia frigida</i>		X	X							
<i>Pseudephebe minuscula</i>		X								X
<i>Psoroma cf. tenue</i>	X					X				
<i>Tephromela atra</i>	X									X
<i>Tephromela disciformis</i>		X								X
<i>Usnea lambii</i>		X								X
<i>Xanthoria elegans</i>		X								X

544

545 Figure captions

546

547 Fig. 1: a: arrow=location of Coal Nunatak on Alexander Island (70°03'S 68°31'W). b:

548 circle=location of the research area on Coal Nunatak.

549

550 Fig. 2: ML-tree of the *Asterochloris* haplotype and the 7 *Trebouxia* haplotypes as shown in

551 Tab. 1.

552 The photobiont sequences are named as follows: abbreviation of the photobiont\_abbreviation

553 of the mycobiont\_number of the haplotype as in Tab.1

554

555 Abbreviations of the mycobiont: Bupa: *Buellia papillata*, Cafl: *Candelariella flava*, Cajo:

556 *Caloplaca johnstonii*, Cale: *Caloplaca lewis-smithii*, Lebo: *Lepraria borealis*, Leca: *Lepraria*

557 *cacuminum*, Lepa: *Lecidella pataviana*, Ocfr: *Ochrolechia frigida*, Psmi: *Pseudephebe*


558 *minuscula*, Pste: *Psoroma cf. tenue*, Teat: *Tephromela atra*, Tedi: *Tephromela disciformis*,


559 Usla: *Usnea lambii*, Xael: *Xanthoria elegans*

560 Abbreviations of the photobionts: Ast: *Asterochloris spec.*, Tja: *Trebouxia jamesii*, Tim:

561 *Trebouxia impressa*







1000

Ast\_Ocfr\_1

Ast\_Lebo\_1

867

Tja\_Lebo\_8

280

Tja\_Psmi\_8

Tja\_Usla\_8

1000

1000

Tim\_Bupa\_2

Tim\_Cale\_5

592

Tim\_Cafl\_3

703

863

Tim\_Pste\_4

990

Tja\_Tedi\_7

Tja\_Lebo\_7

115

Tja\_Xael\_7

45

Tja\_Teat\_7

33

Tja\_Leca\_7

41

Tja\_Cajo\_6

89

FTja\_Lepa\_7



0.06