

Article (refereed) - postprint

Vane, C.H.; Kim, A.W.; Cave, M.R.; Li, N.. 2014 Equal abundance of odd and even n-alkanes from cycad leaves: can the carbon preference index (CPI) faithfully record terrestrial organic matter input at low latitudes? *Encephalartos* (118).

Copyright © 2014 Cycad Society of South Africa

This version available <http://nora.nerc.ac.uk/509715/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

*This is a pre-copy-editing, author-produced PDF of an article accepted for publication in *Encephalartos* following peer review. The definitive publisher-authenticated version is available online at http://cycadsociety.org/?page_id=623*

Contact BGS NORA team at
norabgs@bgs.ac.uk

Equal abundance of odd and even *n*-alkanes from cycad leaves: Can the carbon preference index (CPI) faithfully record terrestrial organic matter input at low latitudes?

C.H. Vane^{1*}, A.W. Kim¹, M.R. Cave¹ and N. Li²

¹British Geological Survey, Environmental Science Centre, Keyworth, Nottingham, NG12 5GG, Tele.: +44(0)115 936 3017: Email: chv@bgs.ac.uk

²Fairy Lake Botanic Garden, Xianhu Road 160, Liatang Shenzhen, Guangdong, 518004, China

Abstract

Long chain *n*-alkanes from the leaves of thirteen extant cycad species within the Cycadaceae, Stangeriaceae, Zamiaceae families were measured by gas chromatography-mass spectrometry (GC-MS). Cycad *n*-alkane patterns ranged from nC_{10} - nC_{37} , were unimodal in distribution, maximised at nC_{27} - nC_{33} and gave average chain length (ACL) values of 23.9 to 31.0. Low carbon preference indices (CPI) in the range of 0.88 to 2.70 were observed which is atypical of lipids from the leaves of terrestrial plants. Analysis of variance between the three families of true cycads showed that there were no significant differences between CPI values. The unusually low CPI values (<1.5) in nine of the thirteen cycad species analysed suggests that caution needs to be exercised in the use of *n*-alkanes distributions as a chemical marker of terrestrial plant input in sediments from tropical and sub-tropical regions.

1. Introduction

Long chain *n*-alkanes from the cuticular waxes of higher plant leaves are characterised by a strong odd predominance, carbon preference index (CPI) values ranging from ~3 to 34 and average chain lengths (ACL) of 26 to 30 (Eglinton & Hamilton, 1967; Rommerskirchen *et al.*, 2006; Stojanovic *et al.*, 2003). In combination, the *n*-alkane distribution pattern and dominance of *n*C₂₇, *n*C₂₉ and *n*C₃₁ over lower molecular weight homologues *n*C₁₅, *n*C₁₇ and *n*C₁₉ have been widely applied as a proxy for terrestrial land plant input to lake, river, estuarine and coastal sediments (Jaffé *et al.*, 2001; Meyers, 2003; Wang *et al.*, 2003). Exceptions to the odd dominated *n*-alkane pattern from higher plants have been shown and on at least one occasion re-examined and corrected (Reddy *et al.*, 2000).

Investigations of the solvent soluble lipid content of Cycadophyta the ancient seed bearing plants found in tropical and subtropical regions reported that *n*-alkanes from the Zamiaceae family varied in a non-systematic manner with respect to genus (Osborne *et al.*, 1993). Epicuticular leaf waxes from the *Ceratozamia* and *Encephalartos* genera ranged from *n*C₁₆ to *n*C₃₃ with maxima at *n*C₁₈ to *n*C₁₉. The same study reported that certain species *n*-alkane distributions were characterised by equal amounts of odd and even alkanes, as well as an absence of significant concentrations of branched chain isomers (Osborne *et al.*, 1993). The aim of this study was to examine the *n*-alkane distribution of Cycad foliage in order to improve our understanding of the utility of the *n*-alkane distributions and associated ratios in ancient sediments.

2. Methods

2.1 Samples and extraction

Fresh leaf material was obtained in October, 2006 from Fairy Lake Botanic Gardens, Shenzhen, China. Leaves were washed with 18 MΩ distilled water, freeze

dried and ground using a freezer mill (SPEX CertiPrep 6850). Powdered leaves ~2.5 g were extracted in an accelerated solvent extraction system ASE 200 (Dionex) using 60 ml dichloromethane (DCM). Extracts were evaporated under a stream of dry nitrogen gas to a constant weight and reconstituted in 3 ml DCM.

2.2 Fractionation and Instrumental Analysis

Lipids were isolated by chromatography on columns containing 6 g Silica (deactivated) eluted with 40 ml *n*-hexane. The hydrocarbons were further fractionated into adduct and non-adduct fractions by urea adduction. Adduct fractions were analysed using a Varian CP3800 series gas chromatograph (GC) directly coupled with a Varian 1200L triple Quadrupole MS/MS system (GC/MS). Peak assignments were made by comparison with mass spectra and retention times of authentic standard compounds.

2. Results and discussion




Figure 1. Gas chromatograms of the saturated hydrocarbon fraction extracted from cycad (*Ceratozamia*, *Dioon*, *Macrozamia*) and tree fern (*Brainea*) leaves.

Alkanes ranging from nC_{10} to nC_{37} were identified in the leaves of 16 specimens from six plant families. For all specimens the major n -alkanes were C_{27} , C_{29} , C_{31} and C_{33} in contrast n -alkanes in the range C_{16} to C_{26} were mainly present at lower relative abundances (Fig. 1). Total amounts of n -alkanes ranged from 0.55 $\mu\text{g/g}$ dry wt for *Cycas panzhihuaensis* to 6.8 $\mu\text{g/g}$ dry wt for *Encephalartos gratus* (Table 1). The average chain length (ACL) ranged from 23.9 for *Cycas panzhihuaensis* to a maximum of 31.0 for *Encephalartos gratus*, confirming the notion that cycad leaves contain a broad range of n -alkanes (Table 1). By contrast long chain n -alkane distributions for non-cycads, namely Equisetacea and Blechnaceae plant families, gave a limited distribution range dominated by C_{27} , C_{28} , C_{29} (Fig 1; Table 1). The carbon preference index (CPI) of the true cycads (Cycadaceae, Stangeriaceae, Zamiaceae) ranged from as low as 0.88 in *Cycas panzhihuaensis* to 2.70 for *Zamia fischeri* Miq. The slight predominance of odd over even n -alkanes was confirmed by the mean CPI of the 12 cycad leaf specimens of 1.47. The cycad CPI values presented here are considerably lower than those usually reported for higher plant sourced n -alkanes which typically vary from about 5 to 30. An earlier study of n -alkane leaf waxes of 28 cycad species from Mexico, Florida and Australia reported that *Macrozamia moorei*, *Zamia fischeri*, *Zamia furfuracea* and *Ceratozamia robusta* yielded odd over even ratios of ~ 1.5 , unimodal in distributions which were maximal at nC_{17} to nC_{21} and showed no significant quantities of branched chain isomers (Osborne *et al.*, 1993). In this current study we report broadly similar CPI values of 1.21 *Macrozamia moorei*, 1.30 *Ceratozamia robusta*, 1.32 *Zamia furfuracea*, but higher CPI values for the cycad *Zamia fischeri* at 2.70 (Table 1). Overall results differ from that of the earlier study of Osborne *et al* (1993) in that here the cycad n -alkanes were generally maximal at C_{29} to C_{31} . Furthermore, branched chain isomers were such a significant component of the

solvent extracts that urea-adduction was required to remove these components in order to achieve baseline separation and facilitating quantification of individual *n*-alkane homologues. One plausible explanation for the difference in *n*-alkane distributions between this and previous studies of the same plant species could be that we analysed the whole leaf as compared to only extracting the surface epicuticular waxes of the leaves. Alternatively it has been speculated that the method of cultivation can modify epicuticular leaf wax *n*-alkane distributions, although this hypothesis remains untested (Osborne *et al.*, 1993).

Table 1. Summary of *n*-alkane data for cycad, horsetail and tree fern leaves

Family	Genus and Species	Common Name	Conc. (µg/g) ^a	CPI ^b	ACL ^c	Odd Max	Even Max
Equisetaceae	<i>Equisetum arvenese</i>	Horsetail	2.21	4.90	28.4	29	30
Blechnaceae	<i>Brainea insignis</i> (Hook.) J.Sm	Cycad Fern	2.47	11.90	28.3	29	28
Cycadaceae	<i>Cycas miquelii</i> Warburg	Unknown	2.16	1.27	28.0	27	28
	<i>Cycas panzhihuaensis</i> L.Zhou & S.Y.Yang	Unknown	0.55	0.88	23.9	27	20
	<i>Cycas revoluta</i> Thunb	Sago Cycad	3.67	1.30	25.3	27	28
	<i>Cycas thouarsii</i> Lehm	Unknown	0.70	1.91	26.7	27	26
Oleandraceae	<i>Nephrolepis auriculata</i> (L.) Trim	Sword Fern	1.22	2.39	28.1	27	28
Stangeriaceae	<i>Stangeria eriopus</i> (Kunze) Baillon	Natal Grass Cycad	2.22	1.69	27.8	31	28
	<i>Bowenia spectabilis</i> Hook.ex J.D.Hook	Unknown	0.93	1.34	26.8	29	28
Zamiaceae	<i>Dioon spinulosum</i> Dyer	Mexican Sago Palm	1.52	1.00	29.4	33	32
	<i>Encephalartos gratus</i> Prain	Unknown	6.80	1.21	31.0	33	32
	<i>Lepidozamia peroffskyana</i> Regel	Pineapple Zamia	3.06	1.88	27.5	29	28
	<i>Macrozamia moorei</i> F.Muell	Unknown	3.79	1.21	29.1	31	30
	<i>Zamia fischeri</i> Miq.	Unknown	2.28	2.70	29.0	31	30
	<i>Zamia furfuracea</i> L.f.	Cardboard Palm	0.57	1.32	27.2	31	30
	<i>Ceratozamia robusta</i> Miq. Belize	Bamboo Cycad	3.67	1.30	28.1	31	30

^aTotal concentration of *n*-alkanes (*n*-C₁₀ to *n*-C₃₆) in µg/g of dry leaf weight.

^bCPI (carbon preference index)=(Σodd C from *n*-C₁₀ to *n*-C₃₆)/ (Σeven C from *n*-C₁₀ to *n*-C₃₆)

^cACL (average carbon chain length)=(Σ%C_n × n)/ 100

The differences between the means of the CPI data (Table 1) for the three true cycad families Cycadaceae, Stangeriaceae, Zamiaceae were analysed using one-way analysis of variance (ANOVA) in the R statistical programming language (R Development Core Team (2008)). Figure 2 shows a comparative box plot of the CPI values for the three cycad families showing all three have similar median values and

interquartile ranges with a possible outlier (*Zamia fischeri* Miq.) in the Zamiaceae family. The analysis of variance gave a non significant p-value of 0.8531 showing that there are no significant differences between the CPI values for the three families. As an additional check, a non-parametric equivalent method to the ANOVA, was also applied (Kruskal-Wallis) which confirmed no significant difference with a p-value of 0.5906.




Figure 2. Boxplot of the CPI values for Cycadaceae (n=4), Stangeriaceae (n=2), Zamiaceae (n=7).

4 Conclusions

This study indicates that soils and sediments accumulating significant amounts of cycad foliage may have low CPI values and that the widely applied *n*-alkane based terrestrial source inference requires refinement particularly in tropical and subtropical soils where cycads grow naturally.

5. Acknowledgements

C.H. Vane is indebted to Nigel Birch at the British Consulate-General, Guangzhou, for logistical and financial assistance. This paper is published by permission of the Executive Director, British Geological Survey.

6. References

- Eglinton, G., Hamilton, J., 1967. Leaf epicuticular waxes. *Science*, 156, 1322-1335.
- Jaffé, R., Mead, R., Hernandez, M.E., Peralba, M.C., DiGuida, O.A., 2001. Origin and transport of sedimentary organic matter in two subtropical estuaries: a comparative, biomarker-based study. *Organic Geochemistry*, 32, 507-526.
- Meyers, P.A., 2003. Applications of organic geochemistry to paleolimnological reconstructions: a summary of examples from the Laurentian Great Lakes. *Organic Geochemistry*, 34, 261-289.
- Osborne, R., Salatino, M.L., Sekiya, C.M., Vazquez Torres, M., 1993. Alkanes of foliar epicuticular waxes from five cycad genera in the Zamiaceae. *Phytochemistry*, 33, 607-609.
- Reddy, C.M., Eglinton, T.I., Palic, R., Benitez-Nelson, B.C., Stojanovic, G., Palic, I., Djordjevic, S., Eglinton, G., 2000. Even carbon number predominance of plant wax *n*-alkanes: a correction. *Organic Geochemistry*, 31, 331-336.
- Rommerskirchen, F., Plader, A., Eglinton, G., Chikaraishi, Y., Rullkötter, J., 2006. Chemotaxonomic significance of distribution and stable carbon isotopic composition of long-chain alkanes and alkan-1-ols in C4 grass waxes. *Organic Geochemistry*, 37, 1303-1332.
- Stojanovic, G., Palic, R., Tarr, C.H., Reddy, C.M., Marinkovic, O., 2003. *n*-Alkanes and fatty acids of *Hypericum perforatum*, *Hypericum maculatum* and *Hypericum olympicum*. *Biochemical Systematics and Ecology*, 31, 223-226.

Wang, X.C., Chen, R.F., Berry, A., 2003. Sources and preservation of organic matter in Plum Island salt marsh sediments (MA, USA): long-chain n-alkanes and stable carbon isotope compositions. *Estuarine, Coastal and Shelf Science*, 58, 917-928.

