



ARTICLE

Received 2 May 2015 | Accepted 24 Feb 2016 | Published 31 Mar 2016

DOI: [10.1038/ncomms11146](https://doi.org/10.1038/ncomms11146)

OPEN

Oxygen depletion recorded in upper waters of the glacial Southern Ocean

Zunli Lu¹, Babette A.A. Hoogakker², Claus-Dieter Hillenbrand³, Xiaoli Zhou¹, Ellen Thomas⁴, Kristina M. Gutchess¹, Wanyi Lu¹, Luke Jones² & Rosalind E.M. Rickaby²

Oxygen depletion in the upper ocean is commonly associated with poor ventilation and storage of respired carbon, potentially linked to atmospheric CO₂ levels. Iodine to calcium ratios (I/Ca) in recent planktonic foraminifera suggest that values less than ~2.5 μmol mol⁻¹ indicate the presence of O₂-depleted water. Here we apply this proxy to estimate past dissolved oxygen concentrations in the near surface waters of the currently well-oxygenated Southern Ocean, which played a critical role in carbon sequestration during glacial times. A down-core planktonic I/Ca record from south of the Antarctic Polar Front (APF) suggests that minimum O₂ concentrations in the upper ocean fell below 70 μmol kg⁻¹ during the last two glacial periods, indicating persistent glacial O₂ depletion at the heart of the carbon engine of the Earth's climate system. These new estimates of past ocean oxygenation variability may assist in resolving mechanisms responsible for the much-debated ice-age atmospheric CO₂ decline.

¹Department of Earth Sciences, Syracuse University, Syracuse, New York 13244, USA. ²Department of Earth Sciences, University of Oxford, Oxford OX1 3AN, UK. ³British Antarctic Survey, Cambridge CB3 0ET, UK. ⁴Department of Geology and Geophysics, Yale University, New Haven, Connecticut, USA. Correspondence and requests for materials should be addressed to Z.L. (email: zunlilu@syr.edu) or to R.E.M.R. (email: rosr@earth.ox.ac.uk).

The Southern Ocean is widely considered to be critical to global nutrient and carbon cycling, including over glacial–interglacial time scales¹. As an area of incomplete nutrient utilization, it is a major source of CO₂ to the atmosphere today². At present, old (CO₂– and nutrient-rich and relatively O₂–depleted) deep waters upwell along most of the Antarctic continental margin^{3,4} (Fig. 1), release CO₂ into, and recharge O₂ from surface waters before they down-well in distinct areas, such as the Weddell and Ross seas, to form Antarctic Bottom Water (AABW). In the glacial Southern Ocean, strengthening of the biological pump due to enhanced iron supply^{5,6}, increased stratification⁷, and expanded sea-ice cover⁸, were among the dominant players in reducing atmospheric CO₂ by ~90 p.p.m.V. Each of these mechanisms could counterbalance the increased O₂ solubility due to lower glacial temperatures, leading to a reduction in the O₂ concentration of the seawater. Since the Southern Ocean is thought to have reduced its CO₂ leakage during glacial periods¹, it provides an ideal location to search for evidence of deoxygenation linked to CO₂ sequestration in the upper ocean.

During the last glacial period, deep waters surrounding Antarctica were less ventilated, and older than today (relative to the atmosphere)⁹. A recent quantitative O₂ proxy study based on benthic foraminiferal δ¹³C indicates that decreased ventilation linked to a reorganization of glacial ocean circulation and a strengthened global biological pump significantly enhanced the ocean storage of respired carbon in the deep North Atlantic¹⁰. Early box-models hypothesized very low-oxygen levels in the high latitude Southern Ocean^{11,12}. Proxies did not paint a clear picture for bottom-water O₂ concentrations in the glacial Southern Ocean¹³. Only a few studies on marine sediment cores south of the APF have found evidence for substantially lowered bottom water O₂ concentrations. There, authigenic uranium concentrations were elevated in sediments deposited during glacial Marine Isotope Stages (MIS) 2 and 6 (refs 14,15). By contrast, another study highlighted a transient stagnation event during the early stage of the last interglacial (MIS 5e)¹⁶.

Bottom water or porewater redox proxies cannot capture upper ocean O₂ levels far from the continental shelf, so there is scant constraint on upper ocean oxygenation conditions in vast tracts of the open ocean¹³. A novel proxy, the I/Ca composition of marine carbonates, especially planktonic and benthic foraminiferal tests, has demonstrated its potential to reconstruct paleo-oxygenation levels in both the upper ocean^{17–20} and bottom waters²¹, respectively. The thermodynamically stable forms of iodine in seawater are iodate (IO₃[–]) and iodide (I[–])²².

The total concentrations of IO₃[–] and I[–] are relatively uniform in the world ocean at around 0.45 μmol l^{–1} due to the residence time of ~300 kyr (ref. 23), supported by a more recent compilation of iodine concentrations in global rivers²⁴. Therefore, the total iodine concentration in the global ocean likely remained invariant over the duration of a glacial termination (~6 kyr).

Iodate is taken up by marine organisms as a micronutrient in surface waters²⁵, but its concentration does not increase during the aging of deep waters^{26,27}, in contrast to those of the major nutrients nitrate and phosphate, probably due to the low I/C_{org} ratio of plankton²⁵. Iodine speciation is strongly redox sensitive. IO₃[–] is completely converted to I[–] when oxygen is depleted²⁸. Because IO₃[–] is the only chemical form of iodine that is incorporated into the structure of carbonate¹⁷, calcareous tests precipitated closer to an oxygen minimum zone (OMZ) will record lower I/Ca and vice versa. An OMZ is defined by O₂ < 20 μmol kg^{–1} in the Pacific Ocean and O₂ < 50 μmol kg^{–1} in the Atlantic Ocean²⁹.

In this paper, we use recent planktonic foraminifera and modern water column data to establish typical I/Ca values for the presence of an OMZ or O₂-depleted water. On the basis of this proxy development, the down-core record of planktonic foraminifera I/Ca obtained at site TC493/PS2547 indicates the persistent presence of oxygen-depletion in the upper waters of high latitude Southern Ocean during the last two glacial periods.

Results

Site selection. We measured I/Ca values on eleven planktonic foraminiferal species in modern to Holocene samples, and in one sample from a previous interglacial (Supplementary Table 1 and Supplementary Figs 1 and 2). We chose sites from well-oxygenated areas (for example, the North and sub-Antarctic South Atlantic), and sites located beneath OMZs, including Ocean Drilling Program (ODP) Sites 658, 709, 720 (Site 720: last interglacial samples), 849 and 1242. First, we use these data to further establish the foundations of the I/Ca proxy. Subsequently, we focus on an I/Ca down-core record on *Neogloboquadrina pachyderma* sinistral deposited during the last two glacial cycles in two sediment cores (PS2547 and TC493) recovered from the same location (71°09' S, 119°55' W, water depth 2,096 m) on a seamount in the Amundsen Sea (Fig. 1)³⁰. The excellent carbonate preservation at this site³⁰ provides a unique window to reconstruct past upper ocean conditions south of the APF. Site




Figure 1 | Hydrographic section of Southern Ocean in the Pacific sector. Dissolved oxygen concentrations showing major water masses⁵⁰ and boundaries, average modern summer (SSI) and winter (WSI) sea-ice extent⁶², and core site PS2547/TC493. The locations of the Antarctic Circumpolar Current (ACC) fronts are marked as SB, Southern Boundary of the ACC; SACCF, Southern ACC Front; APF, Antarctic Polar Front; SAF, Sub-Antarctic Front. AABW, Antarctic Bottom Water; AAIW, Antarctic Intermediate Water; AASW, Antarctic Surface Water; CDW, Circumpolar Deep Water; PDW, Pacific Deep Water. This graph is generated in Ocean Data View, using the Southern Ocean Atlas data set⁶³.





Figure 2 | I/Ca and modern OMZs. (a) Modern and late Holocene I/Ca in planktonic foraminiferal tests versus minimum O₂ concentrations in the upper 500 m of the water column (Note: I/Ca at Site 720 is from a MIS 5 sample). Error bars for y axis indicate the s.d. (1 s.d.) of triplicate measurements. Blue squares show down-core interglacial (IG) I/Ca data on *N. pachyderma* (s) from site TC493/PS2547 plotted against minimum O₂ concentrations in the modern water column, indicating well-oxygenated conditions. I/Ca for glacial *N. pachyderma* (s) tests are marked as red squares, indicating O₂ depletion. (b) Compilation of modern ocean surface water IO₃⁻ concentrations compared with minimum O₂ concentrations^{26–28,47–48,64–69}. Brown dashed line indicates the surface water IO₃⁻ concentration of ~0.25 μmol l⁻¹ as a threshold value for differentiating OMZ-type and normal open ocean type of IO₃⁻ depth profiles. (c) O₂ depth profiles. Yellow shading marks 20–70 μmol kg⁻¹ O₂ concentration as the threshold for complete iodate reduction. (d) IO₃⁻ depth profiles at OMZ sites from the Eastern Equatorial Pacific (EEP)²⁸ and the Arabian Sea (station N8)⁴⁸ and at a well-oxygenated high-latitude site near the Weddell Sea (Station PS71/179-1)⁵⁴.

TC493/PS2547 is currently bathed by Circumpolar Deep Water (CDW), which is overlain by a layer of Antarctic Surface Water (AASW), or Winter Water^{31–33}, and is located on the edge of the average modern summer sea-ice limit³⁴ (Fig. 1). During the Last Glacial Maximum (LGM), the sea-ice boundaries within the Southern Ocean shifted significantly northwards^{35,36}. Thus, it is highly likely that site TC493/PS2547 was located within the permanent sea-ice zone during past glacial periods³⁴.

Age model and glacial polynyas. The sediments of core TC493/PS2547 consist mainly of foraminiferal ooze and sandy mud, with *N. pachyderma* (s) tests forming the primary carbonate component³⁰. The age model of the record is based on

magnetostratigraphy combined with benthic foraminiferal (*Cibicides cf. wuellerstorfi*) oxygen isotope (δ¹⁸O) stratigraphy³⁰, tuned to the global benthic δ¹⁸O stack³⁷. Continuous deposition of foraminifera³⁰ indicates at least episodic opening of polynyas during glacial periods³⁴, because of its seamount location^{38,39}. This scenario is consistent with the occurrence of the benthic foraminifera species *Epistominella exigua*, which is adapted to highly episodic phytodetritus supply⁴⁰.

I/Ca in foraminifera. I/Ca values in the modern and late Holocene samples are lower than ~2.5 μmol mol⁻¹ at sites with O₂ minima <70 μmol kg⁻¹ in the upper ocean (0–500 m) (Fig. 2a).



In contrast, recent planktonic foraminifera at sites with O_2 minima $> 100 \mu\text{mol kg}^{-1}$ have $\text{I/Ca} > 4 \mu\text{mol mol}^{-1}$, regardless of species (Fig. 2a). At site TC493/PS2547, the *N. pachyderma* (s) I/Ca ratio is high ($5\text{--}7 \mu\text{mol mol}^{-1}$) during the Holocene and MIS 5 relative to the lowest values ($< 2 \mu\text{mol mol}^{-1}$) during glacial MIS 2 and 6 (Fig. 3 and Supplementary Table 2).

Discussion

A tremendous amount of work has been devoted to developing foraminiferal proxies for temperature and pH, using global calibrations derived from core-top samples (for example, the Mg/Ca seawater temperature proxy⁴¹). Low I/Ca ratios of planktonic foraminifera unambiguously reveal the presence of low-oxygen waters, but a global calibration approach cannot establish planktonic foraminifera I/Ca as a linearly quantitative proxy for the continuum of dissolved O_2 concentration. Due to the stepwise nature of redox reactions⁴², quantitative IO_3^- reduction does not occur before the dissolved oxygen is depleted to a certain threshold value, triggering nitrate reduction⁴³. IO_3^- concentrations at water depths matching planktonic foraminiferal habitats are often not in equilibrium with the *in situ* O_2 concentrations, and O_2 contents which are sufficiently low to initiate major IO_3^- reduction may be detrimental to many species⁴⁴. Instead, the I/Ca (recording the *in situ* IO_3^- concentration) is determined by the depth habitat of the foraminifera and the upper ocean IO_3^- mixing gradient. This mixing gradient is largely controlled by the surface water IO_3^- concentration and the depth of the IO_3^- reduction zone²⁸. Nonetheless, a planktonic foraminifera proxy that semi-quantitatively approximates dissolved O_2 concentrations, indicative of the presence of an OMZ, can still be highly valuable for the paleoceanography community.

Before interpreting the down-core record from site TC493/PS2547, we identify the characteristic I/Ca signals for modern OMZs. IO_3^- depth profiles in the open ocean generally fall into two types (Fig. 2d): (1) the OMZ-type, with low surface water values and near-zero subsurface values in the OMZ; and (2) the normal open ocean type (for example, in a well-oxygenated water column), with relatively high surface water values and even higher subsurface values. A threshold O_2 concentration will cause complete IO_3^- reduction in the subsurface, and there may be a surface water IO_3^- threshold concentration below which complete IO_3^- reduction is likely to happen in the water column. Combined with modern water column IO_3^- and O_2 data, the I/Ca values measured on modern and late Holocene planktonic foraminifera consistently indicate that $\text{I/Ca} < 2.5 \mu\text{mol mol}^{-1}$ is equivalent to a surface water IO_3^- concentration of $< 0.25 \mu\text{mol l}^{-1}$, thus providing a marker for the presence of oxygen-depleted water with a subsurface O_2 concentration $< 20\text{--}70 \mu\text{mol kg}^{-1}$ (Fig. 2a–c).

Figure 3 | Down-core records of studied sites. (a,b) Stable carbon and oxygen isotopes measured on benthic (*C. cf. wuellerstorfi*) and planktonic (*N. pachyderma*) foraminiferal tests³⁰. It is well documented that $\delta^{13}\text{C}$ of *N. pachyderma* (s) is offset by -1.0‰ south of the APF⁴⁹ and the values plotted here are uncorrected. (c) Atmospheric pCO_2 record at EPICA Dome C is plotted for comparison, with dashed line indicating the long-term mean value of CO_2 (50–270 ka) following Luethi *et al.*⁵⁶. (d) I/Ca measured on *N. pachyderma* (s) tests from cores PS2547 and TC493. (e) Bulk sediment CaCO_3 content from PS2547. Grey columns show the abundance of the benthic foraminifera *E. exigua* as number of tests per cm^3 in core PS2547. Yellow shading highlights peak interglacial periods (including deglaciations), and blue shading marks glacial maxima and cooling intervals.

Modern surface water IO_3^- concentrations are influenced by productivity and the presence of a subsurface OMZ^{25,28}. To visualize this relationship, we compiled surface water IO_3^- concentrations from the literature and plotted them against the minimum O_2 concentrations in the subsurface water (Fig. 2b). The IO_3^- concentration broadly increases with the minimum O_2 concentration when the surface water IO_3^- concentration is $>0.25 \mu\text{mol l}^{-1}$ (Fig. 2b). This correlation is likely a reflection of surface productivity versus subsurface respiration, because lower productivity leads to lower iodine uptake in surface water and less oxygen consumption by subsurface organic matter decomposition. In areas with a strong OMZ and near-zero O_2 values, the surface water IO_3^- concentrations are below $0.25 \mu\text{mol l}^{-1}$ (Fig. 2b). A partition coefficient K_d ($K_d = [\text{I}/\text{Ca}]/[\text{IO}_3^-]$ with units of $[\mu\text{mol mol}^{-1}]/[\mu\text{mol l}^{-1}]$) of ~ 10 was reported from abiological calcite synthesis experiments^{17,20}. Using this K_d value, an IO_3^- concentration $<0.25 \mu\text{mol l}^{-1}$ results in I/Ca values $< \sim 2.5 \mu\text{mol mol}^{-1}$ in calcite. This estimate is consistent with modern I/Ca at OMZ Sites 658, 849 and 1242, as well as the last interglacial I/Ca value at Site 720 (Fig. 2a). Therefore, a surface water I/Ca value $<2.5 \mu\text{mol mol}^{-1}$ indicates that a pronounced subsurface O_2 minimum exerted the dominant control on the upper ocean IO_3^- profile. This I/Ca threshold value does not seem to depend on foraminiferal species (Fig. 2a).

The O_2 threshold for maintaining an OMZ-type IO_3^- profile is useful for the paleoceanographic application of the planktonic I/Ca proxy. At O_2 concentrations $<20 \mu\text{mol kg}^{-1}$, microbial processes become dominant²⁹, and IO_3^- likely would be completely reduced to I^- since the reaction is biologically mediated⁴⁵ (for example, ODP Sites 1242, 720 and 849 in Fig. 2a,c). ODP Site 658 is located at the northern edge of a shallow pocket of distinctively low-oxygen water with mean O_2 concentrations of $\sim 70 \mu\text{mol kg}^{-1}$ in the upper 200 m (ref. 46), which may be sufficiently low to generate an OMZ-type iodate profile. Three species of planktonic foraminifera analysed at ODP Site 1242 show exceptionally low I/Ca ratios around $0.5 \mu\text{mol mol}^{-1}$, corresponding to an IO_3^- concentration of $\sim 0.05 \mu\text{mol l}^{-1}$. Such a low IO_3^- concentration is comparable to that reported for a location where an extreme hypoxic event occurred⁴⁷. Moreover, this low IO_3^- concentration implies that IO_3^- reduction should occur shallower than at Site 849 and at two sites with classic OMZ-type IO_3^- profiles (Eastern Equatorial

Pacific²⁸ and Arabian Sea⁴⁸; Fig. 2c). A comparison of the O_2 profiles of these sites reveals that the O_2 threshold needs to be $>50 \mu\text{mol kg}^{-1}$ to achieve a shallower IO_3^- reduction at Site 1242. Therefore, we suggest that I/Ca values lower than $\sim 2.5 \mu\text{mol mol}^{-1}$ indicate O_2 minima $<20\text{--}70 \mu\text{mol l}^{-1}$. This O_2 range cannot be further narrowed down with the available information, and we refer to this range as the O_2 threshold for an OMZ-type IO_3^- profile. However, the threshold behaviour of IO_3^- reduction (relative to O_2) in subsurface waters does not necessarily lead to step changes in down-core records of planktonic I/Ca . This is because planktonic foraminifera typically record the IO_3^- mixing gradient in the top part of water column, above the O_2 -depleted zone where rapid step changes in IO_3^- concentrations occur. Low planktonic I/Ca values may be driven by shoaling of O_2 -depleted water, and/or by increasing productivity, both of which could change gradually over time.

The available data from modern and late Holocene planktonic foraminifera suggest that the I/Ca ratio acts as a robust (paleo-) proxy for determining the signature of O_2 -depletion in the upper ocean (Fig. 2). At site TC493/PS2547, I/Ca was high ($5\text{--}7 \mu\text{mol mol}^{-1}$) during the Holocene and interglacial MIS 5 when compared with the lowest values ($<2 \mu\text{mol mol}^{-1}$) characterizing peak glacial periods MIS 2 and 6 (Fig. 3). Changes in salinity, temperature and foraminiferal habitat, most likely, are not the main drivers for this record (Supplementary Discussion). The glacial I/Ca values of *N. pachyderma* (s) are best explained by the presence of a water mass with a dissolved O_2 content $<70 \mu\text{mol kg}^{-1}$ close to, i.e., above or near, this site (Figs 2 and 3). We reiterate that the low I/Ca does not necessarily imply O_2 -depleted seawater within the foraminiferal habitat.

At present, CDW wells up to a water depth of approximately 250–300 m in the Amundsen Sea³¹ and has O_2 concentrations notably lower than the top 200 m of the water column (Fig. 2c). Although the interpretation of absolute values of planktonic $\delta^{13}\text{C}$ is far from straightforward in the seasonal ice zone (for example, disequilibrium from seawater⁴⁹), it is reasonable to assume that CDW had a strong influence on the local water column during glacial periods, as its upwelling along the continental margin was probably responsible for the opening of the glacial polynyas. The CDW upwelling at site TC493/PS2547 today partly originates from Pacific Deep Water (PDW) moving southward from the equator, with a low-oxygen and high nutrient signature




Figure 4 | Conceptual illustration of paleo-environmental changes. Upper ocean IO_3^- and O_2 profiles were influenced by circulation, productivity and polynyas over glacial cycles. (a) Well-oxygenated interglacial condition; (b) Relatively oxygen-depleted glacial conditions with expanded sea-ice cover; (c) Episodic polynya opening during glacials.

(Fig. 1)^{50,51}. $\delta^{30}\text{Si}$ data from fossil diatoms and sponges indicate higher silicic acid concentrations in the Pacific sector of the Southern Ocean during the LGM, which further imply that either the southward transport of PDW was more efficient or PDW was less ventilated than today⁵². So glacial CDW was likely more O_2 depleted than during interglacials, and upwelling of this water contributed to the glacial I/Ca signal at site TC493/PS2547.

The oxidation of I^- to IO_3^- is thought to take from a few months up to 40 years⁵³. Long-distance transport of well-oxygenated deep water with low IO_3^- concentrations ($<0.25\ \mu\text{mol l}^{-1}$) has not been documented in the modern ocean, but this scenario should be tested with further work on I^- oxidation kinetics. Today our site is bathed in CDW transported from a Pacific OMZ and the interglacial I/Ca values at site TC493/PS2547 do not show any remnant signal of the OMZ from the Pacific Ocean. On the basis of the knowledge about iodine speciation change in modern ocean, we interpret the observed glacial I/Ca values as a local signal, in principle, indicating the presence of a water mass with low O_2 and low IO_3^- vertically or horizontally close to the planktonic foraminiferal habitat.

In the setting of site TC493/PS2547 a coherent conceptual model for *N. pachyderma* (s) recording the presence/absence of O_2 -depletion needs to integrate changes in productivity, sea-ice extent and the opening/closing of polynyas on time scales of glacial to seasonal cycles (Fig. 4). Although the polynyas complicate the interpretation of the proxy data, their presence arguably provides the only window for sufficient accumulation of planktonic microfossils to record upper ocean conditions during glacial periods at such high latitudes.

The modern O_2 profile at site TC493/PS2547 is defined by equilibration with the atmosphere at 0–250 m, and CDW influence below 250 m, as shown by the distinctively low O_2 concentrations (Fig. 2c). With O_2 above the threshold for complete IO_3^- reduction in the entire water column, the IO_3^- profile at site TC493/PS2547 should be similar to those at other high latitude locations, for example, site PS71/179–1 at $69^\circ 31' \text{ S}$ and $0^\circ 3' \text{ W}$ in the Weddell Sea⁵⁴ (Fig. 2d). An interglacial scenario of relatively high seasonal productivity, high O_2 and surface water IO_3^- ($>0.3\ \mu\text{mol l}^{-1}$) concentrations (Fig. 4a), is well described for the modern Atlantic sector of Southern Ocean⁵⁴.

Relative to the interglacial periods, the Southern Ocean experienced expanded sea-ice cover during glacial periods, and was less ventilated^{9,36}. A more dynamic seasonal sea-ice cycle during ice ages would have increased water column stratification. Increased winter sea-ice formation (spatially and volumetrically) may have generated waters dense enough to sink ultimately to the bottom of the ocean⁵⁵. On the other hand, melting of thicker sea ice during glacial-time summers in the seasonal sea-ice zone would have strengthened the halocline (not considering the influence of polynyas). So, the glacial seasonal stratification was likely stronger than today. These factors overall should have lowered the glacial O_2 concentrations in the Southern Ocean (Fig. 4b). At site TC493/PS2547, glacial I/Ca demonstrate that the IO_3^- profile was OMZ-like with complete IO_3^- reduction near the foraminiferal habitat (Fig. 2d). However, the dynamics of polynyas must be considered when interpreting the location of the low O_2 water mass, and the means by which the signal was recorded by *N. pachyderma* (s).

Without a polynya above site TC493/PS2547, glacial phytoplankton productivity under perennial sea-ice cover would have been relatively low due to the scarcity of light³⁴, and planktonic foraminifera depending on algae could not flourish. The water column would have been relatively poorly ventilated and strongly stratified during these times, creating the ideal environment for developing low O_2 conditions and an OMZ-type IO_3^- profile

(Fig. 4b). The episodic opening of a polynya re-established primary production (mainly by diatoms) and thus a planktonic foraminiferal habitat, vertical mixing and oxygenation in, at least, the uppermost part of the water column (Fig. 4c). While overall glacial-time production was reduced^{30,34}, the planktonic foraminifera preserved in the glacial sediments probably recorded transient I/Ca changes in the water column associated with polynya-induced peaks in glacial productivity. Modern open ocean productivity pulses do not lower IO_3^- concentrations to $<0.25\ \mu\text{mol l}^{-1}$ in oxygenated water (Supplementary Discussion)⁵⁴, thus the glacial I/Ca signal is most likely driven by changes in O_2 and not productivity.

The likely short-lived nature of glacial polynyas makes it difficult to envisage that very brief plankton blooms alone could produce a utilization-driven O_2 depletion in a cold, well-oxygenated Southern Ocean. For the same reason, it is difficult to imagine that the vertical mixing cells restricted by the size of the polynya could rapidly oxygenate voluminous nearby waters outside of the polynya, if most of the sea-ice covered areas were O_2 -depleted. The more likely scenario is that the O_2 concentrations in the deep and abyssal Southern Ocean were generally lower during glacial periods than during interglacial periods. Upwelling of a more O_2 -depleted CDW in the generally stratified upper ocean was mainly responsible for the IO_3^- reduction at site TC493/PS2547 (Fig. 4b), while the episodic opening of polynyas created habitable conditions for planktonic foraminifera to record the deoxygenation in the upper ocean (Fig. 4c). We suggest that the I/Ca proxy should be used as a local proxy, in principle. However, it is probably a reasonable speculation that this record (Fig. 3) shows oxygenation changes integrated over a regional volume of water (e.g. CDW).

The timing of glacial deoxygenation and deglacial reoxygenation at site PS2547 shows potential linkages to global climate changes (Fig. 3). The appearance of OMZ-type I/Ca values ($<\sim 2.5\ \mu\text{mol mol}^{-1}$) during past glacial periods coincided with the lowering in atmospheric pCO_2 level below the long-term mean value⁵⁶. Identical timing was reported for a strongly stratified Antarctic Zone coincident with pCO_2 decrease under the same threshold value (225 p.p.m.) in the Atlantic sector of the Southern Ocean⁵⁷. Stronger stratification may be the common driving force for the productivity change (ODP Site 1094) and oxygenation change (PS2547/TC493) in the Antarctic zone. Furthermore, during the last interglacial period, the recovery of *N. pachyderma* (s) I/Ca values is offset from the $\delta^{18}\text{O}$ trend, with peak I/Ca occurring about 10 kyr after the peak $\delta^{18}\text{O}$ (Fig. 3), an observation worthy of future investigation.

Our I/Ca results build on other evidence^{52,58,59} to make a stronger case for lower oxygen concentrations in CDW (and very likely PDW) during glacial periods. Altogether with the reconstructed O_2 content of deep waters in the glacial North Atlantic¹⁰, these observations seem to allude to large scale deoxygenation in the glacial global ocean interior⁶⁰. Future work providing quantitative reconstructions of bottom water O_2 concentrations in the Southern Ocean, especially south of the APF, and in other major ocean basins will shed new light on the mechanisms of sequestering atmospheric CO_2 during ice ages.

Methods

Foraminifera cleaning. Sediments were sampled from the split core sections and wet sieved. Approximately 40 tests of *N. pachyderma* sinistral were picked from the 200–250 μm size fraction of each sample. The cleaning procedure followed the Mg/Ca protocol of Barker *et al.*⁶¹. Cleaned glass slides were used to gently crack open all chambers. Clay particles were removed in an ultrasonic water bath. After adding NaOH-buffered 1% H_2O_2 solutions the samples were heated in boiling water for 10–20 min to remove organic matter. Calcareous microfossils were then

thoroughly rinsed with de-ionized water. Reductive cleaning was not applied because contribution of iodine from Mn-oxides is deemed negligible¹⁹.

ICP-MS measurements. The cleaned samples were dissolved in 3% nitric acid, and diluted to solutions with 50 p.p.m. Ca for analyses. Iodine calibration standards were freshly prepared also with 50 p.p.m. Ca. 0.5% tertiary amine solution (Spectrasol, CFA-C) was added to stabilize iodine within a few minutes after the sample dissolution. The measurements were performed immediately after that to minimize potential iodine loss. The sensitivity of iodine was tuned to above 80 kcps for a 1 p.p.b. standard. The precision for ¹²⁷I is typically better than 1%. The long-term accuracy is guaranteed by frequently repeated analyses of the reference material JCP-1 (ref. 17). The detection limit of I/Ca is on the order of 0.1 μmol mol⁻¹. The I/Ca measurements were performed using a quadrupole ICP-MS (Bruker M90) at Syracuse University.

References

- Sigman, D. M., Hain, M. P. & Haug, G. H. The polar ocean and glacial cycles in atmospheric CO₂ concentration. *Nature* **466**, 47–55 (2010).
- Marinov, I., Gnanadesikan, A., Toggweiler, J. R. & Sarmiento, J. L. The Southern Ocean biogeochemical divide. *Nature* **441**, 964–967 (2006).
- Orsi, A. H., Whitworth, T. & Nowlin, W. D. On the meridional extent and fronts of the Antarctic Circumpolar Current. *Deep Sea Res. I* **42**, 641–673 (1995).
- Schmidtko, S., Heywood, K. J., Thompson, A. F. & Aoki, S. Multidecadal warming of Antarctic waters. *Science* **346**, 1227–1231 (2014).
- Martin, J. H. GLACIAL-interglacial CO₂ change: the iron hypothesis. *Paleoceanography* **5**, 1–13 (1990).
- Martinez-Garcia, A. *et al.* Iron fertilization of the subantarctic ocean during the last ice age. *Science* **343**, 1347–1350 (2014).
- Toggweiler, J. R. Variation of atmospheric CO₂ by ventilation of the ocean's deepest water. *Paleoceanography* **14**, 571–588 (1999).
- Stephens, B. B. & Keeling, R. F. The influence of Antarctic sea ice on glacial-interglacial CO₂ variations. *Nature* **404**, 171–174 (2000).
- Skinner, L. C., Fallon, S., Waelbroeck, C., Michel, E. & Barker, S. Ventilation of the deep southern ocean and deglacial CO₂ rise. *Science* **328**, 1147–1151 (2010).
- Hoogakker, B. A. A., Elderfield, H., Schmiedl, G., McCave, I. N. & Rickaby, R. E. M. Glacial-interglacial changes in bottom-water oxygen content on the Portuguese margin. *Nat. Geosci.* **8**, 40–43 (2015).
- Knox, F. & McElroy, M. B. Changes in atmospheric CO₂ - influence of the marine biota at high-latitude. *J. Geophys. Res. Atmos.* **89**, 4629–4637 (1984).
- Archer, D. E. *et al.* Model sensitivity in the effect of Antarctic sea ice and stratification on atmospheric pCO₂. *Paleoceanography* **18** (2003).
- Jaccard, S. L. & Galbraith, E. D. Large climate-driven changes of oceanic oxygen concentrations during the last deglaciation. *Nat. Geosci.* **5**, 151–156 (2012).
- Ceccaroni, L. *et al.* Late Quaternary fluctuations of biogenic component fluxes on the continental slope of the Ross Sea, Antarctica. *J. Mar. Syst.* **17**, 515–525 (1998).
- Frank, M. *et al.* Similar glacial and interglacial export bioproductivity in the Atlantic sector of the Southern Ocean: Multiproxy evidence and implications for glacial atmospheric CO₂. *Paleoceanography* **15**, 642–658 (2000).
- Hayes, C. T. *et al.* A stagnation event in the deep South Atlantic during the last interglacial period. *Science* **346**, 1514–1517 (2014).
- Lu, Z., Jenkyns, H. C. & Rickaby, R. E. M. Iodine to calcium ratios in marine carbonate as a paleo-redox proxy during oceanic anoxic events. *Geology* **38**, 1107–1110 (2010).
- Hardisty, D. S. *et al.* An iodine record of Paleoproterozoic surface ocean oxygenation. *Geology* **42**, 619–622 (2014).
- Zhou, X. L., Thomas, E., Rickaby, R. E. M., Winguth, A. M. E. & Lu, Z. L. I/Ca evidence for upper ocean deoxygenation during the PETM. *Paleoceanography* **29**, 964–975 (2014).
- Zhou, X. *et al.* Upper ocean oxygenation dynamics from I/Ca ratios during the Cenomanian-Turonian OAE 2. *Paleoceanography* **30** (2015).
- Glock, N., Liebetrau, V. & Eisenhauer, A. I/Ca ratios in benthic foraminifera from the Peruvian oxygen minimum zone: analytical methodology and evaluation as a proxy for redox conditions. *Biogeosciences* **11**, 7077–7095 (2014).
- Wong, G. T. F. The marine geochemistry of iodine. *Rev. Aquat. Sci.* **4**, 45–73 (1991).
- Broecker, W. S. & Peng, H. T. *Tracers in the Sea* 690 (Lamont-Doherty Geological Observatory, 1982).
- Moran, J. E., Oktay, S. D. & Santschi, P. H. Sources of iodine and iodine 129 in rivers. *Water Resources Research* **38** (2002).
- Elderfield, H. & Truesdale, V. W. On the biophilic nature of iodine in seawater. *Earth Planet. Sci. Lett.* **50**, 105–114 (1980).
- Nakayama, E., Kimoto, T., Ishiki, K., Sohrin, Y. & Okazaki, S. Determination and distribution of iodide-iodine and total-iodine in the North Pacific Ocean - by using a new automated electrochemical method. *Mar. Chem.* **27**, 105–116 (1989).
- Waite, T. J., Truesdale, V. W. & Olafsson, J. The distribution of dissolved inorganic iodine in the seas around Iceland. *Mar. Chem.* **101**, 54–67 (2006).
- Rue, E. L., Smith, G. J., Cutter, G. A. & Bruland, K. W. The response of trace element redox couples to suboxic conditions in the water column. *Deep Sea Res. I* **44**, 113–134 (1997).
- Gilly, W. F., Beman, J. M., Litvin, S. Y. & Robison, B. H. in *Annual Review of Marine Science, Annual Review of Marine Science* Vol. 5 (eds Carlson, C. A. & Giovannoni, S. J.) 393–420 (Annual Reviews, 2013).
- Hillenbrand, C. D., Futterer, D. K., Grobe, H. & Frederichs, T. No evidence for a Pleistocene collapse of the West Antarctic Ice Sheet from continental margin sediments recovered in the Amundsen Sea. *Geo-Mar. Lett.* **22**, 51–59 (2002).
- Wahlin, A. K. *et al.* Some Implications of Ekman layer dynamics for cross-shelf exchange in the Amundsen sea. *J. Phys. Oceanogr.* **42**, 1461–1474 (2012).
- Nakayama, Y., Schroder, M. & Hellmer, H. H. From circumpolar deep water to the glacial meltwater plume on the eastern Amundsen Shelf. *Deep Sea Res. I* **77**, 50–62 (2013).
- Jacobs, S. *et al.* Getz Ice Shelf melting response to changes in ocean forcing. *J. Geophys. Res. Oceans* **118**, 4152–4168 (2013).
- Thatje, S., Hillenbrand, C. D., Mackensen, A. & Larter, R. Life hung by a thread: Endurance of antarctic fauna in glacial periods. *Ecology* **89**, 682–692 (2008).
- Collins, L. G., Pike, J., Allen, C. S. & Hodgson, D. A. High-resolution reconstruction of southwest Atlantic sea-ice and its role in the carbon cycle during marine isotope stages 3 and 2. *Paleoceanography* **27** (2012).
- Gersonde, R., Crosta, X., Abelmann, A. & Armand, L. Sea-surface temperature and sea ice distribution of the Southern Ocean at the EPILOG Last Glacial Maximum—a circum-Antarctic view based on siliceous microfossil records. *Quart. Sci. Rev.* **24**, 869–896 (2005).
- Lisiecki, L. E. & Raymo, M. E. A Pliocene-Pleistocene stack of 57 globally distributed benthic δ¹⁸O-18 records. *Paleoceanography* **20** (2005).
- Bersch, M., Becker, G. A., Frey, H. & Koltermann, K. P. Topographic effects of the Maud Rise on the stratification and circulation of the Weddell gyre. *Deep Sea Res. A* **39**, 303–331 (1992).
- Martin, S. in *Encyclopedia of ocean sciences*. (eds Steele, H., Turekian, K. K. & Thorpe, S. A.) 2241–2247 (Academic Press, 2001).
- Smith, J. A., Hillenbrand, C. D., Pudsey, C. J., Allen, C. S. & Graham, A. G. C. The presence of polynyas in the Weddell Sea during the Last Glacial Period with implications for the reconstruction of sea-ice limits and ice sheet history. *Earth Planet. Sci. Lett.* **296**, 287–298 (2010).
- Elderfield, H. & Ganssen, G. Past temperature and δ¹⁸O-18 of surface ocean waters inferred from foraminiferal Mg/Ca ratios. *Nature* **405**, 442–445 (2000).
- Stumm, W. & Morgan, J. J. *Aquatic Chemistry*, 2nd edn 477 (Wiley, 1981).
- Luther, G. W. *et al.* Simultaneous measurement of O₂, Mn, Fe, I₂, and S(-II) in marine pore waters with a solid-state voltammetric microelectrode. *Limnol. Oceanogr.* **43**, 325–333 (1998).
- Hull, P. M., Osborn, K. J., Norris, R. D. & Robison, B. H. Seasonality and depth distribution of a mesopelagic foraminifer, *Hastigerinella digitata*, in Monterey Bay, California. *Limnol. Oceanogr.* **56**, 562–576 (2011).
- Kuepper, F. C. *et al.* Commemorating two centuries of iodine research: an interdisciplinary overview of current research. *Angew. Chem. Int. Ed.* **50**, 11598–11620 (2011).
- Stramma, L., Huttel, S. & Schafstall, J. Water masses and currents in the upper tropical northeast Atlantic off northwest Africa. *J. Geophys. Res. Oceans* **110** (2005).
- Truesdale, V. W. & Bailey, G. W. Dissolved iodate and total iodine during an extreme hypoxic event in the Southern Benguela system. *Estuar. Coast. Shelf Sci.* **50**, 751–760 (2000).
- Farrenkopf, A. M. & Luther, G. W. Iodine chemistry reflects productivity and denitrification in the Arabian Sea: evidence for flux of dissolved species from sediments of western India into the OMZ. *Deep Sea Res. II* **49**, 2303–2318 (2002).
- Kohfeld, K. E., Anderson, R. F. & Lynch-Stieglitz, J. Carbon isotopic disequilibrium in polar planktonic foraminifera and its impact on modern and Last Glacial Maximum reconstructions. *Paleoceanography* **15**, 53–64 (2000).
- Talley, L. D. Closure of the global overturning circulation through the Indian, Pacific, and Southern Oceans: schematics and transports. *Oceanography* **26**, 80–97 (2013).
- McCave, I. N., Carter, L. & Hall, I. R. Glacial-interglacial changes in water mass structure and flow in the SW Pacific Ocean. *Quart. Sci. Rev.* **27**, 1886–1908 (2008).

52. Ellwood, M. J., Wille, M. & Maher, W. Glacial Silicic Acid Concentrations in the Southern Ocean. *Science* **330**, 1088–1091 (2010).
53. Chance, R., Baker, A. R., Carpenter, L. & Jickells, T. D. The distribution of iodide at the sea surface. *Environ. Sci. Process. Impacts* **16**, 1841–1859 (2014).
54. Bluhm, K., Croot, P. L., Huhn, O., Rohardt, G. & Lochte, K. Distribution of iodide and iodate in the Atlantic sector of the southern ocean during austral summer. *Deep Sea Res. II* **58**, 2733–2748 (2011).
55. Mackensen, A. Strong thermodynamic imprint on Recent bottom-water and epibenthic delta C-13 in the Weddell Sea revealed: Implications for glacial Southern Ocean ventilation. *Earth Planet. Sci. Lett.* **317**, 20–26 (2012).
56. Luthi, D. *et al.* High-resolution carbon dioxide concentration record 650,000–800,000 years before present. *Nature* **453**, 379–382 (2008).
57. Jaccard, S. L. *et al.* Two modes of change in southern ocean productivity over the past million years. *Science* **339**, 1419–1423 (2013).
58. Moy, A. D., Howard, W. R. & Gagan, M. K. Late Quaternary palaeoceanography of the Circumpolar Deep Water from the South Tasman Rise. *J. Quat. Sci.* **21**, 763–777 (2006).
59. Galbraith, E. D. & Jaccard, S. L. Deglacial weakening of the oceanic soft tissue pump: global constraints from sedimentary nitrogen isotopes and oxygenation proxies. *Quart. Sci. Rev.* **109**, 38–48 (2015).
60. Jaccard, S. L., Galbraith, E. D., Martínez-García, A. & Anderson, R. F. Covariation of deep Southern Ocean oxygenation and atmospheric CO₂ through the last ice age. *Nature* **530**, 207–210 (2016).
61. Barker, S., Greaves, M. & Elderfield, H. A study of cleaning procedures used for foraminiferal Mg/Ca paleothermometry. *Geochem. Geophys. Geosyst.* **4** (2003).
62. Comiso, J. C. *Large-Scale Characteristics and Variability of the Global Sea Ice Cover* 112–142 (Blackwell, 2003).
63. Olbers, D., Gouretski, V., Seiss, G. & Schroeter, J. *Hydrographic Atlas of the Southern Ocean* (Alfred-Wegener Institute (AWI), 1992).
64. Truesdale, V. W., Bale, A. J. & Woodward, E. M. S. The meridional distribution of dissolved iodine in near-surface waters of the Atlantic Ocean. *Progress Oceanogr.* **45**, 387–400 (2000).
65. Campos, M., Farrenkopf, A. M., Jickells, T. D. & Luther, G. W. A comparison of dissolved iodine cycling at the Bermuda Atlantic Time-Series station and Hawaii Ocean Time-Series Station. *Deep Sea Res. II* **43**, 455–466 (1996).
66. Wong, G. T. F. Dissolved iodine across the Gulf Stream Front and in the South Atlantic Bight. *Deep Sea Res. I* **42**, 2005–2023 (1995).
67. Farrenkopf, A. M., Dollhopf, M. E., NiChadhain, S., Luther, G. W. & Neelson, K. H. Reduction of iodate in seawater during Arabian Sea shipboard incubations and in laboratory cultures of the marine bacterium *Shewanella putrefaciens* strain MR-4. *Mar. Chem.* **57**, 347–354 (1997).
68. Smith, J. D., Butler, E. C. V., Airey, D. & Sandars, G. Chemical-properties of a low-oxygen water column in port-hacking (Australia) - arsenic, iodine and nutrients. *Mar. Chem.* **28**, 353–364 (1990).
69. Emerson, S., Cranston, R. E. & Liss, P. S. Redox species in a reducing fjord - equilibrium and kinetic considerations. *Deep Sea Res. A* **26**, 859–878 (1979).

Acknowledgements

Z.L. thanks NSF OCE 1232620. B.A.A.H. is supported by Natural Environment Research Council (NERC) grant NE/1020563/1. C.-D.H. is supported by NERC. Z.L. and R.E.M.R. were supported by NERC NE/E018432/1 during the initial development of the I/Ca proxy. We thank the captains, officers, crews and shipboard scientists of expeditions JR179 and ANT-XI/3 for recovering cores TC493 and PS2547, Professor David Hodell and Professor Andreas Mackensen for analysing stable isotopes on these cores, and Dr Hannes Grobe for providing samples from core PS2547. We acknowledge Professor Nick McCave for providing some of the core top materials and Mr Thomas Williams provided assistance for Fig. 1. This research used samples provided by the Ocean Drilling Program (ODP). ODP is sponsored by the US National Science Foundation and participating countries under management of the Joint Oceanographic Institutions (JOI). This manuscript greatly benefited from comments by Dr Nicolaas Glock and two anonymous reviewers.

Author contributions

Z.L., X.Z., K.M.G. and W.L. carried out the I/Ca analysis. C.-D.H. provided all samples from core TC493 and CaCO₃ and stable isotope data for cores TC493 and PS2547. B.A.A.H. and L.J. contributed the core top samples. E.T. identified *E. exigua*. All authors participated in data interpretation and manuscript preparation.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Lu, Z. *et al.* Oxygen depletion recorded in upper waters of the glacial Southern Ocean. *Nat. Commun.* 7:11146 doi: 10.1038/ncomms11146 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>